

Technical note: A procedure to clean, decompose and aggregate time series

François Ritter¹

¹Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette 91191, France

Correspondence to: François Ritter (ritter.francois@gmail.com)

Abstract. Errors, gaps and outliers complicate and sometimes invalidate the analysis of time series. While most fields have developed their own strategy to clean the raw data, no generic procedure has been promoted to standardize the pre-processing. This lack of harmonization makes the inter-comparison of studies difficult, and leads to screening methods that can be arbitrary or case-specific. This study provides a generic pre-processing procedure implemented in R (**ctbi**, for cyclic/trend decomposition using bin interpolation) dedicated to univariate time series. **Ctbi** is based on data binning and decomposes the time series into a long-term trend and a cyclic component (quantified by a new metric, the Stacked Cycles Index) to finally aggregate the data. Outliers are flagged with an enhanced boxplot rule called Logbox that corrects biases due to the sample size and that is adapted to non-Gaussian residuals. Three different Earth Science datasets (contaminated with gaps and outliers) are successfully cleaned and aggregated with **ctbi**. This illustrates the robustness of this procedure that can be valuable to any discipline.

1 Introduction

In any discipline, raw data need to be evaluated during a pre-processing procedure before performing the analysis. Errors are removed, values that deviate from the rest of the population are flagged (outliers, see Aguinis et al., 2013), in some cases gaps are filled. Because the raw data are altered, pre-processing is a delicate and time-consuming task that can be neglected due to cognitive biases deflecting our understanding of reality ('I see what I want to see'), or due to our impatience to obtain results. The fate of extreme values is crucial as they usually challenge scientific or economic theories (Reiss et al., [19972007](#)).

Time series are particularly difficult to pre-process (Chandola et al., 2009). A value can or cannot be considered as an outlier just depending on its timestamp (e.g., a freezing temperature in summer), large data gaps are common, abrupt changes can occur and a background noise covers the true signal. In Earth Science, in-situ or remote measurements routinely produce time series that first need to be visually inspected. The expert-knowledge of the researcher, technician or engineer is essential to

flag suspicious periods of possible instrument failure (e.g., a rain gauge blocked by snowflakes), violation of the experimental conditions (e.g., a passing car during CO₂ measurements in a forest), or human error (e.g., calibration of the wrong sensor). Once these suspicious periods have been flagged, a pre-processing algorithm is necessary to evaluate the quality of the remaining portion of the measurements. However, there currently is no consensus on which procedure to use even in the simple univariate case: a recent review (Ranjan et al., 2021) covered more than 37 preprocessing methods for univariate time series, and Aguinis et al. (2013) listed 14 different outlier definitions that are mutually exclusive. Despite this (overwhelming) abundance of methods and conventions, there are surprisingly few R packages that offer a pre-processing function. It is worth mentioning *hampel* (package **pracma**, Borchers, 2021) that applies a Hampel filter (Pearson, 2002) to time series and flags outliers based on the Mean Absolute Deviation (MAD), which is a robust approximation of the standard deviation defined as $MAD(x) = 1.4826 \times M(|x - M(x)|)$ with M the median operator. However, the *hampel* function is not robust to missing values and the scaling factor of 1.4826 is not adapted to non-Gaussian residuals. Another option is the function *tsoutliers* (package **forecast**, Hyndman et al., 2020) that applies a Seasonal and Trend decomposition using Loess (STL, Cleveland et al., 1990) to data showing a seasonal pattern, complemented by a smoothing function to estimate the trend of non-seasonal time series (Friedman's super smoother, Friedman, 1984). The residuals obtained can be transformed to follow a Gaussian distribution (Cox-Box method, Box & Cox, 1964), and then outliers are flagged using the boxplot rule (Tukey, 1977). This method will be proved in this study to work well with data associated with nearly-Gaussian residuals, but to show poor performance otherwise.

This study offers an alternative pre-processing procedure (implemented in R) called **ctbi** for cyclic/trend decomposition using bin interpolation. The time series is divided into a sequence of non-overlapping time intervals of equal period (called bins), and outliers are flagged with an enhanced version of the boxplot rule (called Logbox) that is adapted to non-Gaussian data for different sample sizes. **Ctbi** fulfils four purposes:

- i) Data cleaning: bins with insufficient data are discarded, and outliers are flagged in the remaining bins. If there is a cyclic pattern within each bin, missing values can be imputed as well.
- ii) Decomposition: the time series is decomposed into a long-term trend and a cyclic component.
- iii) Cyclicity analysis: the mean cycle of the stacked bins is calculated, and the strength of the cyclicity is quantified by a novel index, the Stacked Cycles Index.
- iv) Aggregation: data are averaged (or summed) within each bin.

This procedure is particularly adapted to univariate time series that are *messy*, with outliers, data gaps or irregular timesteps. The inputs offer a large flexibility in terms of imputation level or outlier cutoff, but also in the timestamp of the bins: a day does not necessarily start at midnight or a year the 1st of January. The timeline is not limited to daily or monthly data but can

vary from milliseconds to millenarians. The outputs keep track of the changes brought to the data: contaminated bins are flagged, as well as outliers and imputed data points.

This paper is divided into two distinct parts. The first part describes the Logbox method and compares its performance with ~~four~~ five other outlier detection methods in the literature based on daily precipitation & temperature data extracted from ~~6307~~ century-old weather stations. The second part describes the **ctbi** procedure, and then applies it to three datasets that have been contaminated beforehand to show the efficiency of the algorithm. A comparison with *tsoutliers* is performed, and, finally, limitations and good practice recommendations are discussed.

2 Part I, outliers

65 2.1 Context

This first part is dedicated to the detection of outliers present in univariate datasets (without the time component). The boxplot (or Tukey's) rule is a commonly used method to flag outliers below a lower boundary l and above an upper boundary u (Tukey, 1977):

$$\begin{cases} l = q(0.25) - \alpha \times (q(0.75) - q(0.25)) \\ u = q(0.75) + \alpha \times (q(0.75) - q(0.25)) \end{cases}$$

70 With q the sample quantile (e.g., $q(0.5)$ is the median) and $\alpha = 1.5$ a constant that corresponds to 99.3% of Gaussian data falling within $[l, u]$. This method is simple and robust to the presence of a maximum of 25% of outliers in the dataset (known as the breakdown point). When a real data point falls outside the $[l, u]$ range, it is considered as an erroneously flagged outlier (or type I error). Conversely, a type II error occurs when a real outlier is not flagged. The type I error is more common for three reasons:

- 75 (i) For small Gaussian samples ($n < 30$), up to 8.6% of data (Hoaglin et al., 1986) can be cut due to the inaccuracy of the sample quantile for small n .
- (ii) For large Gaussian samples ($n > 10^3$), $\alpha = 1.5$ is inappropriate because the number of erroneously flagged outliers increases linearly with n due to the 99.3% of data captured by $[l, u]$.
- 80 (iii) For non-Gaussian populations, $\alpha = 1.5$ is generally too restrictive. For example, ~4.8 % of data following an Exponential distribution would be cut.

~~While studies~~ Studies have corrected biases in the detection of outliers in small samples (see Carling, 2000; Schwertman et al., 2004), ~~they have ignored these emerging in-~~ and large samples. Concerning the last issue (Barbato et al., 2011), ~~but these methods were adapted to Gaussian populations. For non-Gaussian populations,~~ Kimber (1990) and Hubert & Vandervieren (2008) have adjusted α to the skewness (related to the asymmetry of a distribution) but did not consider the kurtosis (related

85 to the tail weight) that will be proven to be a key variable in this study. Therefore, there currently is no generic procedure that
can be used when the population is non-Gaussian ~~and/or the sample size is large~~.

To understand how to address this problem, two sets of common distributions with known skewness S , kurtosis excess κ_{ex}
and quantile function Q are used (Fig. 1). The first set is the Pearson family composed of light-tailed distributions that represent
90 any theoretically possible residuals with moderate S & κ_{ex} . Pearson originally worked to create distributions that cover the
entire (S, κ_{ex}) space (Pearson, 1895; 1901 & 1916), but they took their modern names later on (Gamma, Inverse-gamma, Beta
prime, Student, Pearson IV). The second set is the Generalized Extreme Value family composed of the Gumbel, Weibull and
Fréchet that are heavy-tailed distributions (high S & κ_{ex}) used in Extreme Value Theory to model the behavior of extrema
(Jenkinson, 1955). Based on this framework, this study finds that $\alpha(n) = A \log(n) + B + \frac{C}{n}$ reasonably addresses all
95 previously mentioned issues, with C fixed as a constant ($C = 36$). The two parameters A and B correspond to the nature of the
distribution and are estimated based on a predictor of the ~~kurtosis modified from maximum tail weight and inspired by~~ Moors
(1988). A comparison between this procedure (called Logbox) and ~~four~~five other existing models is performed on residuals
obtained from 6307 weather stations with more than 100 years of daily temperature and precipitation measurements (Fig. 2).
Finally, Logbox is implemented in part II to clean the residuals obtained after fitting the univariate time series with a robust
100 and nonparametric method.

2.2 Method

2.2.1 Distributions

Residuals with moderate κ_{ex} & S are represented in this study with 4999 light-tailed distributions from the Pearson family
(Pearson, 1895; 1901 & 1916) composed of the Gaussian, Gamma (196 distributions, including the Exponential), Inverse
105 gamma (170), Beta prime (1135), Pearson IV (3377) and Student (120) distributions (Fig. 1a). These distributions cover the
entire (κ_{ex}, S^2) space without overlap, except for the Beta distribution that has been discarded due to a bounded support
(unrealistic residuals). The shape parameters of each distribution have been chosen to produce regularly-spaced points with a
mean distance of 0.05 in the (κ_{ex}, S^2) space and with a range between the Gaussian and the Exponential: $\kappa_{ex} \in [0,6]$ and $S \in$
[0,2]. Heavy-tailed residuals are represented with 368 distributions from the Generalized Extreme Value (GEV) family (Fig.
110 1d) composed of the Gumbel, Weibull (244 distributions) and Fréchet (123). Their shape parameters cover a larger range:
 $\kappa_{ex} \in [0,500]$ and $S \in [0,15]$.

2.2.2 The Logbox model

Based on the boxplot rule, α can be defined as:

$$\alpha(n) = \frac{Q\left(1 - \frac{f(n)}{2n}\right) - Q(0.75)}{Q(0.75) - Q(0.25)}$$

With n the sample size, Q the population quantile function and f a function that gives the number of erroneously flagged outliers. In the original boxplot rule, $Q = \Phi^{-1}$ (with Φ the cumulative distribution function of the Gaussian) and $f(n) = 0.007 \times n^{0.07n}$ which leads to $\alpha = 1.5$. As explained in the introduction, this choice of f is not valid for large sample sizes due to the linear dependence on n . A flat number of erroneously flagged outliers ($f(n) = b$) or a logarithmic relationship

($f(n) = b \log(n)$) would not be appropriate either, because $\alpha(n)$ could take arbitrary large values as $1 - \frac{f(n)}{2n}$ would approach 1 too rapidly ($Q(1) = \infty$). This study suggests instead $f(n) = 0.001\sqrt{n}$ as a compromise. For example, for a sample of size $n = 10^2, 10^4$ or 10^6 ; respectively 0.01, 0.1 or 1 point would be erroneously flagged as outlier (instead of 0.7, 70 or 7000 points with the original boxplot rule). To characterize the relationship $\alpha(n)$ versus n , $\alpha(n)$ is derived with high accuracy (Q implemented in R) for each distribution of the Pearson and GEV family for 5 sample sizes ($n_i = 10^i$ with $i \in [2,6]$). It appears that $\alpha(n) = A \log(n) + B$ is an accurate model for both the Pearson family (mean of $r^2 = 0.994 \pm 0.005$) and the GEV family ($r^2 = 0.99 \pm 0.01$). [Barbato et al. \(2011\) found the same law for the Gaussian distribution based on empirical considerations only, with reported values of \$A = 0.15\$ and \$B = 1.15\$. For comparison, this study finds \$A = 0.08\$ and \$B = 2\$ for the Gaussian distribution \(\$r^2 = 0.999\$ \).](#)

[The relationship \$\alpha\(n\) = A \log\(n\) + B\$ now needs to be extended to small or non-Gaussian samples.](#) To account for biases emerging at small sample size, an additional ~~constant~~ term is added following Carling (2000): $\alpha(n) = A \log(n) + B + \frac{C}{n}$. The parameter $C = 36$ has been numerically determined with a Monte-Carlo simulation on the distributions of the Pearson family to ensure that the percentage of erroneously flagged outliers corresponds to $\sim 0.1\%$ for $n = 9$ (supplementary material). ~~For the particular case of the~~ [To account for non-Gaussian \(ubiquitous in nature\) populations, \$A = 0.08\$ and \$B = 2\$ \(\$r^2 =\$ will be estimated with a new robust predictor sensitive to the tail weight. ~~0.999~~\).](#)

2.2.3 The Logbox procedure

Let (m_-, m_+) be two ~~positive~~ functions defined as $m_- = (E_3 - E_1)/(E_6 - E_2)$ and $m_+ = (E_7 - E_5)/(E_6 - E_2)$ with $E_i = q(i/8)$ the sample octile. The centered Moors $m = m_- + m_+ - 1.23$ is a [known](#) robust predictor of the kurtosis excess with a breakdown point of 12.5% (Moors 1988, Kim & White 2004). ~~This~~ [However, this](#) study introduces a modified version defined as $m_* = \max(m_-, m_+) - 0.6165$. The parameter m_* is more appropriate than m to determine if a sample is light-tailed or heavy-tailed. For example, a Gaussian distribution ($m_- = m_+ \approx 0.6165$; ~~$m = m_* \approx 0$~~) and a right-skewed distribution with one heavy tail ($m_- = 0.1$ and $m_+ = 1.13$) will share identical m but different m_* . [The Logbox procedure is the following for an unknown sample of size \$n\$:](#)

- 145 (i) ~~If $n \in [3, 9]$, the outlier threshold $[l, u]$ is $l = q(0.50) - \beta \times MAD$ relationships shown in Fig. 1 are $A = g_A(m_*)$ and $u = q(0.50) + \beta \times MAD$ with $\beta = 12.5$.~~
(ii) ~~If $n \geq 9$, $m_* B = g_B$ is computed (bounded by $[0, 2]$) and the boxplot rule is used with $\alpha(n) = g_A(m_*) \log(n) + g_B(m_*) + \frac{36}{n^2}$.~~

150 For very small sample sizes (case i), the median and MAD are preferred, with (Leys et al., 2013) over the boxplot rule because a single outlier would break the estimator m_* . The value of $\beta = 12.5$ has been numerically determined with a Monte Carlo simulation and corresponds to $\sim 0.1\%$ of erroneously flagged outliers in the Pearson family for $n = 9$ (supplementary material). For larger sample sizes (case ii), $g_A(x) = 0.2294e^{2.9416x - 0.0512x^2 - 0.0684x^3}$ ($r^2 = 0.999$) and $g_B(x) = 1.0585 + 15.6960x - 17.3618x^2 + 28.3511x^3 - 11.4726x^4$ ($r^2 = 0.999$) for $x \in [0, 2]$. Each function has been parametrized based on the Pearson and GEV family together (Fréchet has been excluded due to a different behavior, see Fig. 1e,f). The coefficients have been determined with a Monte-Carlo simulation that minimizes the root-mean square error ($N \sim 10^8$).

155 ~~For an unknown sample of size $n \geq 9$, the Logbox procedure is finally the following: m_* is computed (bounded by $[0, 2]$) and the boxplot rule is used with $\alpha(n) = g_A(m_*) \log(n) + g_B(m_*) + \frac{36}{n^2}$.~~

160 2.42.3 Former models

Logbox is compared with four other models (Kimber, 1990; Hubert & Vandervieren, 2008; Schwertman et al., 2004; Leys et al., 2013; Barbato et al., 2011). The first two models (Kim. and Hub.) adjust the boxplot method with respect to the skewness:

$$\begin{cases} l_{Kim.} = q(0.25) - 3 \times (q(0.50) - q(0.25)) \\ u_{Kim.} = q(0.75) + 3 \times (q(0.75) - q(0.50)) \end{cases}$$

165 And

$$\begin{cases} l_{Hub.} = q(0.25) - 1.5 \times h(-MC) \times (q(0.75) - q(0.25)) \\ u_{Hub.} = q(0.75) + 1.5 \times h(MC) \times (q(0.75) - q(0.25)) \end{cases}$$

With the function h defined as $h(MC) = e^{4MC}$ for $MC < 0$ and $h(MC) = e^{3MC}$ for $MC \geq 0$. The Medcouple $MC \in [-1, 1]$ is a robust estimator of S , with an algorithm complexity of $O(n \log n)$ and a breakdown point of 25% (Brys et al., 2004). The third model (Sch.) constructs the lower and upper boundary around the median:

$$\begin{cases} l_{Sch.} = q(0.50) - \frac{Z}{k_n} \times 2(q(0.50) - q(0.25)) \\ u_{Sch.} = q(0.50) + \frac{Z}{k_n} \times 2(q(0.75) - q(0.50)) \end{cases}$$

170

Mis en forme : Police : Times New Roman

Mis en forme : Police : + Corps (Times New Roman)

With k_n a function of the sample size n to adjust for small samples (given as a table in Schwertman et al., 2004) and Z a constant related to the percentage of data captured by $[l_{Sch.}, u_{Sch.}]$, here picked as $Z = 3$ (Gaussian case for the $\pm 3\sigma$ window).

Finally, the last model (Ley.) uses the MAD around the median:

$$\begin{cases} l_{Ley.} = q(0.50) - 3 \times MAD \\ u_{Ley.} = q(0.50) + 3 \times MAD \end{cases}$$

175 Finally, the last model (Bar.) is similar to the Logbox procedure but parametrized on the Gaussian distribution only:

$$\begin{cases} l_{Bar.} = q(0.25) - (0.15 \times \log(n) + 1.15) \times (q(0.75) - q(0.25)) \\ u_{Bar.} = q(0.75) + (0.15 \times \log(n) + 1.15) \times (q(0.75) - q(0.25)) \end{cases}$$

2.2.54 Comparison between models

180 The comparison between models is performed on two sets of residuals obtained from weather stations part of the Global Historical Climatology Network (GHCN-daily) with at least 100 years of daily temperature (2693 stations, 9.4×10^7 days) or daily precipitation (6277 stations, 5.8×10^7 wet days, dry days are excluded). Because this network is used to calibrate products that are remote-sensing based and because suspicious values are routinely flagged (Menne et al., 2012; Xungang et al., 2012), the risk of errors in these century-old stations can be considered small. The residuals are extracted with the robust method described in part II based on non-overlapping bins (bins with less than 80% of data are discarded). To reduce the impact of the extraction method on the analysis, three bin intervals (5, 10 and 20 days) are used to obtain three replicas for each station. The sensitivity of each outlier detection method to the sample size has also been estimated. For each station and for each sample size $l_i = 10i$ (i varying from 1 to 10), $N_i = \frac{100}{i}$ samples are randomly selected and the number of flagged outliers is summed over all the N_i samples (the total number of points is constant, $N_i \times l_i = 1000$).

190 For the ~~four~~five models (Ley., Hub., Kim., Sch., Bar.), the percentage of flagged outliers is computed for each station, and then the mean (± 1 SD) is calculated over all stations. For the Logbox model, this method is not appropriate because the expected number of erroneously flagged outlier per station is less than one ($0.001\sqrt{\sim 10^8} \sim 0.1$). Instead, the percentage of flagged outliers is calculated over the total number of points: $\rho = (\sum n_j^{flagged}) \times 100 / (\sum n_j)$ with j a station. The variability is estimated by subsampling the total number of stations N_s : $\sqrt{N_s}$ sets of $\sqrt{N_s}$ random stations are selected without replacement. The parameter ρ is computed for each set, and the associated variability is calculated on all ρ values (± 1 SD in Fig. 2f and quantiles in Fig. 2c).





Fig. 1. Location of the 4999 light-tailed distributions of the Pearson family (panel a) and the 368 heavy-tailed distributions of the GEV family (panel d) in the (κ_{ex}, S^2) space (kurtosis excess, squared skewness). The coefficients A and B correspond to $\alpha(n) = A \times \log(n) + B + \frac{36}{n}$ used to replace $\alpha = 1.5$ in the boxplot rule. For the Pearson family, they are shown in the (κ_{ex}, S^2) space (panels b and c). For the GEV family (panels e and f), they are shown against a predictor of κ_{ex} the maximum tail weight, defined for right-skewed distributions as $m_* = (E_7 - E_5)/(E_6 - E_2) - 0.6165$ for right-skewed distributions (E_i with $E_i = q(i/8)$ the sample octile).

Mis en forme : fontstyle01

2.3 Results and discussion

The parameter $\alpha = 1.5$ used in the boxplot rule is sensitive to the sample size n , and the relationship $\alpha(n) = A \log(n) + B + \frac{C}{n}$ corrects for this effect for both light-tailed distributions (Pearson family, Fig. 1a) and heavy-tailed distributions (GEV family, Fig. 1d). The value of A, B and C depends on the outlier threshold level and the nature of the distribution. The convention in this study is to set the expected number of erroneously flagged outliers to $f(n) = 0.001\sqrt{n}$, which corresponds to a percentage of type I error of $0.1/\sqrt{n}\%$. This leads to homogeneous A and B values among the Pearson family ($A = 0.8 \pm 0.3, B = 3 \pm 1$, Fig. 1b,c) used to numerically determine $C = 36$ (supplementary material). Because the value of A and B rapidly diverges

for heavy-tailed distributions, a model adapted to the shape of the residuals is required (Fig. 1e,f). To keep this model simple, the asymmetry of a distribution (i.e., the skewness) is ignored in this study in order to only focus on the weight of the heavier tail. Possible outliers might not be flagged on the light tail of an asymmetric distribution (risk of type II error), but residuals with strong asymmetry are usually produced when the range of possible values is semi-bounded (e.g., precipitation in $[0, +\infty[$), which makes the detection of errors trivial (negative precipitation). For this purpose, the parameter m_* is a robust predictor of the kurtosis excess (maximum tail weight with a breakdown point of 12.5%) that has been slightly modified from Kim & White (2004).²²⁰ Finally, $\alpha(n) = g_A(m_*) \log(n) + g_B(m_*) + \frac{36}{n}$ for $n \geq 9$ and $m_* \in [0, 2]$, with the functions g_A and g_B parametrized on both families (Fig. 1e,f). The Fréchet distribution has been excluded because its tails are decaying too rapidly (the A and B coefficients are bounded despite an extreme kurtosis).






Fig. 2. Comparison between [five](#) outlier detection methods performed on two sets of residuals (temperature and precipitation) obtained from weather stations with daily measurements over at least 100 years. The two histograms (panels **a** and **b**) represent aggregated residuals from all stations (for visualization purpose only) and show counts with at least 100 daily occurrences, with the median of the lower/upper threshold displayed for each method. For the methods [Ley.](#) (Leyset al., 2013), Kim. (Kimber, 1990), Sch. (Schwertman et al., 2004), Hub. (Hubert & Vandervieren, 2008) and [Ley.](#) (Ley&Bar. (Barbató et al., 2013, 2011)), the mean percentage ($\pm 1SD$) of flagged data is shown for sample sizes varying from 10 to 100, and for all available points per station ($n \sim 4 \times 10^4$ for the temperature and $n \sim 10^4$ for the precipitation, panels **d** and **e**). For Logbox (panels **c** and **f**), this percentage is calculated by pooling all points, and the variability is estimated with a random resampling of stations (see method). The theoretical threshold is the expected percentage of erroneously flagged outliers ($P_{theo} = f(n) \times \frac{100}{n} = \frac{0.1}{\sqrt{n}}$ %).

The Logbox procedure is tested and compared with [four](#) other models on daily precipitation and temperature residuals from century-old weather stations (Fig. 2). It is firstly visually striking that the outlier threshold from the [four](#) traditional methods cut too many data points not only for the precipitation but also for the temperature residuals (Fig. 2a,b). The percentage of flagged data points per station varies around 2 ± 0.5 to 1.7 ± 1 % for the temperature (Fig. 2d, median of $36634 \approx 4 \times 10^4$ days per station), and from 5.5% (Sch) to 4.1% (Bar.) to 10.5% (Ley.) for the precipitation (Fig. 2e, median of $8352 \approx 10^4$ wet days per station).

The reason for the large discrepancy between observed and expected percentage of flagged outliers ($\sim 0.7\%$ based on the boxplot rule) is that these [four](#) methods have been designed for *nearly*-Gaussian residuals. Even daily temperatures are diverging from normality because the fitting model used to extract residuals from the time series minimizes the root-mean-

Mis en forme : Police :Times New Roman, Non Gras

Mis en forme : Police :NonGras

Mis en forme : Police :Times New Roman

Mis en forme : Police :Italique

square-error. The anomalies are therefore more concentrated around 0 than those produced by a Gaussian, but with larger extremes (Fig. 2a, leptokurtic distribution). ~~For small samples of~~ Only the *Bar.* model correctly captures outliers present in the temperature residuals (0.17% data points flagged) as it accounts for large sample size effects (logarithmic law in α similar to Logbox). However, *Bar.* fails at capturing outliers in the precipitation residuals because this method has been parametrized on the Gaussian only. For small samples, the type I error is even higher in all traditional methods due to the inaccuracy of the quantiles: from 1.4% (*Bar.*) to 4.25% (*Sch.*) to 7% (*Hub.*) of points temperature residuals are cut for $n = 20100$ (Fig. 2d). The precipitation is less affected by the sample size effect because the type I error was already high in large samples. This analysis proves that none of the traditional former methods is suitable to the outlier detection detect outliers in non-Gaussian residuals.

Mis en forme : Police :Italique

In comparison, the Logbox procedure shows a percentage of flagged outliers close to the expected values for large sample sizes (Fig. 2c), with 0.0006 ± 0.0003 % for the temperature (expected value of 0.0005%) and 0.0017 ± 0.0009 % for the precipitation (expected value of 0.001%). These results are surprisingly accurate knowing that 12.5% of the extreme values are disregarded for robustness reasons (m_*), and also knowing that Logbox has only been parametrized on theoretical distributions (Pearson & GEV family). For smaller sample sizes ($n < 30$ in Fig. 2f), the precipitation residuals are cut too frequently (~ 0.2 — 0.325 %) compared to the expected threshold (~ 0.03 %), but the temperature residuals are not cut enough. The constant parameter used to correct for a sample size effect ($C = 36$) is only adapted to nearly-Gaussian residuals, and it cannot be better estimated because any predictor (such as m_*) becomes inaccurate at smaller sample sizes. However, the percentage of flagged outliers remains within one order of magnitude of the expected threshold, which is a reasonable compromise between type I errors (precipitation) and type II errors (temperature).

Mis en forme : Police :NonItalique

To summarize, Logbox enhances the boxplot rule by considering the sample size effect and by adapting the cutting thresholds to the data. This method has been implemented in the function *ctbi.outlier* (in the R package **ctbi**) that will be used to flag potential outliers in the residuals obtained by the aggregation procedure described in part II.

3 Part II, the **ctbi** procedure

3.1 Context

This second part is dedicated to the pre-processing, partial imputation and aggregation of univariate time series. In order to flag outliers, one first needs to produce residuals that represent the variability around the *signal*. In its simplest form, the time series y_t is represented with the following additive decomposition (Hyndman & Athanasopoulos, 2018, 2021): $y_t = T_t + S_t + \epsilon_t$, with T_t a long-term trend, S_t a cyclic component (originally, *seasonal component* but the term cyclic is preferred here as it

is more generic) with period τ ($\forall t, S_t = S_{t+\tau}$) and ε_t the residuals that are considered to be stationary. A popular algorithm that performs this decomposition is the Seasonal and Trend decomposition using Loess (or **STL**, Cleveland et al., 1990), that is robust to the presence of outliers. The enhanced version of the algorithm, **STLplus** (Hafen, 2016), is also robust to the presence of missing values and data gaps. Unfortunately, there are three major drawbacks to using **STLplus** in the general case: (i) This algorithm has specifically been designed for signals showing seasonal patterns, which makes it less relevant for other types of data; (ii) The long-term trend based on loess needs several input parameters (s.window, s.degree,...) and the decomposition is therefore not unique; (iii) The algorithm has a complexity of $O(n^2)$ due to the loess, which is resource intensive and not adapted to long time series ($n > 10^7$). In particular, the first point explains why the function *tsoutliers* needs to use a smoothing function (Friedman, 1984) to complement the **STL** procedure.

A new robust and nonparametric procedure (**ctbi**) is proposed instead to calculate T_t and S_t using non-overlapping bins. Outliers are flagged in the residuals ε_t with the Logbox method described in part I, and imputation is performed using $T_t + S_t$ if the cyclic pattern is strong enough, which is quantified by a new index introduced in this study (the Stacked Cycles Index or SCI). Bins with sufficient data can finally be aggregated, while other bins are discarded. The procedure is simple (entirely described in Fig. 3), the long-term trend T_t is unique and non-parametrized (based on linear interpolations crossing each bin), the cyclic component S_t is simply the mean stack of bins using detrended data (equivalent to STL for periodic time series). The algorithm complexity is of the order of $O(n \log(n))$ because the loess is not necessary anymore. In the following, the procedure is first described more in details and then applied to three case studies (a temperature, precipitation and methane dataset) that have been contaminated with outliers, missing values and data gaps. Comparison with the raw data demonstrates the reliability of the **ctbi** procedure, whose performance is compared to *tsoutliers*.

3.2 Method

3.2.1 Definitions

Bin: a bin is a time window characterized by a left *side* (inclusive), a right *side* (exclusive), a *center* and a *period* (e.g., 1 year in Fig. 3a). Any univariate time series can be decomposed in a sequence of non-overlapping bins, with the first and last data point contained in the first and last bin, respectively (Fig. 3a). The *bin size* n_{bin} is the rounded median of the number of points (including NA values) present in each non-empty bin. A bin is *accepted* when its number of non-NA data points is above $n_{bin}(1 - f_{NA})$ with $f_{NA} \in [0,1]$ the maximum fraction of NA values per bin (input left to the user). Otherwise, the bin is *rejected* and all its data points are set to NA (Fig. 3a, bin 4).

Long-term trend: the *long-term trend (median based)* is a linear interpolation of the median values associated with each side (calculated between two consecutive centers, see Fig. 3a). A side value is set as missing if the number of non-NA data points

(between the two nearest consecutive centers) is below $n_{bin}(1 - f_{NA})$. To solve for boundaries issues and missing sides values, the interpolation is extended using the median value associated with each center (bin 1, 3 & 5 in Fig. 3a). Once the outliers have been quarantined, the *long-term trend (mean based)* will be calculated following the same method but using the mean instead of the median (Fig. 3c).

Cycle: the *cycle (median based)* is composed of n_{bin} points that are the medians of the stack of all accepted bins with the long-term trend (median based) removed (Fig. 3b1). Once the outliers have been quarantined, the *cycle (mean based)* will be the mean stack of accepted bins with the long-term trend (mean based) removed (bin 2, 3 & 5 in Fig. 4a). The cyclic component S_t is the sequence of consecutive cycles.




Fig. 3. Example of the aggregation procedure with the following inputs: bin side = 2020-06-01, bin period = 1 year, $f_{NA} = 0.2$ (minimum of 10 months of data for a bin to be accepted) and $SCI_{min} = 0.6$ (cyclic imputation level). The bin 4 has been rejected because it contains only 6 months of data (panel a). Two outliers have been flagged in the residuals (detrended and deseasonalized data, panel b.2). After the outliers have been replaced with NA values, the bin 1 has been rejected (9 months of data), and the long-term trend and cycle have been updated using the mean instead of the median (panel c). A point in bin 3 has been imputed based on the cyclicity ($SCI_{min} \leq SCI = 0.61$).

Stacked Cycles Index: $SCI \leq 1$ is an adimensional parameter quantifying the strength of a cycle based on the variability around the mean stack (Fig. 4). Its structure is similar to another index developed in a former study (Wang et al., 2006), however a factor of N_{bin}^{-1} has been added to correct for a bias emerging at a small number of bins (N_{bin} is the number of accepted bins). This correcting factor has been calculated based on stationary time series of Gaussian noise (with therefore a null cyclicity per definition, see supplementary material).




Fig. 4. The Stacked Cycles Index ($SCI \leq 1$) quantifies the strength of the cyclicity associated with the period of a bin. The long-term trend (mean based) is first removed to compute the total sum of squares (panel a). Then the cyclic component (mean based) is also removed to compute the sum of squared residuals (panel b). SCI is the coefficient of determination minus N_{bin}^{-1} to correct for a bias emerging at a small number of bins, with N_{bin} the number of accepted bins (here $N_{bin} = 3$, panel c).

3.2.2 Ctbi procedure

Inputs.

1. The univariate time series (1st and 2nd column: time and raw data, respectively).
2. One bin center or one bin side (e.g., 2020-06-01).
3. The period of the bin (e.g., 1 year).
4. The aggregation operator (mean, median or sum).
5. The range of possible values (default value $y_{lim} \in] - \infty, +\infty[$).
6. The maximum fraction of NA values per bin (default value $f_{NA} = 0.2$).
7. The A, B, C coefficients used in the Logbox method (automatically calculated by default, *coeff.outlier* = 'auto').

8. The minimum SCI for imputation (default value $SCI_{min} = 0.6$).

Outputs.

1. The original dataset, with 9 columns: (i) time; (ii) outlier-free and imputed data; (iii) index of the bins associated with each data points (the index is negative if the bin is rejected); (iv) long-term trend; (v) cyclic component; (vi) residuals (including the outliers); (vii) quarantined outliers; (viii) value of the imputed data points; (ix) relative position of the data points in their bins, between 0 (the point falls on the left side) and 1 (the point falls on the right side).
350
 2. The aggregated dataset, with 10 columns: (i) aggregated time (center of the bins); (ii) aggregated data; (iii) index of the bin (negative value if the bin is rejected); (iv) start of the bin; (v) end of the bin; (vi) number of points per bin (including NA values); (vii) number of NA values per bin, originally; (viii) number of outliers per bin; (ix) number of imputed points per bin; (x) variability associated with the aggregation (standard deviation for the mean, MAD for the median and nothing for the sum).
355
 3. The mean cycle, with 3 columns: (i) time boundary of the first bin with n_{bin} points equally spaced; (ii) the mean value associated with each point; (iii) the standard deviation associated with the mean value.
360
 4. A summary of the bins: the Stacked Cycle index (SCI), the representative number of data points per bin (n_{bin}) and the minimum number of data points for a bin to be accepted ($n_{bin min}$).
 5. A summary of the Logbox output: the coefficients A , B and C , m_* , the number of points used, the lower/upper outlier threshold.
- 365 Step 1, data screening. The bin size n_{bin} is calculated; values above or below y_{lim} are set to NA; the number of accepted bins N_{bin} is assessed; all data points within rejected bins are set to NA; the long-term trend and cycle (both median based) are calculated (Fig. 3a,b1).
- Step 2, outliers. Outliers are flagged in the residuals (detrended and deseasonalized data) using Logbox (Fig. 3b2); outliers are quarantined and their values are set to NA; the number of accepted bins N_{bin} is updated; all data points within newly rejected
370 bins are set to NA (bin 1 in Fig. 3c).
- Step 3, long-term trend and cycle (mean based): The long-term trend and the cycle are calculated using the mean instead of the median (Fig. 3c); SCI is calculated (Fig. 4).
- Step 4, imputation: If $SCI > SCI_{min}$, all NA values in accepted bins are imputed with the long-term trend + the mean cycle (imputation bounded by y_{lim}). Repeat Step 3 and Step 4 three times to reach convergence.
- 375 Step 5, aggregation: Accepted bins are aggregated around their center.

3.2.3 Case studies

Three univariate datasets are chosen to illustrate the potential of the aggregation procedure (Fig. 5, first column). The first dataset is an in-situ temperature (in °C) measured during summer in the canopy of an Oak woodland of California (month of August, temporal resolution of 5 min), and provided by the National Ecological Observatory Network (NEON 2021, site SJER). The second dataset is an in-situ daily precipitation record (in mm) measured at the station of Cape-Leeuwin (South westerly coast of Australia) from 1990 to 2020 and available on the Global Historical Climatology Network (Menne et al., 2012; Xungang et al. 2012). The last dataset is a Methane proxy record (in ppbv) published in Louergue et al. (2008) that covers 800000 years with irregular timesteps (varying from 1 to 3461 years, with a median of 311 years). None of the datasets contain obvious outliers or large data gap.


3.2.4 Contamination of the datasets

To test for the robustness of the aggregation procedure, the three raw datasets are contaminated by 30% (Fig 5, second column) with the use of three data gap (20% of the dataset), random NA values (9.5% of the dataset) and outliers (0.5% of the dataset). The three data gaps are picked with random length and position. The position of the outliers and the NA values follows a Poisson law. The value of the outliers is picked equal to $y_{min} - \frac{1}{2}(\mu - y_{min})$ or $y_{max} + \frac{1}{2}(y_{max} - \mu)$ with y_{min} , y_{max} and μ respectively the minimum, maximum and mean of the dataset (temperature and methane datasets). The precipitation is supposed to follow a heavy-tail distribution (extremes are more frequent), and negative values are impossible, which is why outlier values are set to $1.6 \times y_{max}$ instead (supplementary material).

3.2.5 Aggregation of the datasets

Each dataset (raw and contaminated version) is consecutively aggregated twice (Fig. 6). The temperature dataset is aggregated (using the mean) every hour ($n_{bin} = 12$) and then every day ($n_{bin} = 24$). The precipitation dataset is aggregated (using the sum) every month ($n_{bin} = 31$) and then every year ($n_{bin} = 12$). The methane dataset is aggregated (using the mean) every 2000 years ($n_{bin} = 4$) and then every 20000 years ($n_{bin} = 10$). For each dataset, the mean cycle of the second level of aggregation is shown in Fig. 6 (second column). The aggregation inputs are chosen as default values. The only exceptions are $coeff.outlier = NA$ and $SCI_{min} = NA$ for the raw data (outliers are not checked, data are not imputed), $f_{NA} = 1$ for the Methane dataset (bins with at least 1 non-NA data point are accepted due to the high irregularity in the sampling frequency) and $y_{lim} = [0, +\infty[$ for the precipitation dataset (negative precipitation are impossible). The number of false positive (real data points flagged as outliers) and false negative (outliers that have not been flagged) are counted during the first level of aggregation (Table 1), and compared with the $tsoutliers$ function with $\lambda = \text{"auto"}$, which means that the residuals have been transformed to follow a Gaussian with the Cox-Box method (Box & Cox, 1964), or $\lambda = \text{NULL}$, which means the original

residuals are not transformed. The boxplot rule in *tsoutliers* uses $\alpha = 3$, and the long-term trend or cyclic component are not available for comparison.



410 **Fig. 5.** Raw and contaminated versions of the three datasets used as case studies: temperature (panel **a**), precipitation (panel **b**) and methane (panel **c**). The sampling frequency is given in parenthesis. The contaminated versions contain three large data gaps (20% of the datasets), random missing values (9.5%) and random outliers (0.5%) set as a constant level.

3.3 Results and discussion

415 The three univariate time series have been chosen as case studies due to their various statistical characteristics that are commonly seen in the scientific or economic field (Fig. 5, 1st column). The long-term trend follows smooth or moderate variations in the temperature and precipitation datasets, but shows a much higher volatility in the methane dataset. The cyclic pattern varies from strong diurnal cycles (temperature) and moderate seasonal cycles (precipitation) to no apparent cyclicality

over a period of 20000 years (methane). The detrended and deseasonalized residuals follow distributions from Gaussian (temperature) or seemingly Exponential (methane) to heavy-tailed (precipitation). Finally, the sampling frequency goes from sub-hourly (temperature) or daily (precipitation) to highly variable (1 to 3461 years, methane). To test the limits of the aggregation procedure, these three datasets are *severely* contaminated by data gaps, outliers and missing values (Fig. 5, 2nd column).

425




Fig. 6. Aggregation of the temperature (panel a), precipitation (panel b) and methane (panel c) in two consecutive levels: 1 (thin lines) and 2 (bold lines). Only the first level of aggregated precipitation is shown for clarity. Black and red colors are associated with the raw and

430 contaminated datasets, respectively. The mean cycles of the second level of aggregation are shown in the second column, with their SCI displayed (the raw and contaminated versions share similar values).

The first level of aggregation recovers most of the destroyed signal with $\sim 80\%$ of the bins being accepted for all three datasets (Fig. 6). In these accepted bins, all outliers have been correctly flagged (Table 1). The mean percentage of difference between the contaminated and raw aggregates (level 1) is virtually zero for the temperature ($0 \pm 0.1\%$), the methane ($-0.1 \pm 2\%$) and the precipitation ($0 \pm 17\%$). For the [Methanemethane](#) dataset, the only false positive (Table 1) is located at the beginning of the time series (modern time), because the anthropogenic change in CH_4 is unprecedented when compared to the geological history (the long-term fit does not capture the abrupt increase due to climate change). In comparison, the function *tsoutliers* successfully flags the outliers in the contaminated temperature and methane datasets (with the Cox-Box method), however it fails with the contaminated daily precipitation dataset (Table 1). This comes from the inability of *tsoutliers* to handle heavy tailed distributions, creating 55 false negatives (all outliers have been missed) with the Cox-Box method and 1125 false positives without it, due to the limitation of the boxplot rule using a constant $\alpha = 3$ (see part I).

Procedure	ctbi			<i>tsoutliers</i> (with/without Cox-Box)		
	T	P	CH ₄	T	P	CH ₄
Number of false positives	0	0	1	0 / 0	0 / 1125	0 / 3
Number of false negatives	0	0	0	3 / 0	55 / 0	0 / 2

445 **Table 1.** Number of false positives (real data points flagged as outliers, type I error) and false negatives (outliers that have not been flagged, type II error) for the contaminated Temperature (n=8952), Precipitation (n=10949) and Methane (n=2103) datasets shown in Fig. 5 with the *ctbi* procedure and the *tsoutliers* function (with/without the Cox-Box method).

450 The second level of aggregation has been performed to test for the cyclicity in the signal (Fig. 6, 2nd column) using the mean cycles and their associated Stacked Cycles Index (Fig. 4). The raw and contaminated mean cycles share similar magnitude within 1 standard deviation on the mean, and their SCI are the same: -0.02 for the methane (no apparent cycles of 20000 years period), 0.65 for the precipitation (moderate seasonality) and 0.88 for the temperature (strong diurnal cycles). The SCI reveals itself being useful when comparing signals of different nature or periodicities, which is not possible for seasonal indices that only focuses on one field (e.g., hydrology) or data format. (e.g., monthly) such as the seasonality index of Feng et al. (2013). The cyclicity seen in the temperature and precipitation is strong enough to impute the missing data in all accepted bins, which further improves the reconstruction of the signal. Because *SCI* has a similar structure than a coefficient of determination, imputations based on high *SCI* (> 0.6) are respecting the original signal, which is sometimes not the case with a linear

460 interpolation. These three case studies demonstrate that **ctbi** is capable of aggregating signals of poor quality that have a
stationary variance in the residuals. The next section explains how to handle more complex time series.

3.4 Limits & recommendations

465 The **ctbi** procedure complements the expert-knowledge related to a dataset, but it does not replace it. In particular, this
procedure is not capable of detecting long periods of instrument failure or human error, and it is essential to flag them manually
and/or visually before running **ctbi**. This procedure also presents difficulties to pre-process signals with a complex seasonality
associated with residuals of non-stationary variance. A typical example is a daily precipitation record with a pronounced
monsoon: several months of droughts (low variability in the signal) are followed by few weeks of severe floods (high
variability). These two periods do not have the same statistical characteristics, and need to be treated separately. In this
470 situation, two pools of bins can be created using the MAD as a robust indicator of variability within each bin. The procedure
is the following: (i) apply **ctbi** with the median operator (do not flag outliers or impute data, *coeff.outlier* = NA and
SCI_{min} = NA) so that each bin will be associated with a specific MAD; (ii) Flag bins with a low MAD ('dry' season) and a
high MAD ('wet' season); (iii) split the raw data into two datasets of bins with a low and high MAD, respectively; (iv) apply
ctbi separately to each dataset to flag outliers and/or impute data; (v) merge the two datasets. This procedure is successfully
475 applied to a soil respiration dataset (supplementary material).

Other issues can usually be addressed by varying the inputs: period of the bin, maximum ratio of missing values per bin (f_{NA}),
and cyclic imputation level (SCI_{min}). It is recommended to pick the period of a bin so that it contains on average between 4
and ~50 data points. Below 4 would decrease the breakdown point to unsafe levels (1 outlier would be enough to contaminate
480 the bin), and above 50 would produce a long-term trend that might not properly capture the variability in the signal. A
maximum of 20% of the bin can be missing by default ($f_{NA} = 0.2$), but when data are sparse and irregularly distributed, a
value of $f_{NA} = 1$ is possible (example with the Methane dataset: bins with only 1 data point were accepted). Finally, the
imputation level (default of $SCI_{min} = 0.6$) can vary between 0 (forced imputation even without cyclic pattern) and 1 (no
imputation).

485 4 Conclusion (Part I & II)

Although univariate time series are the simplest type of temporal data, this study reveals a lack of consensus in the literature
on how to objectively flag outliers especially in raw data of poor quality. In part I, a comparison between outlier detection
methods is performed on daily residuals from century-old weather stations (precipitation & temperature data). All traditional
outlier detection methods flag extreme events as outliers too frequently (type I error). The alternative procedure developed in
490 this study (Logbox) improves the boxplot rule by replacing the original $\alpha = 1.5$ with $\alpha = A \log(n) + B + \frac{36}{n}$, with A and B

determined with a predictor of the ~~kurtosis excess~~ [maximum tail weight](#) (m_*). Logbox is parametrized on two families of distributions (Pearson & Generalized Extreme Value), and the theoretical percentage of type I error decreases with the sample size ($p_{theo} = \frac{0.1}{\sqrt{n}}\%$). Logbox therefore produces cutting thresholds that are tailored to the shape and size of the data, with a good match between observed and expected type I errors in the precipitation and temperature residuals.

495

In part II, a pre-processing procedure (**ctbi** for cyclic/trend decomposition using bin interpolation) cleans, decomposes, imputes and aggregates time series based on data binning. The strength of the cyclic pattern within each bin is assessed with a novel and adimensional index (the Stacked Cycles Index) inspired by the coefficient of determination. The **ctbi** procedure is able to filter contaminated data by selecting bins with sufficient data points (input: f_{NA}) which are then cleaned from outliers (input: *coeff.outlier*). The cyclic pattern within each bin is evaluated (*SCI*) and missing data are imputed in accepted bins if the cyclicity is strong enough (input: SCI_{min}). Most of the signal can be retrieved from univariate time series with diverse statistical characteristics, illustrated in this study with a temperature, precipitation and methane datasets that have been contaminated with gaps and outliers. Limits in the use of **ctbi** ~~have been~~ [are](#) acknowledged for signals with a long-period of instrument failure, but also for signals presenting a complex seasonality. The last situation can be handled by splitting the raw data into two (or more) datasets containing bins with similar variability quantified by the Mean Absolute Deviation (MAD). The pre-processing procedure is then separately applied to each dataset to correctly identify outliers. It is strongly recommended to examine the data before and after using **ctbi** to ensure that rejected bins and flagged outliers seem reasonable, and to be transparent about the inputs used in your future study.

510

Author contribution

F.R.: Design, writing, coding.

Competing interests

515 The author declares no competing interests.

Acknowledgement

The author would like to warmly thank Dr. Rob Hyndman for his advice, as well as the ~~three~~ reviewers of this study. Dr. Jens Schumacher has particularly stimulated the improvement of the Logbox method. ~~The author is also grateful to the Fonds de~~
520 ~~Dotations O that funded this project.~~

Data availability

The GHCN dataset is available on <https://www1.ncdc.noaa.gov/pub/data/ghcn/daily/>. The Methane dataset is available on <https://doi.org/10.1038/nature06950>. The temperature dataset is available on <https://doi.org/10.48443/2nt3-wj42>.

525

Code availability

The **ctbi** package is available on the comprehensive R Archive Network (CRAN). The code and data used in the study & the supplementary material are available on https://github.com/fritte2/ctbi_article.

530 References

- Aguinis, [Herman, Ryan K.H.](#), Gottfredson, [R. K.](#), and [Harry-Joo](#), H.: Best-practice recommendations [Practice Recommendations](#) for defining-identifying [Defining, Identifying, and handling outliers](#). [Handling Outliers](#), *Organizational Research Methods* 16, no. 2 (<https://doi.org/10.1177/1094428112470848>, 2013): 270-304.
- Barbato, G., Barini, E. M., Genta, G., and Levi, R.: Features and Performance of Some Outlier Detection Methods, *Journal of Applied Statistics*, <https://doi.org/10.1080/02664763.2010.545119>, 2011.
- Borchers, [Hans W.](#), and Maintainer [Hans W. Borchers](#). "H.: Package 'pracma'." (Pracma', R Package Version 2.4.2, 2021).
- Box, [George E. P.](#), and [David R. Cox](#). "D. R.: An analysis of transformations." [Transformations](#), *Journal of the Royal Statistical Society: Series B (Methodological)* 26, no. 2 (1964): 211-243 <https://doi.org/10.1111/j.2517-6161.1964.tb00553.x>, 1964.
- Brys, [Guy](#), [Mia G.](#), [Hubert, M.](#), and [Anja Struyf](#). "A robust measure: A Robust Measure of skewness." [Skewness](#), *Journal of Computational and Graphical Statistics* 13, no. 4 (<https://doi.org/10.1198/106186004X12632>, 2004): 996-1017.
- Carling, [Kenneth](#), K.: Resistant outlier rules [Outlier Rules](#) and the non-Gaussian case. [Case](#), *Computational Statistics & Data Analysis* 33, no. 3 ([https://doi.org/10.1016/S0167-9473\(99\)00057-2](https://doi.org/10.1016/S0167-9473(99)00057-2), 2000): 249-258.
- Chandola, [Varun](#), [Arindam V.](#), [Banerjee, A.](#), and [Vipin Kumar](#), V.: Anomaly detection [Detection: A survey](#). [Survey](#), *ACM computing surveys (CSUR)* 41, no. 3 (<https://doi.org/10.1145/1541880.1541882>, 2009): 1-58.
- Cleveland, [Robert B.](#), [William S.](#), [Cleveland, Jean E. W. S.](#), [McRae, J. E.](#), and [Irma Terpenning](#), I.: STL: A seasonal trend decomposition. *J. Off. Stat* [Seasonal-Trend Decomposition Procedure Based on Loess \(with Discussion\)](http://bit.ly/stl1990), *Journal of Official Statistics* 6, no. 1 (<http://bit.ly/stl1990>, 1990): 3-73.
- Feng, [Xue](#), [Amilcare X.](#), [Porporato, A.](#), and [Ignacio Rodriguez-Iturbe](#), I.: Changes in rainfall seasonality [Rainfall Seasonality in the Tropics](#). *Tropics*, *Nature Climate Change* 3, no. 9 (2013): 811-815-15. <https://doi.org/10.1038/nclimate1907>, 2013.
- Friedman, [Jerome J. H.](#): A Variable Span Smoother. October. Vol. [variable span smoother](#). Stanford Univ CA lab for computational statistics, 1984. <https://doi.org/10.2172/1447470>, 1984.
- Hafen, [Ryan](#). "Stlplus: enhanced seasonal decomposition of time series by Loess." R-package version.: Package 'Stlplus', R Package Version 0.5-1 (<https://doi.org/10.1080/01621459.1986.10478363>, 2016).
- Hoaglin, [David D. C.](#), [Boris Iglewicz, B.](#), and [John W. Tukey](#), J. W.: Performance of some resistant rules [Some Resistant Rules](#) for outlier labeling. [Outlier Labeling](#), *Journal of the American Statistical Association* 81, no. 396 (1986): 991-999 <https://doi.org/10.1080/01621459.1986.10478363>, 1986.
- Hubert, [Mia M.](#), and [Ellen Vandervieren](#), E.: An adjusted boxplot [Adjusted Boxplot](#) for skewed distributions. [Skewed Distributions](#), *Computational statistics & data analysis* *Statistics and Data Analysis* 52, no. 12 (<https://doi.org/10.1016/j.csda.2007.11.008>, 2008): 5186-5204.
- Hyndman, R. J., & Athanasopoulos, G. (2018). *OTexts*: Forecasting: principles and practice, 2nd/3rd edition, OTexts: Melbourne, Australia. OTexts.com/[fpp2](https://otexts.com/fpp2). Accessed on [fpp3](https://otexts.com/fpp3), 2021/10/13, last access: 21 December 2022.
- Hyndman, Rob J., George Athanasopoulos, Christoph Bergmeir, Gabriel Caceres, Leanne Chhay, Mitchell O'Hara Wild, Fotios Petropoulos, Slava Razbash, and Eazo Wang. "Package 'forecast'." [Online]. <https://cran.r-project.org/web/packages/forecast/forecast.pdf> (2020).

Mis en forme : Police : NonItalique

Mis en forme : Police : NonItalique

Mis en forme : Accentuation

Hyndman, R. J., and Khandakar, Y.: Automatic Time Series Forecasting: The Forecast Package for R, *Journal of Statistical Software* 27, <https://doi.org/10.18637/jss.v027.i03>, 2008.

Jenkinson, Arthur A. F.: The frequency distribution of the annual maximum (or minimum) values of meteorological elements. "Minimum Values of Meteorological Elements, *Quarterly Journal of the Royal Meteorological Society* 81, no. 348 (<https://doi.org/10.1002/qj.49708134804>), 1955): 158-174.

Kim, T. H., and White, H.: On More Robust Estimation of Skewness and Kurtosis, *Finance Research Letters* 1, [https://doi.org/10.1016/S1544-6123\(03\)00003-5](https://doi.org/10.1016/S1544-6123(03)00003-5), 2004.

Kimber, A. C.: Exploratory data analysis for possibly censored data from skewed distributions. "Journal of the Royal Statistical Society: Series C (Skewed Distributions, Applied Statistics)" 39, no. 4 (<https://doi.org/10.2307/2347808>), 1990): 21-30.

Kim, Tae Hwan, and Halbert White. "On more robust estimation of skewness and kurtosis." *Finance Research Letters* 1, no. 1 (2004): 56-73.

Leys, Christophe, Olivier C. Ley, Philippe O. Klein, Pierre Bernard, and Laurent Licata. "Detecting outliers: Do not use standard deviation around the mean, use absolute deviation around the median." *Journal of experimental social psychology* 49, no. 4 (2013): 764-766 (<https://doi.org/10.1016/j.jesp.2013.03.013>), 2013.

Loulergue, Laetitia, Adrian L. Schilt, Renato A. Spahni, Valérie R. Masson-Delmotte, Thomas V. Blunier, Bénédicte T. Lemieux, Jean-Marc B. Barnola, Dominique J. M. Raynaud, Thomas F. D. Stocker, T. F., and Jérôme Chappellaz. "Orbital and millennial scale features of atmospheric CH₄ over the past 800,000 years." *Nature* 453, no. 7193 (2008): 383-386 (<https://doi.org/10.1038/nature06950>), 2008.

Menne, Matthew J., Inke Durre, Russell S. L. Vose, Byron E. R. S. Gleason, B. E., and Tamara G. Houston. "An overview of the global historical climatology network daily database." *Global Historical Climatology Network-Daily Database, Journal of atmospheric and oceanic technology* 29, no. 7 (2012): 897-910-910. <https://doi.org/10.1175/JTECH-D-11-00103.1>, 2012.

Moors, J. J. A. "A quantile alternative for kurtosis." *Journal of the Royal Statistical Society: Series D (Kurtosis, The Statistician)* 37, no. 1 (<https://doi.org/10.2307/2348376>), 1988): 25-32.

NEON (National Ecological Observatory Network). Single aspirated air temperature, RELEASE-2021 (DP1.00002.001). <https://doi.org/10.48443/2nt3-wj42>. Dataset accessed from <https://data.neonscience.org> on October 13, 2021 (<https://doi.org/10.48443/2nt3-wj42>), 2021, last access: 13 October 2021.

Pearson, Karl. "X. Contributions to the mathematical theory of evolution. - II. Skew variation in homogeneous material." *Philosophical Transactions of the Royal Society of London, Series A*, 186 (1895): 343-414. <https://doi.org/10.1098/rsta.1895.0010>, 1895.

Pearson, Karl. "XI. Mathematical contributions to the theory of evolution. - X. Supplement to a memoir on skew variation." *Philosophical Transactions of the Royal Society of London, Series A, Containing Papers of a Mathematical or Physical Character* 197, no. 287-299 (1901): 443-459 (<https://doi.org/10.1098/rsta.1901.0023>), 1901.

Pearson, Karl. "IX. Mathematical contributions to the theory of evolution. - XIX. Second supplement to a memoir on skew variation." *Philosophical Transactions of the Royal Society of London, Series A, Containing Papers of a Mathematical or Physical Character* 216, no. 538-548 (1916): 429-457 (<https://doi.org/10.1098/rsta.1916.0009>), 1916.

Pearson, Ronald R. K. "Outliers in process modeling and identification." *IEEE Transactions on control systems technology* 10, no. 1 (<https://doi.org/10.1109/87.974338>), 2002): 55-63.

Ranjan, Kumar, Gaurav, B. K. G., Rajanarayan Prusty, B. P., and Debashisha Jena. "Review of preprocessing methods for univariate volatile time series in power system applications." *Electric Power Systems Research* 191, (<https://doi.org/10.1016/j.epsr.2020.106885>), 2021): 106885.

Reiss, Rolf Dieter, Michael Thomas, and R. D. Reiss. *Statistical analysis of extreme values* (Birkhäuser Basel): *Statistical Analysis of Extreme Values: With Applications to Insurance, Finance, Hydrology and Other Fields*, 3rd Edition, <https://doi.org/10.1007/978-3-7643-7399-3>, 2007.

Vol. 2. Basel: Birkhäuser, 1997.

Mis en forme : Police : NonItalique

Mis en forme : Police : NonItalique

Schwertman, Neil N. C., Margaret Ann Owens, M. A., and Robiah Adnan. "A simple more general boxplot method. R: A Simple More General Boxplot Method for identifying outliers." *Identifying Outliers, Computational statistics & data analysis* 47, no. 1 (<https://doi.org/10.1016/j.csda.2003.10.012>, 2004): 165-174.

Tukey, John J. W. (Pearson). *Exploratory data analysis. Vol. 2. Data Analysis*, 1st edition, 1977.

620 Wang, Xiaozhe, Kate X. Smith, K., and Rob Hyndman. "R. J.: Characteristic-based clustering Based Clustering for time series data." *Time Series Data Mining, Data Mining and Knowledge Knowledge Discovery* 13, no. 3 (<https://doi.org/10.1007/s10618-005-0039-x>, 2006): 335-364.

625 Xungang Yin, Steven Anthony, Ron Ray, Russell S. Vose, Byron E. Gleason, and Tamara G. Houston (2012): Global Historical Climatology Network Daily (GHCN Daily), Version 3. NOAA National Climatic Data Center. doi:10.7289/V5D21VHZ (2021/10/13).