

1 **Response of active catchment water storage capacity**
2 **to the prolonged meteorological drought and**
3 **asymptotic climate variation**

4

5 **Jing Tian¹, Zhengke Pan^{1,2}, Shenglian Guo^{1*}, Jiabo Yin¹, Yanlai Zhou¹, Jun Wang¹**

6

7 ¹State Key Laboratory of Water Resources and Hydropower Engineering Science,

8 Wuhan University, Wuhan 430072, China

9

10 ²Changjiang Institute of Survey, Planning, Design and Research, Wuhan, 430010,

11 China

12

13 *Corresponding author. Email: slguo@whu.edu.cn

14 **Abstract:** Studies on the hydrological response to continuous extreme and asymptotic
15 climate change can improve our ability to cope with the intensified water-related
16 problems. Most of the literature focused on the runoff response to climate change, while
17 neglecting the impacts of the potential variation in the [active catchment water storage](#)
18 [capacity \(ACWSC\)](#) that plays an essential role in the transfer of climate input to the
19 catchment runoff. This study aims to systematically identify the response of the
20 ACWSC to the long-term meteorological drought and asymptotic climate change.
21 Firstly, the time-varying parameter is derived to reflect the ACWSC periodic/abrupt
22 variations in both drought and non-drought periods. Secondly, the change points and
23 varying patterns of the ACWSC are analyzed based on the Bayesian change point
24 analysis with multiple evaluation criteria. Finally, various catchment properties and
25 climate characteristics are used to explore the possible relationship between these
26 variables and the temporal variation characteristics of the ACWSC. The catchments that
27 suffered from the prolonged meteorological drought in southeast Australia were
28 selected as the case study. Results indicate that: (1) the increase of amplitude change in
29 the ACWSC is observed in 83/92 catchments during the prolonged drought period, and
30 significant shifts in the mean value of the ACWSC are detected in 77/92 catchments;
31 (2) the average response time of the ACWSC for all 92 catchments with significant
32 changes is 641.3 days; (3) the values of the ACWSC are changed significantly in the
33 catchments with small areas, low elevations, small slope ranges, large forest coverage,
34 and high soil water holding capacities. This study could enhance our understanding of

35 the variations in catchment property under climate change.

36 **Keywords:** catchment water storage capacity; prolonged meteorological drought;

37 extreme and asymptotic climate change; southeast Australia

38 **1. Introduction**

39 Climate change has been one of the most important drivers influencing the mechanism
40 of runoff generation and the confluence process of catchments (Jung et al.,
41 2012; Changnon and Gensini, 2019). Depending on the extent and duration of climate
42 change, it could be classified into extreme (e.g., from prolonged meteorological drought
43 to extremely wet conditions in a period) and asymptotic changes (climate change in
44 different seasons in a normal year). For instance, significant variations (i.e., less runoff
45 than expected) in hydrological behavior have been reported during the decade-long
46 millennium drought of many catchments in south-eastern Australia compared with the
47 previous wet period (Saft et al., 2016). In addition, seasonally asymptotic variations
48 have been identified in many catchments in America due to the seasonal growth and
49 die-off of vegetation (Deng et al., 2018; Pan et al., 2019a), Asia (Deng et al., 2016) and
50 Australia (Pan et al., 2019b). Studies on the hydrological response of catchments to
51 different climate change scenarios not only can improve our understanding of the
52 hydrological variation mechanism of the catchment, but also enhance our ability to
53 prevent unpredictable extreme events.(Kusangaya et al., 2014; Kundu et al., 2017).

54 Accordingly, studies on the hydrological response to the changing environments
55 generally included two main approaches, i.e., statistical analysis and hydrological

56 modeling. Statistical analysis methods can be used to detect trend changes of prolonged
57 hydrological and meteorological data series (Costa et al., 2003; Siriwardena et al.,
58 2006); nevertheless, they usually lack sufficient physical explanations for the potential
59 variation in catchment hydrological response (Lin et al., 2015; Liu et al., 2018).
60 Hydrological models that can comprehensively consider the spatial heterogeneity and
61 physical process of the catchment are broadly used to quantify the hydrological
62 response under multiple climate conditions (Abbaspour et al., 2007; Tu, 2009; Chen et
63 al., 2019; Tian et al., 2021). For example, Chawla and Mujumdar (2015) adopted the
64 Variable Infiltration Capacity (VIC) model to evaluate the runoff response in the upper
65 Ganga basin. Shen et al. (2018) adopted the Hydrological Model of École de
66 Technologies Supérieure (HMETS) to estimate the uncertainty of runoff response to
67 climate change. Tian et al. (2021) applied the Soil and Water Assessment Tool (SWAT)
68 model to assess the effects of climate change on future runoff in the Han River basin,
69 China. However, most of the previous studies on hydrologic response mainly focused
70 on the variations in runoff response to climate change, without paying attention to the
71 causality between the varying climates (i.e., extreme, and asymptotic changes of
72 climates) and variation in catchment properties.

73 Many previous studies (McNamara et al., 2011; Melsen et al., 2016; Carrer et al.,
74 2019) indicated that the catchment water storage capacity (ACWSC) is one of the most
75 significant parameters influencing the mechanism of hydrological response of
76 catchments. The ACWSC is defined as ‘the active water storage capacity refers to the

77 maximum volume of water stored within a catchment and its distribution among
78 groundwater, soil moisture, vegetation, surface water, and snowpack, which are the
79 variables that ultimately characterize the state of the hydrological system' (McNamara
80 et al., 2011). The root zone storage capacity is defined as "the maximum amount of soil
81 moisture that can be accessed by vegetation for transpiration" (Gao et al., 2014; Nijzink
82 et al., 2016; Singh et al., 2020; Laurène, 2021). For a given catchment, the value of the
83 ACWSC should be greater than or equal to the root zone storage capacity.

84 Our previous study identified the impact of meteorological drought on the
85 ACWSC by investigating the changes in hydrological model parameters before and
86 after drought events (Pan et al., 2020). Results showed that significant shifts in the
87 ACWSC were identified in almost two-thirds of the catchments in south-eastern
88 Australia during the prolonged meteorological drought period. Two subsets of
89 catchments with opposite response directions were identified in the study area, i.e., the
90 subsets of catchments with reduced and increased runoff generation rates, respectively.
91 The main potential reasons may be the difference in the proportion of evergreen
92 broadleaf forests in these catchments. We only considered the average shifts from the
93 non-drought period to the drought period and treated the ACWSC of each period as a
94 constant while neglecting the time-varying characteristics of the ACWSC of each
95 catchment due to the periodic climate change, and thus were unable to reflect variation
96 in catchment characteristics under asymptotic climate.

97 Recently, studies of the potential time-varying ACWSC characteristics based on

98 the simulation of the temporal variations of hydrological parameters have attracted a
99 lot of attention (Coron et al., 2012; Brigode et al., 2013; Patil and Stieglitz, 2015; Deng
100 et al., 2018), and provided a new approach for better-representing changes in catchment
101 characteristics (Deng et al., 2016). Accordingly, the selected model parameters that
102 refer to the ACWSC in the model structure were constructed as multiple hypothetical
103 functions based on physical covariates (e.g., time covariates and catchment attributes),
104 and their simulation results were evaluated and compared with observations through
105 specific criteria. Thus, the functional form that achieved the best simulation
106 performance would be recognized as the best item to represent the potential changes in
107 the catchment property (Jeremiah et al., 2013; Westra et al., 2014; Pan et al., 2019a;
108 Pan et al., 2019b).

109 In this study, we systematically explore the response of the ACWSC to both
110 extreme climate changes (i.e., prolonged meteorological drought) and asymptotically
111 periodic climate changes. Three scientific questions will be investigated as follows.

112 (1) What are the change characteristics of the ACWSC under the conditions of
113 prolonged meteorological drought and asymptotic climate variation?

114 (2) Which catchment features and climate factors are more likely to relate to the
115 change of the ACWSC?

116 (3) What is the difference in the ACWSC when both extreme climate variation and
117 asymptotic climate variation are considered compared with extreme climate variation?

118 **2. Materials**

119 **2.1. Study area**

120 In this study, south-eastern Australia was selected as the initial study area. To minimize
121 the impact of human activities, 398 catchments that were not disturbed by reservoirs or
122 irrigation systems are selected in this study. The study area extends from southern
123 Victoria to New South Wales and Queensland. The study area and the locations of the
124 398 initial catchments are illustrated in **Fig. 1**. Saft et al. (2015) and Pan et al. (2019b)
125 indicated that these catchments had experienced about ten years of meteorological
126 drought near the millennium, which had a significant impact on the stability of local
127 ecosystems and the development of society, economy, and politics (Nicholls, 2004; Hunt,
128 2009; Potter et al., 2011; Hughes et al., 2012; van Dijk et al., 2013; Saft et al., 2015).

129 The essential climate characteristics include the large proportion of arid areas, the
130 semi-annular distribution of annual precipitation, and the terrain, geology, land cover,
131 and climate conditions are differentiated between various state catchments. The annual
132 mean precipitation and temperature range from 507 mm to 1814 mm and 8.26°C to
133 19.52°C, respectively. From the perspective of spatial and temporal distribution, the
134 precipitation in the catchments of Victoria state is mainly concentrated in winter. In
135 contrast, the northern catchments in New South Wales and Queensland states have more
136 rain in summer than in winter. The potential reason for this phenomenon is ENSO (El
137 Niño-Southern Oscillation). In terms of runoff, runoff in summer is dominant in

138 northern catchments, while runoff in winter is more likely to occur in southern
139 catchments.

140 **2.2. Data set**

141 **Table 1** summarized the description and source of the three types of data sets, which
142 include (1) meteorological data (daily precipitation and potential evapotranspiration
143 (PET)), (2) hydrological data (daily runoff), and (3) catchment characteristics
144 (catchment area, mean elevation, mean slope, forest coverage percentage, etc.).

145 398 catchments were selected by Zhang et al. (2013), with catchment areas ranging
146 from 50 km² to 17000 km². The collection period of observations of these catchments
147 ranges from 1976 to 2011. It is noted that the historical meteorological observations of
148 all catchments in the data sets were complete. However, the daily runoff observations
149 of 125 catchments were incomplete with the integrity of the time series being less than
150 80%. Thus, these catchments were excluded, and the remaining 273 catchments were
151 used for meteorological drought identification. Finally, 145 catchments were identified
152 through a long-term meteorological drought with a drought period longer than seven
153 years. The drought periods corresponding to those 145 catchments are exhibited in
154 **Fig.2**. Based on the identification criteria of the prolonged drought period, all the
155 drought periods in these catchments lasted more than seven years. In addition, the
156 drought periods of 35% of the catchments spanned over thirteen years. It can be found
157 that the prolonged meteorological drought of most catchments started after 1990 and

158 ended before 2009. In particular, the meteorological drought of 34 catchments began in
159 1997, and 37 catchments began in 2001.

160 The characteristics of the 145 catchments with prolonged meteorological drought
161 (**Table 2**) demonstrate that there are significant differences in physical properties
162 among different catchments. For example, the catchment area, mean elevation, and
163 mean slope range from 54 to 6818 km², from 47 to 1351m, and from 0.3 to 13.6°,
164 respectively. The interval of forest coverage is [15%, 92%]. These catchment features
165 were selected as potential impact factors and analyzed further in Section 4.3.

166 **3. Methodology**

167 The proposed methodology and procedures are sketched in **Fig.3**. To investigate the
168 response of the ACWSC to the prolonged meteorological drought and asymptotic
169 climate variation, the study scheme is conducted with the following four procedures:
170 (1) identification of prolonged meteorological drought; (2) derivation of the response
171 of the ACWSC to long-term meteorological drought and asymptotic climate variation
172 based on the Bayesian change point analysis and the hydrological modeling approach;
173 and (3) analysis of potential factors (i.e., properties of the catchments and climate
174 characteristics) that may be related to the potential changes of the ACWSC and the
175 response time (defined as the time interval between the occurrence of the prolonged
176 meteorological drought and the abrupt shift of the ACWSC).

177 **3.1. Identification of prolonged meteorological drought**

178 There are many methods/indexes, such as the Standardized Precipitation Index (SPI)
179 (Bayat et al., 2015), Rainfall Departure Analysis (Kumar et al., 2020), and Standardized
180 Precipitation-Evapotranspiration Index (SPEI) (Das et al., 2021), have been used to
181 identify the prolonged meteorological drought. Saft et al. (2015) introduced a drought
182 definition algorithm that was based on the annual rainfall only and proved to have a
183 lower degree of dependence and more robustness than other selected approaches in the
184 south-eastern Australia catchments. It is mentioned that the prolonged drought period
185 should be longer than 7 years according to the defined algorithm. For more detailed
186 information about this method, please refer to Saft et al. (2015) and Pan et al. (2019b).

187 **3.2. Hydrological model**

188 The GR4J hydrological model (modèle du Génie Rural à 4 paramètres Journalier) was
189 used to simulate the potential change characteristics of the ACWSC before and after
190 the prolonged meteorological drought. The GR4J model is a daily lumped rainfall-
191 runoff model developed by Perrin et al. (2003) and improved by Le Moine et al. (2008),
192 and it has been used in more than 400 regions with various climatic characteristics
193 around the world, such as China (Zeng et al., 2019), France (Perrin et al., 2003), North
194 America (Pan et al., 2019a), and Australia (Coron et al., 2012). Its validity in the
195 simulation of the rainfall-runoff relationship and reflection of potential changes in
196 catchment properties has been verified by Le Moine et al. (2008) and Simonneaux et al.

197 (2008).

198 **3.2.1 Model structure**

199 The original GR4J model framework proposed by Perrin et al. (2003) only contains
200 four parameters, and its structure is shown in **Fig.4**. The meanings of the four model
201 parameters are introduced as follows: θ_1 is the maximum capacity of the soil moisture
202 accounting storage, which is used to represent the ACWSC (mm) in this study; θ_2 is the
203 groundwater exchange coefficient (mm); θ_3 represents the one-day-ahead maximum
204 capacity of the routing storage (mm); and θ_4 is the time base of unit hydrograph UH1
205 (day). All model parameters are real values, θ_1 , θ_3 and θ_4 are positive, and θ_2 can be
206 positive, negative, or 0.

207 Based on the existing data and catchment attributes, it is almost impossible to
208 obtain the real value of the ACWSC with current technology. However, the hydrological
209 simulation method provides a new perspective for revealing the potential changes of
210 the ACWSC, i.e., we can use a specific parameter (θ_1) in the GR4J model to represent
211 the ACWSC and characterize its variation in the real catchment. Similar studies can be
212 found in Westra et al. (2014), and Deng et al. (2016). Hence, the simulated values of
213 parameter θ_1 and its time-varying characteristics are used to represent the change of the
214 real ACWSC. It should be noted that θ_2 , θ_3 and θ_4 are assumed to remain constant;
215 similar parameter settings can be found in previous studies (Westra et al., 2014; Pan et
216 al., 2020).

217 **3.2.2 Periodicity of the ACWSC**

218 As explained, parameter θ_l in the GR4J model was used to represent the real ACWSC
219 according to its implications. Our previous work (Pan et al., 2020) verified that the
220 ACWSC (i.e., parameter θ_l) had an “abrupt” point after the prolonged meteorological
221 drought, which assumes that the offset of the estimated θ_l represents the change of the
222 ACWSC. Meanwhile, θ_l in each period is recognized as a constant value and does not
223 include the periodicity of the ACWSC that were outlined by many previous works
224 (Nepal et al., 2017; Kunnath-Poovakka and Eldho, 2019; Sezen and Partal, 2019).
225 However, Westra et al. (2014) and Pan et al. (2020) indicated that the ACWSC had
226 periodic variability that may be due to the seasonal growth and wiling of catchment
227 vegetation.

228 In this study, the potentially periodic variation characteristics of the ACWSC
229 (represented by GR4J model parameter θ_l) were included to reflect the asymptotic
230 change within different periods (i.e., periods before and after the change-point), which
231 was described by the sine function. The sine function is one of the most fundamental
232 functional forms to represent the periodic change of variables (Westra et al., 2014; Pan
233 et al., 2019a; Pan et al., 2019b). Furthermore, the potentially extreme change of the
234 ACWSC between the two periods was denoted by the variations between Equations (1)
235 and (2). The time-varying functions of θ_l during two periods are presented as follows:

236 Before the change-point:

$$\theta_1 = \alpha_1 \sin(\beta_1 t + \gamma_1) + \delta_1 \quad (1)$$

237 After the change-point:

$$\theta_1' = \alpha_2 \sin(\beta_2 t + \gamma_2) + \delta_2 \quad (2)$$

238 where, $\alpha_1, \beta_1, \gamma_1, \delta_1$ and $\alpha_2, \beta_2, \gamma_2, \delta_2$ are regression parameters for the time-varying
239 function; α_1 and α_2 signify the amplitude of the sine function; β_1 and β_2 represent
240 the frequency of the sine function; γ_1 and γ_2 denotes the remainder in the sine function;
241 δ_1 and δ_2 refer to the intercept.

242 3.2.3 Likelihood function and parameter estimation

243 (1) Likelihood function

244 In this study, the likelihood function for catchment i from Thiemann et al. (2001)
245 was adopted, which is shown as follows.

$$p_i(\theta(i) / \xi(i), q(i), r) \propto \left[\frac{w(r)}{\sigma} \right]^T \exp \left[-i(r) \sum_{t=1}^T \left| \frac{e_t(\theta(i))}{\sigma} \right|^{2/(1+r)} \right] \cdot p(\theta(i)) \quad (3)$$

$$\omega(r) = \frac{\{\Gamma[3(1+r)/2]\}^{1/2}}{(1+r) \left\{ \Gamma[(1+r)/2] \right\}^{3/2}}, \beta(r) = \left\{ \frac{\Gamma[3(1+r)/2]}{\Gamma[(1+r)/2]} \right\}^{1/(1+r)} \quad (4)$$

246 where p means the probability of likelihood. $\theta(i) = (\theta_1, \theta_2, \theta_3, \theta_4)$; $\Gamma(\cdot)$ denotes the
247 gamma function; T is the number of time steps; q represents the measured runoff; ξ
248 denotes the climate variable input into the hydrological model; e_t refers to the residual
249 error at time step t ; and r is the type of the residual-error model (in this study, r is
250 represented by Gaussian distribution). When verifying the model type of the residual,

251 parameters $\omega(r), \beta(r)$ are constant values as r is certain. In addition, the prior
252 distribution of all unknown quantities is the uniform distribution.

253 **(2) Parameter estimation**

254 The posterior distribution of all unknown variables was estimated using the
255 Shuffled complex evolution metropolis (SCEM-UA) algorithm, which was based on
256 the Markov chain Monte Carlo method (Vrugt et al., 2003; Ajami et al., 2007). For the
257 convergence of parameters, the Gelman-Rubin convergence value was selected as the
258 evaluation standard, and the convergence threshold was 1.2. The pre-set ranges of all
259 parameters are shown in **Table 3**.

260 **3.3 Change point analysis of ACWSC**

261 **3.3.1 Bayesian change point analysis**

262 The Bayesian change point analysis is one of the strongest ways available to explore
263 the possible change time of the ACWSC (Carlin et al., 1992; Cahill et al., 2015). The
264 likelihood probability was used to evaluate the possibility of each potential change
265 point. The most likely time point of each potential scheme is regarded as the ultimate
266 change point of that catchment.

267 **3.3.2 Criteria for evaluating significant changes in ACWSC**

268 To evaluate whether the ACWSC changed significantly under climate change, the
269 following three criteria were adopted.

270 **(1) The Nash-Sutcliffe efficiency coefficient**

271 To guarantee the reasonable simulation results of the GR4J model, the Nash-
272 Sutcliffe efficiency (NSE) coefficient values before and after the change point should
273 be greater than 0.6. Furthermore, the difference in NSE values between the two periods
274 should be less than $|\pm 20\%|$.

275 **(2) The minimum requirements for significant changes in storage capacity**

276 The change rate of the estimated parameter θ_l (θ'_l) before and after the change
277 point should exceed $|\pm 20\%|$. i.e., $\left| \frac{\theta'_l - \theta_l}{\theta_l} \right| \times 100\% \geq 20\%$.

278 **(3) Robustness requirements of the results**

279 The initial values of model parameters were created three times to reduce their
280 impacts on the final simulation results. Moreover, only the catchments that have
281 significant changes in computation results will be taken as the final change items. If the
282 simulation results meet such robustness requirements, the results would have the lowest
283 dependency and the strongest stability on the adopted algorithm and model.

284 **3.4. Response time of a catchment**

285 Van Lanen et al. (2013) and Huang et al. (2017) showed that the recharge between the
286 groundwater and surface runoff would alleviate the hydrological response under short-
287 term meteorological drought. In other words, groundwater would buffer the surface
288 runoff during the drought period. If the duration of the meteorological drought was

289 longer than several years or even decades, the hydraulic connection between the surface
290 runoff and the underground runoff would be weak due to the gradual decrease of
291 groundwater level. For example, Pan et al. (2020) indicated that the ACWSC may
292 change with the occurrence of the prolonged meteorological drought, and the potential
293 reasons were the difference in soil composition and the extensive death of vegetation
294 during the drought period. It also should be noted that the ACWSC would not change
295 immediately after the occurrence of the meteorological drought but respond after a
296 period due to the existence of catchment elasticity (e.g., the existence of the hydraulic
297 connection between surface runoff and groundwater). Thus, the time interval between
298 the occurrence of the meteorological drought and the change point of the ACWSC is
299 named the catchment response time.

300 **3.5 Potential factors associated with the changes in ACWSC**

301 The process that leads to the change of the ACWSC cannot be measured directly, so
302 some measurable factors are used to probe the lurking correlation between the change
303 of the ACWSC and the catchment response time. We select 33 potential factors of
304 catchments and list them in **Table 4**, which includes 9 catchment features and 24 local
305 climate variables. It is noted that because of the limitation of available data for
306 catchment characteristics, only one static/constant value of the catchment features (A1-
307 A9) was used for the correlation analysis. Furthermore, climate variables in four-time
308 scales were used, including daily (B1-B4), monthly (B5-B7), seasonal (B8-B15), and
309 annual (B16-B24) variables.

310 **4. Results**

311 **4.1 Change pattern of the ACWSC**

312 The most likely change point was confirmed when three criteria had been satisfied. The
313 changing pattern of the ACWSC was determined by Equations (1) and (2). In other
314 words, Equation (1)/Equation (2) reflects the potential periodic/asymptotic feature
315 during the period before/after the change point. It is obvious that α_1 (α_2) and δ_1 (δ_2)
316 are the most important parameters in the regression function, which refer to the
317 amplitude and intercept of the time-varying parameter θ_1 , respectively. Furthermore, the
318 variation between δ_1 and δ_2 denotes the average difference between θ_1 and θ_1' ,
319 reflecting the potential change between the ACWSC of periods before and after the
320 change point.

321 **Table 5** presents the variation characteristics (amplitude α and mean value δ of
322 the ACWSC in the 145 studied catchments with meteorological drought in south-
323 eastern Australia. The results showed that 36.6% of the catchments (55 of 145
324 catchments) were identified to violate the criteria of the maximum performance
325 degradation and result robustness, and thus were removed from further analysis. The
326 remaining 92 catchments were retained as the-set of catchments that satisfied the basic
327 criteria of NSE performance and resultant robustness. As presented in Equations (1)
328 and (2), amplitude α represents the range of variation in the ACWSC, a larger $|\alpha|$
329 implies a greater variation interval of the ACWSC during the specific period.
330 Significant changes in amplitude α were found in 60.0% of the catchments (87 of 145

331 catchments) during the drought period, in which 57.2% of the catchments (83 of 145
332 catchments) experienced a significantly increased change in amplitude α while 2.8%
333 of the catchments (4 of 145 catchments) had significantly decreased variation during
334 the drought period. In addition, only 3.4% of the catchments (5 of 145 catchments)
335 experienced a non-significant change in amplitude α , in which 3 (2) catchments had a
336 slight increase (decrease) trend. It means that most of the catchments (87 of 92
337 catchments) experienced a significant increase trend in the range of variation during the
338 prolonged drought period (Table 5), indicating an increased dramatic cyclical variation
339 magnitude of the ACWSC during the transformation from the non-drought period to
340 the prolonged drought period.

341 The regression parameter δ , which refers to the intercept/mean value of the
342 ACWSC during the specific period, was used to evaluate the average difference
343 between the ACWSC during the two periods. As Table 5 indicated: a significant
344 increase in mean value δ was identified in 84% of the catchments (77 of 145
345 catchments) after the change point, but no catchment was found to experience a
346 significant decrease of δ during the drought period. In addition, the number of
347 catchments with non-significant changes in δ was 15, and 6.9% of the catchments (10
348 of 145 catchments) and 3.5% of the catchments (5 of 145 catchments) were identified
349 to have a non-significant increase and decrease trend during the drought period,
350 respectively. These results illustrated that most catchments (77 of 92 catchments)
351 experienced a significant increase trend in the average ACWSC during the

352 transformation from the non-drought period to the prolonged drought period, indicating
353 a mainstream trend of increased ACWSC during the latter period.

354 The spatial distribution of the 92 catchments that satisfied the criteria of NSE
355 performance and resultant robustness is presented in **Fig. 5**. Obvious convergence was
356 found in the spatial distribution of the catchments with different change forms in the
357 amplitude of the periodic change and the average variation level of the two periods. For
358 instance, catchments with non-significant change in δ were mainly concentrated in the
359 middle part of the south region of Australia. The reason for this phenomenon may be
360 the similar physical features and climatic characteristics of adjacent catchments, which
361 may result in the relatively consistent change direction of catchments in a region.

362 **Fig.6** illustrates the statistical results of the change of amplitude α and mean value
363 δ between two periods (before and after the change point) in all catchments in south-
364 eastern Australia. **Figs.6(a) and 6(b)** show the absolute and relative change percentage
365 of amplitude α between two periods, indicating that the absolute differences in the
366 amplitude between two periods, i.e., $|\alpha_2 - \alpha_1|$ are concentrated within the interval of
367 $[0, 75]$ for 80.4% of the catchments while the relative changes $(\alpha_2 - \alpha_1) / \alpha_1$ are mostly
368 concentrated within the interval of $[0, 400\%]$ for 69.6% of the catchments. The fitting
369 curves in **Figs.6(a) and 6(b)**, which were based on the kernel smoother method (Yandell,
370 1996), had significant positive biases, indicating that much more catchments
371 experienced an increased tendency in the variation range of periodic changes of the
372 ACWSC during the drought period. **Figs.6 (c) and 6(d)** show the absolute and relative

373 change percentage of the mean value δ , respectively, indicating that the absolute
374 change of the mean value, i.e., $|\delta_2 - \delta_1|$, are concentrated within the interval of [50, 150]
375 for 75% of the catchments while the relative change, i.e., $(\delta_2 - \delta_1) / \delta_1$, are mostly
376 concentrated within the interval of [0, 50%] for 65.2% of the catchments. Similarly, the
377 fitting curves in **Figs.6(c) and 6(d)** had remarkable positive biases as well, indicating
378 that much more catchments experienced an increased tendency in the mean value of the
379 ACWSC after the change point.

380 Among the catchments with significant variation in θ_t , two types of typical
381 catchments were taken as examples to present the specific changes of the ACWSC
382 (shown in **Fig.7**). In catchment #222206, both α_2 and δ_2 increased significantly after
383 the change point compared with α_1 and δ_1 . Based on the posterior probability of each
384 possible change point, it was found that the change probability of the ACWSC was the
385 greatest on 2002/12/27. Changes in θ_t indicate that the ACWSC of catchment #222206
386 tends to increase after the change point. In catchment #421042, the amplitude α_2
387 decreases significantly while the mean value δ_2 increases significantly after the
388 change point. The time corresponding to the change point was 2001/7/30, which refers
389 to the moment when θ_t changes. Therefore, the above results of the two example
390 catchments suggest that the ACWSC of various catchments may experience different
391 magnitudes of change under a sustained reduction in rainfall. In addition, a time lag
392 phenomenon clearly occurred between the onset of the meteorological drought and the
393 change in θ_t .

394 **4.2 Response time of catchments with significant change in the** 395 **ACWSC**

396 As mentioned in Section 3.4, the response time refers to the time interval between the
397 occurrence of the meteorological drought and the change point of the ACWSC. The
398 magnitude distribution of response time in the 92 catchments that satisfied the basic
399 criteria of NSE performance and robustness of results was manifested in **Fig.8**, which
400 indicates that the response time in nearly one-third of the catchments (27/92) fell within
401 the range of 800-1000 days, followed by the response time of 17 catchments fell within
402 the range of 600-800 days. Furthermore, as shown in **Table 6**, the average and median
403 response times of the catchments with significant changes in δ are 660.7 days and
404 750.6 days, respectively. Since no significantly decreased variation in δ was found, the
405 catchments with significant changes in δ after the change point all realized a
406 significantly increased trend. In the catchments with a significant increase in amplitude
407 α , the average and median estimates of the response time are 660.4 and 750.6 days,
408 respectively; while those of the catchments with a significant decrease in α are 391.9
409 and 422 days, respectively. According to the results shown in **Table 6**, a significant
410 difference was identified in the length of the response time between two sets of
411 catchments with a significant increase and decrease in amplitude α . However, it is not
412 clear whether the difference between the groups of catchments with significant
413 increase/decrease change of the amplitude α is real or just sampling fluctuations.

414 **4.3 Factors for shifts in the ACWSC**

415 To provide a better understanding of the response of the variation pattern of the
416 ACWSC to the prolonged meteorological drought and the variation characteristics
417 under asymptotic climate change, we investigated whether the change in the ACWSC
418 (especially in the amplitude α and mean value δ) was associated with particular
419 catchment features and/or climate inputs, i.e., are variation in the ACWSC more likely
420 to occur in the catchments with certain characteristics? Thus, 9 multiple catchment
421 features and 24 climate variables that may drive the shifts in the variation of the
422 amplitude α and mean value δ were analyzed in this part.

423 **4.3.1 Difference analysis of factors**

424 **4.3.1.1 Difference between groups of catchments with significant and** 425 **non-significant change in α**

426 To explore the potential differences in catchment properties and climate inputs
427 between catchments with different variation patterns, the 92 selected catchments were
428 divided into two groups (namely $g_{\alpha}(S)$ (catchments with significant change in α)
429 and the $g_{\alpha}(NS)$ group (catchments with non-significant change in α)) according to
430 the significance level of the variation in amplitude α between the periods before and
431 after the change point. As illustrated in Table 5, $g_{\alpha}(S)$ and $g_{\alpha}(NS)$ groups included
432 87 and 5 catchments, respectively. 94.6% (87/92) of studied catchments experienced a
433 significant shift in amplitude α which indicated that the long-term drought in these

434 catchments resulted in a remarkable change in the variation range of the ACWSC. The
435 left two columns in each sub-figure of **Fig.9** referred to the statistical features of
436 catchments within the $g_{\alpha}(S)$ and $g_{\alpha}(NS)$ groups. There was a significant
437 difference in the mean and median estimate of catchment area between these two groups,
438 with their difference ratio reaching 21.2% and 25.1%, respectively, i.e., the $g_{\alpha}(NS)$
439 group indicated a notably larger catchment area than the $g_{\alpha}(S)$ group. However, no
440 other features (mentioned in Table 1) showed similarly significant variation between
441 $g_{\alpha}(S)$ and $g_{\alpha}(NS)$ groups. Among the adopted 9 catchment features, the results
442 indicated the difference in the catchment area may be one of the most important factors
443 for influencing the variation degree of the amplitude α of the ACWSC. However, due
444 to the limited number of catchments in the $g_{\alpha}(NS)$ group (only 5.4% of the adopted
445 92 catchments), it is still not clear whether the statistical values of this group were real
446 or just sampling fluctuation.

447 The right two columns in **Fig.9** referred to catchment subsets with a significant
448 increase pattern in amplitude α after the change point, namely the $s_{\alpha}(IS)$ and the
449 $s_{\alpha}(DS)$ subsets, which denoted the catchment aggregation that experienced
450 significantly increased and decreased changes after the change point, respectively. It
451 should be noted that the two subsets were extracted from the $g_{\alpha}(S)$ group. Most
452 catchments (95.4% of catchments) experienced a significantly increased change in the
453 amplitude α of the ACWSC after the change point, while only 4.6% (4 in 87
454 catchments) of catchments went through a significantly decreased change after the

455 change point. The increased variation range of the ACWSC that occurred during the
456 prolonged drought led to a higher fluctuation range of the ACWSC and more intense
457 variation in runoff generation rate. Thus, the significantly increased pattern in
458 amplitude α and more intense variation in runoff generation rate were the mainstream
459 change direction in the studied catchment dataset.

460 Significant differences have been found in both the mean and median estimate of
461 features of catchment area and mean elevation between the $s_\alpha(IS)$ and the $s_\alpha(DS)$
462 subsets (see right two columns in **Fig.9**), with the difference ratio reaching 46.7% and
463 58.5%, respectively. The $s_\alpha(DS)$ subset had a significantly larger catchment area
464 than the $s_\alpha(IS)$ subset. Meanwhile, there was a significant difference in the median
465 estimate of the Ks of subsoil between the two subsets, with the difference ratio reaching
466 27.7%, however, it was non-significant in the mean estimate of the Ks of subsoil. Due
467 to the limited number of catchments within the $s_\alpha(DS)$ subset (only included 4
468 catchments), it was inadequate to judge whether it was popular findings or just the
469 uniqueness of the sample.

470 Overall, it was likely that catchments with small areas, low elevations, small slope
471 ranges, large forest coverage, and high AWHC of soil may change more significantly
472 in amplitude α after the interference of the meteorological drought. Generally, small
473 areas of large forest cover will require considerable (partitioning of) soil water storage.
474 After experiencing persistent meteorological drought, the pressure on water resources
475 in the catchment increased and tree cover was lost in large quantities due to withering.

476 Canopy retention and uptake by the forest is an important part of ACWSC, and the
477 dieback of trees in the forest may result in a significant change in ACWSC (Adams et
478 al., 2012). Therefore, these catchments are more vulnerable under prolonged drought
479 due to competition for moisture uptake than catchments with low forest cover and large
480 areas.

481 **4.3.1.2 Difference between groups of catchments with significant and** 482 **non-significant change in δ**

483 Similarly, we also analyzed the potential relationship between the change in the mean
484 value δ of the ACWSC and the catchment features/climate characteristics. According
485 to the significance level of the change in mean value δ , the 92 catchments were also
486 segmented into two groups, denoted as $g_{\delta}(S)$ (catchments with significant change
487 in δ) and the $g_{\delta}(NS)$ groups (catchments with non-significant change in δ).

488 As illustrated in **Table 5** and **Fig.10**, 77 in 92 catchments were found to experience
489 a significantly increased change in the mean value δ after the change point, while no
490 catchment went through a significantly decreased pattern after the change point. The
491 non-significant change in the mean value δ occurred in 15 studied catchments. The
492 significant increase in the mean value δ indicated the increased mean ACWSC after
493 the change point due to the long-term meteorological drought, resulting the even less
494 runoff (on average) than the historical relationship suggested. In other words, the low
495 runoff caused by the reduced rainfall was expected as the previous rainfall-runoff
496 relationship showed, the increase in the ACWSC may imply an even lower runoff

497 generation rate than expected.

498 The two left columns in each sub-figure of **Fig.10** presented the comparison of
499 catchment features between $g_{\delta}(S)$ and $g_{\delta}(NS)$ groups. There was a significant
500 difference in the mean (median) estimate of the catchment area, AWHC of the subsoil,
501 Ks of the subsoil, mean slope and slope range between these two groups, with their
502 difference ratio reaching 50.3% (33.8%), 34.2% (54.4%), 20.6% (57.1%), 38.8%
503 (91.1%) and 24.4% (37.4%) respectively. In the other words, the $g_{\delta}(S)$ group had a
504 notably smaller catchment area, Ks of the subsoil, mean slope and slope range, and
505 larger AWHC of the subsoil than the $g_{\delta}(NS)$ group. Meanwhile, there was a
506 significant difference in the median estimate of the Ks of topsoil between the two
507 groups, with the difference ratio reaching 29.6%, however, it was non-significant in the
508 mean estimate of the Ks of topsoil.

509 **4.3.2 Association analysis of factors**

510 **Fig.11** presented the Pearson correlation between the change of amplitude α of θ_1
511 with 9 catchment features and 24 climate variables that were listed in **Table 4**. A
512 positive association has been identified between the absolute change of amplitude α
513 and two catchment features (i.e., mean elevation and Ks of subsoil), while negative
514 relationship between the former and other catchment features (see **Fig.11(a)**). Similarly,
515 the relative change of amplitude α was positively associated with only one catchment
516 feature, i.e., the AWHC of the topsoil (see **Fig.11(b)**). However, no strong correlation
517 was found between the change of amplitude α (including both absolute and relative

518 changes) and both catchment features. **Fig.11(c) and (d)** illustrated the possible
519 correlations between the changes (absolute and relative changes) in the amplitude α of
520 the ACWSC and 24 climate variables. Generally, a weak positive correlation was found
521 between the absolute change of amplitude α and all climate variables, with the highest
522 Correlation Coefficient (CC) reaching 0.203 that occurred with the B6 feature (i.e., Cv
523 of monthly runoff). Similarly, there was no strong correlation between the relative
524 change of amplitude α and all climate variables (see **Fig.11(d)**), with the highest CC
525 only reaching 0.19 that occurred with B17 feature (i.e., the mean annual potential
526 evapotranspiration). Since no strong correlation was found between the variation in the
527 amplitude α and a single factor, we speculated that the potential change of the variation
528 range of the ACWSC after the change point was the result of the combination of various
529 catchment properties and climate characteristics.

530 **Fig.12** illustrates the Pearson correlation between the changes (absolute change
531 and relative change) of the mean value δ of the ACWSC and catchment features
532 between the periods before and after the change point. The absolute change of the mean
533 value δ was negatively correlated with both catchment features (see **Fig. 12(a)**), with
534 the highest CC reaching -0.362 that occurred with the Ks of topsoil, subsequently
535 followed by the AWHC of the subsoil (CC=-0.341), the Ks of subsoil (CC=-0.267), and
536 the forest percentage (CC=-0.242). Similar to **Fig. 12(a)**, the relative change of the
537 mean value δ was negatively correlated with most of catchment features (**Fig. 12(b)**),
538 except for A3 (slope range) and A8 (AWHC of topsoil), with the largest CC reaching -

539 0.362 that occurred with the K_s of topsoil, followed by AWHC of the subsoil (CC=-
 540 0.341), and forest coverage (CC=-0.242). It is obvious that the soil and forest-related
 541 features had the largest relationship with the relative change of the mean value δ
 542 among both catchment features. The potential reasons may lie that the water holding
 543 capacities of various soil types were different due to the dissimilarity of void and
 544 adhesion in different soil types, which directly affected the ability of the catchment to
 545 absorb and store water, thereby influencing the magnitude of the ACWSC of the
 546 catchment (Leblanc et al., 2009). Furthermore, the coverage of various forest
 547 percentages would affect the water holding capacity and water assumption ability
 548 (Fohrer et al., 2005), resulting in potential changes in the ACWSC. **Figs.12(c) and 12(d)**
 549 illustrate the association between the changes (absolute and relative change) of the
 550 mean value δ and 24 climate variables before and after the change point. As **Figs.12(c)**
 551 indicates: the absolute change of the mean value δ had positive correlations with B19
 552 (Annual aridity index, CC=0.421), followed by B9 (mean summer precipitation,
 553 CC=0.306), while it had negative correlations with B8 and B21. **Fig.12 (d)** shows that
 554 the relative change of the mean value δ had the largest negative correlation with B24
 555 (Annual base flow ratio, CC=-0.279), followed by B20 (Mean annual runoff index,
 556 CC=-0.215). No correlation (CC< 0.2) has been found in the relative change of the
 557 mean value δ with other climate variables.

558 In total, $g_\alpha(S)$ and $g_\delta(S)$ groups had a significantly smaller catchment area
 559 than those of the $g_\alpha(NS)$ and $g_\delta(NS)$ groups, indicating the reduced possibility

560 that the ACWSC would change significantly (including changes in both amplitude α
561 and mean value δ) along with the increased catchment area. Furthermore, the
562 catchments with a smaller hydraulic conductivity of the soil may be more prone to change
563 in statistical significance to experience a significant variation on the average level of
564 the ACWSC during a prolonged meteorological drought.

565 **4.3.3 Trend analysis within the significantly changed group**

566 As our findings in **Table 5**, most of the studied catchments experienced a
567 significantly increased variation after the change point, the $s_{\alpha}(IS)$ and $s_{\delta}(IS)$
568 subsets of catchments were further used as typical samples for the trend analysis
569 between the variation in the ACWSC and certain characteristics. According to the
570 results in sections 4.3.1 and 4.3.2, four catchment properties, i.e., catchment area, mean
571 elevation, forest coverage, and soil characteristics, were adopted for the trend analysis.
572 As illustrated in **Fig.13**, the absolute changes in α and δ both show an increasing
573 trend with the increase in catchment area, the catchment group with the mean elevation
574 within the interval of [300, 600] had the largest absolute change in both the amplitude
575 α and mean value δ among all groups with different elevation interval, implying the
576 potentially most suitable elevation range for the occurrence of the variation of ACWSC.
577 Furthermore, the decreased variation of the estimated value of α and δ has been
578 identified along with the increase in the forest coverage of catchments. In addition,
579 **Fig.13** indicated that the changes in α and δ were both negatively associated with the
580 increase in forest coverage percentage of the catchment, implying the positive

581 contribution of high forest coverage to the potential change in the ACWSC during the
582 meteorological drought. A similar relationship was observed in changes of δ with the
583 AWHC subsoil.

584 **4.4 Factors for the response time of catchments**

585 The Pearson correlation coefficient between the response time with both
586 catchment features and climate variables was presented in **Fig.14**. Positive correlations
587 were identified between the response time with A6 (AWHC of the topsoil, CC=0.249)
588 and A2 (mean elevation, CC=0.239). While a negative correlation was found between
589 the response time and A5 (forest coverage, CC=-0.225). The potential reasons for this
590 finding may lie that the larger ACWSC indicated a higher ability of the soil to retain
591 water and make it more sufficiently available for plant use, thus resulting in an
592 increased response time in the catchment (Lawes et al., 2009; Leenaars et al., 2018).
593 Meanwhile, the increased catchment elevation may promote changes in forest
594 architecture (i.e., decreases in tree stature and stem diameter, trends in stem
595 deformation, hard, thick, and smaller leaves) and enhance the dominant position of
596 plants with less water assumption (Lenoir et al., 2008; Oke and Thompson., 2015), and
597 thus relatively enlarge the response time. In addition, the persistent decline of the
598 groundwater level and storage has been observed in catchments of South-eastern
599 Australia (Leblanc et al., 2009), resulting in the gradual reduction of the interactions
600 between the surface water and groundwater (Van et al., 2013). Thus, the increased forest
601 coverage of the catchment may result in larger water demand for the ecosystem (Adams

602 et al., 2012), and thus caused a shorter response time of the ACWSC to the
603 meteorological drought.

604 As for the relationship between the response time and the climate variables
605 mentioned in **Table 4**, the absolute variations of many climate variables (i.e., B1-B4,
606 B9, B13, B17) had negative correlations with the response time (**Fig.14(b)**), with their
607 correlation coefficient between 0.20 and 0.31. The highest CC in **Fig.14(b)** was 0.31
608 which reached with B2 (mean daily potential evapotranspiration). As shown in
609 **Fig.14(c)**, the response time was negatively correlated with the absolute change of B2
610 (mean daily potential evapotranspiration), B3 (mean T_{\max}), and B13 (mean summer
611 runoff), with the CC were -0.313, -0.263, and -0.27, respectively. It also should be noted
612 that only a weak association has been identified between the response time and these
613 climate variables. In addition, no positive correlation ($CC > 0.2$) has been identified
614 between the response time with the absolute and relative changes of both climate
615 variables.

616 Similarly, the potential connections between the response time and several
617 catchment properties were further analyzed in the significantly changed subsets. As
618 shown in **Fig.15**, negative associations have been found between the length of response
619 time with the size of the catchment area and forest coverage. Furthermore, the
620 catchment group with the mean elevation within the interval of [300, 600] had the
621 smallest response time within all range groups of catchments.

622 **5. Discussions**

623 **5.1 Possible reasons for different changes in the ACWSC**

624 The results showed that most catchments were identified to have an increasing trend in
625 both the amplitude α and the mean value δ of the ACWSC after a prolonged
626 meteorological drought. According to our findings, soil type and forest coverage are
627 the variables the most related to the ACWSC. The soil water holding capacities of
628 various soil types were different due to the dissimilarity of void and adhesion in
629 different soil types, which directly affects the ability of the catchment to absorb/store
630 water, thereby affecting the ACWSC of the catchment. Saft et al. (2015) showed that
631 the annual rainfall-runoff relationships of many catchments changed in southeastern
632 Australia during the millennium drought (1997-2009). The prolonged meteorological
633 drought led to the continuous decrease of the groundwater level as well as a significant
634 change in soil properties. Leblanc's study for southeastern Australia showed that only
635 two years after the 2001 drought, soil moisture and surface water storage lost 80 and
636 12 km³, respectively, and the rapid drying up reached near-steady low levels (Leblanc
637 et al., 2009). Years of drought led to an almost complete drying up of surface water
638 resources, and the hydrological drought continued even after rainfall resumed. In
639 addition, the soil types in the study area include silt loam, loam, silt, sand, sandy loam,
640 clay and loamy sand, among which silt loam accounts for more than 45% of the total
641 study area (Pan et al., 2020). Moreover, the silt loam possessed a strong field capacity
642 and large adhesion property. The silt loam may maintain the original soil structure state

643 even if the soil pore space increases due to the declined groundwater level, which may
644 partly explain the increase in the ACWSC of the catchments.

645 Furthermore, the variation of forest coverage and composition would affect the
646 water holding capacity and water assumption ability, resulting in potential changes in
647 the ACWSC. Previous studies (Fensham et al., 2009; Allen et al., 2010) showed that
648 the increased frequency, duration of drought, and heat stress associated with climate
649 change are strong factors contributing to changes in vegetation dynamics that may
650 fundamentally alter forest composition and structure in many areas. Drought-induced
651 vegetation dieback was more likely to occur in regions with relatively high densities of
652 local woody cover. Adams et al. (2012) combined the extensive literature on the
653 ecohydrological effects of tree harvesting with existing studies to propose a new and
654 relevant hypothesis. For most forests, evapotranspiration would be dramatically
655 reduced after the significant dieback of the tree cover due to drought. According to Pan
656 et al. (2020), the main land use types throughout the study area are evergreen broadleaf
657 forest, grassland, woodland, and cropland. As the evergreen broadleaf forest and
658 woodland occupied most of the study region, the notable loss of tree cover caused by
659 the prolonged meteorological drought may dramatically reduce the evapotranspiration
660 in catchments. Catchments with large coverage of evergreen broadleaf forest processed
661 the large water demand per unit area (Adams et al., 2012). For comparison, the water
662 consumption of catchments with other land use types (grassland and farmland) was less,
663 and their drought resistance ability was relatively stronger. It can be hypothesized that

664 in catchments with large coverage of vegetation, the occurrence of the prolonged
665 drought may intensify the competition for water demand between different varieties of
666 vegetation, promoting the survival of the vegetation types with less water consumption
667 but with higher water adoption ability. Therefore, the catchments with high forest cover
668 may lead to an increase in the ACWSC.

669 **5.2 The limitations of the hydrological model**

670 The GR4J model was used to address the response of the ACWSC to the prolonged
671 meteorological drought. The model processes a relatively simple structure with
672 relatively low requirements for input data, and it has been widely used in the rainfall-
673 runoff simulation for small and medium-sized catchments (Dhemi et al., 2010; Demirel
674 et al., 2013; Sezen et al., 2019; Kunnath et al., 2019). However, the GR4J model is
675 implemented subject to restrictions and limitations due to the inadequate description of
676 the runoff generation and flow confluence processes in large catchments (e.g., larger
677 than 10,000 km²). Conceptual models usually consider the entire catchment to be one
678 entity, then use empirical functional relationships or conceptual simulations to describe
679 the runoff generation and flow confluence processes, and consequently adopt certain
680 parameters with physical meanings to characterize the inhomogeneity of the spatial
681 distribution of catchment characteristics. It has been argued that conceptual lumped
682 rainfall-runoff models are far from being able to tackle the challenging problem of
683 assessing the impacts of land use or forest variation. The GR4J model lacks a physical
684 foundation but seems to best detect changes in a basin behavior (Perrin et al., 2003).

685 According to Westra et al. (2014), θ_1 is the most sensitive parameter in the GR4J
686 model and therefore was used to represent the ACWSC in this study. The sine function
687 was used to reflect the periodic change of the ACWSC. Further studies are necessary
688 to explore the impacts of different forms of functions on the identification and
689 simulation of the periodic variation of the ACWSC.

690 **6. Conclusions**

691 This study focused on the response of the ACWSC to the long-term meteorological
692 drought and asymptotic climate change systematically based on the hydrological
693 simulation method. Firstly, the time-varying parameter (the most sensitive model
694 parameter in the adopted GR4J model) was derived to reflect the ACWSC
695 periodic/abrupt variations in drought and non-drought periods. Secondly, the change
696 points and varying patterns of the ACWSC during the transformation from non-drought
697 to drought periods were analyzed based on the Bayesian change point analysis with
698 multiple evaluation criteria. Finally, a variety of catchment features and climate
699 characteristics were used to explore the possible relationship between these variables
700 and the temporal variation characteristics of the ACWSC. Catchments that suffered
701 from the prolonged meteorological drought in southeast Australia were selected as the
702 case study. The main conclusions were summarized as follows.

703 (1) The increase of ACWSC amplitude change was observed in 83/92 catchments
704 during the prolonged drought period, and significant shifts in the mean value of the
705 ACWSC were detected in 77/92 catchments.

706 (2) The average response time of the ACWSC for all 92 catchments with
707 significant changes was 641.3 days. Specifically, the response time in 27 and 17
708 catchments fell within the ranges of 800-1000 days and 600-800 days, respectively.

709 (3) The ACWSC changed significantly in the catchments with small areas, low
710 elevations, small slope ranges, large forest coverage, and high soil water holding
711 capacities.

712 In this study, the response characteristics of the ACWSC to the prolonged
713 meteorological drought in southeastern Australia were analyzed. It was found that the
714 catchment response time and mode are greatly different. However, only the correlations
715 between the changes of parameter θ_1 , response time, and single-factor of catchment
716 features and climate variables were considered in this study. Subsequent studies could
717 be conducted by combining data from multiple sources to carry out multi-factor
718 regression analysis. Nevertheless, this study could enhance our understanding of the
719 variations in catchment property under climate change.

720 **Acknowledgments**

721 This study was supported by the National Natural Science Foundation of China (Grant
722 No. U20A20317) and the National Key Research and Development Program of China
723 (2021YFC3200303). The numerical calculations were done on the supercomputing
724 system in the Supercomputing Center of Wuhan University. The authors would like to
725 thank the editor and anonymous reviewers for their comments, which helped improve
726 the quality of the paper.

727 **Author contributions**

728 All the authors helped to conceive and design the analysis. Jing Tian and Zhengke Pan
729 performed the analysis and wrote the paper. Shenglian Guo, Jun Wang, Jiabo Yin and
730 Yanlai Zhou contributed to the writing of the paper and made comments.

731 **Compliance with ethical standards**

732 **Conflict of interest:** The authors declare that they have no conflict of interest.

733 **References**

- 734 Abbaspour, K. C., Yang, J., Maximov, I., Siber, R., Bogner, K., Mieleitner, J., Zobrist, J., and Srinivasan,
735 R.: Modelling hydrology and water quality in the pre-alpine/alpine Thur watershed using SWAT, *J.*
736 *Hydrol.*, 333, 413-430, 10.1016/j.jhydrol.2006.09.014, 2007.
- 737 Adams, H. D., Luce, C. H., Breshears, D. D., Allen, C. D., Weiler, M., Hale, V. C., Smith, A. M. S., and
738 Huxman, T. E.: Ecohydrological consequences of drought- and infestation- triggered tree die-off:
739 insights and hypotheses, *Ecohydrology*, 5, 145-159, 10.1002/eco.233, 2012.
- 740 Ajami, N. K., Duan, Q. Y., and Sorooshian, S.: An integrated hydrologic Bayesian multimodel
741 combination framework: Confronting input, parameter, and model structural uncertainty in
742 hydrologic prediction, *Water Resour. Res.*, 43, 10.1029/2005wr004745, 2007.
- 743 Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger,
744 T., Rigling, A., Breshears, D. D., Hogg, E. H., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J.,
745 Demidova, N., Lim, J. H., Allard, G., Running, S. W., Semerci, A., and Cobb, N.: A global overview
746 of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *For.*
747 *Ecol. Manage.*, 259, 660-684, 10.1016/j.foreco.2009.09.001, 2010.
- 748 Bayat, B., Nasserli, M., and Zahraie, B.: Identification of long-term annual pattern of meteorological
749 drought based on spatiotemporal methods: evaluation of different geostatistical approaches, *Nat.*
750 *Hazards*, 76, 515-541, 10.1007/s11069-014-1499-3, 2015.
- 751 Brigode, P., Oudin, L., and Perrin, C.: Hydrological model parameter instability: A source of additional
752 uncertainty in estimating the hydrological impacts of climate change?, *J. Hydrol.*, 476, 410-425,
753 10.1016/j.jhydrol.2012.11.012, 2013.
- 754 Cahill, N., Rahmstorf, S., and Parnell, A. C.: Change points of global temperature, *Environ. Res. Lett.*,
755 10, 10.1088/1748-9326/10/8/084002, 2015.
- 756 Carlin, B. P., Gelfand, A. E., and Smith, A. F. M.: Hierarchical bayesian-analysis of changepoint
757 problems, *J. R. Stat. Soc. C-Appl.*, 41, 389-405, 10.2307/2347570, 1992.
- 758 Carrer, G. E., Klaus, J., and Pfister, L.: Assessing the Catchment Storage Function Through a Dual-
759 Storage Concept, *Water Resour. Res.*, 55, 476-494, 10.1029/2018wr022856, 2019.
- 760 Changnon, D., and Gensini, V. A.: Changing Spatiotemporal Patterns of 5-and 10-Day Illinois Heavy

761 Precipitation Amounts, 1900-2018, *J. Appl. Meteorol. Clim.*, 58, 1523-1533, 10.1175/jamc-d-18-
762 0335.1, 2019.

763 Chawla, I., and Mujumdar, P. P.: Isolating the impacts of land use and climate change on streamflow,
764 *Hydrol. Earth Syst. Sc.*, 19, 3633-3651, 10.5194/hess-19-3633-2015, 2015.

765 Chen, Q. H., Chen, H., Wang, J. X., Zhao, Y., Chen, J., and Xu, C. Y.: Impacts of Climate Change and
766 Land-Use Change on Hydrological Extremes in the Jinsha River Basin, *Water*, 11,
767 10.3390/w11071398, 2019.

768 Coron, L., Andreassian, V., Perrin, C., Lerat, J., Vaze, J., Bourqui, M., and Hendrickx, F.: Crash testing
769 hydrological models in contrasted climate conditions: An experiment on 216 Australian catchments,
770 *Water Resour. Res.*, 48, 10.1029/2011wr011721, 2012.

771 Costa, M. H., Botta, A., and Cardille, J. A.: Effects of large-scale changes in land cover on the discharge
772 of the Tocantins River, Southeastern Amazonia, *J. Hydrol.*, 283, 206-217, 10.1016/s0022-
773 1694(03)00267-1, 2003.

774 Das, S., Das, J., and Umamahesh, N. V.: Identification of future meteorological drought hotspots over
775 Indian region: A study based on NEX-GDDP data, *Int. J. Climatol.*, 41, 5644-5662,
776 10.1002/joc.7145, 2021.

777 Demirel, M. C., Booij, M. J., and Hoekstra, A. Y.: Effect of different uncertainty sources on the skill of
778 10 day ensemble low flow forecasts for two hydrological models, *Water Resour. Res.*, 49, 4035-
779 4053, 10.1002/wrcr.20294, 2013.

780 Deng, C., Liu, P., Guo, S. L., Li, Z. J., and Wang, D. B.: Identification of hydrological model parameter
781 variation using ensemble Kalman filter, *Hydrol. Earth Syst. Sc.*, 20, 4949-4961, 10.5194/hess-20-
782 4949-2016, 2016.

783 Deng, C., Liu, P., Wang, D. B., and Wang, W. G.: Temporal variation and scaling of parameters for a
784 monthly hydrologic model, *J. Hydrol.*, 558, 290-300, 10.1016/j.jhydrol.2018.01.049, 2018.

785 Fensham, R. J., Fairfax, R. J., and Ward, D. P.: Drought-induced tree death in savanna, *Global Change*
786 *Biol.*, 15, 380-387, 10.1111/j.1365-2486.2008.01718.x, 2009.

787 Fohrer, N., Haverkamp, S., and Frede, H. G.: Assessment of the effects of land use patterns on hydrologic
788 landscape functions: development of sustainable land use concepts for low mountain range areas,
789 *Hydrol. Process.*, 19(3), 659-672, 10.1002/hyp.5623, 2005.

790 Gao, H., Hrachowitz, M., Schymanski, S. J., Fenicia, F., Sriwongsitanon, N., and Savenije, H. H. G.:
791 Climate controls how ecosystems size the root zone storage capacity at catchment scale, *Geophys.*
792 *Res. Lett.*, 41, 7916-7923, 10.1002/2014gl061668, 2014.

793 Huang, S. Z., Li, P., Huang, Q., Leng, G. Y., Hou, B. B., and Ma, L.: The propagation from meteorological
794 to hydrological drought and its potential influence factors, *J. Hydrol.*, 547, 184-195,
795 10.1016/j.jhydrol.2017.01.041, 2017.

796 Hughes, J. D., Petrone, K. C., and Silberstein, R. P.: Drought, groundwater storage and stream flow
797 decline in southwestern Australia, *Geophys. Res. Lett.*, 39, 10.1029/2011gl050797, 2012.

798 Hunt, B. G.: Multi-annual dry episodes in Australian climatic variability, *Int. J. Climatol.*, 29, 1715-1730,
799 10.1002/joc.1820, 2009.

800 Jeremiah, E., Marshall, L., Sisson, S. A., and Sharma, A.: Specifying a hierarchical mixture of experts
801 for hydrologic modeling: Gating function variable selection, *Water Resour. Res.*, 49, 2926-2939,
802 10.1002/wrcr.20150, 2013.

803 Jung, I. W., Moradkhani, H., and Chang, H.: Uncertainty assessment of climate change impacts for
804 hydrologically distinct river basins, *J. Hydrol.*, 466, 73-87, 10.1016/j.jhydrol.2012.08.002, 2012.

805 Kumar, A., Panda, K. C., Nafil, M., and Sharma, G.: Identification of meteorological drought
806 characteristics and drought year based on rainfall departure analysis, *J. Appl. Sci. Technol.*, 51-59,
807 2020.

808 Kundu, S., Khare, D., and Mondal, A.: Individual and combined impacts of future climate and land use
809 changes on the water balance, *Ecol. Eng.*, 105, 42-57, 10.1016/j.ecoleng.2017.04.061, 2017.

810 Kunnath-Poovakka, A., and Eldho, T. I.: A comparative study of conceptual rainfall-runoff models GR4J,
811 AWBM and Sacramento at catchments in the upper Godavari river basin, India, *J. Earth Syst. Sci.*,
812 128, 10.1007/s12040-018-1055-8, 2019.

813 Kusangaya, S., Warburton, M. L., van Garderen, E. A., and Jewitt, G. P. W.: Impacts of climate change
814 on water resources in southern Africa: A review, *Phys. Chem. Earth*, 67-69, 47-54,
815 10.1016/j.pce.2013.09.014, 2014.

816 Laurène, J. E., Bouaziz, Aalbers, E. E., Weerts, A.H., Hegnauer, M., and Hrachowitz, M.: The importance
817 of ecosystem adaptation on hydrological model predictions in response to climate change, *Hydrol.*
818 *Earth Syst. Sc.*, 2021.

819 [Lawes, R. A., Oliver, Y. M., Robertson, M. J.: Integrating the effects of climate and plant available soil](#)
820 [water holding capacity on wheat yield, *Field Crop. Res.*, 113\(3\), 297-305, 10.1016/j.fcr.2009.06.008,](#)
821 [2009.](#)

822 Le Moine, N., Andreassian, V., and Mathevet, T.: Confronting surface- and groundwater balances on the
823 La Rochefoucauld-Touvre karstic system (Charente, France), *Water Resour. Res.*, 44,
824 10.1029/2007wr005984, 2008.

825 Leblanc, M. J., Tregoning, P., Ramillien, G., Tweed, S. O., and Fakes, A.: Basin-scale, integrated
826 observations of the early 21st century multiyear drought in southeast Australia, *Water Resour. Res.*,
827 45, 10.1029/2008wr007333, 2009.

828 [Leenaars, J. G. B., Claessens, L., Heuvelink, G. B. M., et al. Mapping rootable depth and root zone plant-](#)
829 [available water holding capacity of the soil of sub-Saharan Africa, *Geoderma*, 324, 18-36,](#)
830 [10.13140/RG.2.1.3950.9209, 2018.](#)

831 [Lenoir, J., Gégout, J. C., Marquet, P. A., de Ruffray, P., and Brisse, H.: A significant upward shift in plant](#)
832 [species optimum elevation during the 20th century, *Science*, 320\(5884\), 1768-1771,](#)
833 [10.1016/j.idairyj.2006.12.007, 2008.](#)

834 Lin, B. Q., Chen, X. W., Yao, H. X., Chen, Y., Liu, M. B., Gao, L., and James, A.: Analyses of landuse
835 change impacts on catchment runoff using different time indicators based on SWAT model, *Ecol.*
836 *Indicators*, 58, 55-63, 10.1016/j.ecolind.2015.05.031, 2015.

837 McNamara, J. P., Tetzlaff, D., Bishop, K., Soulsby, C., Seyfried, M., Peters, N. E., Aulenbach, B. T., and
838 Hooper, R.: Storage as a metric of catchment comparison, *Hydrol. Process.*, 25, 3364-3371,
839 10.1002/hyp.8113, 2011.

840 Melsen, L., Teuling, A., Torfs, P., Zappa, M., Mizukami, N., Clark, M., and Uijlenhoet, R.:
841 Representation of spatial and temporal variability in large-domain hydrological models: case study
842 for a mesoscale pre-Alpine basin, *Hydrol. Earth Syst. Sc.*, 20, 2207-2226, 10.5194/hess-20-2207-
843 2016, 2016.

844 Nepal, S., Chen, J., Penton, D. J., Neumann, L. E., Zheng, H. X., and Wahid, S.: Spatial GR4J

845 conceptualization of the Tamor glaciated alpine catchment in Eastern Nepal: evaluation of GR4JSG
846 against streamflow and MODIS snow extent, *Hydrol. Process.*, 31, 51-68, 10.1002/hyp.10962, 2017.

847 Nicholls, N.: The changing nature of Australian droughts, *Clim. Change*, 63, 323-336,
848 10.1023/B:CLIM.0000018515.46344.6d, 2004.

849 Nijzink, R., Hutton, C., Pechlivanidis, I., Capell, R., Arheimer, B., Freer, J., Han, D., Wagener, T.,
850 McGuire, K., Savenije, H., and Hrachowitz, M.: The evolution of root-zone moisture capacities
851 after deforestation: a step towards hydrological predictions under change?, *Hydrol. Earth Syst. Sc.*,
852 20, 4775-4799, 10.5194/hess-20-4775-2016, 2016.

853 Oke, O. A., Thompson, K. A.: [Distribution models for mountain plant species: the value of elevation](#),
854 *Ecol. Model.*, 301, 72-77, 10.1016/j.ecolmodel.2015.01.019, 2015.

855 Pan, Z. K., Liu, P., Gao, S. D., Cheng, L., Chen, J., and Zhang, X. J.: Reducing the uncertainty of time-
856 varying hydrological model parameters using spatial coherence within a hierarchical Bayesian
857 framework, *J. Hydrol.*, 577, 10.1016/j.jhydrol.2019.123927, 2019a.

858 Pan, Z. K., Liu, P., Gao, S. D., Xia, J., Chen, J., and Cheng, L.: Improving hydrological projection
859 performance under contrasting climatic conditions using spatial coherence through a hierarchical
860 Bayesian regression framework, *Hydrol. Earth Syst. Sc.*, 23, 3405-3421, 10.5194/hess-23-3405-
861 2019, 2019b.

862 Pan, Z. K., Liu, P., Xu, C. Y., Cheng, L., Tian, J., Cheng, S. J., and Xie, K.: The influence of a prolonged
863 meteorological drought on catchment water storage capacity: a hydrological-model perspective,
864 *Hydrol. Earth Syst. Sc.*, 24, 4369-4387, 10.5194/hess-24-4369-2020, 2020.

865 Patil, S. D., and Stieglitz, M.: Comparing Spatial and temporal transferability of hydrological model
866 parameters, *J. Hydrol.*, 525, 409-417, 10.1016/j.jhydrol.2015.04.003, 2015.

867 Perrin, C., Michel, C., and Andreassian, V.: Improvement of a parsimonious model for streamflow
868 simulation, *J. Hydrol.*, 279, 275-289, 10.1016/s0022-1694(03)00225-7, 2003.

869 Potter, N. J., Petheram, C., and Zhang, L.: Sensitivity of streamflow to rainfall and temperature in south-
870 eastern Australia during the Millennium drought, 19th International Congress on Modelling and
871 Simulation (MODSIM), Perth, Australia, 2011, WOS:000314989303087, 3636-3642, 2011.

872 Saft, M., Western, A. W., Zhang, L., Peel, M. C., and Potter, N. J.: The influence of multiyear drought on
873 the annual rainfall-runoff relationship: An Australian perspective, *Water Resour. Res.*, 51, 2444-
874 2463, 10.1002/2014wr015348, 2015.

875 Saft, M., Peel, M. C., Western, A. W., and Zhang, L.: Predicting shifts in rainfall-runoff partitioning
876 during multiyear drought: Roles of dry period and catchment characteristics, *Water Resour. Res.*,
877 52, 9290-9305, 10.1002/2016wr019525, 2016.

878 Sezen, C., and Partal, T.: The utilization of a GR4J model and wavelet-based artificial neural network for
879 rainfall-runoff modelling, *Water Supply*, 19, 1295-1304, 10.2166/ws.2018.189, 2019.

880 Shen, M. X., Chen, J., Zhuan, M. J., Chen, H., Xu, C. Y., and Xiong, L. H.: Estimating uncertainty and
881 its temporal variation related to global climate models in quantifying climate change impacts on
882 hydrology, *J. Hydrol.*, 556, 10-24, 10.1016/j.jhydrol.2017.11.004, 2018.

883 Simonneaux V, H. L., Boulet G, et al.: Modelling runoff in the Rheraya Catchment (High Atlas, Morocco)
884 using the simple daily model GR4J., Trends over the last decades [C]//13th IWRA World Water
885 Congress, Montpellier, France., 2008.

886 Singh, C., Wang-Erlandsson, L., Fetzer, I., Rockstrom, J., and van der Ent, R.: Rootzone storage capacity

887 reveals drought coping strategies along rainforest-savanna transitions, *Environ. Res. Lett.*, 15,
888 10.1088/1748-9326/abc377, 2020.

889 Siriwardena, L., Finlayson, B. L., and McMahon, T. A.: The impact of land use change on catchment
890 hydrology in large catchments: The Comet River, Central Queensland, Australia, *J. Hydrol.*, 326,
891 199-214, 10.1016/j.jhydrol.2005.10.030, 2006.

892 Thiemann, M., Trosset, M., Gupta, H., and Sorooshian, S.: Bayesian recursive parameter estimation for
893 hydrologic models, *Water Resour. Res.*, 37, 2521-2535, Doi 10.1029/2000wr900405, 2001.

894 Tian, J., Guo, S. L., Deng, L. L., Yin, J. B., Pan, Z. K., He, S. K., and Li, Q. X.: Adaptive optimal
895 allocation of water resources response to future water availability and water demand in the Han
896 River basin, China, *Sci. Rep.*, 11, 10.1038/s41598-021-86961-1, 2021.

897 Tu, J.: Combined impact of climate and land use changes on streamflow and water quality in eastern
898 Massachusetts, USA, *J. Hydrol.*, 379, 268-283, 10.1016/j.jhydrol.2009.10.009, 2009.

899 van Dijk, A., Beck, H. E., Crosbie, R. S., de Jeu, R. A. M., Liu, Y. Y., Podger, G. M., Timbal, B., and
900 Viney, N. R.: The Millennium Drought in southeast Australia (2001-2009): Natural and human
901 causes and implications for water resources, ecosystems, economy, and society, *Water Resour. Res.*,
902 49, 1040-1057, 10.1002/wrcr.20123, 2013.

903 Van Lanen, H. A. J., Wanders, N., Tallaksen, L. M., and Van Loon, A. F.: Hydrological drought across
904 the world: impact of climate and physical catchment structure, *Hydrol. Earth Syst. Sc.*, 17, 1715-
905 1732, 10.5194/hess-17-1715-2013, 2013.

906 Vrugt, J. A., Gupta, H. V., Bouten, W., and Sorooshian, S.: A Shuffled Complex Evolution Metropolis
907 algorithm for optimization and uncertainty assessment of hydrologic model parameters, *Water*
908 *Resour. Res.*, 39, 10.1029/2002wr001642, 2003.

909 Westra, S., Thyer, M., Leonard, M., Kavetski, D., and Lambert, M.: A strategy for diagnosing and
910 interpreting hydrological model nonstationarity, *Water Resour. Res.*, 50, 5090-5113,
911 10.1002/2013wr014719, 2014.

912 Yandell, B. S.: Kernel Smoothing, *Technometrics*, 38, 75-76, 1996.

913 Zeng, L., Xiong, L. H., Liu, D. D., Chen, J., and Kim, J. S.: Improving Parameter Transferability of GR4J
914 Model under Changing Environments Considering Nonstationarity, *Water*, 11, 10.3390/w11102029,
915 2019.

916 Zhang, Y. Q., Viney, N., Frost, A., Oke, A., Brooks, M., Chen, Y., and Campbell, N.: Collation of
917 Australian modeller's streamflow dataset for 780 unregulated Australian catchments, CSIRO: Water for
918 a healthy country national research flagship, 115 pp, 2013.

919

920 **Tables**

921 **Table 1.** Description of the dataset adopted in this study.

922

Data type	Description	Data source
Meteorological data	Daily precipitation, potential evapotranspiration	
Runoff data	Daily runoff data from hydrological stations	Australian Water Resources Assessment system
Catchment features	Catchment area, elevation, slope, forest coverage percentage, AWHC of the soil, K_s of the soil	

923 **Note:** AWHC denotes the available soil water holding capacity; K_s refers to the saturated hydraulic
924 conductivity.

925

926

927 **Table 2.** Summary of the characteristics of the 145 catchments that had the prolonged
928 meteorological drought, including the mean, median, minimum, and
929 maximum estimates of 9 catchment features.

930

Number	Catchment features	Mean	Median	Minimum	Maximum
A1	Area (km ²)	711.17	363.0	54.0	6818.0
A2	Mean elevation (m)	542.57	468.0	47.0	1351.0
A3	Slope range (°)	22.18	22.6	2.1	49.9
A4	Mean slope (°)	5.49	5.0	0.3	13.6
A5	Forest coverage (%)	55.00	57.0	15.0	92.0
A6	AWHC of the topsoil (mm)	41.26	42.0	22.0	64.0
A7	AWHC of the subsoil (mm)	88.66	87.5	27.0	188.0
A8	K _s of topsoil (mm/h)	157.52	160.0	31.0	283.0
A9	K _s of subsoil (mm/h)	62.10	53.0	4.0	216.0

931

932

933
 934
 935

Table 3. Ranges of the initial values of GR4J model parameters.

Parameters	Meaning	Unit	Min	Max	
α_1, α_2	Amplitude of the sine function	/	-200	200	
θ_1	β_1, β_2	Frequency of the sine function	/	0	1
	γ_1, γ_2	Remainder in the sine function	/	-200	200
	δ_1, δ_2	Intercept of the sine function	/	-300	300
θ_2	Groundwater exchange coefficient	mm	-5.0	5.0	
θ_3	Capacity of catchment reservoir	mm	1.0	200.0	
θ_4	Unit line confluence time	day	0.1	10.0	

936
 937

938 **Table 4. Category of the** selected variables that may be associated with the changes in
 939 the ACWSC. The selected variables was divided into two parts, i.e., cathment features
 940 (9 variables) and climate variables (24 variables).

941

Category	Catchment features	Category	Climate variables
A1	Area (km ²)	A6	AWHC of the topsoil (mm)
A2	Mean elevation (m)	A7	AWHC of the subsoil (mm)
A3	Slope range (°)	A8	K _s of topsoil (mm/h)
A4	Mean slope (°)	A9	K _s of subsoil (mm/h)
A5	Forest coverage (%)		
Category	Climate variables	Category	Climate variables
B1	Mean daily precipitation (mm)	B13	Mean summer runoff(mm)
B2	Mean daily potential evapotranspiration(mm)	B14	Mean autumn runoff(mm)
B3	Mean Daily T _{max} (°C)	B15	Mean winter runoff(mm)
B4	Mean Daily T _{min} (°C)	B16	Mean annual precipitation (mm)
B5	C _v of monthly precipitation	B17	Mean annual potential evapotranspiration(mm)
B6	C _v of monthly runoff	B18	Mean annual runoff(mm)
B7	Mean monthly runoff index	B19	Mean annual aridity ratio
B8	Mean spring precipitation (mm)	B20	Mean annual runoff index
B9	Mean summer precipitation (mm)	B21	C _v of annual precipitation
B10	Mean autumn precipitation (mm)	B22	C _v of annual runoff
B11	Mean winter precipitation (mm)	B23	Mean annual base flow (mm)
B12	Mean spring runoff(mm)	B24	Annual base flow ratio

942

943 **Table 5.** Summary of catchments with different change patterns in the amplitude α
 944 and mean value δ in the regression function of the ACWSC due to a prolonged
 945 meteorological drought.

946

Factors	Magnitude	Change direction	Number of catchments	Percentage	
Amplitude (α)	Significant change	Increased	83	57.24%	
		Decreased	4	2.76%	
	Non-significant change	Increased	3	2.07%	
		Decreased	2	1.38%	
	Catchments that do not meet the criteria for the maximum performance degradation and result robustness			53	36.55%
	Catchments with a prolonged meteorological drought			145	100%
Mean value (δ)	Significant change	Increased	77	53.10%	
		Decreased	0	0	
	Non-significant change	Increased	10	6.90%	
		Decreased	5	3.45%	
	Catchments that do not meet the criteria of the maximum performance degradation and result robustness			53	36.55%
	Catchments with a prolonged meteorological drought			145	100%

947

948

949 **Table 6.** Response times of different groups of catchments with significant increase/
950 decrease in regression parameters α and δ .


951

Catchment type	Average (day)	Median (day)	Minimum (day)	Maximum (day)
Catchments with significant increase in δ	691.1	781.0	92.2	1082.0
Catchments with significant decrease in δ	/	/	/	/
Catchments with significant increase in α	690.8	781.0	92.2	1082.0
Catchments with significant decrease in α	422.3	452.4	122.6	661.9

952

953 **Figures**


954



955

956 **Fig.1.** Spatial distribution of the original 398 catchments in south-eastern Australia


957 that were selected from Zhang et al. (2013).



959

960


961 **Fig.2.** The drought periods correspond to 145 catchments with prolonged
 962 meteorological drought in the south-eastern Australia.



963

964


Fig.3. Flowchart of the proposed methodology and procedures.




965

966

Fig.4. Diagram of the GR4J model proposed by Perrin et al. (2003).



(a) Amplitude α




(b) Mean value δ


967
968

969
970


971 **Fig.5.** Spatial distribution of catchments with different change patterns in the ACWSC
 972 after the prolonged drought. Subfigures (a) and (b) illustrate the spatial
 973 distribution of catchments with different variation forms in the amplitude α and
 974 mean value δ during the drought period, respectively.




(a) Magnitude of absolute change in estimated parameter α



(b) Magnitude of the relative change percentage in estimated parameter α




(c) Magnitude of absolute change in estimated parameter δ




(d) Magnitude of the relative change percentage in estimated parameter δ

975

976 **Fig.6.** The magnitudes of change in the amplitude and mean value of the ACWSC
 977 between the periods before and after the change point. Sub-figures (a) and (b) illustrate
 978 the magnitude of absolute and relative percentage changes in estimated parameter α ,
 979 respectively. Sub-figures (c) and (d) refer to magnitude of absolute and relative
 980 percentage changes in the estimated mean value of parameter δ .




(a) Catchment 222206



(b) Catchment 421042

981


982 **Fig.7.** Time-varying patterns of model parameter θ_l in two example catchments (i.e.,
 983 catchment 222206 and 421042).



984

985 **Fig.8.** Magnitude distribution of the response time in 92 catchments that satisfied the
986 basic criteria of NSE performance and result robustness.

987



988

989 **Fig.9.** Comparison of physical features between the $g_{\alpha}(S)$ and $g_{\alpha}(NS)$ groups and

990 $g_{\alpha}(SI)$ and $g_{\alpha}(SD)$ subsets for the study catchments. The orange and green boxes (left

991 two columns) denote the physical characteristics of the $g_{\alpha}(S)$ and $g_{\alpha}(NS)$ groups

992 which was divided according to the significance level of the variation in the amplitude


993 after the change point. The purple and yellow columns (right two columns) denote the

994 catchment features of the $g_{\alpha}(SI)$ and $g_{\alpha}(SD)$ subsets with significantly increased and

995 decreased change patterns in the amplitude after the change point, respectively.

996

997




998

999 **Fig.10.** Comparison of catchment characteristics between the groups of catchments

1000 with significant and non-significant changes in mean value δ , i.e., $g_\delta(S)$ and


1001 the $g_\delta(NS)$ groups.



1002


1003 **Fig.11.** The Pearson correlation coefficient between the variation in the amplitude α
 1004 with multiple catchments features and climate variables. (a) Correlation between the
 1005 absolute variation of amplitude α and catchment features; (b) Correlation between the
 1006 relative variation of amplitude α and catchment features; (c) Correlation between the
 1007 absolute variation of amplitude α and absolute variation of climate variables; (d)
 1008 Correlation between the relative variation of amplitude α and relative variation of
 1009 climate variables.

1010




1011

1012 **Fig.12.** The Pearson correlation coefficient between the variation in the mean value δ
1013 with multiple catchment features and climate variables. (a) Correlation between the
1014 absolute variation of mean value δ and catchment features; (b) Correlation between
1015 the relative variation of mean value δ and catchment features; (c) Correlation between
1016 the absolute variation of mean value δ and absolute variation of climate variables; (d)
1017 Correlation between the relative variation of mean value δ and relative variation of
1018 climate variables.



1019 Fig.13. Trend analysis between the variation in the ACWSC and catchment properties.

1020



1021

1022


1023

1024

1025

1026

Fig.14. The Pearson correlation coefficient between the response time with catchment features and variation in climate variables before and after the change point. (a) Correlation between the response time and catchment features; (b) Correlation between the response time and absolute change of climate variables; (c) Correlation between the response time and relative change of climate variables.



1027

1028

Fig.15. The potential connections between the response time and catchment properties.