



## Flood generation: process patterns from the raindrop to the ocean

Günter Blöschl<sup>1,\*</sup>

5 <sup>1</sup> Institute of Hydraulic Engineering and Water Resources Management, Vienna University of Technology, Karlsplatz  
13/223, 1040 Vienna, Austria

\* Invited contribution by Günter Blöschl, recipient of the EGU Dalton Medal 2019.

*Correspondence to:* Günter Blöschl (bloeschl@hydro.tuwien.ac.at)

10

**Abstract.** This article reviews river flood generation processes and flow paths across space scales. The scale steps include the pore, profile, hillslope, catchment, regional and continental scales, representing a scale range of a total of 10 orders of magnitude. Although the processes differ between the scales, there are notable similarities. At all scales, there are media  
15 patterns that control the flow of water, and are themselves influenced by the flow of water. The processes are therefore not spatially random (as in thermodynamics) but organised, and preferential flow is the rule rather than the exception. Hydrological connectivity, i.e. the presence of coherent flow paths, is an essential characteristic at all scales. There are similar controls on water flow and thus on flood generation at all scales, however, with different relative magnitudes. Processes at lower scales affect flood generation at the larger scales not simply as a multiple repetition of pore scale  
20 processes, but through interactions, which cause emergent behaviour of process patterns. For this reason, when modelling these processes, the scale transitions need to be simplified in a way that reflects the relevant structures (e.g. connectivity) and boundary conditions (e.g. groundwater table) at each scale. In conclusion, it is argued that upscaling as the mere multiple application of small scale process descriptions will not capture the larger scale patterns of flood generation. Instead, there is a need to learn from observed patterns of flood generation processes at all spatial scales.

25

### 1 Introduction

30 The importance of research on flood generation processes cannot be overestimated. Numerous recent floods in the last years in Europe and worldwide have highlighted that floods continue to be one of the most serious threats to humankind (Merz et al., 2021). Major recent events include the July 2021 flood in Germany with more than 200 fatalities; the July 2020 flood in Bangladesh where the monsoon season was one of the worst on record; the more local June 2016 flood in Simbach, Bavaria;



and the June 2013 flood in central Europe, all with enormous economic damage. While, previously, much of the flood  
35 planning and management has been based on empirical approaches, these are no longer sufficient. This is because better  
accuracy is required than simple extrapolations from observations permit, and because any changes in the flood  
characteristics, be it through climate, land use or river works, can be much more reliably predicted when explicitly  
accounting for the flood generation processes (Blöschl et al., 2015).

40 The purpose of this article is to review our present understanding of flood generation processes across space scales. The  
focus is on the land phase of the processes, i.e. on what happens to extreme rainfall when it falls on the ground. At each  
scale, the mechanisms of runoff generation are discussed, as are the processes that control the flow paths and thus the  
magnitude of the floods. A sound understanding of these processes will assist in building more reliable models of floods and  
their probabilities, and ultimately enhance the robustness of flood management methods.

45

## 2 How do raindrops aggregate to form a flood?

Let us take one step back, and conceptually envisage the fate of one raindrop when it falls on the ground. Obviously, its flow  
50 path will depend on *where* it falls. Depending on the surface characteristics, it may infiltrate into the soil, remain on the soil  
surface until it evaporates, or else run off on the surface. The raindrop may in fact not even reach the ground and get  
intercepted in the canopy of the vegetation, or it may fall as snow when it is cold. The raindrop, if not evaporated, will start  
its journey to the ocean, either at low speed in the subsurface where resistance is high, or on the surface where it quickly  
reaches a stream, or – is often the case – as a mix of the two.

55

So far, so good. In flood generation one is interested in the effect of many raindrops on the flow of water in the streams and  
on the land surface so, again conceptually thinking, the task seems easy. There is ‘only’ a need to follow the path of each and  
every raindrop to the ocean and the cumulative effect of all of them will give an exact representation of the water cycle,  
including flooding. Alas, there are too many drops. For example, the 2013 Danube flood at Vienna had a volume (including  
60 baseflow) of 9.5 billion m<sup>3</sup> (Blöschl et al., 2013) which, assuming a rain drop diameter of 1 mm, gives a total of  $2 \cdot 10^{19}$  drops.  
If one needed only one second to simulate the path of a drop on a computer, the total computation time would be  $6 \cdot 10^{11}$   
years, i.e., the simulations would be finished at a time when the state of the universe will be rather uncertain. So this is  
clearly not feasible, apart from observational difficulties. Alternatives are needed and they involve some kind of aggregation,  
which leads us to the question: “How do raindrops aggregate to form a flood?”

65

Hydrology has first fully realised this to be a problem with the advent of digital computers (Sivapalan and Blöschl, 2017).  
The classical treatment of this question is Allan Freeze’s “Blueprint for a physically-based model” (Freeze and Harlan, 1969)



70 where they conclude: “The level of development is not adequate to permit the construction of complete physically-based, hydrologic response models at this time.” and “simplification of the model is needed [...] while maintaining physical relevance of the controlling hydrological parameters.” (Freeze and Harlan, 1969, pp. 255-256). Since then the aggregation problem has also become known as the “scale issue” and has given rise to numerous initiatives and publications, among them a series of workshops on the topic in Caracas, Venezuela; Princeton, USA; Robertson, Australia; and Krumbach, Austria (Rodríguez-Iturbe and Gupta, 1983; Gupta et al., 1986; Kalma and Sivapalan, 1995; Blöschl et al., 1997).

75 A key element in understanding how best to scale from the point scale to large areas is the nature of hydrological variability. Again a classic is the paper of Dooge (1986) who, drawing on Weinberg (1975), suggests that variability may fall in three types: (i) organized simplicity (which can be treated analytically, e.g. by the methods of mechanics); (ii) unorganized complexity (which can be treated statistically, e.g. by the methods of thermodynamics) and (iii) organized complexity (the treatment of which is difficult). Dooge (1986, p. 49S) notes that “Most problems arising in catchment hydrology fall in the category of complex systems with some degree of organization”. Which means the difficult one. The discussion on the degree to which spatial organisation and patterns need to be represented in catchment and subsurface hydrology (as opposed to a purely statistical treatment as in thermodynamics has since then not subsided (Western et al., 2001; Rajaram, 2016).

85 A related issue is the interpretation of variability on which, too, no last word has been spoken in hydrology. Specifically, the question is whether randomness and thus uncertainty are inherent in the processes (so there is no way of reducing them and a probabilistic treatment is unavoidable) or simply a reflection of a lack of knowledge (so more accurate measurements will allow a deterministic treatment). Two contrasting opinions on the subject are the following. The first, in the context of a new hydrological blueprint, suggests that uncertainty is inherent in the processes: “a probability based theoretical scheme for building process-based models of uncertain hydrological systems, thereby unifying hydrological modeling and uncertainty assessment” (Montanari and Koutsoyiannis, 2012; p. 1). The second, more pragmatic, interprets it as a lack of knowledge: “It should be noted that the possibility of using, in the treatment of a real process, schemes of well determined or of only stochastically definite processes stands in no relation to the question whether the real process is itself determined or random.” (Kolmogoroff, 1931; p. 417).

95 While these considerations seem to highlight the challenges of connecting the scales, there is perhaps also a positive side to it, drawing on the opportunities of scale research. As Blöschl (2001, p. 711) remarked: “Given that scale issues are common to all of the hydrologic subdisciplines (and indeed far beyond them), scaling work may perhaps reveal its greatest potential as an umbrella under which a rich spectrum of concepts, tools, and measurement techniques covering a range of areas can be unified.”

100



We now turn to the more concrete question of how floods are generated. Given that the scale issues are of paramount importance, we will deal with flood generation separately at different space scales, starting from the pore scale (at a characteristic length of 1 mm) and moving up to the profile, hillslope, catchment, regional and continental scales, covering a scale range of a total of 10 orders of magnitude (Fig. 1). We will review the mechanisms of flood generation at each scale with the ultimate purpose of linking them across scales. Fortunately, there is a very rich body of hydrological literature on the processes at each scale, although their linkages have been studied less.




Figure 1: Scale transitions across 10 orders of magnitude.

110

### 3 The pore scale: $10^{-3}$ m

115 At the pore scale we are in the domain of soil physics which has a mature theory of water movement in unsaturated soils under the influence of gravity and capillary forces between water and the soil grains (e.g. Jury and Horton, 2004) as well as of infiltration (Smith et al., 2002). Much has been said about the overwhelming spatial variability of soil characteristics (e.g. Bierkens et al., 2000) but, even at one single point, there is enormous complexity. Clay minerals for example, important components of many soils, may have diverse shapes and tend to change their form and size very significantly as they become wet or dry out (Tessier, 1990), leading to soil swelling and shrinking. The implication of this is that traditional theories of water flow no longer apply, but attempts at accounting for swelling and shrinking exist. For example, Tuller and Or (2003) have modelled the changes in volume and pore space induced by the swelling and shrinking behaviour of clay minerals for predicting the permeability of clayey soils as a function of soil suction. An even more taunting task is the understanding of water – soil – water feedbacks, i.e. the phenomenon that the moisture state will modify the hydraulic characteristics of the soil, which will modify the flow and thus moisture, which will modify the hydraulic characteristics and so forth. If plants interact with the water and the soil characteristics, these feedbacks become even more complex (Alaoui et al., 2018)

125



The swelling and shrinking processes of the soil, while operative at the scale of micrometers, can become very relevant at larger scales, visible to the naked eye, and a major determinant of infiltration and flood generation. An example is shown in  
130 Fig. 1 where water flowing overland infiltrates into the shrinkage cracks, giving rise to preferential water movement in the soil, and reducing the amount of runoff generated locally. One would expect that, for small storms, these cracks will reduce flood peaks while, when they are filled with water, flood discharges may abruptly increase.




135 **Figure 2: Shrinkage cracks in the Hydrological Open Air Laboratory (HOAL), Lower Austria, which will lead to preferential infiltration. Photo: A. Eder.**

#### 140 **4 The profile scale: $10^0$ m**

At the soil profile scale, there is, again, established theory and the dominant paradigm is Richards equation (Richards, 1931) which combines Darcy's law with mass balance for representing the vertical flow of water in the soil matrix, and thus infiltration on the soil surface. Its physical basis (i.e. Darcy's law) is sometimes used as a justification for a widespread use  
145 in hydrology, and one could even say, it is occasionally idolised as a "sacred cow". Notwithstanding all its favourable properties, one has to remember that it is only applicable to some soils and, perhaps more importantly, it represents only one out of many processes controlling runoff generation. Not only the issues discussed at the pore scale also apply here, but also additional phenomena occur at the profile scale not captured by Richards equation.



- 150 Based on dye-tracing experiments carried out at 14 field sites in Switzerland, Flury et al. (1994, p. 1945) noted: “The spatial structure of flow patterns showed remarkable differences. In most soils, water bypassed the soil matrix.” Preferential flow is thus the rule rather than the exception (Fig. 3). Macropores and soil layers tend to contribute to the preferential nature of the flow as does the varying moisture state. Based on simulations, Zehe et al. (2007) explored the effect of initial soil moisture on the occurrence of vertical preferential flow and found that at intermediate soil moisture, when a transition between matrix and preferential flow occurs, flow becomes very sensitive to small-scale soil moisture variations. Thus, “it seems that small uncertainties can easily amplify under certain conditions and will limit predictability.” (Blöschl and Zehe, 2005, p. 3923).
- 155



- 160 **Figure 3: Preferential infiltration. Dye flow patterns observed at the Weiherbach, Germany, irrigation site. From Zehe et al. (2007).**



## 5 The hillslope scale: $10^2$ m

165

The analysis at the profile scale considers one point in the landscape at a time, and thus treats runoff generation as a local phenomenon, usually referred to as the infiltration excess runoff generation mechanism attributable to Robert Horton (Horton, 1933). In reality, spatial connections of the processes are often important, so a treatment at the hillslope scale is appropriate. The classical contribution here is that of Dunne and Black (1970) who have highlighted the relevance of saturation excess runoff generation based on field observations in the Sleepers river watershed in Vermont. They note: "... overland flow generated by the mechanism described by Horton did not occur. [...] When the water table rose to the surface of the ground, however, overland flow was generated on small areas of the hillside." Dunne and Black (1970, p. 478). Since the water table is controlled, among other factors, by the lateral subsurface redistribution, this is no longer a local mechanism. A similarly spatial process was observed on the Panola hillslope in Georgia: "When connectivity was achieved, the instantaneous subsurface stormflow rate increased more than fivefold compared to before the subsurface saturated areas were connected to the trench face." (Tromp-van Meerveld and McDonnell, 2006, p. 1). In this case, the critical process is saturation near the bedrock-soil interface, as opposed to surface saturation in the case of saturation excess runoff.

170

175

A term mentioned in the Panola study is "flow connectivity", representing the presence of coherent flow paths along the hillslope. Traditionally, connectivity had mainly been studied in fractured rocks (e.g. Andersson and Dverstorp, 1987) and it is only in the 1990s that it has received increased attention in catchment hydrology (e.g. Western et al., 1998, 2001; Grayson and Blöschl, 2001). Of key relevance has been the recognition in hydrology (although long known to pedology, Milne, 1935; Jenny, 1941), that the soil characteristics are organised along a hillslope, and that this organisation is critically important for runoff generation. Here, again, feedbacks can occur between the soil structure and flow, but this time modulated by animal activity. An example is the role of earthworms. In some climates, there is a tendency for larger soil moisture at the foot of a hillslope, which favours the presence of earthworms and thus increases the number of earthworm holes, which in turns increases the flow conductivity, with repercussions on soil moisture (Zehe et al., 2010).

185

Flow connectivity and feedbacks can also be very relevant in the context of sediment movement. As overland flow concentrates along rills, in particular in agricultural landscapes, erosion is enhanced (because shear stress is proportional to flow depth, Biron et al., 2004), which leads to a deeper incision of rills and therefore more flow concentration. While traditional stream gauge measurements are not very good at capturing these connectivity patterns, novel techniques such as terrestrial or aerial video images can provide detailed pattern information (Silasari et al., 2017; Deng et al., 2020). As an example, Fig. 4 shows the saturation patterns in the HOAL. As the saturation area exceeds  $80 \text{ m}^2$ , the flow paths connect and runoff from the hillslope starts abruptly.

190

195




Figure 4: Connectivity of overland flow on 2 March 2015 in the Hydrological Open Air Laboratory (HOAL), Lower Austria. Left: Observations of precipitation (at a raingauge), saturation area (from an analysis of the videos), and runoff (at a flume). Right:

200 Video images of the saturation patterns. From Silasari et al. (2017)

## 6 The catchment scale: $10^3$ m

205

A catchment is of course composed of many hillslopes, but how the flood generation responses of many hillslopes combine exactly, has been the subject of a long debate in hydrology. At one end of the spectrum is again the blueprint of Freeze and Harlan (1969), which describes the processes on each hillslope in a detailed way, to subsequently combine them to catchment response, an approach sometimes called “upward”. At the other end of the spectrum is the “downward” approach that views catchments as organised complex systems in the sense of Dooge (1986), that co-evolve (Sivapalan and Blöschl, 2015), and “develop features of spatial organization, including surface or sub-surface drainage patterns, as a by-product of this evolution.” (Savenije and Hrachowitz, 2017, p. 1107). If the latter is true, flood generation should no longer be treated as a boundary value problem (as in Freeze and Harlan, 1969) because “models that fail to account for patterns and the associated feedbacks miss a critical element of how systems at the interface of atmosphere, biosphere and pedosphere function.” (Savenije and Hrachowitz, 2017, p. 1107). So how do we reconcile these views? Sivapalan (2003, p. 1040) suggests the following: “One way to achieve this reconciliation is to focus on common concepts, features or patterns that have physical meanings that transcend the range of scales in question, and which are easily scalable.” Common concepts are, for example, travel time distributions. Additionally, instead of analysing a single catchment in detail, comparing many





catchments with contrasting characteristics may be useful for understanding the controls in a holistic way, an approach  
220 known as comparative hydrology (Gaál et al., 2012).

Experimental basins provide evidence of how the hillslope responses combine to that of an entire catchment. One example is  
the 66 ha Hydrological Open Air Laboratory (HOAL), where streamflow is observed at 13 flumes, along with numerous  
other hydrological variables. The moisture observations of Fig. 5 show that the spatial patterns differ between winter and  
225 summer with a stronger spatial organisation in winter because of the higher relevance of lateral redistribution of water, in  
line with studies in the Tarrawarra, Australia, for example (Grayson et al., 1997). The HOAL is particularly interesting as it  
features different runoff generation mechanisms within a rather limited area, including runoff generation through overland  
flow (partly infiltration excess during high intensity storms in summer, partly saturation excess during winter), runoff  
generation via tile drains (with very fast response times of less than an hour even though the drain pipes are about 60 cm  
230 below ground) and runoff from permanently saturated areas (local wetlands) (Blöschl et al., 2016). Soil moisture is a major  
control on runoff generation in this catchment, as illustrated in Fig. 6, but the peak flow – soil moisture relationship depends  
very much on the runoff generation mechanism: threshold type behaviour for the overland flow mechanism, more gradual  
for the other mechanisms, and the total outflow of the catchment is a mix of these processes. The threshold behaviour  
observed in the HOAL is not an exception. For example, a similar relationship was found by Penna et al. (2011) in a  
235 catchment in the Italian Dolomites. In order to explain the non-linear runoff generation in a spatial context, various  
theoretical studies have explored the aggregation of hillslope response in the spirit of these observations, accounting for the  
space-time covariance of precipitation and soil moisture by an upward approach (Woods and Sivapalan, 1999; Viglione et  
al., 2010). These studies assist in generalising the understanding of the scaling behaviour beyond individual case studies.

240





Figure 5: Soil moisture patterns in the Hydrological Open Air Laboratory (HOAL), Lower Austria. Top: a summer event (9 and 10 August, 2014). Bottom: a winter event (6 and 7 December, 2014). Note the more pronounced spatial organisation of soil moisture in winter, in particular during the event. Bottom left shows the flumes used in Fig. 6: O (Overland flow), T (Tile drains), S (Saturation area), E (Entire catchment). See Blöschl et al. (2016) for details.

245



250 **Figure 6: Non-linear runoff generation in the Hydrological Open Air Laboratory (HOAL), Lower Austria. Scaled peak runoff of events plotted against scaled soil moisture at the beginning of these events. Each point represents one event. The panels refer to the flumes shown in Fig. 5, which represent different runoff generation mechanisms. From Vreugdenhil et al. (2022).**

## 255 **7 The regional scale: $10^5$ m**

At the regional scale, the comparative, diagnostic approach to identifying flood generation processes is particularly powerful. For example, Thompson et al. (2011, p. 1) “suggest that the spatial patterns of vegetation may offer a lens through which to investigate scale dependence of hydrology “. Analysing various regions in the US, they found that the organisation of

260 vegetation imposes spatial dependence in areally averaged hydrologic variables such as soil moisture and the partitioning of flow paths. There is an example of the downward approach that harmonises with the identification of spatial organisation in catchments viewed as complex systems (Savenije and Hrachowitz, 2017). The findings are directly relevant to scale effects in flood generation, given that soil moisture is a key control at all scales (Western et al., 2002). A similar comparative approach, albeit with a stronger focus on climatic controls, identifies flood types such as snow melt floods, rain-on-snow

265 floods, flash floods and synoptic floods. An example is shown in Fig. 7. The patterns of the frequency of types could not easily be inferred from other variables such as topography. For example, rain-on-snow floods are most frequent in the north of Austria, and this is related to the seasonal interplay of snowmelt processes and soil moisture that differs from other regions. The more frequent flash floods in the East are related to the higher frequency of convective storms and the faster catchment response times (Gaal et al., 2012). The typology approach has recently been extended to also include processes

270 related to infiltration and runoff routing (Tarasova et al., 2019, 2020). The importance of these typologies for understanding



flood generation derives from the process similarity of events pertaining to the same class, e.g., similarity in terms of the transition to extreme events and in terms of changes of flood processes with time (Merz et al., 2021; Blöschl et al., 2015).

Similar to the other scales, the aggregation processes at the regional scale are complex. Fig. 8 shows the spatial flood hydrograph pattern of the June 2013 flood in the Danube basin as an example. The flood response of the Bavarian Danube in the northwest of the basin is delayed and the runoff coefficient at Hofkirchen is only 0.27 because of the highly permeable subsurface, while the Inn in the southwest exhibits a faster response and the runoff coefficient at Schärding is around 0.5 because of shallower soils, steeper terrain and higher rainfall (Blöschl et al., 2013). The Upper Inn shows very little flood runoff, and the flood wave built up through tributaries in Bavaria. The confluence of the Inn at Schärding with the Bavarian Danube at Hofkirchen resulted in a characteristic, combined shape of the flood wave at Achleiten, where the fast and slow contributions of the Inn and Danube are clearly visible. During the propagation of the flood wave along the Austrian Danube, it changed shape due to retention in the flood plains. It can be appreciated that the spatial patterns of runoff controlled by precipitation, some snowmelt, soil characteristics and soil moisture are aggregated and modulated as the flood wave propagates through the stream network. Understanding these patterns is extremely important both for flood forecasting and for flood risk mapping (e.g. Vorogushyn et al., 2010; Viglione et al., 2013; Merz et al. 2014; Nester et al., 2015).





Figure 7: Frequency of regional flood process types in Austria based on observed maximum annual peak discharges of 12000 events classified according to their type. Each polygon represents a gauged catchment. From Merz and Blöschl (2003).

290



**Figure 8:** Spatial flood hydrograph pattern of the 2013 event in the Danube Basin upstream of Vienna, Austria. Red circles indicate the stream gauges. The scale shown on the bottom right relates to all hydrographs (light blue areas). From Blöschl et al. (2013).

295

### 300 **8 The continental scale: $10^7$ m**

As one moves up in scale to an entire continent, one could – similarly as at other aggregate scales – view flood generation simply as the sum of the processes in many pores, profiles, hillslopes, catchments and regions. This upward view (in the spirit of Freeze's blueprint) has given rise to the hyperresolution modelling paradigm at continental and global scales (Wood et al., 2011; Bierkens et al., 2015; O'Neill et al., 2021). There is much to be said in favour of these models as they codify existing system knowledge, and modern computer resources permit simulations at resolutions of hundreds or even tens of meters globally. On the other hand, Sivapalan (2018, p. 1674) noted that “[upward] reductionist type models are best suited to represent ‘known’ knowledge, and are at their weakest to reveal ‘unknown’ knowledge [...]; vice versa for [downward]

305



[...] models". Complexities such as preferential flow, connectivity and media-flow interactions are rarely codified at each  
310 scale they can be observed, let alone their summary effect at the continental scale modulated by cross-scale feedbacks.  
Hydrologists have been sorely aware of these scale issues since the 1980s. Ways forward may include downward  
diagnostics, i.e. learning from observed patterns not only at small scales, but also at the continental scale in order to capture  
the summary effect of all these interactions on runoff generation.

315 Meteorology has a longer tradition of continental (and global) scale diagnostics. At the continental scale, the structure of the  
stream network tends to become less important and, instead, the atmospheric situation controls the movement and direction  
of cyclones, the intensity of the rainfall they produce, and thus the flood characteristics. Classifying cyclone tracks based on  
air pressure and other data allows identifying those cyclones that are most prone to producing regional floods. For example,  
Vb tracks (van Bebber, 1891) that propagate from northern Italy to Poland, tend to produce the highest precipitation rates in  
320 Central Europe, such as those responsible for the June 2003 flood (Fig. 8). Fig. 9 illustrates that, in the Ore mountains, the 20  
year maximum annual daily precipitation during Vb events is 62 mm/d, while it ranges between 20 and 35 mm/d for other  
event types (Hofstätter et al., 2016, 2018). Superposition of the polar and the subtropical jet stream over the Western  
Mediterranean has been identified as a main feature triggering Vb cyclones (Hofstätter et al., 2019). A classification of flood  
generation mechanisms similar to that of Fig. 7 can also be performed at the continental scale. For example, Kemter et al.  
325 (2020) showed that synoptic rainfall is particularly important in the Alps and the Carpathians, soil moisture excess in the  
Atlantic climate of Western Europe, and snowmelt in the north and east of Europe. Rain-on-snow has some relevance in the  
midlands of central Europe.

A simple, but very efficient, diagnostic of these runoff generation processes at the continental scale is the timing of the year  
330 floods occur, which is illustrated in Fig. 10a. In Europe, the timing of the floods varies gradually from winter floods in the  
west to spring floods in the east because of increasing continentality (distance from the Atlantic), and from winter floods in  
the south to spring floods in the north because of the increasing influence of snowmelt. The effect of snow storage and melt  
in the Alps and the Carpathians (summer floods) is superimposed on this spatial pattern. The spatial pattern in Fig. 10a is  
quite different from that of the individual drivers, as illustrated for the case of extreme precipitation (Fig. 10b). This is  
335 because of the interplay between the seasonality of event precipitation, snowmelt, soil moisture and evaporation.

Hydrologists are not commonly used to interpret observed patterns at the continental scale, but there is a lot to be learned  
from them, also for small scale studies. For example, one notes that the west coasts of the European islands and peninsulas  
(e.g. UK, Denmark, Iberia, Italy, Greece) all tend to have earlier flood occurrence by one month than the respective east  
340 coasts, a finding that suggests similarities between the relative magnitudes of flood generation processes also at the  
catchment scale. Comparative hydrology (Sivapalan, 2018) exploits these similarities and helps avoid being "frogs in the  
well" (as the Chinese saying goes), and instead see the bigger picture that is also valuable in a local context.




Figure 9: Storm tracks of cyclone types and associated heavy precipitation in the Ore mountains, 2500 cyclone events. From information in Hofstätter et al. (2018).  
345




Figure 10: (a) Observed average timing of river floods in Europe based on maximum annual discharge series 1960–2020 from 4062 stream gauges. Light blue: winter floods; green to yellow: spring floods; orange to red: summer floods; purple to dark blue: autumn floods. (b) Similar maps but for 7-day maximum annual precipitation based on E-OBS data. Redrawn from Blöschl et al. (2017).  
350



## 9 Conclusions

355

This paper has reviewed flood generation processes across scales with a focus on spatial process patterns. Understanding these processes is not only of scientific interest but has enormous practical importance for risk management purposes, including flood forecasting and flood risk assessment in a changing climate.

360

There is remarkable similarity of the flood generation mechanisms between the scales from the pores to the continent. At all scales, processes are not random (as in thermodynamics) but they exhibit organised structures that control the flow of water. There is often an interplay between the structure (e.g. soil characteristics, topography) and the flow characteristics (e.g. flow velocity). For example, overland flow is controlled by the rills on the surface, but they themselves are created by overland flow. Remarkably, there is preferential water flow at all scales; flow in pores and macropores in the soil, rills on the land surface, preferential flow paths in aquifers, rivers in the landscape, and atmospheric storm tracks that transport moisture in a preferential way. Connectivity is important as it controls whether flow paths connect and thus the overall behaviour of the system. While the controls on flood generation are similar at all scales, their relative magnitudes differ between scales.

365

370

Given all this process complexity, an obvious question is, how these scales can be linked, e.g. through upscaling. From the discussion it seems to be clear that processes *at all scales* affect flood generation at the continental scale in some way. Additionally, large scale processes possess emergent behaviour. They are not simply the multiple re-occurrence of pore scale processes. This is reminiscent of the very different behaviour of groups of people compared to that of individuals (studied respectively in sociology and psychology). Groups mainly behave differently because people interact, talk to each other. In the domain of hydrology, these interactions are feedbacks: feedbacks between processes (such as flow and structure), feedbacks between places (such as the subsurface and the surface), and feedbacks between scales. Often, a small number of people plays a disproportionately large role in influencing the group's opinion, e.g. influencers in social networks (Langner, 2013), and this phenomenon is similar to the effect of preferential flow on flood generation at higher scales. Without capturing preferential flow it will thus be difficult representing runoff generation across scales.

375

380

In modelling these processes, simplifications of the scale transitions are needed that reflect the respective structures (e.g. connectivity) and the boundary conditions (e.g. groundwater table). Upscaling by million-fold repetition of pore scale process will unlikely give the right patterns at the continental scale because of emergent behaviour. On the other hand, we can learn from observed patterns at all scales in a downward way. As Dunne (1998, p. 24) put it: "When formulating a particular problem, the scale must be chosen judiciously to maximize its utility. This challenges the belief that only progressively finer-scale studies are really 'scientific' and 'rigorous' and that 'scaling up' from some presumably

385





fundamental understanding is the only way to solve problems in hydrology.” It is hoped that scientific curiosity will push hydrologists further to learn about processes from observed patterns *at all scales*, to better understand how floods are generated as water moves from the raindrop to the ocean.

390

### Acknowledgements

This article is an outgrowth from the John Dalton Medal Lecture I gave at the General Assembly of the European Geosciences Union in April 2019. Over the years, my own ideas I have presented here, have benefitted immensely from interactions with numerous friends and collaborators which I would like to thank most cordially. These include the co-authors with which I have published the most: J. Parajka, A. Viglione, R. Merz, M. Sivapalan, J. Komma, R. Kirnbauer, A. Montanari, M. Rogger, R. B. Grayson, D. Gutknecht, J. L. Salinas, H. H. G. Savenije, G. Laaha, J. Szolgay, B. Merz, C. Reszler, A. W. Western, M. Borga, W. Wagner, G. Carr, T. Nester, J. O. Skoien, A. P. Blaschke, A. Castellarin, A. H. Farnleitner, S. Kohnová, E. Zehe, G. B. Chirico, G. DiBaldassarre, P. Strauss, M. Zessner, M. Bertola, A. Eder, K. Haslinger, M. Hofstätter, A. Kiss, D. Lun, B. Szeles, M. Vreugdenhil, J. Waser, B. Arheimer, T. Blume, X. Chen, P. Hogan, L. Kuil, R. A. P. Perdigão, W. Schöner, E. Szolgayová, S. Vorogushyn, C. Bucher, C. Cudennec, L. Gaál, J. Hall, D. Koutsoyiannis, P. Stadler, A. Bárdossy, H. Bormann, J. Derx, M. Exner-Kittridge, A. Fürnkranz-Prskawetz, H. V. Gupta, K. Hlavcová, J. Jansa, T. R. Kjeldsen, D. P. Loucks, H. Pirkel, E. Toth, T. Wagener, R. Woods, M. H. Barendrecht, L. Brandimarte, K. Breinl, A. Buttinger-Kreuzhuber, W. Buytaert, X. Cai, P. Claps, U. Ehret, H. Flühler, R. Fuchs, E. Gaume, A. Gelfan, E. Gröller, D. Gustafsson, P. Haas, B. Hansjürgens, S. Hasenauer, L. Holko, H. M. Holländer, H. Hölzel, Z. Horváth, M. Hrachowitz, G. Humer, B. Kohl, P. Kraft, H. Kreibich, U. Lall, L. R. Leung, N. Macdonald, D. S. Mackay, A. T. A. McMahon, L. Mediero, M. Michalak, F. Nobilis, M. Oismüller, T. Picciafuoco, L. Plavcová, H. Rajaram, G. H. Reischer, G. Sander, E. Sauquet, R. Sommer, V. Srinivasan, R. Tong, P. Valent, E. Yeshaneh, A. Alaoui, E. N. Anagnostou, H. Apel, S. A. Archfield, A. Becker, K. Beven, A. R. Blanch, D. Borchardt, R. Brázdil, M. Broer, S. Ceola, V. Chaplot, F. H. S. Chiew, M. Clark, J.-D. Creutin, A. Domeneghetti, M. Doubková, A. Dumitrescu, G. Glatzel, D. C. Goodrich, S. Harrigan, K. Hlavková, D. Hughes, G. Jewitt, D. Kollanur, Z. Kostka, Z. W. Kundzewicz, G. Kuschnig, C. Leibundgut, J. J. McDonnell, N. McIntyre, P. Miklánek, P. Molnar, V. Naeimi, L. Nemeth, M. Pfaundler, U. Piock-Ellena, H. Rechberger, H. Ribicic, D. Rosbjerg, B. Schindler, S. Schymanski, K. Scipal, R. Silasari, M. Šraj, H. Stadler, G. Stancalie, A. H. Thieken, S. E. Thompson, P. A. Troch, R. Vertessy, J. Vierheilig, D. J. Wilson, H. C. Winsemius

**Competing interests.** The author declares that he has no conflict of interest.



420

## References

- Alaoui, A., Rogger, M., Peth, S. and Blöschl, G.: Does soil compaction increase floods? A review. *Journal of Hydrology*, 557, 631-642, 2018. <https://doi.org/10.1016/j.jhydrol.2017.12.052>.
- Andersson, J., and Dverstorp, B.: Conditional simulations of fluid flow in three - dimensional networks of discrete fractures. 425 *Water Resources Research*, 23(10), 1876-1886, 1987.
- Bierkens, M. F., Bell, V. A., Burek, P., Chaney, N., Condon, L. E., David, C. H., ... and Wood, E. F.: Hyper - resolution global hydrological modelling: what is next? “Everywhere and locally relevant”. *Hydrological Processes*, 29(2), 310-320, 2015).
- Bierkens, M., Finke, P. and De Willigen, P.: *Upscaling and downscaling methods for environmental research*. Kluwer 430 Academic, 2000.
- Biron, P. M., Robson, C., Lapointe, M. F. and Gaskin, S. J.: Comparing different methods of bed shear stress estimates in simple and complex flow fields. *Earth Surface Processes and Landforms: The Journal of the British Geomorphological Research Group*, 29(11), 1403-1415, 2004.
- Blöschl, G. and Zehe, E.: On hydrological predictability. *Hydrological Processes*, 19 (19), pp. 3923-3929, 2005.
- Blöschl, G., Blaschke, A. P., Broer, M., Bucher, C., Carr, G., Chen, X., ... and Zessner, M.: The hydrological open air 435 laboratory (HOAL) in Petzenkirchen: A hypothesis-driven observatory. *Hydrology and Earth System Sciences*, 20(1), 227-255, 2016.
- Blöschl, G., Gaál, L., Hall, J., Kiss, A., Komma, J., Nester, T., ... and Viglione, A.: Increasing river floods: fiction or reality?. *Wiley Interdisciplinary Reviews: Water*, 2(4), 329-344, 2015.
- Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A., Merz, B., Arheimer, B., ... and Živković, N.: Changing climate shifts timing 440 of European floods. *Science*, 357(6351), 588-590, 2017.
- Blöschl, G., Nester, T., Komma, J., Parajka, J., & Perdigão, R. A.. The June 2013 flood in the Upper Danube Basin, and comparisons with the 2002, 1954 and 1899 floods. *Hydrology and Earth System Sciences*, 17(12), 5197-5212, 2013.
- Blöschl, G., Sivapalan, M., Gupta, V. K., Beven, K. and Lettenmaier, D.: Special section on scale problems in hydrology. *Water Resources Research*, 33(12), 2881-2999, 1997.
- 445 Blöschl, G.: Scaling in hydrology. *Hydrological Processes*, 15, 709-711, 2001.
- Deng, J., Yin, H., Kong, F., Chen, J., Dronova, I. and Pu, Y.: Determination of runoff response to variation in overland flow area by flow routes using UAV imagery. *Journal of Environmental Management*, 265, 109868, 2020.



- Dooge, J. C.: Looking for hydrologic laws. *Water Resources Research*, 22(9S), 46S-58S, 1986.
- Dunne, T. and Black, R. D.: An experimental investigation of runoff production in permeable soils. *Water Resources Research*, 6(2), 478-490, 1970.
- 450 Dunne, T.: Wolman Lecture: hydrologic science in landscapes on a planet in the future. *Hydrologic Sciences: Taking Stock and Looking Ahead*, 10-43, 1998.
- Flury, M., Flühler, H., Jury, W. A., and Leuenberger, J.: Susceptibility of soils to preferential flow of water: A field study. *Water Resources Research*, 30(7), 1945-1954, 1994.
- 455 Freeze, R. A. and Harlan, R. L.: Blueprint for a physically-based, digitally-simulated hydrologic response model. *Journal of Hydrology*, 9(3), 237-258, 1969.
- Gaál, L., Szolgay, J., Kohnová, S., Parajka, J., Merz, R., Viglione, A. and Blöschl, G.: Flood timescales: Understanding the interplay of climate and catchment processes through comparative hydrology. *Water Resources Research*, 48, W04511, 2012. doi:10.1029/2011WR011509.
- 460 Grayson, R. and Blöschl, G.: Spatial processes, organisation and patterns. In: *Spatial patterns in catchment hydrology*, edited by: Grayson, R. and Blöschl, G., Cambridge University Press, pp. 3-16, 2001.
- Grayson, R. B., Western, A. W., Chiew, F. H. and Blöschl, G.: Preferred states in spatial soil moisture patterns: local and non-local controls. *Water Resources Research*, 33 (12), 2897-2908, 1997.
- Gupta, V. K., Rodríguez-Iturbe I. and Wood, E. F.: Scale Problems in Hydrology, Runoff Generation and Basin Response. 465 D. Reidel Publ., 246 pp., 1986.
- Hofstätter, M. and Blöschl G.: Vb cyclones synchronized with the Arctic-/North Atlantic Oscillation. *Journal of Geophysical Research: Atmospheres*, 124, pp. 3259-3278, 2019. <https://doi.org/10.1029/2018JD029420>
- Hofstätter, M., Chimani, B., Lexer, A. and Blöschl, G.: A new classification scheme of European cyclone tracks with relevance to precipitation. *Water Resources Research*, 52, 2016. 7086–7104, doi:10.1002/2016WR019146.
- Hofstätter, M., Lexer, A., Homann, M. and Blöschl, G.: Large-scale heavy precipitation over central Europe and the role of atmospheric cyclone track types. *International Journal of Climatology*, 38, pp. e497–e517, 2018. <https://doi.org/10.1002/joc.5386>
- Horton R.E.: The role of infiltration in the hydrologic cycle. *Transactions, American Geophysical Union*, 14, 446–460, 1933.
- Jenny, H.: Factors of soil formation—a system of quantitative pedology. McGraw-Hill, New York, 281 pp., 1941.
- 470 Jury, W. A. and Horton, R.: *Soil physics*. John Wiley & Sons, 2004.
- Kalma, J. D. and Sivapalan, M.: *Scale issues in hydrological modelling*. John Wiley and Sons, 489 pp, 1995.



- Kemter, M., Merz, B., Marwan, N., Vorogushyn, S. and Blöschl, G.: Joint trends in flood magnitudes and spatial extents across Europe. *Geophysical Research Letters*, 47(7), 2020. e2020GL087464. <https://doi.org/10.1029/2020GL087464>
- Kolmogoroff, A.: Über die analytischen Methoden in der Wahrscheinlichkeitsrechnung. *Mathematische Annalen*, 104, 415-475, 1931.
- Langner, S., Hennigs, N. and Wiedmann, K.-P.: Social persuasion: targeting social identities through social influencers, *Journal of Consumer Marketing*, 30(1), 31-49, 2013. <https://doi.org/10.1108/07363761311290821>
- Merz R. and Blöschl, G.: A process typology of regional floods. *Water Resources Research*, 39 (12), article number 1340, 2003.
- 480 Merz, B., Aerts, J. C. J. H., Arnbjerg-Nielsen, K., Baldi, M., Becker, A., Bichet, A., ... and Nied, M.: Floods and climate: emerging perspectives for flood risk assessment and management. *Natural Hazards and Earth System Sciences*, 14(7), 1921-1942, 2014.
- Merz, B., Blöschl, G., Vorogushyn, S., Dottori, F., Aerts, J. C., Bates, P., ... and Macdonald, E.: Causes, impacts and patterns of disastrous river floods. *Nature Reviews Earth & Environment*, 2(9), 592-609, 2021.
- 485 Milne, G. Some suggested units of classification and mapping, particularly of East African soils. *Soil Research. Supplements to the Proceedings of the International Society of Soil Science*, (3), 183-198, 1935.
- Montanari, A. and Koutsoyiannis, D.: A blueprint for process-based modeling of uncertain hydrological systems, *Water Resour. Res.*, 48, W09555, 2012. doi:10.1029/2011WR011412.
- Nester, T., Komma, J. and Blöschl, G.: Real time flood forecasting in the Upper Danube basin. *Journal of Hydrology and Hydromechanics*, 64 (4), 404-414, 2015. doi: 10.1515/johh-2016-0033
- O'Neill, M. M. F., Tijerina, D. T., Condon, L. E. and Maxwell, R. M.: Assessment of the ParFlow-CLM CONUS 1.0 integrated hydrologic model: evaluation of hyper-resolution water balance components across the contiguous United States, *Geosci. Model Dev.*, 14, 7223-7254, 2021. <https://doi.org/10.5194/gmd-14-7223-2021>.
- 490 Penna, D., Tromp-van Meerveld, H. J., Gobbi, A., Borga, M. and Dalla Fontana, G. : The influence of soil moisture on threshold runoff generation processes in an alpine headwater catchment. *Hydrology and Earth System Sciences*, 15(3), 689-702, 2011.
- 495 Rajaram, H.: Debates - Stochastic subsurface hydrology from theory to practice: Introduction, *Water Resour. Res.*, 52, 9215-9217, 2016. doi:10.1002/2016WR020066.
- Richards, L. A.: Capillary conduction of liquids through porous mediums. *Physics*, 1(5), 318-333, 1931.
- Rodriguez-Iturbe, I. and Gupta, V.K.: Scale problems in hydrology. *Journal of Hydrology*, 65(1-3), 1-257, 1983.



- Savenije, H. H. and Hrachowitz, M.: HESS Opinions Catchments as meta-organisms—a new blueprint for hydrological  
500 modelling. *Hydrology and Earth System Sciences*, 21(2), 1107-1116, 2017.
- Silasari, R., Parajka, J., Ressler, C., Strauss, P. and Blöschl, G.: Potential of time - lapse photography for identifying saturation  
area dynamics on agricultural hillslopes. *Hydrological Processes*, 31(21), 3610-3627, 2017. doi: 10.1002/hyp.11272
- Sivapalan, M. and Blöschl, G.: The growth of hydrological understanding: Technologies, ideas, and societal needs shape the  
field. *Water Resources Research*, 53, 8137–8146, 2017. doi: 10.1002/2017WR021396
- 505 Sivapalan, M. and Blöschl, G.: Time scale interactions and the coevolution of humans and water. *Water Resour. Res.*, 51,  
6988–7022, 2015. doi:10.1002/2015WR017896.
- Sivapalan, M.: From engineering hydrology to Earth system science: milestones in the transformation of hydrologic science.  
*Hydrology and Earth System Sciences*, 22(3), 1665-1693, 2018.
- Sivapalan, M.: Process complexity at hillslope scale, process simplicity at the watershed scale: is there a connection?.  
510 *Hydrological Processes*, 17(5), 1037-1041, 2003.
- Smith, R. E., Smettem, K. R., Broadbridge, P., Woolhiser, D.A.: *Infiltration theory for hydrologic applications*. Water  
Resources Monograph Series Vol. 15., American Geophysical Union, 2002.
- Tarasova, L., Basso, S., Wendi, D., Viglione, A., Kumar, R. and Merz, R.: A process-based framework to characterize and  
classify runoff events: The event typology of Germany. *Water Resources Research*, 56(5), 2020. e2019WR026951.
- 515 Tarasova, L., Merz, R., Kiss, A., Basso, S., Blöschl, G., Merz, B., ... and Wietzke, L.: Causative classification of river flood  
events. *Wiley Interdisciplinary Reviews: Water*, 6(4), e1353, 2019.
- Tessier, D.: Behaviour and microstructure of clay minerals. In: *Soil colloids and their associations in aggregates*, pp. 387-  
415. Springer, Boston, MA, 1990.
- Thompson, S. E., Harman, C. J., Troch, P. A., Brooks, P. D. and Sivapalan, M.: Spatial scale dependence of  
520 ecohydrologically mediated water balance partitioning: A synthesis framework for catchment ecohydrology. *Water  
Resources Research*, 47(10), 2011.
- Tromp-van Meerveld, H. J. and McDonnell, J. J.: Threshold relations in subsurface stormflow: 2. The fill and spill  
hypothesis. *Water Resources Research*, 42(2), 2006.
- Tuller, M. and Or, D.: Hydraulic functions for swelling soils: pore scale considerations. *Journal of Hydrology*, 272(1-4), 50-  
525 71, 2003.
- van Bebbber, W. J.: Die Zugstrassen der barometrischen Minima nach den Bahnenkarten der Deutschen Seewarte für den  
Zeitraum 1875–1890, *Meteorol. Z.*, 8, 361–366, 1891.



- Viglione, A., Chirico, G. B., Woods, R. and Blöschl, G.: Generalised synthesis of space–time variability in flood response: An analytical framework. *Journal of Hydrology*, 394(1-2), 198-212, 2010.
- 530 Viglione, A., Merz, R., Salinas, J. L. and Blöschl, G.: Flood frequency hydrology: 3. A Bayesian analysis. *Water Resources Research*, 49(2), 675-692, 2013. doi:10.1029/2011WR010782
- Vorogushyn, S., Merz, B., Lindenschmidt, K. E. and Apel, H.: A new methodology for flood hazard assessment considering dike breaches. *Water Resources Research*, 46(8), 2010.
- Vreugdenhil, M., Szeles, B., Silasari, R., Salinas, J.L., Strauss, P. Oismueller, M., Hogan, P. Wagner, W. and Blöschl, G.:  
535 Nonlinear controls on event runoff generation in a small agricultural catchment. In preparation, 2022.
- Weinberg, G. M.: An Introduction to General Systems Thinking, Wiley-Interscience, New York, 1975.
- Western, A. W., Blöschl, G. and Grayson, R. B.: How well do indicator variograms capture the spatial connectivity of soil moisture?. *Hydrological Processes*, 12(12), 1851-1868, 1998.
- Western, A. W., Blöschl, G. and Grayson, R. B.: Toward capturing hydrologically significant connectivity in spatial patterns.  
540 *Water Resources Research*, 37(1), 83-97, 2001.
- Western, A. W., Grayson, R. B. and Blöschl, G.: Scaling of soil moisture: A hydrologic perspective. *Annual Review of Earth and Planetary Sciences*, 30(1), 149-180, 2002.
- Wood, E. F., Roundy, J. K., Troy, T. J., Van Beek, L. P. H., Bierkens, M. F., Blyth, E., ... and Whitehead, P.:  
545 Hyperresolution global land surface modeling: Meeting a grand challenge for monitoring Earth's terrestrial water. *Water Resources Research*, 47(5), 2011.
- Woods, R. and Sivapalan, M.: A synthesis of space - time variability in storm response: Rainfall, runoff generation, and routing. *Water Resources Research*, 35(8), 2469-2485, 1999.
- Zehe, E., Blume, T. and Blöschl, G. The principle of ‘maximum energy dissipation’: a novel thermodynamic perspective on rapid water flow in connected soil structures. *Philosophical Transactions of the Royal Society B: Biological Sciences*,  
550 365(1545), 1377-1386, 2010.
- Zehe, E., Elsenbeer, H., Lindenmaier, F., Schulz, K. and Blöschl, G.: Patterns of predictability in hydrological threshold systems. *Water Resources Research*, 43(7), article number W07434, 2007.