

Revisiting the Hydrological Basis of the Budyko Framework With the Hydrologically Similar Groups Principle

Yuchan Chen¹, Xiuzhi Chen¹, Meimei Xue¹, Chuanxun Yang^{2,3}, Wei Zheng¹, Jun Cao⁴, Wenting Yan¹, Wenping Yuan¹

5 ¹Guangdong Province Key Laboratory for Climate Change and Natural Disaster Studies, School of Atmospheric Sciences, Sun Yat-sen University, Zhuhai, 519082, China;

²Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, 510640, China;

³University of Chinese Academy of Sciences, Beijing, 100049, China;

⁴Guangdong provincial Academy of Environmental Science, Guangzhou, 510635, China;

10 *Correspondence to: Xiuzhi Chen (chenxzh73@mail.sysu.edu.cn)*

Abstract. The Budyko framework is a simple and effective tool for estimating the water balance of watershed. Quantification of the watershed-characteristic-related parameter (Pw) is critical for accurate water balance simulations with the Budyko framework. However, there is no universal method for calculating Pw as the interactions between hydrologic, climatic, and watershed characteristic factors differ greatly across watersheds. To fill this research gap, this study
15 introduced the hydrologically similar groups principle into the Budyko framework for quantifying the Pw of watersheds in similar environments. We first classified the selected 366 watersheds worldwide into six hydrologically similar groups based on watershed attributes, including climate, soil, and vegetation. Results show that soil moisture (SM) and fractional vegetation cover (FVC) are two controlling factors of the Pw in each group. The SM exhibits a power-law relationship with the Pw values, with increasing SM leading to higher Pw values in dry watersheds ($SM \leq 20$ mm) and lower Pw values
20 in humid watersheds ($SM > 20$ mm). Additionally, the FVC shows to be linearly correlated with the Pw values in most hydrologically similar groups, except in that group with moist soil and no strong rainfall seasonality ($SM > 20$ and $SI \leq 0.4$). Multiple non-linear regression models between Pw and the controlling factors (SM and FVC) were developed to individually estimate the Pw of six hydrologically similar groups. Cross-validations using the bootstrap sampling method ($R^2 = 0.63$) and validations of time-series Global Runoff Data Centre (GRDC) data ($R^2 = 0.89$) both indicate that the
25 proposed models perform satisfactorily in estimating the Pw parameter in the Budyko framework. Overall, this study is a new attempt to quantify the unknown watershed characteristic-related parameter in the Budyko framework using the hydrologically similar groups method. The results will be helpful in improving the applicability of the Budyko framework for estimating the annual runoff of watersheds in diverse climates and with different characteristics.

1 Introduction

30 There has been an increasing interest in estimating the water balance of watersheds with a simple and effective tool—the Budyko framework. Unlike process-based models that typically require a large number of parameters as inputs for accurate simulations (Caracciolo et al., 2018; Lei et al., 2014), the Budyko framework is a top-down approach that is rooted on a firm physical basis, relating a catchment’s long-term evaporative ratio (ratio between actual evapotranspiration and precipitation) to its aridity index (ratio between potential evapotranspiration and precipitation) (Vora and Singh, 2021; 35 Sivapalan, 2003; Wang and Tang, 2014). Currently, the Budyko framework has been widely used for assessing linkages and feedbacks between climate forcing and land surface characteristics on water and energy cycles (Zhang et al., 2001; Milly and Shmakin, 2002; Li et al., 2013; Xu et al., 2013), prompting a great deal of empirical, theoretical, and process-based studies (Chen and Sivapalan, 2020; Roderick and Farquhar, 2011; Rau et al., 2018; Goswami and Goyal, 2022).

The original Budyko equation assumes that evapotranspiration is mainly controlled by precipitation (representing the 40 availability of water) and potential evapotranspiration (representing the availability of energy) (Budyko, 1974; Wang et al., 2022). Despite its solid performance, the original Budyko equation still produces a bias between modeled and measured evapotranspiration or runoff because it does not consider the effects of watershed characteristics other than mean annual climatic conditions on water balance (Kim and Chun, 2021; Zhang et al., 2001). As a result, hydrologists have invested considerable efforts to improve model performance by introducing parameters related to watershed characteristics 45 (watershed-characteristic-related parameter, P_w) into the original Budyko equation. The popular parametric equations of the Budyko framework are presented in Table 1.

Table 1. Parametric formulations of the Budyko framework (P_w - watershed-characteristic-related parameter; ET - actual evaporation, R - runoff, P - precipitation, PET - potential evapotranspiration, all in mm yr⁻¹).

Reference	Formulation	P_w (Theoretical range)	Reference values of P_w
Budyko (1974)	$\frac{ET}{P} = \left[\frac{PET}{P} \tanh \left(\frac{PET}{P} \right)^{-1} \left(1 - \exp \left(-\frac{PET}{P} \right) \right) \right]^{0.5}$	0.5	0.5
Zhang et al. (2001)	$\frac{ET}{P} = \frac{1 + w \frac{PET}{P}}{1 + w \frac{PET}{P} + \left(\frac{PET}{P} \right)^{-1}}$	w (0, ∞)	Trees – 2.0, Plants – 0.5

Turc (1954), Mezentsev (1955), Choudhury (1999), Yang et al. (2008)	$\frac{ET}{P} = \frac{1}{\left[1 + \left(\frac{P}{PET}\right)^n\right]^{\frac{1}{n}}}$	n (0, ∞)	Field – 2.6, River basins – 1.8
Wang and Tang (2014)	$\frac{ET}{P} = \frac{1 + \frac{PET}{P} - \sqrt{\left(1 + \frac{PET}{P}\right)^2 - 4\varepsilon(2 - \varepsilon)\frac{PET}{P}}}{2\varepsilon(2 - \varepsilon)}$	ε (0,1)	0.55 - 0.58
Tixeront (1964), Fu (1981), Zhou et al. (2015)	$\frac{R}{P} = \left[1 + \left(\frac{P}{PET}\right)^{-m}\right]^{\frac{1}{m}} - \left(\frac{P}{PET}\right)^{-1}$	m (1, ∞)	Forest – 2.83, Shrub – 2.33, Grassland or cropland – 2.28, Mixed land – 2.12

50 From the hydrological point of view, P_w controls the fraction of precipitation diverted into runoff for a given aridity index (Caracciolo et al., 2018). Watersheds with higher P_w values partition more precipitation to evapotranspiration and consequently less to runoff than those with lower P_w values; some studies defined P_w as the water retention capacity of a watershed (Fu, 1981; Zhou et al., 2015). Overall, P_w denotes the adjustment of water-energy partitioning by various watershed characteristics (Yao et al., 2017; Li et al., 2013).

55 During the past decades, researchers have done lots of work to quantify P_w for the accurate simulation of evapotranspiration or runoff using the Budyko framework (Wang et al., 2022; Yao et al., 2017; Guo et al., 2019; Yu et al., 2021) and considerably improved the estimation of P_w by taking into account the influence of watershed characteristics (Fu, 1981; Liu and Liang, 2015; Guan et al., 2022; Yang et al., 2008). Although there is agreement that P_w represents the integrated effects of various environmental factors (Wang et al., 2022; Liu et al., 2022b; Yu et al., 2021; Gan et al., 2021), studies
60 still differed greatly as to what factors and effects should relate to P_w and failed to give a general framework for quantifying it. For instance, whether the P_w in the Budyko framework is controlled by vegetation or not has been much debated. Ning et al. (2017) found that P_w generally correlated positively with vegetation cover. Zhang et al. (2018) obtained the sensitivity of P_w to changes in LAI by taking a derivative of the P_w function with respect to LAI, implying a crucial role of vegetation cover in impacting P_w . However, other studies indicated that most regions or watersheds show no significant influences of
65 vegetation indices or cover on P_w (Li et al., 2013; Liu et al., 2021). For example, Li et al. (2013) noted that the variations in the P_w values are not entirely controlled by vegetation cover in small catchments. Another study by Liu et al. (2021) also found a weak correlation between the vegetation leaf area index and P_w . Therefore, more in-depth studies are needed for revisiting the hydrological basis of P_w in the Budyko framework.

Here, we hypothesize that watersheds with similar climatic, hydrologic, and watershed characteristics have consistent
70 controlling factors of Pw in the Budyko framework. Classifying watersheds into groups that are hydrologically similar may
help us identify how Pw responds to different watershed characteristic factors. However, to date, few studies have been
conducted on classifying watersheds based on the highly variable hydro-climate-Pw relationships in the Budyko framework.
This may be an important reason why researchers disagree about the factors and extent of the influence on Pw.

This study proposes a new approach to address the research gap in accurately estimating the Pw parameter in the
75 Budyko framework by classifying watersheds into hydrologically similar groups and developing a framework for
estimating Pw (PwM) in each group to simulate global runoff. More specifically, we collected 726 hydrological records in
366 watersheds from published literature for analyses. These 726 samples were classified into six hydrologically similar
groups according to the hydrologically homogenous attributes of watersheds using the Decision Tree Regressor method.
Then, we identified the controlling factors of Pw from various environmental factors in each hydrologically similar group
80 and developed multiple non-linear regression models for estimating Pw in the Budyko framework. We expect that
classifying watersheds into hydrologically similar groups can help explore the effect of watershed characteristics on their
water balance and interpret the physical meaning of the Pw in the Budyko framework. This study highlights the need to
account for the interactions among hydrologic, climatic, and watershed characteristic factors for explaining Pw in the
Budyko framework.

85 **2 Fu's formula**

This study employed Fu's formula (Zhou et al., 2015) to analyze Pw in the Budyko framework. Fu's equation is a
commonly used parametric equation in Budyko-type formulas due to its versatility and adaptability (Zhou et al., 2015).
The formula is expressed as:

$$\frac{R}{P} = \left(1 + \left(\frac{P}{PET} \right)^{-Pw} \right)^{\frac{1}{Pw}} - \left(\frac{P}{PET} \right)^{-1} \quad (1)$$

90 where R/P is a dimensionless annual water yield coefficient; P/PET is an aridity index; Pw is a dimensionless constant
varying from 1 to infinity and represents water retention capacity for evapotranspiration. When Pw=1, all the precipitation
becomes flow and the residence time is 0. When Pw tends to infinity, the runoff approaches to the difference between
precipitation and potential evapotranspiration. In this scenario, all precipitation remains in the watershed and all available
water is lost through evapotranspiration. The duration of water residence equals to the time for converting all precipitation

95 to evapotranspiration. However, in natural watersheds, it may be difficult to observe P_w approaching infinity since it is nearly impossible for all precipitation to be retained in the watershed. The natural watersheds with a high P_w value may be “non-conservative” (i.e., precipitation is not the sum of streamflow and evapotranspiration), as a portion of the water that remains in the watershed may not be solely from precipitation but may include groundwater flow and other difficult to measure flows. As a result, it may be challenging to accurately estimate the water balance, especially in regions with complex hydrological systems (De Lavenne and Andréassian, 2018; Goswami and O'connor, 2010). As a precautionary measure, this study sets an empirical upper limit of 10 for P_w to ensure that the watersheds in question remain conservative.

3 Data


3.1 Hydrological data

Hydrological data for modelling, including runoff and corresponding precipitation data, were collected from published literature (726 samples listed in Supplement 1, Fig. 1). Potential evapotranspiration data were downloaded from version 4.05 of the CRU TS (Climatic Research Unit gridded Time Series) climate dataset (<https://doi.org/10.6084/m9.figshare.11980500>), which is produced by the CRU at the University of East Anglia. For consistency, we used potential evapotranspiration values extracted from the CRU TS dataset of all watersheds listed in Supplement 1, even for studies with potential evapotranspiration values reported. The potential evapotranspiration values were extracted based on the coordinate points of watersheds. Using collected and extracted the annual average runoff, precipitation and potential evapotranspiration data for the observation period, we calculated the annual water yield coefficient (R/P) and aridity index (P/PET) for each sample. Then, we derive the annual average P_w value of each sample for the corresponding period according to Equation 1.

Observed river discharge data for validation were obtained from the Global Runoff Data Centre (GRDC, https://www.bafg.de/GRDC/EN/02_srvcs/21_tmsrs/riverdischarge_node.html). Only the GRDC stations meeting the following criteria were selected for further analysis: (1) The sites with continuous time-series runoff observations during the period 2000–2016 and corresponding surface soil moisture (SM), fractional vegetation cover (FVC) and seasonal index (SI) data were also available during such a period; (2) The drainage area reports can be found in the original data to provide area parameters for converting original flow volumes to runoff rates; (3) The geographical coordinates reports can be found in the original data and the shape of the drainage can be found in the GRDC Watershed Boundaries (2011); (4) The watersheds with “non-conservative” ($P_w > 10$) and unrealistic runoff rates ($P_w < 1$) are removed. Based on these criteria,

545 GRDC stations were selected for validation (Fig. 1). Then, the flow volumes of selected sites were converted to runoff rates (Ghiggi et al., 2019).

125 We used the GRDC Watershed Boundaries (2011) to extract the average values of potential evapotranspiration and precipitation from grid datasets for each watershed. The potential evapotranspiration values were extracted from the CRU TS dataset. The precipitation values for runoff reconstruction were extracted from the Global Precipitation Climatology Centre (GPCC) Precipitation Total Full V2018 data provided by the NOAA/OAR/ESRL (PSL, Boulder, Colorado, USA) because these were found to agree better with observations in previous research compared to the CRU TS precipitation dataset (Ahmed et al., 2019; Degefu et al., 2022; Fiedler and Döll, 2007; Hu et al., 2018; Salaudeen et al., 2021).



130 **Figure 1.** Location of observation sites for modeling (green dots) ($n = 726$) and GRDC (Global Runoff Data Centre) observation sites (orange triangles) ($n = 545$) for validation. Background colors represent UNEP (1997) climate classification for P/PET values (Hyper Arid: $P/PET < 0.03$; Arid: $0.03 \leq P/PET < 0.2$; Semi-Arid: $0.2 \leq P/PET < 0.5$; Dry sub-humid: $0.5 \leq P/PET < 0.65$; Humid: $P/PET \geq 0.65$). The globe was divided into nine geographic regions: North America (west, southwest, midwest, northeast, southeast, except of the USA), South America, Africa, and Europe. Due to the limited availability of GRDC observation data in Asia and Australia, these regions were absent in the division of global geographic regions.

135

3.2 Watershed characteristic-related data

140 The watershed characteristic-related factors mainly include SM (0-10 cm underground), FVC and SI of Walsh and Lawler (1981). For the collected watersheds from published literature without boundary files, these three datasets were extracted from grid data according to the coordinate points of these watersheds. For the GRDC watersheds, records of these three fields were extracted from grid data based on the boundary files provided by GRDC Watershed Boundaries (2011). The sources of datasets are summarized in Table 2.

Table 2. Data sources for watershed characteristic factors

Watershed characteristic factors	Data source/version	Units	Reference
Surface soil moisture (0-10cm underground, SM)	GLDAS Noah Land Surface Model L4	mm	Rodell et al. (2004)
Fractional vegetation cover (FVC)	GLASS FVC V4	m ² m ⁻²	Liang et al. (2021)
Seasonal index (SI)	CRU TS dataset version 4.03, global maps of seasonality indices	dimensionless	Walsh and Lawler (1981);Feng (2019)

145

4 Methods

4.1 Classification of watersheds into hydrologically similar groups using watershed attributes

150

A hydrologically similar group (i.e., hydrologically homogeneous region) is defined as a group of drainage basins whose hydrologic responses are similar (Kanishka and Eldho, 2020). Therefore, the relationship between Pw and any watershed characteristic variable does not change substantially in a hydrologically similar group. However, when that relationship between Pw and the variable changes as certain boundaries are crossed, the corresponding watersheds are divided into different groups by these boundaries.

155

We used SM, SI and FVC for classification. For SM and FVC, the bounded intervals of the variables were given by the Decision Tree Regressor (DTR) from the Scikit-learn library (Pedregosa et al., 2011) in Python. The locations of splits in DTR were used as dividing intervals. The criterion for measuring the quality of the split was set to “poisson”, which uses a reduction in Poisson deviance to find splits. The “random” strategy was used to choose local optimal splitting at each node. The results and performances of DTR are shown in Supplement 2. Based on the criteria used by Walsh and Lawler (1981), we divided the SI into three parts ($SI \leq 0.4$, $0.4 < SI \leq 0.8$, $SI > 0.8$) to represent three hydroclimatic seasonality (precipitation spread throughout the year, marked seasonality with a short drier season, extreme seasonality with a long drier season). Finally, six hydrologically similar groups were classified (Table 3).

160

Table 3. Classification of watersheds

Soil moisture classifier	Water soil regime	Seasonality index classifier	Seasonality precipitation regime	Fractional vegetation cover classifier	vegetation cover regime	Name of the group
SM \leq 20	Dry soil	—	—	—	—	IN _D
		SI \leq 0.4	Seasonless	—	—	IN _{WP}
SM $>$ 20	Wet soil	0.4 < SI \leq 0.8	Marked seasonality	FVC \leq 0.2 0.2 < FVC \leq 0.5 FVC > 0.5	Low density Middle density High density	IN _{WMS} IN _{WMM} IN _{WML}
		SI > 0.8	Extreme seasonality	—	—	IN _{WE}

4.2 Setup of proposed Pw simulation model (PwM)

4.2.1 PwM with the classification of hydrologically similar groups

We performed regression analysis between the Pw and watershed characteristic variables to determine the input variables of the PwM. The variables whose R^2 of the regression model was greater than 0.1 were selected as input variables. We used a polynomial as the basic model form. Each term of the polynomial depends on the regression model of the corresponding variable and the Pw. For each hydrological group, the Pw value is modeled as the function:

$$Pw = \sum Coef_n \times f(Var_n) \quad (2)$$

where Pw represents the value of Pw; Var_n represents the input variable that passes the regression test; f corresponds to the function derived from the regression of Pw on Var_n ; $Coef_n$ represents the empirical coefficient fitted by multiple non-linear regression (MNR).

4.2.2 PwM without classification of hydrologically similar groups

For comparison, we estimated Pw without the hydrologically similar groups, defined as non_PwM. The non_PwM was defined as follows:

$$non_Pw = a_1 \times SM^2 + a_2 \times SM + b_1 \times FVC^2 + b_2 \times FVC \quad (3)$$

where non_Pw is the annual value of Pw simulated by non_PwM ; SM is the annual average value of surface soil moisture (0-10 cm underground); FVC is the annual average value of fractional vegetation cover; a_1 , a_2 , b_1 and b_2 represent the empirical coefficients fitted by the least square method.

4.3 Model validation

180 4.3.1 Performance metrics

Three performance metrics were used to assess the accuracy of PwM. The variable N is the number of observations, i is the ith value to be simulated, and y_s and y_o are the simulated and observed series, respectively.

The relative bias (RelBIAS) represents systematic errors. A positive value indicates a general overestimation, while a negative one indicates an underestimation. The perfect agreement is achieved when RelBIAS equals zero. RelBIAS is
185 defined as:

$$RelBIAS = \frac{mean(y_s - y_o)}{mean(y_o)} \quad (4)$$

The coefficient of determination (R²) assesses the linear relationship between the simulated and observed time series data and is defined as:

$$R^2 = \frac{\sum_{i=1}^N (y_o^i - \bar{y}_o)(y_s^i - \bar{y}_s)}{[\sum_{i=1}^N (y_o^i - \bar{y}_o)^2]^{0.5} [\sum_{i=1}^N (y_s^i - \bar{y}_s)^2]^{0.5}} \quad (5)$$

190 The Nash–Sutcliffe efficiency (NSE) (Nash and Sutcliffe, 1970), a goodness-of-fit index, is usually used to assess the accuracy of the model. When NSE = 1, the model predictions perfectly match the observed data. A value higher than 0 indicates that the modeled mean is a good predictor compared to the observed value. It is defined as:

$$NSE = 1 - \frac{\sum_{i=1}^N (y_s^i - y_o^i)^2}{\sum_{i=1}^N (y_o^i - \bar{y}_o)^2} \quad (6)$$

4.3.2 Cross-validations using the bootstrap sampling method

195 We used cross-validation to test the stability of the proposed PwM using the bootstrap sampling method. The collected public data were split into two parts, one for model training and the other for model validation. A subset of 60% of the data was randomly selected using the bootstrap sampling method for training PwM. The remaining 40% of the data was used to evaluate the model performance using the validation metrics in Sect. 4.3.1. For each metric, the variable N is the number of test sets, i is the ith value to be simulated by the trained PwM, and y_s and y_o are the simulated and observed series of test
200 sets, respectively. The process was repeated randomly 10000 times. We documented the cross-validation result of each bootstrapping and showed them in the violin plot (Fig. 3).

4.3.3 Validations of GRDC time-series runoff reconstruction results

205 To further assess the model performance, we applied the proposed PwM to Fu's model to reconstruct the time-series runoff data of GRDC from 2000 to 2016. Finally, the time-series runoff data from 545 GRDC stations, which were selected by Sect. 3.1, were used to evaluate the model performance using the validation metrics in Sect. 4.3.1. For each metric, the terms y_s and y_o represent the simulated and observed time-series runoff data, respectively.

5 Results and discussion

5.1 The new proposed model for estimating Pw in Fu's formula

210 The regressions between Pw in Fu's formula and watershed characteristic variables collected from globally published datasets are shown in Fig. 2.




Figure 2. Regression between Pw in Fu’s formula and (a) SM ($SM \leq 20$ mm), (b) SM ($SM > 20$ mm), (c) FVC (IN_D), (d) FVC (IN_{WP}), (e) FVC (IN_{WMS}), (f) FVC (IN_{WMM}), (g) FVC (IN_{WML}), and (h) FVC (IN_{WE}). Symbol shapes indicate SM (dots) and FVC (squares).

As shown in Fig. 2a-b, the relationship between Pw and SM conforms to a power function, consistent with prior findings reported by Chen and Sivapalan (2020). The important finding here is that there is a critical soil moisture threshold at 20 mm that separates watersheds with two different water balances. In watersheds characterized by arid conditions ($SM \leq 20$ mm), as shown in Fig. 2a, the Pw values have an upward trend as SM values increase. On the other side, in watersheds characterized by humid conditions ($SM > 20$ mm), as shown in Fig. 2b, the Pw values exhibit a decreasing trend as SM

values increase. This is likely because transpiration usually increases as soil water increases in relatively dry conditions (Jiao et al., 2019; Bierhuizen, 1958; Wang et al., 2012; Yao et al., 2016; Schwarzel et al., 2020). However, once the soil moisture exceeds the threshold (20 mm in this study), the acceleration of transpiration from soil moisture slows down quickly (Havranek and Benecke, 1978; Verhoef and Egea, 2014; Metselaar and De Jong Van Lier, 2007). These findings are very in line with previous studies (Havranek and Benecke, 1978; Jiao et al., 2019; Cavanaugh et al., 2011; Ducharme et al., 1998), although the threshold of soil moisture varies slightly in these studies (e.g., 0.25 m³ m⁻³ in Ducharme et al. (1998), 0.10 m³ m⁻³ in Cavanaugh et al. (2011) and 0.20 m³ m⁻³ in Jiao et al. (2019)).

As shown in Fig. 2c-h, the FVC is linearly correlated with the Pw values of watersheds in most hydrologically similar groups but differs greatly between different groups. In dry watersheds (IN_D), the relationship between Pw and FVC followed a positive linear function (Fig. 2c). This finding is consistent with the majority view that vegetation transpiration increases (reflected by the increased Pw) with increasing vegetation cover in regions with insufficient soil moisture (Wang et al., 2012; Yao et al., 2016; Schwarzel et al., 2020). For those small and wet watersheds, vegetation-related factors are considered to be weakly correlated with Pw (Liu et al., 2021; Padrón et al., 2017; Yang et al., 2014). However, our study reveals a positive linear correlation between Pw and FVC in the IN_{WMS} (Fig. 2e) and IN_{WE} groups (Fig. 2h), whereas a negative linear correlation is observed in the IN_{WMM} (Fig. 2f) and IN_{WML} groups (Fig. 2g). Only in the IN_{WP} group, the relationship between Pw and FVC is not significant. These results indicate that the relationship between Pw and FVC may be stronger than what was previously believed, and this relationship varies across different groups characterized by specific combinations of FVC and SI. This confirms that climate, soil moisture, and vegetation cover are not independent factors affecting the water balance (Gan et al., 2021; Yang et al., 2009). Coupling vegetation with other catchment properties resulted in greater Pw variations (Gan et al., 2021).

Based on the results of the regression analysis illustrated in Fig. 2, the proposed PwM employs SM and FVC as input variables (i.e., Var_n) for all groups, except for the IN_{WP} group, for which FVC was not chosen. The formula in PwM for calculating the Pw is modeled as a sum of a power function of SM and a linear function of FVC, given by Equation 7:

$$Pw = \begin{cases} 0.91 \times SM^{0.38} + 1.48 \times FVC & (IN_D, SM \leq 20) \\ 28.72 \times SM^{-0.76} & (IN_{WP}, SM > 20, SI \leq 0.4) \\ 39.03 \times SM^{-0.96} + 11.82 \times FVC & (IN_{WMS}, SM > 20, 0.4 < SI \leq 0.8, FVC \leq 0.2) \\ 33.76 \times SM^{-0.71} - 1.47 \times FVC & (IN_{WMM}, SM > 20, 0.4 < SI \leq 0.8, 0.2 < FVC \leq 0.5) \\ 20.41 \times SM^{-0.42} - 4.221 \times FVC & (IN_{WML}, SM > 20, 0.4 < SI \leq 0.8, FVC > 0.5) \\ 3078 \times SM^{-2.43} + 3.53 \times FVC & (IN_{WE}, SM > 20, SI > 0.8) \end{cases} \quad (7)$$

where Pw is the annual value of Pw; SM is the annual average value of surface soil moisture (0-10cm underground); FVC is the annual average value of fractional vegetation cover.

245 **5.2 Cross-validations based on data collected from globally published literature**

The performances of PwM and non_PwM were cross-validated based on the data collected from globally published literature using the bootstrap sampling method (Fig. 3). On average, the ensemble RelBIAS of Pw simulated by PwM is slightly negative (Fig. 3a), indicating a weak tendency to underestimate the values of Pw, but with a maximum relative bias less than 0.1. The interquartile range of R^2 for PwM is from 0.35 to 0.40, with a median of 0.37. The scores of R^2 are higher than 0.3 in more than 95% of the bootstrap sampling events. The NSE skill scores show that in most bootstrap samplings, the estimation error estimated variance for PwM is less than the variance of the observations ($NSE > 0$), with an interquartile range from 0.33 to 0.39. In comparison, the maximum relative bias of the Pw simulated by the non_PwM is 0.12, the median of R^2 is 0.13, and the median of NSE is 0.13. Overall, cross-validations show that the performance of the PwM with the hydrologically similar groups is better and more stable than that of the non_PwM.

255 Grouping watersheds based on their hydrological similarities ensures that watersheds within the same category exhibit similar behaviors in settings with comparable climate, soil and vegetation characteristics (Kanishka and Eldho, 2017; Sinha et al., 2019). The model developed based on the principle of hydrologically similar groups considers the unique hydrological characteristics of different watersheds and can more accurately simulate the hydrological response in complex watershed systems (Santra et al., 2011; Jin et al., 2017; Kouwen et al., 1993; Gao et al., 2018; Kanishka and Eldho, 2017). As a comparison, in the non_PwM, all watersheds were lumped into a single category and showed a similar hydrological response to changes in watershed characteristics. That non_PwM, as the similar model used in previous studies (Zhang et al., 2018; Liang et al., 2015; Xu et al., 2013), may overlook and oversimplify the intricate interplay between climate, watershed characteristics and hydrology, thereby potentially resulting in less precise predictions of Pw across diverse watersheds.




Figure 3. Cross-validation results of (a) PwM and (b) non_PwM. A violin represents the distribution of the considered skill scores. The white dot on the violin plot represents the median. The black bar in the center of the violin represents the interquartile range. Colors distinguish three performance metrics: Red (RelBIAS), yellow (R^2) and blue (NSE).

270

The skill scores of cross-validations for the six groups are shown in Fig. 4. Though its overall RelBIAS is negative, PwM tends to overestimate values of Pw in the IN_{WP} group (the median of RelBIAS is positive). The IN_{WMS} group scores highest in R², with a median of 0.73, while the IN_{WP} group scores the lowest, with a median of 0.16. The grouped NSE scores show more uncertainty than the overall, especially in the IN_{WMS}: the lower adjacent value (LAV) larger than zero indicates more skill than the mean of observations; however, the outliers are far below zero. The low NSE value may be due to the low number of watersheds sampled in this interval, which increased the inconclusive results.

275




Figure 4. Cross-validation results of PwM for (a) IN_D, (b) IN_{WP}, (c) IN_{WMS}, (d) IN_{WMM}, (e) IN_{WML}, and (f) IN_{WE}.

280

Figure 5 shows the simulated R/P by PwM in comparison to site observations. The R² between the observed and the simulated values is 0.63 (Fig. 5a). The model performs well in humid regions with P/PET ≥ 1 in southeast America, Europe, middle China and southeast Australia. However, PwM likely underestimated the runoff in the arid (P/PET < 0.2) and semi-arid regions (0.2 ≤ P/PET < 0.5), mainly in western America and northwest China (Fig. 5b).




Figure 5. Simulated R/P using PwM in comparison with the observations collected from published literatures. (a) Scatter plots between R/P (yellow: simulation; green: observations) and P/PET; (b) Difference between simulated R/P from PmM and observations from the published datasets.

285

5.3 Validations of reconstructing the time-series GRDC runoff

For the selected 545 GRDC watersheds, the annual runoff estimated by PwM ranges from 229.84 to 320.34 mm, which is slightly lower than the observed range of GRDC (265.82 ~ 345.50 mm yr⁻¹) (Fig. 6a). Overall, the temporal evolution of runoff is captured well in the period 2000-2010. However, since 2011, the consistency between reconstructed runoff and GRDC runoff decreases, and the reconstruction results are consistently lower than the GRDC observations. The scatter plot between simulated and observed R/P also shows a slight underestimation of reconstructed global long-term mean runoff (Fig. 6b). The spatial patterns of long-term mean runoff reconstruction are shown in Fig. 6c-f. The estimated time-series runoff shows lower values in the west of the United States and south of Africa, and shows higher values in the northeastern United States and the European Mediterranean area, in comparison with the GRDC time series.

290

295





Figure 6. Time-series runoff reconstruction results in the selected GRDC stations. (a) Time-series annual mean runoff of the selected 545 GRDC watersheds; (b) Scatterplot between the modeled runoff and observed runoff; The spatial distribution of annual mean runoff in (c) North America, (d) South America, (e) Africa, and (f) Europe.

300

Figure 7 displays the skill scores of the reconstructed runoff by PwM in comparison with the GRDC ensemble from 2000-2016. It can be seen that, generally, the result of reconstruction by PwM is satisfactory, as indicated by the RelBIAS close to 0. The underestimation of runoff mainly occurs in the high mountains of the western United States (Fig. 7a), where the runoff is much smaller. Humid regions such as the northeastern United States and the European Mediterranean area have quite high R^2 values, while lower values are observed in the semi-arid ($0.2 \leq P/PET < 0.5$) and the dry sub-humid ($0.5 \leq P/PET < 0.65$) regions, which are mainly located in the western and midwestern United States (Fig. 7e-h). There are low NSE scores in the watersheds where runoff is unusually under-estimated or over-estimated (Fig. 7i-l), especially in the western United States.


305



310 **Figure 7.** Spatial distribution of the skill scores of the reconstructed time-series runoff.

We classified the GRDC data into nine geographic regions (Fig. 1) and further evaluated the performance of PwM in each sub-region individually. In general, the simulated time-series runoff is consistent with the time-series observations (Fig. 8-9), except in the western United States, where runoff was consistently underestimated (Fig. 8a). Spatially, there is an underestimation of runoff in sub-regions like the western United States (Fig. 8a) and high latitudes in North America (Fig. 8f). The runoff underestimation is more severe in the arid areas of the western United States (Fig. 9a) than in the relatively wet areas of northwest North America (Fig. 9f). The reconstructed time-series runoff in the Milk River watershed (GRDC station number: 4220501) and Near Lethbridge watershed (GRDC station number: 4213111) both show an underestimation of annual runoff in arid areas. The Milk River and Near Lethbridge are two adjacent watersheds with similar drainage areas located on the border of the United States and Canada. However, the underestimation is more serious in the Milk River watershed (RelBIAS=-0.32, annual mean P/PET=0.52) than in the Near Lethbridge watershed (RelBIAS=-0.27, annual mean P/PET=0.55). Interestingly, the spatial pattern of runoff underestimation almost coincides with that of the glaciers. Therefore, we considered that glacial meltwater might be the probable cause of runoff underestimation in glacier-covered areas (Li et al., 2021), where glacial snowmelt plays a more important role as a water input in arid regions than in wet ones. Therefore, the underestimation of runoff in the western United States is greater than

325 in northwest North America. Temporally, the runoff was mostly underestimated by PwM in the year 2011, when the world experienced abnormally high temperatures (Frölicher et al., 2018; NOAA/NOAA, 2011) and glacier melting was thus accelerated and increased runoff (Du et al., 2022; Liu et al., 2022a).



330 **Figure 8.** Observed time-series runoff versus reconstructed time-series runoff. Nine geographic sub-regions were in Fig. 1: North America ((a) west, (b) southwest, (c) midwest, (d) northeast, (e) southeast, (f) except of the USA), (g) South America, (h) Africa, and (i) Europe.




Figure 9. Scatterplots between observed annual mean runoff and reconstructed annual mean runoff. Nine geographic sub-regions were in Fig. 1: North America ((a) west, (b) southwest, (c) midwest, (d) northeast, (e) southeast, (f) except of the USA), (g) South America, (h) Africa, and (i) Europe.

335

340

In this paper, we selected the new Fu's equation and developed a universal framework for estimating Pw. Our results show that, to a large extent, the Pw in Budyko equation can be well estimated by the PwM using only soil moisture and fractional vegetation cover parameters. This indicates that soil moisture and fractional vegetation cover strongly control the water balance of watersheds (Gan et al., 2021; Chen and Sivapalan, 2020; Yang et al., 2009; Wang et al., 2021). The better performance of PwM than non_PwM supports our hypothesis that watersheds with similar climatic, hydrologic, and watershed-related characteristics have consistent controlling factors of Pw in the Budyko Framework, and suggest that the classification of watersheds can reduce uncertainty and improve the accuracy of Pw and runoff predictions.

345 **6 Conclusion**

This study developed a new framework for estimating the Pw in the Budyko framework for watersheds in similar environments based on the principle of hydrologically similar groups. The proposed method not only represented runoff observations in 366 watersheds from published literature but also reconstructed the time-series runoff in 545 GRDC stations. The findings indicated that Pw is closely related to SM and FVC, and the relationship varies across specific hydrologically similar groups. However, due to the complexity of hydrological processes, the new framework could not fully account for the impacts of all other factors, which might result in an underestimation of runoff in regions with glaciers or under climates with temperature anomalies. Overall, our findings lay a sound basis for estimating Pw in the Budyko framework, provide references for calibrating hydrological models, and will be helpful in improving global runoff estimations.

355 *Code availability.* The pieces of code that were used for all analyses are available from the authors upon request.

Data availability. All data used in this study are publicly available. Potential evapotranspiration data are available from CRU TS (<https://doi.org/10.6084/m9.figshare.11980500>), precipitation data used to model validation are available from GPCC (<https://psl.noaa.gov/data/gridded/data.gpcc.html>), observed river discharge data are available from GRDC (https://www.bafg.de/GRDC/EN/02_srvcs/21_tmsrs/riverdischarge_node.html), SM data are available from GLDAS (https://disc.gsfc.nasa.gov/datasets/GLDAS_NOAH025_M_2.1/summary?keywords=GLDAS), FVC data are available from GLASS (<http://www.glass.umd.edu/05D/FVC/>), and SI data are available from HydroShare (<http://www.hydroshare.org/resource/ff287c90c9e947a78e351c8d07d9d3f3>)

Author contributions. XC designed the study and proposed the scientific hypothesis. YC implemented the experiments, conducted the analysis and wrote the paper. MX helped with data collection and checked the technical adequacy of the experiments. CY and WZ helped with data processing. WPY provided guidance on the SI. CY, WZ, CJ, WTY and WPY reviewed and edited the manuscript. XC oversaw the study and conducted manuscript revision as a mentor.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Financial support. This study was financed by the National Natural Science Foundation of China (grant numbers 31971458, 41971275), the Innovation Group Project of Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), grant number 311021009 and the Special High-level Plan Project of Guangdong Province (grant number 2016TQ03Z354).

References

- Ahmed, K., Shahid, S., Wang, X., Nawaz, N., and Khan, N.: Evaluation of gridded precipitation datasets over arid regions of Pakistan, *Water*, 11, 210, <https://doi.org/10.3390/w11020210>, 2019.
- Bierhuizen, J.: Some observations on the relation between transpiration and soil moisture, *Netherlands Journal of Agricultural Science*, 6, 94-98, <https://doi.org/10.18174/njas.v6i2.17713>, 1958.
- Budyko, M. I.: *Climate and life*, Academic press 1974.
- Caracciolo, D., Pumo, D., and Viola, F.: Budyko's based method for annual runoff characterization across different climatic areas: an application to United States, *Water Resources Management*, 32, 3189-3202, <https://doi.org/10.1007/s11269-018-1984-7>, 2018.
- Cavanaugh, M. L., Kurc, S. A., and Scott, R. L.: Evapotranspiration partitioning in semiarid shrubland ecosystems: a two-site evaluation of soil moisture control on transpiration, *Ecohydrology*, 4, 671-681, <https://doi.org/10.1002/eco.157>, 2011.
- Chen, X. and Sivapalan, M.: Hydrological basis of the Budyko curve: Data-guided exploration of the mediating role of soil moisture, *Water Resources Research*, 56, e2020WR028221, <https://doi.org/10.1029/2020WR028221>, 2020.
- Choudhury, B.: Evaluation of an empirical equation for annual evaporation using field observations and results from a biophysical model, *Journal of Hydrology*, 216, 99-110, [https://doi.org/10.1016/S0022-1694\(98\)00293-5](https://doi.org/10.1016/S0022-1694(98)00293-5), 1999.
- de Lavenne, A. and Andréassian, V.: Impact of climate seasonality on catchment yield: A parameterization for commonly-used water balance formulas, *Journal of Hydrology*, 558, 266-274, <https://doi.org/10.1016/j.jhydrol.2018.01.009>, 2018.
- Degefu, M. A., Bewket, W., and Amha, Y.: Evaluating performance of 20 global and quasi-global precipitation products in representing drought events in Ethiopia I: Visual and correlation analysis, *Weather and Climate Extremes*, 35, 100416, <https://doi.org/10.1016/j.wace.2022.100416>, 2022.
- Du, X., Silwal, G., and Faramarzi, M.: Investigating the impacts of glacier melt on stream temperature in a cold-region watershed: Coupling a glacier melt model with a hydrological model, *Journal of Hydrology*, 605, 127303, <https://doi.org/10.1016/j.jhydrol.2021.127303>, 2022.
- Ducharne, A., Laval, K., and Polcher, J.: Sensitivity of the hydrological cycle to the parametrization of soil hydrology in a GCM, *Climate dynamics*, 14, 307-327, <https://doi.org/10.1007/s003820050226>, 1998.
- Feng, X.: Global maps of seasonality indices, HydroShare [dataset], 2019.
- Fiedler, K. and Döll, P.: Global modelling of continental water storage changes—sensitivity to different climate data sets, *Advances in Geosciences*, 11, 63-68, <https://doi.org/10.5194/adgeo-11-63-2007>, 2007.
- Frölicher, T. L., Fischer, E. M., and Gruber, N.: Marine heatwaves under global warming, *Nature*, 560, 360-364, <https://doi.org/10.1038/s41586-018-0383-9>, 2018.
- Fu, B.: On the calculation of the evaporation from land surface, *Chinese Journal of Atmospheric Sciences*, 5, 23-31, <https://doi.org/10.3878/j.issn.1006-9895.1981.01.03>, 1981.
- Gan, G., Liu, Y., and Sun, G.: Understanding interactions among climate, water, and vegetation with the Budyko framework, *Earth-Science Reviews*, 212, 103451, <https://doi.org/10.1016/j.earscirev.2020.103451>, 2021.

- Gao, M., Chen, X., Liu, J., and Zhang, Z.: Regionalization of annual runoff characteristics and its indication of co-dependence among hydro-climate-landscape factors in Jinghe River Basin, China, *Stochastic Environmental Research and Risk Assessment*, 32, 1613-1630, [10.1007/s00477-017-1494-9](https://doi.org/10.1007/s00477-017-1494-9), 2018.
- 410 Ghiggi, G., Humphrey, V., Seneviratne, S. I., and Gudmundsson, L.: GRUN: an observation-based global gridded runoff dataset from 1902 to 2014, *Earth System Science Data*, 11, 1655-1674, <https://doi.org/10.5194/essd-11-1655-2019>, 2019.
- Goswami, M. and O'Connor, K. M.: A “monster” that made the SMAR conceptual model “right for the wrong reasons”, *Hydrological Sciences Journal*, 55, 913-927, <https://doi.org/10.1080/02626667.2010.505170>, 2010.
- 415 Goswami, U. P. and Goyal, M. K.: Relative Contribution of Climate Variables on Long-Term Runoff Using Budyko Framework, in: *Water Resources Management and Sustainability*, Springer, 147-159, https://doi.org/10.1007/978-981-16-6573-8_7, 2022.
- GRDC: Watershed Boundaries of GRDC Stations / Global Runoff Data Centre, Federal Institute of Hydrology (BfG) [dataset], 2011.
- 420 Guan, X., Zhang, J., Yang, Q., and Wang, G.: Quantifying the effects of climate and watershed structure changes on runoff variations in the Tao River basin by using three different methods under the Budyko framework, *Theoretical and Applied Climatology*, 1-14, <https://doi.org/10.1007/s00704-021-03894-5>, 2022.
- Guo, A., Chang, J., Wang, Y., Huang, Q., Guo, Z., and Li, Y.: Uncertainty analysis of water availability assessment through the Budyko framework, *Journal of Hydrology*, 576, 396-407, <https://doi.org/10.1016/j.jhydrol.2019.06.033>, 2019.
- 425 Havranek, W. M. and Benecke, U.: The influence of soil moisture on water potential, transpiration and photosynthesis of conifer seedlings, *Plant and Soil*, 49, 91-103, <https://doi.org/10.1007/BF02149911>, 1978.
- Hu, Z., Zhou, Q., Chen, X., Li, J., Li, Q., Chen, D., Liu, W., and Yin, G.: Evaluation of three global gridded precipitation data sets in central Asia based on rain gauge observations, *International Journal of Climatology*, 38, 3475-3493, <https://doi.org/10.1002/joc.5510>, 2018.
- 430 Jiao, L., Lu, N., Fang, W., Li, Z., Wang, J., and Jin, Z.: Determining the independent impact of soil water on forest transpiration: a case study of a black locust plantation in the Loess Plateau, China, *Journal of Hydrology*, 572, 671-681, <https://doi.org/10.1016/j.jhydrol.2019.03.045>, 2019.
- Jin, Y., Liu, J., Lin, L., Wang, A., and Chen, X.: Exploring hydrologically similar catchments in terms of the physical characteristics of upstream regions, *Hydrology Research*, 49, 1467-1483, <https://doi.org/10.2166/nh.2017.191>, 2017.
- 435 Kanishka, G. and Eldho, T.: Watershed classification using isomap technique and hydrometeorological attributes, *Journal of Hydrologic Engineering*, 22, 04017040, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0001562](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001562), 2017.
- Kanishka, G. and Eldho, T.: Streamflow estimation in ungauged basins using watershed classification and regionalization techniques, *Journal of Earth System Science*, 129, 1-18, <https://doi.org/10.1007/s12040-020-01451-8>, 2020.
- 440 Kim, D. and Chun, J. A.: Revisiting a Two-Parameter Budyko Equation With the Complementary Evaporation Principle for Proper Consideration of Surface Energy Balance, *Water Resources Research*, 57, e2021WR030838, <https://doi.org/10.1029/2021WR030838>, 2021.
- Kouwen, N., Soulis, E. D., Pietroniro, A., Donald, J., and Harrington, R. A.: Grouped Response Units for Distributed Hydrologic Modeling, *Journal of Water Resources Planning and Management*, 119, 289-305, [https://doi.org/10.1061/\(ASCE\)0733-9496\(1993\)119:3\(289\)](https://doi.org/10.1061/(ASCE)0733-9496(1993)119:3(289)), 1993.
- 445 Lei, H., Yang, D., and Huang, M.: Impacts of climate change and vegetation dynamics on runoff in the mountainous region of the Haihe River basin in the past five decades, *Journal of Hydrology*, 511, 786-799, <https://doi.org/10.1016/j.jhydrol.2014.02.029>, 2014.

- Li, D., Pan, M., Cong, Z., Zhang, L., and Wood, E.: Vegetation control on water and energy balance within the Budyko framework, *Water Resources Research*, 49, 969-976, <https://doi.org/10.1002/wrcr.20107>, 2013.
- 450 Li, Y., Li, F., Shangguan, D., and Ding, Y.: A new global gridded glacier dataset based on the Randolph Glacier Inventory version 6.0, *Journal of Glaciology*, 67, 773-776, <https://doi.org/10.1017/jog.2021.28>, 2021.
- Liang, S., Cheng, J., Jia, K., Jiang, B., Liu, Q., Xiao, Z., Yao, Y., Yuan, W., Zhang, X., and Zhao, X.: The global land surface satellite (GLASS) product suite, *Bulletin of the American Meteorological Society*, 102, E323-E337, <https://doi.org/10.1175/BAMS-D-18-0341.1>, 2021.
- 455 Liang, W., Bai, D., Wang, F., Fu, B., Yan, J., Wang, S., Yang, Y., Long, D., and Feng, M.: Quantifying the impacts of climate change and ecological restoration on streamflow changes based on a Budyko hydrological model in China's Loess Plateau, *Water Resources Research*, 51, 6500-6519, <https://doi.org/10.1002/2014WR016589>, 2015.
- Liu, J., You, Y., Zhang, Q., and Gu, X.: Attribution of streamflow changes across the globe based on the Budyko framework, *Science of The Total Environment*, 794, 148662, <https://doi.org/10.1016/j.scitotenv.2021.148662>, 2021.
- 460 Liu, J., Long, A., Deng, X., Yin, Z., Deng, M., An, Q., Gu, X., Li, S., and Liu, G.: The Impact of Climate Change on Hydrological Processes of the Glacierized Watershed and Projections, *Remote Sensing*, 14, 1314, <https://doi.org/10.3390/rs14061314>, 2022a.
- Liu, Q. and Liang, L.: Impacts of climate change on the water balance of a large nonhumid natural basin in China, *Theoretical and Applied Climatology*, 121, 489-497, <https://doi.org/10.1007/s00704-014-1255-3>, 2015.
- 465 Liu, S., Wang, X., Zhang, L., Kong, W., Gao, H., and Xiao, C.: Effect of glaciers on the annual catchment water balance within Budyko framework, *Advances in Climate Change Research*, 13, 51-62, <https://doi.org/10.1016/j.accre.2021.10.004>, 2022b.
- Metselaar, K. and de Jong van Lier, Q.: The shape of the transpiration reduction function under plant water stress, *Vadose Zone Journal*, 6, 124-139, <https://doi.org/10.2136/vzj2006.0086>, 2007.
- 470 Mezentsev, V.: Back to the computation of total evaporation, *Meteorologia i Hidrologia*, 5, 24-26, 1955.
- Milly, P. and Shmakin, A.: Global modeling of land water and energy balances. Part II: Land-characteristic contributions to spatial variability, *Journal of Hydrometeorology*, 3, 301-310, [https://doi.org/10.1175/1525-7541\(2002\)003<0301:Gmolwa>2.0.Co;2](https://doi.org/10.1175/1525-7541(2002)003<0301:Gmolwa>2.0.Co;2), 2002.
- Nash, J. E. and Sutcliffe, J. V.: River flow forecasting through conceptual models part I—A discussion of principles, *Journal of hydrology*, 10, 282-290, [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6), 1970.
- 475 Ning, T., Li, Z., and Liu, W.: Vegetation dynamics and climate seasonality jointly control the interannual catchment water balance in the Loess Plateau under the Budyko framework, *Hydrology and Earth System Sciences*, 21, 1515-1526, <https://doi.org/10.5194/hess-21-1515-2017>, 2017.
- NOAANCEI: Monthly National Climate Report for July 2011, NOAA National Centers for Environmental Information, 2011.
- 480 Padrón, R. S., Gudmundsson, L., Greve, P., and Seneviratne, S. I.: Large-scale controls of the surface water balance over land: Insights from a systematic review and meta-analysis, *Water Resources Research*, 53, 9659-9678, <https://doi.org/10.1002/2017WR021215>, 2017.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., and Dubourg, V.: Scikit-learn: Machine learning in Python, the *Journal of machine Learning research*, 12, 2825-2830, 2011.
- 485 Rau, P., Bourrel, L., Labat, D., Frappart, F., Ruelland, D., Lavado, W., Dewitte, B., and Felipe, O.: Hydroclimatic change disparity of Peruvian Pacific drainage catchments, *Theoretical and applied climatology*, 134, 139-153, <https://doi.org/10.1007/s00704-017-2263-x>, 2018.

- 490 Rodell, M., Houser, P., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C.-J., Arsenault, K., Cosgrove, B., Radakovich, J.,
and Bosilovich, M.: The global land data assimilation system, *Bulletin of the American Meteorological society*, 85,
381-394, <https://doi.org/10.1175/BAMS-85-3-381>, 2004.
- Roderick, M. L. and Farquhar, G. D.: A simple framework for relating variations in runoff to variations in climatic
conditions and catchment properties, *Water Resources Research*, 47, <https://doi.org/10.1029/2010wr009826>, 2011.
- 495 Salaudeen, A., Ismail, A., Adeogun, B. K., Ajibike, M. A., and Zubairu, I.: Evaluation of ground-based, daily, gridded
precipitation products for Upper Benue River basin, Nigeria, *Engineering and Applied Science Research*, 48, 397-
405, <https://doi.org/10.14456/easr.2021.42>, 2021.
- Santra, P., Das, B. S., and Chakravarty, D.: Delineation of hydrologically similar units in a watershed based on fuzzy
classification of soil hydraulic properties, *Hydrological Processes*, 25, 64-79, <https://doi.org/10.1002/hyp.7820>, 2011.
- 500 Schwarzel, K., Zhang, L., Montanarella, L., Wang, Y., and Sun, G.: How afforestation affects the water cycle in drylands:
A process-based comparative analysis, *Global Change Biology*, 26, 944-959, <https://doi.org/10.1111/gcb.14875>, 2020.
- Sinha, J., Jha, S., and Goyal, M. K.: Influences of watershed characteristics on long-term annual and intra-annual water
balances over India, *Journal of Hydrology*, 577, 123970, <https://doi.org/10.1016/j.jhydrol.2019.123970>, 2019.
- Sivapalan, M.: Process complexity at hillslope scale, process simplicity at the watershed scale: is there a connection?,
505 *Hydrological Processes*, 17, 1037-1041, <https://doi.org/10.1002/hyp.5109>, 2003.
- Tixeront, J.: Prediction of streamflow, IAHS publication, 118-126, 1964.
- Turc, L.: The water balance of soils: relation between precipitation, evaporation and flow, *Ann. Agron*, 5, 491-569, 1954.
- UNEP: World atlas of desertification, United Nations Environment Programme [dataset], 1997.
- Verhoef, A. and Egea, G.: Modeling plant transpiration under limited soil water: Comparison of different plant and soil
510 hydraulic parameterizations and preliminary implications for their use in land surface models, *Agricultural and Forest
Meteorology*, 191, 22-32, <https://doi.org/10.1016/j.agrformet.2014.02.009>, 2014.
- Vora, A. and Singh, R.: Satellite based Budyko framework reveals the human imprint on long-term surface water
partitioning across India, *Journal of Hydrology*, 602, 126770, <https://doi.org/10.1016/j.jhydrol.2021.126770>, 2021.
- Walsh, R. and Lawler, D.: Rainfall seasonality: description, spatial patterns and change through time, *Weather*, 36, 201-
515 208, <https://doi.org/10.1002/j.1477-8696.1981.tb05400.x>, 1981.
- Wang, D. and Tang, Y.: A one-parameter Budyko model for water balance captures emergent behavior in Darwinian
hydrologic models, *Geophysical Research Letters*, 41, 4569-4577, <https://doi.org/10.1002/2014gl060509>, 2014.
- Wang, F., Xia, J., Zou, L., Zhan, C., and Liang, W.: Estimation of time-varying parameter in Budyko framework using
long short-term memory network over the Loess Plateau, China, *Journal of Hydrology*, 607, 127571,
520 <https://doi.org/10.1016/j.jhydrol.2022.127571>, 2022.
- Wang, H., Lv, X., and Zhang, M.: Sensitivity and attribution analysis of vegetation changes on evapotranspiration with the
Budyko framework in the Baiyangdian catchment, China, *Ecological Indicators*, 120, 106963,
<https://doi.org/10.1016/j.ecolind.2020.106963>, 2021.
- Wang, Y., Bredemeier, M., Bonell, M., Yu, P., Feger, K.-H., Xiong, W., and Xu, L.: Comparison between a statistical
525 approach and paired catchment study in estimating water yield response to afforestation, *Revisiting Experimental
Catchment Studies in Forest Hydrology:(Proceedings of a Workshop held during the XXV IUGG General Assembly
in Melbourne, June–July 2011, 3-11,*
- Xu, X., Liu, W., Scanlon, B. R., Zhang, L., and Pan, M.: Local and global factors controlling water-energy balances within
the Budyko framework, *Geophysical Research Letters*, 40, 6123-6129, <https://doi.org/10.1002/2013gl058324>, 2013.

- 530 Yang, D., Shao, W., Yeh, P. J. F., Yang, H., Kanae, S., and Oki, T.: Impact of vegetation coverage on regional water balance in the nonhumid regions of China, *Water Resources Research*, 45, <https://doi.org/10.1029/2008WR006948>, 2009.
- Yang, H., Yang, D., Lei, Z., and Sun, F.: New analytical derivation of the mean annual water-energy balance equation, *Water resources research*, 44, <https://doi.org/10.1029/2007wr006135>, 2008.
- 535 Yang, H., Qi, J., Xu, X., Yang, D., and Lv, H.: The regional variation in climate elasticity and climate contribution to runoff across China, *Journal of hydrology*, 517, 607-616, <https://doi.org/10.1016/j.jhydrol.2014.05.062>, 2014.
- Yao, J., Mao, W., Yang, Q., Xu, X., and Liu, Z.: Annual actual evapotranspiration in inland river catchments of China based on the Budyko framework, *Stochastic Environmental Research and Risk Assessment*, 31, 1409-1421, <https://doi.org/10.1007/s00477-016-1271-1>, 2017.
- 540 Yao, W., Xiao, P., Shen, Z., Wang, J., and Jiao, P.: Analysis of the contribution of multiple factors to the recent decrease in discharge and sediment yield in the Yellow River Basin, China, *Journal of Geographical Sciences*, 26, 1289-1304, <https://doi.org/10.1007/s11442-016-1227-7>, 2016.
- Yu, K.-x., Zhang, X., Xu, B., Li, P., Zhang, X., Li, Z., and Zhao, Y.: Evaluating the impact of ecological construction measures on water balance in the Loess Plateau region of China within the Budyko framework, *Journal of Hydrology*, 545, 601, 126596, <https://doi.org/10.1016/j.jhydrol.2021.126596>, 2021.
- Zhang, L., Dawes, W., and Walker, G.: Response of mean annual evapotranspiration to vegetation changes at catchment scale, *Water resources research*, 37, 701-708, <https://doi.org/10.1029/2000WR900325>, 2001.
- Zhang, S., Yang, Y., McVicar, T. R., and Yang, D.: An analytical solution for the impact of vegetation changes on hydrological partitioning within the Budyko framework, *Water Resources Research*, 54, 519-537, 550 <https://doi.org/10.1002/2017wr022028>, 2018.
- Zhou, G., Wei, X., Chen, X., Zhou, P., Liu, X., Xiao, Y., Sun, G., Scott, D. F., Zhou, S., and Han, L.: Global pattern for the effect of climate and land cover on water yield, *Nature communications*, 6, 1-9, <https://doi.org/10.1038/ncomms6918>, 2015.