

Sources of skill in lake temperature, discharge and ice-off seasonal forecasting tools

François Clayer¹, Leah Jackson-Blake¹, Daniel Mercado^{2,3}, Muhammed Shikhani⁴, Andrew French⁵,
Tadhg Moore^{6‡}, James Sample¹, Magnus Norling¹, Maria-Dolores Frias⁷, Sixto Herrera⁷, Elvira de Eyto⁵,
Eleanor Jennings⁶, Karsten Rinke⁴, Leon van der Linden⁸, Rafael Marcé^{2,3}.

¹Norwegian Institute for Water Research (NIVA), Oslo, Norway

²Catalan Institute for Water Research (ICRA), Girona, Spain

³Universitat de Girona, Girona, Spain

⁴Department of Lake Research, Helmholtz Centre for Environmental Research, Magdeburg, Germany

⁵Foras na Mara - Marine Institute, Furnace, Newport, Co. Mayo, Ireland

⁶Dundalk Institute of Technology, Dundalk, Co. Louth, Ireland

⁷Grupo de Meteorología. Dpto. de Matemática Aplicada y Ciencias de la Computación. Universidad de Cantabria, Santander, Spain

⁸SA Water, Adelaide SA 5000, Australia

15 *Correspondence to:* François Clayer (francois.clayer@niva.no)

‡ Present address: Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA

Abstract. Despite high potential benefits, the development of seasonal forecasting tools in the water sector has been slower than in other sectors. Here we assess the skill of seasonal forecasting tools for lake and reservoir set up at four sites in Australia and Europe. These tools consist of coupled hydrological catchment and lake models forced with seasonal meteorological forecast ensembles to provide probabilistic predictions of seasonal anomalies in water discharge, temperature and ice-off. Successful implementation requires a rigorous assessment of the tools' predictive skill and an apportionment of the predictability between legacy effects and input forcing data. To this end, models were forced with two meteorological datasets from the European Centre for Medium Range Weather Forecasts (ECMWF), the seasonal forecasts SEAS5 with three-month lead times and the ERA5 reanalysis. Historical skill was assessed by comparing both model outputs, i.e., seasonal lake hindcasts (forced with SEAS5) and pseudo-observations (forced with ERA5). The skill of the seasonal lake hindcasts was generally low although higher than the reference hindcasts, i.e., pseudo-observations. In addition, the SEAS5 meteorological predictions showed less skill than the lake hindcasts. In fact, skilful lake hindcasts identified for selected seasons and variables were not always synchronous with skilful SEAS5 meteorological hindcasts, raising questions on the source of the predictability. A set of sensitivity analyses showed that most of the forecasting skill originates from legacy effects, although during winter and spring in Norway some skill was coming from SEAS5 over the three-month target season. When SEAS5 hindcasts were skillful, additional predictive skill originates from the interaction between legacy and SEAS5 skill. We conclude that a climatology driven forecast is currently likely to yield higher quality forecasts.

1. Introduction

Freshwater provides essential services for food and energy production, manufacturing, cultural heritage, and natural habitats.

35 However, it is threatened by more frequent extreme events (Jeppesen et al., 2021), climate change (Labrousse et al., 2020), anthropogenic water depletion (Yi et al., 2016) and agricultural pressures (Wuijts et al., 2021). Implementation of mitigation measures can help preserve freshwater resources, although they come with trade-offs between production from economic sectors with related social benefits, and availability of good quality freshwater. Hence, successful implementation of measures requires capacity at the local-regional level for cross-sectoral decision-making (Wuijts et al., 2021). Seasonal forecasting tools 40 for water quality can help facilitate the decision-making process by refining optimal actions over the next season, e.g., magnitude and timing of reservoir drawdowns. Indeed, they can supply knowledge on the impacts of future climatic conditions on freshwater over a realistic time frame enabling implementation with reduced negative effects on economic activities. Nevertheless, the use and access to forecasting tools is still very limited for water managers (Lopez & Haines, 2017; Soares et al., 2018). The probabilistic nature of seasonal forecasts can be a key barrier coupled with the lack of reliability and credibility 45 of these predictions in most regions out of tropics. Hence, a better access to seasonal forecasting tools as well as increased comprehension and description of these tools are required prior to their successful implementation in the decision-making process within the water sector.

Seasonal meteorological predictions provide a probabilistic description of the weather over the next few months, e.g., an 80% chance of the weather being wetter than normal. Seasonal climate predictability mainly originates from ocean–atmosphere 50 interactions (Troccoli, 2010). In fact, the ocean inertia, given its volume and the heat capacity of liquid water, exerts an influence on the atmosphere on the scale of months which allows us to estimate its future effect on weather. Given that ocean–atmosphere interactions are relatively strong in the equatorial region (Troccoli, 2010), seasonal meteorological predictions typically show stronger predictive skill, or prediction performance, around the tropics (Johnson et al., 2019; Manzanas et al., 2014). Under higher latitudes, skills from seasonal meteorological predictions are patchy and less consistent among variables 55 and seasons. Hence, the boundary conditions, e.g., seasonal air temperature forecasts used to force a hydrological model, is usually not the main source of predictability outside the tropics, at least for stream flow (Greuell et al., 2019; Harrigan et al., 2018; Wood et al., 2016). Nevertheless, climate models producing seasonal meteorological forecasts are constantly improving and it is reasonable to expect that forecast opportunities will expand in the future (Mariotti et al., 2020). Developing seasonal forecasting workflows, quantifying the skill and investigating the source of predictability represent a necessary and essential 60 step towards reliable water quality seasonal forecasting.


While some of the first forecasting tools were originally developed for flood warnings (e.g., Pagano et al., 2014; Werner et al., 2009), applications to other sectors are becoming more frequent. In the agricultural sector, for example, a recent study shows that flowering time can be reliably predicted from seasonal meteorological forecasts in central and eastern Europe, enabling 65 early variety selection and planning of farm management (Ceglar & Toreti, 2021). Seasonal meteorological forecasts were also shown to provide useful information for the wind energy sector (Lledó et al., 2019), and to avoid significant economic losses

from hydropower generation during droughts (Portele et al., 2021). Nevertheless, the use of seasonal meteorological forecasts for water temperature in lakes and reservoirs has been limited so far, where the focus has been on water quantity (Arnal et al., 2018; Giuliani et al., 2020; Greuell et al., 2019; Pechlivanidis et al., 2020). Studies forecasting water temperature, a fundamental water quality variable, are rare in the literature (Mercado-Bettin et al., 2021; Zhu et al., 2020; Baracchini et al., 70 2020), despite the diverse influence of this variable on lake ecosystem structure and functioning (Dokulil et al., 2021). Nevertheless, a simple lumped model (*air2water*; Piccolroaz et al., 2013), previously developed to estimate surface lake water temperature as a function of air temperature, has been applied to predict water temperature in thousands of lakes (Zhu et al., 2021). While this hybrid approach yielded skillful surface lake water temperature forecasts (Piccolroaz et al., 2018; Toffolon et al., 2014), it doesn't allow forecasting any other lake variable, such as bottom temperature or ice-off, i.e., the first ice-free 75 day after a ice-covered period.

Research on seasonal forecasting in hydrology has started more than a decade ago (Troin et al., 2021) and now represents a source of knowledge for other research fields. When forecasting river flow, for example, predictability can originate from two main sources: (i) initial conditions such as catchment water stores of initial soil moisture, groundwater, and snowpack, which are directly linked to the water residence time; and (ii) boundary conditions, i.e., meteorological forecasts used to force the 80 hydrological model (Greuell et al., 2019). Throughout the many studies of river flow seasonal forecasting in Europe, it appears that initial conditions form the dominant source of skill in run-off (Greuell et al., 2019; Harrigan et al., 2018; Wood et al., 2016) and predictability can be extended up to a year ahead in case of very low flow (Staudinger & Seibert, 2014). When dealing with standing water bodies, antecedent conditions are also likely to provide significant predictability, given that the water storage in lakes and reservoirs is large compared to river channels, providing higher inertia. Water residence time is thus 85 expected to exert a strong influence on discharge predictability. Water temperature, on the other hand, is influenced by multiple meteorological variables, e.g., wind, air temperature and radiation, in addition to water stores which can affect the source of its predictability.

Here, we further investigate the performance and in particular the source of this prediction performance, also referred to as predictive or forecasting skill, of lake seasonal forecasting tools first described by Mercado-Bettin et al. (2021) and Jackson-90 Blake et al. (2022). These tools integrate hydrological catchment and physical lake models forced with seasonal meteorological forecasts with three-month lead times at four case study sites in Europe and Australia (Fig. 1). In fact, the model forcing meteorological variables as well as output catchment and lake variables are a set of retrospective seasonal forecasts for past dates, hereafter referred to as hindcasts, that can be compared to historical records. The objective of this study is to assess whether seasonal meteorological hindcast ensembles with three-month lead time used as inputs to catchment and lake process-95 based models provide some predictive skill to seasonal lake hindcasts. To this end, the forecasting skill of the tools was assessed for combinations of season and freshwater variables, i.e., discharge, water temperature or ice-off, and for each tercile. In parallel, we quantified the forecasting skill of each meteorological variable of the seasonal meteorological prediction at each site. Both assessments were carried out following aggregation of model outputs from daily to seasonal temporal resolution, i.e., seasonal means or sums. When a hindcast was found to perform significantly better than a reference hindcast, e.g.,

100 climatology from pseudo-observations as defined in the Methods, for a combination of a given season, variable and tercile, this latter combination was defined as a “window of opportunity”. This terminology is introduced to emphasize the fact that these forecasts can be used in the decision-making processes by water managers but only for specific variable and season. A set of sensitivity analyses was performed to identify input-output relationships and to partition the source of the prediction skill for each window of opportunity among warm-up, first lead-month and seasonal meteorological predictions. The 105 comparison in hindcasts with the aim of isolating the contributions of different sources to skill has been applied before on streamflow hindcasts (e.g., Arnal et al., 2018; Greuell et al., 2019), but this is, to our knowledge, the first study investigating the origin of seasonal hindcast ensemble skills on water discharge, temperature and ice-off in lakes and reservoirs. The implications for lake forecasting tools are discussed.



110 **Figure 1: Location of the four case studies in Europe and Australia along with climate type and coordinates. Map is modified from Jackson-Blake et al. (2022). Detailed catchment maps are given in Jackson-Blake et al. (2022).**

2. Methods

2.1 Description of the forecasting tools

The forecasting tools consist of a coupled catchment runoff model to a one-dimensional water column lake model, forced by seasonal meteorological predictions, to simulate three output variables at daily resolution: inflow discharge, and lake surface and bottom temperature. For Lake Vansjø in Norway, the timing of ice melt (ice-off) was also included in the output variables in spring. The workflow consisted in running the catchment models first, providing inflow water discharge and water temperature to the lake models.

2.1.1 Case study sites

Lake forecasting tools were developed for four regulated water lakes/reservoirs in Europe and Australia which have been described earlier (Mercado-Bettin et al., 2021; Table 1; Fig. 1). Briefly, Sau (Spain) and Mount Bold (Australia) reservoirs large water supplies for the cities of Barcelona and Adelaide, respectively. Lake Vansjø (Norway), is a drinking water source for three municipalities and the Wupper reservoir (Germany) is used for flood control, environmental flows, and recreation.

Table 1: Characteristics of the study sites. Mixing timing refers to boreal seasons only.

Case study (Country)	Catchment area (km ²)	Surface area (ha)	Volume (hm ³)	Water retention time (yrs)	Max. Depth (m)	Mixing regime	Mixing timing
Sau (Spain)	1680	575	165	0.2	60	monomictic	Winter
Mt Bold (Australia)	357	254	46.4	0.2-0.6	44.5	monomictic	Summer
Vansjø (Norway)	690	3600	252	1.1	19	dimictic	Spring Fall
Wupper (Germany)	215	211	26	0.2	31	dimictic	Spring Fall

2.1.2 Meteorological input data

We used two different meteorological datasets to force the catchment hydrological and lake physical models in our tools: a climate reanalysis (ERA5) and a seasonal forecasting product (SEAS5) which both offer a global spatial and continuous temporal coverages to ensure future transferability of our workflows and easy comparison between our case-studies (Johnson et al., 2019). ERA5 is the latest reanalysis at 0.25° spatial resolution (Hersbach et al., 2020) produced by the European Centre for Medium Range Weather Forecasts (ECMWF; <https://www.ecmwf.int>) within the Copernicus Climate Change Service (C3S, <https://climate.copernicus.eu>). ERA5 data (1988-2016) were used (i) to correct for bias in the SEAS5 data using the quantile mapping technique as described below; (ii) to provide meteorological pseudo-observations for retrospective skill evaluation of SEAS5 hindcasts, (iii) to force catchment hydrological and lake physical models to produce pseudo-observations of the output variables, (iv) to force our catchment and lake models to produce antecedent/warm-up period data preceding

135 seasonal hindcast periods (i.e., combined one lead-month and three-month target season). SEAS5 is the latest seasonal
forecasting system from the ECMWF at 1° spatial resolution and provides operational seasonal forecasts and retrospective
seasonal forecasts for past years (hindcasts). We used hindcasts (1994-2016) in this study. A hindcast with 25 members was
considered for the period 1994-2016 for the three-month boreal seasons (spring: March through May; summer: June through
August; autumn: September through November; winter: December through February), with one month as lead time. A
140 dedicated R package (climate4R; Iturbide et al., 2019) was used for ERA5 and SEAS5 meteorological data pre-processing.
SEAS5 members were pre-processed using the quantile mapping technique (Gutiérrez et al., 2019) to correct for systematic
bias relative to pseudo-observations (ERA5 reanalysis). We used the empirical quantile mapping approach (EQM) due to its
ability to deal with multivariate problems (Wilcke et al., 2013). EQM adjusts 99 percentiles and linearly interpolates inside
this range every two consecutive percentiles; outside this range, a constant extrapolation (using the correction obtained for the
145 1st or 99th percentile) is applied (Déqué, 2007). In the case of precipitation, we applied the wet-day frequency adaptation
proposed by Themeßl et al. (2011). The resulting bias-corrected data were used for hydrologic and lake models meteorological
forcing, noting that we implemented bias-correction using leave-one-(year)-out cross-validation. Therefore, for each year,
seasonal climate hindcast member predictions were adjusted with the bias correction parameters derived from training with all
150 other years; after which all bias-corrected data were appended to obtain a corrected (i.e., locally calibrated) time series of
seasonal meteorological hindcasts for the full period for each case study. Finally, to use the bias-corrected data as
meteorological forcing for hydrologic and lake models, we used bilinear interpolation (akima method), whereby we specified
lake/reservoir coordinates from which seasonal meteorological hindcast data from surrounding pixels were interpolated.
155 Meteorological datasets include daily average 2-meter air temperature, u and v components of wind, surface air pressure,
relative humidity (or dew-point temperature), cloud cover, short-wave radiation, downwelling long-wave radiation and daily
sum of precipitation.

2.1.3 Observations

Daily inflow discharge and daily to monthly lake water temperature observations (Table S1) were used for catchment and lake
model calibration and validation, as well as quantification of forecasting skills. For Lake Vansjø, daily measurements of
discharge over 1994–2016 were taken from the gauging station at Høgfoss (Station 3.22.0.1000.1; Norwegian Water Resources
160 and Energy Directorate). Lake temperature data were gathered from the Vansjø-Hobøl monitoring program dataset, conducted
by the Norwegian Institute for Bioeconomy Research and by the Norwegian Institute for Water Research (Haande et al., 2016).
These data are available freely on the Norwegian national database (<https://vannmiljo.miljodirektoratet.no>). For Sau reservoir,
daily measurements of discharge into Sau Reservoir were provided by the Catalan water agency (Agència Catalana de l'Aigua,
ACA) while lake temperature and weather data are part of a long-term monitoring program (Marce et al., 2010). Discharge,
165 water temperature and weather observations at the two other reservoir sites were collected from the water reservoir operators
(Wupperverband for Wupper and SA Water for Mt Bold). Lake water temperature data are discontinuous and covered only
part of the modelled time-period (1994–2016) because of limited funding for monitoring programs. In addition, precipitation,

temperature, short-wave radiation, humidity, and wind daily records at nearby meteorological stations were obtained for each case study from the local meteorological institutes. For Lake Vansjø, this included ice-off dates from the Norwegian Meteorological Institute station 1,715 (Rygge) located on the lake shore (59° 38'N, 10° 79' E).

2.1.4 Catchment-lake process-based model setup and calibration

A catchment-lake process-based model chain was setup at each site to predict daily inflow discharge into the lake/reservoir and daily lake water temperature. Given the specificity of each catchment regarding flow dynamics and water management, different models were used at each site (Fig. 2). While this disparity prevents us from an in-depth comparison among case-studies, the common methods and code established to manipulate input and output data enable us to quantify forecast performance and the source of the predictability at each site in a consistent and comparable way.

Inflow water temperature and discharge for Sau and Vansjø was modelled with the mesoscale Hydrologic Model (mHM v5.9: <http://www.ufz.de/mhm>) and SimplyQ (hydrological module of SimplyP; Jackson-Blake et al. 2017), respectively. Inflow water temperature and discharge for Wupper and Mt Bold was modelled with the *Génie Rural* (GR) suite of models implemented within the R package airGR (Coron et al., 2017), GR6J and GR4J, respectively. mHM and SimplyQ hydrologic models were forced with ERA5 daily precipitation and daily average surface air temperature, and the GR models were forced with daily precipitation and daily potential evapotranspiration (Hargreaves-Samani potential evapotranspiration, derived from daily minimum and maximum temperature, implemented in drought4R; Iturbide et al., 2019). All hydrological models were calibrated and validated against local observations using the Nash–Sutcliffe efficiency coefficient (NSE) as the objective function.




Figure 2: Description of the forecasting workflow—Calibrated hydrologic and lake models are used to produce seasonal lake hindcasts with 25 members

The General Ocean Turbulence Model (GOTM, <http://gotm.net>) was used to simulate the water temperature profile of Sau Reservoir and Lake Vansjø. The General Lake Model (GLM, Hipsey et al., 2019) was used to simulate water temperature in

the Mt. Bold and Wupper reservoirs. Lake models were forced with ERA5 surface air temperature, u and v wind components, surface air pressure, relative humidity (or dew-point temperature), cloud cover, short-wave radiation, precipitation and, in some cases, also downwelling long-wave radiation, and calibrated and validated against observations using the Root-Mean-Square Error (RMSE) and NSE as objective functions.

195 For Lake Vansjø, the water level was set to constant given that observed fluctuations are < 1 m which are not critical for the lake heat and water budgets. The three reservoirs, on the other hand, experience much larger water level fluctuations because of complex water pumping patterns and/or water scarcity. It was thus critical to allow for water level fluctuations and parametrize the outflows to avoid dry outs. For Wupper Reservoir, a statistical model was developed to calculate the reservoir's outflow based on the inflow using the timeseries over the warm-up period for each discharge simulation of the catchment 200 model. Such an approach allowed mimicking the outflow decision and approximately resembling the observed water-level to avoid the cases of dry-outs or exceedingly low volumes of water due to inflow/outflow misestimation. More details on the performance of the linear regression are given in the supplementary information. For Sau, historical observations of outflow and pumping volumes were used to force the model. For Mt. Bold Reservoir, an average annual cycle was calculated from historical observations and then replicated throughout the entire timeseries. While this assumption does not allow for inter- 205 annual variation, it allowed for simulation of water level fluctuation each year that represented the seasonal cycle apparent within Mt. Bold and avoided dry outs.

The lake energy budget includes exchanges through the air-water interface, i.e., downward short-wave radiation, downward and upward long-wave radiation, latent and sensible heat fluxes, and by lateral fluxes of water, i.e., inflow and outflow of water (Schmid and Read 2022). The energy fluxes at the air-water interface are accounted for in the GLM or GOTM lake 210 model, however, the lateral fluxes caused by throughflow (inflow-outflow balance) need to be parametrized through the addition of water temperature to the inflow provided by the catchment model. Inflow temperature was estimated based on the assumption that water temperatures follow the air temperatures closely with some time lag (Stefan & Preud'homme 1993; Ducharne 2008). Hence, water temperature was predicted with a linear model of the form $A + B * \text{AirTemperature}$ where A and B were optimized against local observations when available. At Sau Reservoir, the values of A and B were 5.12 and 0.799, 215 respectively, while for Mt Bold reservoir and Lake Vansjø, the values of A and B were 5 and 0.75, respectively. The validation of this model for Wupper Reservoir, as an example, is described in the supplementary information (SI).

Most common verification statistics, e.g., Kling-Gupta efficiency (KGE), NSE and RMSE, for hydrological and lake modeling were calculated. Details on calibration and validation periods as well as statistics are shown in Table 4 and Table S2.

2.1.5 Pseudo-observations (Lake_PO)

220 Following calibration, lake and hydrologic models were forced with ERA5 over 1994-2016 to produce daily pseudo-observations of river discharge, daily surface and bottom temperature, as well as presence or absence of ice (for Lake Vansjø only). The output of this simulation is hereafter referred to as lake pseudo-observations (Lake_PO). Theoretical prediction skill of seasonal forecasts is commonly evaluated against pseudo-observations (Greuell et al., 2019; Harrigan et al., 2018;

Wood et al., 2016). In contrast to lake real observations, Lake_PO have the advantages of being complete and allow to disregard changes in skill related to model errors or biases (Harrigan et al., 2018), and to focus on skill originating from initial and boundary conditions. In contrast to the theoretical prediction skill, the total prediction skill includes any error or bias introduced by the model. Here, the total prediction skill of seasonal lake hindcasts (discharge, water temperature and ice-off) was also evaluated against real observations, when those were available and covering a representative time period.


2.1.6 Seasonal forecasts (Lake_F)

For each of the 92 three-month hindcast seasons (11/1993 to 11/2016), we simulated ensemble predictions of daily river discharge, daily surface and bottom water temperature as well as presence or absence of ice (for Lake Vansjø only; Fig. 3). Catchment and lake models were forced with ERA5 data over the 1-year warm-up period followed by a set of 25 members of SEAS5 data covering an first lead month (M0) and the 3-month long target season (M1–M3). The first lead month is defined in agreement with Gruell et al. (2019) as the month following the date on which the forecast would have been issued. Over the first lead month, the 25 members of SEAS5 progressively diverge from ERA5 to their respective SEAS5 member. Model outputs for the final 3 months, i.e., the target season, were aggregated into three month (M1–M3) seasonal averages or sums (i.e., average surface and bottom water temperature and cumulative seasonal inflow discharge). The output of this simulation is hereafter referred to as lake forecasts (Lake_F).

2.1 Assessment of modeling performance and source of forecasting skills

2.1.1 Model and forecast verification

A complete assessment of the modelling and forecasting performance of our workflow was performed through several verifications (Table 2). The first verification (Verification 1 in Table 2) consisted in evaluating the performance of the models forced with ERA5 by comparing model outputs (Lake_PO) to observations at daily temporal resolution, as described in section 2.1.4. This verification step included the reporting of traditional verification statistics for modelling, i.e., NSE; KGE and RMSE. The second and third verifications (Verifications 2 and 3 in Table 2) consisted in quantifying the lake forecast (Lake_F) performance compared to climatology from pseudo-observations (Lake_PO) or from observations, respectively. These steps allowed to quantify the forecasting skill of a perfect model and the total forecasting skill, respectively. Forecast verification 2 and 3 were performed using model output data at seasonal temporal resolution, i.e., daily model outputs over the target season (M1–M3), were aggregated in seasonal averages or sums.



250 **Figure 3: Time series of the air temperature (a), precipitation (b), discharge (c), surface (d) and bottom (e) water temperature over the warm-up, first lead month (M0) and target season (M1–M3) for Autumn 2000. The black lines indicate ERA5 (a and b) and Lake_PO (c–e) data, the light and dark blue lines are, respectively, the 25 members and the mean of SEASS5 (a and b) and Lake_F (c–e).**

255

Forecast performance was quantified with two skill scores: the Ranked Probability Skill Score (RPSS) and the Relative Operating Characteristic Skill Score (ROCSS). Skill scores are a measure of the relative improvement of the forecast compared to a reference forecast which here is the climatology based on either Lake_PO or observations. ROCSS values were calculated against climatology from real observations ($ROCSS_{obs}$), in addition to pseudo-observations ($ROCSS_{original}$), only when

260

observations covered the whole season. Indeed, $ROCSS_{original}$ was calculated only if there was at least one observation point in each month of the season and observations for at least 70% of the seasons. Observations that met these criteria only included inflow discharge at Vansjø, Sau and Wupper for all seasons, surface and bottom temperature at Vansjø in summer only, surface and bottom temperature at Wupper for all seasons, surface temperature at Sau for all seasons and ice-off at Vansjø.

RPSS and ROCSS are commonly used as evaluation measures of probabilistic forecasting skill (Jolliffe & Stephenson, 2012; Müller et al., 2005). The visualizeR package (Frías et al., 2018) was used to compute the RPSS and ROCSS for Lake_PO and Lake_F. Briefly, the RPSS provides a relative performance measure on how well the probabilistic ensemble is distributed over the lower, middle and upper terciles, while the ROCSS provides a relative measure of discriminative skill for each category. A $RPSS > 0$ is associated with a better forecast than the reference (1 being a perfect score), while $RPSS \leq 0$ indicates no improvement compared to the reference. The ROCSS value ranges from -1 (perfectly bad forecast) to 1 (perfect forecast) and a zero value indicates no skill compared to the reference. The RPSS has been shown to be sensitive to the ensemble size, but this effect can be corrected for using the Fair (or unbiased) RPSS (Ferro et al., 2014). To allow for comparison with other forecasting systems, we have used the fair RPSS (FRPSS) forecast verification. In this study, the FRPSS is calculated for tercile events. The statistical significance of the FRPSS and ROCSS is computed based on the 95% confidence level from a one-tailed Z test. When a forecast for a given season, variable and tercile was associated with a ROCSS value that was statistically significant, we referred to it as a window of opportunity (i.e., a combination of season, variable and tercile for which forecast performance was significantly better than the reference). In our case, the threshold values above which a ROCSS was considered significant typically range between 0.47 and 0.55.

Table 2: Comparison carried out to evaluate model and forecast performance

Verification	Outputs used	Reference forecast data	Purpose	Statistics
1	Lake_PO	Observations	Assess lake model skill	KGE NSE RMSE
2	Lake_F	Lake_PO	Assess the transfer of meteorological forecast skill through process-based models – Perfect model forecasting skill	$ROCSS_{original}$
3	Lake_F	Observations	Assess total forecasting skill	$ROCSS_{obs}$

280 **2.1.2 Sensitivity analyses to initial conditions and meteorological forcing**

Several sensitivity analyses (SA), summarized in Table 3, were performed to identify the origin of the forecasting skill for a given window of opportunity, i.e., a combination of season, variable and tercile for which forecast performance was significantly better than the reference. Results of the SA are only reported for sites having a substantial number of windows of opportunity for conciseness. These SA allowed quantifying the sensitivity of hindcast performance to forcing data over specific periods: the target season (M1–M3; SEAS5), the first lead month (M0) and the warm-up period (ERA5). It was thus possible to quantify the proportion of skills originating from each of these periods.

The SA consisted of replacing the forcing data of interest, i.e., over the target season, the first lead month or the warm-up period, by data from an equivalent season/period but from a randomly selected year. For example, for the target season SA (S-SA), the SEAS5 forcing data covering the 3-month target season was replaced by SEAS5 data from a randomly selected equivalent season. Furthermore, the SA for the warm-up period (W-SA) consisted in replacing the ERA5 data covering the warm-up period by ERA5 data from a randomly selected equivalent time-period. The last SA covered warm-up and first lead month (W+M0-SA) and consisted in replacing ERA5 data over the warm-up, as in W-SA, but also SEAS5 data over the first lead month. To ensure that the randomly sampled forcing data are representative of the whole SEAS5 or ERA5 datasets, we introduce two levels of repetitions for all experiments. First, we randomly selected a year for each of the 25 members of SEAS5, meaning that the data selected to replace the original SEAS5 forcing data is extremely likely to be from a different year for each SEAS5 member. Second, we repeated the analysis 25 times, for each season. Sensitivity analyses were only carried out for Spain and Norway because of the low number of windows of opportunity at the two other sites and considering the resources needed to execute these hindcast experiments.

The outputs of each of the sensitivity analysis were used to calculate ROCSS values against the climatology based on Lake_PO, as for Lake_F in the Verification 2 described above (Table 2). The ROCSS values obtained through this procedure were, respectively, $ROCSS_S$, $ROCSS_W$ and $ROCSS_{W+M0}$ for S-SA, W-SA, and W+M0-SA. The ROCSS values obtained for the various SAs were compared to the original Lake_F ROCSS values ($ROCSS_{original}$) to investigate the sources of prediction skill. An estimation of the proportion of prediction skill originating from the SEAS5 data over the target season (P_{season}) was expressed as follows:

$$305 \quad P_{season} = ROCSS_{original} - ROCSS_S \quad (1)$$

Similarly, the proportions of prediction skill originating from the ERA5 data over the warm-up ($P_{warm-up}$) and from the SEAS5 data over the first lead month (P_{M0}) can be respectively estimated as:

$$P_{warm-up} = ROCSS_{original} - ROCSS_W \quad (2)$$

$$P_{M0} = ROCSS_W - ROCSS_{W+M0} \quad (3)$$

310 In Eq. 1–3, prediction skill was assumed to linearly scale with ROCSS values and skill from any interaction effect was neglected. While we admit that Eq. 1–3 are not necessarily statistically correct, they are useful to quantify the relative importance of the sources of skill. Hence, the values of P_{season} , $P_{warm-up}$ and $P_{transition}$ should be interpreted with care.

2.2.3 Sensitivity analyses to individual input variables

To further investigate through which process forecasting skill is transferred from input to output variables, a one-at-a-time sensitivity analysis (OAT-SA) was performed for Lake_PO and the Pearson partial correlation coefficients (PPCC) between each variable of Lake_PO, i.e., surface temperature, bottom temperature, discharge, ice-off, and a set of relevant input variables were determined (Table 3). The OAT-SA consisted in replacing the data for a specific input meteorological variable by data

from an equivalent target season but from a randomly selected year. The seasonal means of OAT-SA outputs were compared to default outputs (Lake_PO) with R^2 . Higher $(1 - R^2)$ values indicate more influence of input variables on Lake_PO.

320 PPCC allowed quantifying the sensitivity of model outputs to a given input variable while removing the effect of the remaining input variables. Note that PPCC were calculated on seasonally aggregated variables. To ensure that PPCC were statistically appropriate, i.e., only when a linear relationship exists between the seasonal means of input factors and those of the output (Pianosi et al., 2016), the linearity assumption was checked through visual inspection of scatter plots between each input and output variables. Partial correlation coefficients are a good alternative to ‘All-At-a-Time’ (or global) SA when the latter is not 325 possible because of the lack of computing resources (Pianosi et al., 2016). To avoid misleading conclusions, correlation between input variables should be minimized (Marino et al., 2008). Hence, only the most relevant input variables were included. Precipitation and air temperature were retained for discharge, while air temperature, precipitation, wind speed (wind speed calculated from u and v components of wind) and short-wave radiation were retained for surface and bottom temperature. In fact, short-wave radiation was retained over relative humidity, cloud cover and air pressure because it was responsible for 330 most of air-water heat fluxes (see SI). Wind was retained because of its impact on thermal stability (Blottiere, 2015).

Table 3: List of sensitivity analyses (SA) performed

SA	Forcing data to be replaced		Model output	Purpose	Sensitivity index
	Period	Variable			
S-SA	Target season (SEAS5)	All	Lake_F	Quantifying the proportion of forecasting skill originating from SEAS5 data over the target season	$ROCSS_S$
W-SA	Warm-up period (ERA5)	All	Lake_F	Quantifying the proportion of forecasting skill originating from ERA5 data over the warm-up season – initial conditions	$ROCSS_W$
W+M0-SA	Warm-up period (ERA5) and first lead month (SEAS5)	All	Lake_F	Quantifying the proportion of forecasting skill originating from SEAS5 data over the first lead month	$ROCSS_{W+M0}$
OAT-SA	Target season (ERA5)	One at a time	Lake_PO	Quantifying the sensitivity of Lake_PO to a specific forcing variable	$1 - R^2$
PPCC	None	None	Lake_PO	Quantifying the sensitivity of Lake_PO to a specific forcing variable while removing the effect of the remaining variables	PPCC

3 Results

3.1 Performance of the calibrated catchment and lake models (Lake_PO)

Catchment and lake models calibrated against local observations performed reasonably well (Table 4). For river discharge,

335 NSE and KGE both ranged between 0.51 and 0.85 over the calibration and validation periods. For surface water temperature, RMSE ranged from 1.10 to 1.63 and NSE from 0.78 to 0.94 over the calibration and validation periods. Over each season, however, Lake_PO showed more heterogeneous performance (Table S2). Discharge simulations were usually worse in summer, except in Australia where performance was poor for most seasons. Surface water temperature modeling typically showed better performance during spring and fall than during summer or winter. There is no clear pattern for bottom water 340 temperature, but overall, it seems more difficult to be accurately simulated compared to surface temperature.

Table 4: Verification statistics of the catchment and lake model for each case study

	Output variable	Calibration				Validation			
		Time	NSE	KGE	RMSE	Time	NSE	KGE	RMSE
Norway	Discharge	2005–2010	0.51	0.56		2011–2015	0.57	0.57	
	Temperature	2005–2010	0.92		1.12	2011–2015	0.93		1.10
Spain	Discharge	1997–2007	0.60	0.66		2008–2018	0.54	0.63	
	Temperature	1997–2007	0.93		1.63	2008–2018	0.94		1.45
Germany	Discharge	1991–2011	0.71	0.85		2012–2016	0.63	0.81	
	Temperature	1993–2010	0.93		1.31	2011–2016	0.91		1.53
Australia	Discharge	2003–2007	0.64–0.80	0.70–0.84		2008–2013	0.65–0.80	0.54–0.75	
	Temperature	2014–2016	0.91		1.17	2016–2018	0.78		1.50

3.2 Skill of the seasonal meteorological (SEAS5) and Lake (Lake_F) hindcasts

345 Table 5 displays the ROCSS values for each combination of Lake_F output variable, season and tercile while Table 6 summarizes the windows of opportunity, i.e., a combination of season, variable and tercile for which forecast performance, or predictive skill, was significantly better than the reference, for SEAS seasonal meteorological hindcasts as well as for Lake_F hindcasts. These windows of opportunity typically had ROCSS values larger than 0.47 to 0.55 (see Methods section for details). For SEAS5 seasonal meteorological hindcasts, only 3 to 10 windows of opportunity were observed for each case study out of 350 the 96 possibilities, i.e., 3 terciles of 8 variables over 4 seasons (Table 6). Regarding Lake_F, larger proportions of the 36–39 possible variable-tercile-season combinations were associated with statistically significant ROCSS values (Table 6). Winter and Spring in Norway, as well as Summer and Autumn in Spain were the seasons associated with the most skillful Lake_F hindcasts. Lake Vansjø in Norway was the only case study where windows of opportunity for SEAS5 and Lake_F were consistently concentrated within the same seasons, i.e., mostly in Spring and to a lesser extent in Winter. For the other case

355 studies, there were fewer windows of opportunity for SEAS5 and those were more randomly distributed over the year. FRPSS values were typically reported for surface water temperature in spring and autumn, except for autumn in Spain. Norway and Germany also showed significant fair RPSS for bottom water temperature in spring and autumn, and summer and autumn, respectively. Note that neither river discharge nor any of the SEAS5 variables had FRPSS values in any case study. Windows of opportunity for bottom temperature represented more than half of the total for all case-studies and variables while those for
360 surface temperature and discharge were more sporadic.

The comparison of SEAS5 and Lake_F skillful hindcasts in Table 6 is already useful for identifying possible transfer of forecasting skill from the SEAS5 seasonal meteorological hindcasts to the catchment and lake models. SEAS5 meteorological hindcasts are skillful over only a very limited number of seasons, variables and terciles (Table 6). However, for Norway, there is a higher number of skillful meteorological and lake hindcasts in spring than in the other seasons. For the other case-studies, 365 such a clear connection between SEAS5 meteorological hindcasts and catchment/lake model outputs is not as apparent. We can thus hypothesize that the skill of catchment and lake model hindcasts in Norway is more inherited from the SEAS5 data than at other case studies. In contrast, skill of the catchment and lake model hindcasts at the other case-studies is hypothesized to originate from the legacy of the warm-up period or from the parametrization of the inflow-outflow water balance.

Verification statistics for Lake_PO seasonal means compared to observations (Table 7) show that the catchment and lake 370 models performed well at the Norwegian and Spanish sites in capturing interannual variability. In Germany and Australia, performance was lower. Note that when observation coverage was below 50%, no statistics were calculated given the low number of seasons represented and the risk of bias when computing seasonal averages. The difference between $ROCSS_{original}$ (comparing Lake_F and Lake_PO) and $ROCSS_{obs}$ (comparing Lake_F and lake observations) did not necessarily scale inversely with the verification statistics (Table 7). In fact, the $ROCSS_{obs}$ reported for the German site were 375 slightly lower or even larger than their respective $ROCSS_{original}$ with differences lower than 0.23. Whereas, for the Spanish site, three $ROCSS_{obs}$ values out of 4 were significantly lower than the $ROCSS_{original}$ with a difference larger than 0.33. Nevertheless, several output variables, e.g., bottom temperature in Germany and ice-off in Norway, are associated with significant $ROCSS_{original}$ and $ROCSS_{obs}$ which provides further confidence in model calibration and low model error. In contrast, even if the verification statistics for discharge were not worse than for the other variables, $ROCSS_{obs}$ values are all 380 below the significance threshold pointing towards some limitations in predicting hydrology.

Table 5: ROCSS_{original} for each combination of season, variable and tercile of lake hindcasts (Lake_F). Color scale range from dark blue (ROCSS = -1, perfectly bad forecast) to dark red (ROCSS = 1, perfect forecast) with white in the middle (ROCSS = 0, no change compared to reference forecast). Windows of opportunity are highlighted by bold, black numbers, i.e., combination of season, variable and tercile associated with a statistically significant ROCSS value.

		Norway			Spain			Germany			Australia		
		Lower	Middle	Upper	Lower	Middle	Upper	Lower	Middle	Upper	Lower	Middle	Upper
Discharge	Spring	0.58	0.18	0.54	0.37	-0.11	0.33	-0.34	-0.03	-0.47	0.01	-0.33	-0.64
	Summer	-0.13	0.23	-0.41	0.73	-0.02	0.23	-0.59	-0.13	-0.34	-0.05	0.28	0.36
	Autumn	0.32	-0.24	0.27	0.17	0.46	0.47	-0.29	-0.33	0.03	0.48	-0.02	0.18
	Winter	0.21	-0.08	-0.1	0.4	0.52	0.08	-0.28	0.17	-0.34	0.16	-0.02	-0.22
Surface Temperature	Spring	0.75	0.14	0.53	0.2	0.18	0.19	0.2	-0.14	0.38	0.19	-0.27	0.26
	Summer	0.33	0.39	-0.13	0.42	-0.13	0.57	-0.25	-0.18	-0.22	0.16	-0.48	0.23
	Autumn	-0.52	-0.24	-0.12	0.22	0.11	0.47	0.23	0.42	0.3	0.13	-0.34	-0.04
	Winter	0.48	-0.12	-0.17	0.32	-0.66	-0.15	-0.22	0.04	-0.09	0.13	-0.14	-0.1
Bottom Temperature	Spring	0.56	0.05	0.68	0.44	0.12	0.86	0.59	0.28	0.6	0.37	0.31	-0.16
	Summer	-0.12	0.14	-0.35	0.53	0.36	0.72	0.48	0.21	0.71	0.6	-0.45	0.18
	Autumn	-0.53	0.03	-0.26	0.5	0.55	0.64	0.27	0.26	0.15	0.28	0.4	0.38
	Winter	0.48	0.01	0.53	-0.02	0.54	0.27	-0.16	-0.3	0.27	0.63	-0.1	0.29
Ice-off	Spring	0.69	0.29	0.75									

Table 6: SEAS5 meteorological and Lake_F lake hindcasts associated with statistically significant FRPSS or ROCSS at each case-study.

Site	Indexes	Numbers of skillful hindcasts: windows of opportunity										
		variable (tercile)										
		Winter		Spring		Summer		Autumn		TOTAL		
		SEAS5	Lake_F	SEAS5	Lake_F	SEAS5	Lake_F	SEAS5	Lake_F	SEAS5	Lake_F	
Norway	FRPSS					ST; BT				ST; BT	0/32	4/12
	ROCSS	3	3	7	8	0	0	0	0	10/96	11/39	
		cc (-)	ST (-)	airP(=,+)	Q (-,+)							
		sw (+)	BT (-,+)	airT (+)	ST (-,+)							
		lw (=)		cc (=)	BT (-,+)							
				hum (-)	Ice-off							
				U (+)	(-,+)							
				V (-)								
Australia	FRPSS				ST				ST	0/32	2/12	
	ROCSS	2	1	1		1	1	4	1	8/96	3/36	
		airP (+)	BT (-)	lw (+)		cc (=)	BT (-)	airP (+)	Q (-)			
		hum (+)						cc (+)				
								airT (=)				
								sw (+)				
Spain	FRPSS				ST					0/32	1/12	
	ROCSS	0	1	2	1	2	5	1	3	5/96	9/36	
		Q (=)	cc (+)	airP (+)	BT (+)	cc (+)	Q (-)	cc (+)	Q (+)			
						hum (+)	ST (+)		BT (-,+)			
							BT (-,+)					
Germany	FRPSS				ST		BT		ST; BT	0/32	4/12	
	ROCSS	2	0	1	2	0	2	0	0	3/96	4/36	
		lw (=)		hum (+)	BT (-,+)		BT (-,+)					
		V (-)										

390 Lake_F variable abbreviations: ST, BT and Q stand for surface-, bottom-temperature and discharge, respectively. SEAS5 meteorological variable abbreviations: airT, airP, cc, hum, sw, lw, U, V stand for surface air temperature, air pressure, cloud cover, relative humidity (or dew-point temperature), short-wave radiation, downwelling long-wave radiation, and u and v components of wind, respectively. -, + and = stands for lower, upper and middle terciles, respectively.

395 **Table 7: Verification statistics (NSE, R², RMSE, RMSE/sd, bias) for Lake_PO seasonal means, as well as comparison of the
ROCSS_{original}-(comparing Lake_F and Lake_PO) and *ROCSS_{obs}* (comparing Lake_F and lake observations).**

Site	Variable	Season	Obs coverage			NSE	R ²	RMSE	RMSE/ sd	bias	<i>ROCSS_{original}</i>			<i>ROCSS_{obs}</i>		
			S	M	D						lower	middle	upper	lower	middle	upper
Norway	Discharge	Spring	100	96	93	0.72	0.80	2.0	0.52	-1.0	0.58*	0.54*	0.36			0.34
	Surface	Winter	0	0	0						0.48*		n.a			
	Temperature	Spring	48	58	5						0.75*	0.53*	n.a			n.a
	Bottom	Winter	0	0	0						0.48*	0.53*	n.a			n.a
	Temperature	Spring	43	52	4						0.56*	0.68*	n.a			n.a
	Ice-on		100	-	-	0.97	0.99	2.2	0.16	1.8				a		
	Ice-off		100	-	-	0.36	0.76	19.3	1.09	-14.7	0.69*	0.75*	0.55*			0.68*
Spain	Discharge	Winter	100	100	99	0.88	0.89	3.9	0.34	-0.6	0.52*				0.18	
		Summer	100	100	98	0.51	0.62	3.5	0.69	-1.6	0.73*			0.40		
		Autumn	100	100	98	0.73	0.74	4.0	0.51	-0.8		0.47*			0.40	
	Surf. Temp.	Summer	78	78	3	0.12	0.40	1.1	0.87	-0.6		0.57*			-0.08	
	Bottom	Spring	48	70	3							0.86*			n.a	
	Temperature	Summer	48	67	2						0.53*	0.72*	n.a			n.a
		Autumn	35	58	3						0.50*	0.64*	n.a			n.a
Ger.	Bottom	Spring	100	96	6	-5.01	0.49	1.2	2.40	1.0	0.59*	0.60*	0.41			0.46*
	Temperature	Summer	100	100	7	-8.63	0.26	3.8	3.04	3.6	0.48*	0.71*	0.51*			0.49*
Australia	Discharge	Autumn	43	100	100	-0.67	0.41	1.61	1.23	-1.27	0.48*			n.a		
	Bottom	Winter	23	100	82	-0.70	0.32	1.98	1.17	1.51	0.63*			n.a		
	Temperature	Summer	17	75	46						0.60*			n.a		

Only output variables associated with statistically significant *ROCSS_{original}* are included. Statistically significant *ROCSS* are highlighted with an asterisk.

“Obs coverage” is the percentage of seasons (S), months (M) and days (D) covered by observations. Spring is March to May, Summer is June to August, Autumn is September to November, and Winter is December to February.

^aIce-on typically occurs between November and December which is the autumn and winter boundary. Therefore, *ROCSS* values could not be calculated for

400 ice-on.

3.3 Sensitivity analyses to initial conditions and meteorological forcing

The *ROCSS_S*, *ROCSS_W* and *ROCSS_{W+M0}* values obtained for each run of S-SA, W-SA and W+M0-SA, respectively, are summarized in boxplots in Figure 4 together with the original *ROCSS* value for each window of opportunity at the Norwegian and Spanish sites. This set of sensitivity analyses (SA) were performed to identify the origin of the forecasting skill for a given window of opportunity and allowed quantifying the sensitivity of hindcasts performance to forcing data over specific periods: the target season (M1–M3; SEAS5), the first lead month (M0) and the warm-up period (ERA5). In general, output variable


sensitivity to SEAS5 data over the target season (S-SA) is small relative to sensitivity to ERA5 data over the warm-up season and/or SEAS5 data over the first lead month. In fact, at Sau, replacing SEAS5 data over the target season with random data (S-SA) does not yield any significant change in the ROCSS values, except for the surface temperature upper tercile (Fig. 4 410 panel 1). However, significant changes in ROCSS values are seen for W-SA compared to $ROCSS_{original}$ indicating high sensitivity to warm-up. The similar ranges in $ROCSS_W$ and $ROCSS_{W+M0}$ values suggest limited or no impact of the SEAS5 data over the first lead month on output variable forecasts.

At Vansjø in Norway, on the other hand, 8 out of 11 windows of opportunity show significant changes in $ROCSS_S$ values, indicating higher sensitivity to SEAS5 data over the target season than at Sau. Furthermore, 3 windows of opportunity are 415 associated with $ROCSS_S$ that are lower than $ROCSS_{original}$ (Fig. 4b, f and g), i.e., suggesting SEAS5 is providing some skill, while 5 have $ROCSS_S$ that are higher than $ROCSS_{original}$ (Fig. 4a, h–k), suggesting the use of SEAS5 is in fact reducing forecasting skill compared to a random forecast. Then, a progressive decrease in ROCSS values is typically observed for all windows of opportunity following W-SA and W+M0-SA, indicating a progressive loss of forecasting skill related to ERA5 data over the warm-up and SEAS5 data over the first lead month.

420 3.4 Sensitivity analyses to specific input variables

Figure 5 and 6 summarizes the results from the two sensitivity analyses to specific input variables: OAT-SA and PPCC. Seasonal means of Lake_PO at Vansjø also showed higher sensitivity to specific input variables than Lake_PO at Sau (Fig. 5). In fact, surface temperature is highly sensitive to surface air temperature over the year while some other input variables have 425 more specific influence. Bottom temperature is also highly sensitive to surface air temperature but wind also plays a large role, especially in summer which is consistent with its expected impact on lake thermal stability (Blottiere, 2015). Finally, as expected, discharge at Vansjø is highly sensitive to precipitation, and to a lesser degree to surface air temperature, except in winter where surface air temperature has a larger influence on discharge.

The PPCC also show similar patterns regarding sensitivity (Fig. 6) where discharge is highly correlated with precipitation at the four sites and surface air temperature plays a secondary role for specific seasons. Once again, surface and bottom 430 temperature at Sau stand out due to their limited sensitivity to input variables while at the three other sites, surface temperature, and to a lesser degree bottom temperature, are generally strongly positively correlated with surface air temperature. Others, like precipitation and short-wave radiation have more of an anecdotal influence on lake temperature, while wind shows a more consistent negative impact on surface temperature at Vansjø, Wupper and Mt Bold. Wind also shows some impact on bottom temperature, although less consistent. At Vansjø and Mt Bold following the coldest season, wind is positively correlated with 435 bottom temperature, while at Wupper during the two coldest seasons, wind is negatively correlated with bottom temperature. Finally, ice-off date in Vansjø shows a strong negative correlation with surface air temperature (Fig. 6m) that can be linked back to the snow content and the intensity of snow melt in the catchment (Fig. 6n and o).



440 **Figure 4: Box plots (n = 25) of $ROCSS_S$, $ROCSS_W$ and $ROCSS_{W+M0}$ from sensitivity analysis runs S-SA (replacing target season
441 SEAS5 data with random data), W-SA (replacing warm-up ERA5 data with random data) and W+M0-SA (replacing warm-up
442 period – ERA5 and first lead month – SEAS5 data with random data) for each window of opportunity at the Norwegian (a–k) and
443 Spanish (l–q) sites. $ROCSS_{original}$ is given by the red line, so $ROCSS_S$, $ROCSS_W$ and $ROCSS_{W+M0}$ below the red line indicate a loss**

of skill and values above the line indicate higher skill than the original forecast. ***, ** and * indicate significant difference between a given group of $ROCSS_S$, $ROCSS_W$ and $ROCSS_{W+M0}$ values and $ROCSS_{original}$ following Mann-Whitney Rank Sum test at a significance level of 0.001, 0.01 and 0.05, respectively. Note that the S-SA, W-SA and W+M0-SA were only performed for Sau Reservoir in Spain and Lake Vansjø in Norway because of the significant resources needed to perform this hindcast experiments.





Figure 5: Relative sensitivity expressed as $1 - R^2$ of Lake_PO seasonal means to specific input variables estimated following the OAT-SA (see section 2.2.3 and Table 3 in the Methods section for details). Circle size represent relative sensitivity on a scale from 0 to 1. Meteorological variable abbreviations: airT, p, wind, cc, hum, and sw stand for surface air temperature, precipitation, wind speed, cloud cover, relative humidity (or dew-point temperature) and short-wave radiation, respectively. Note that the relatively larger sensitivity of Lake_PO to specific input variables over the whole year can be larger compared to over a given season because of the strong seasonal cyclicity. Note that the OAT-SA was only performed for Sau Reservoir in Spain and Lake Vansjø in Norway.



455

460

Figure 6: Pearson partial correlation coefficients (PPCC) between Lake PO seasonal means and seasonal means of selected input variables. Circle color and size represent PPCC value (from -1 to 1) and significance, respectively. Meteorological variable abbreviations: airT, P, wind, cc, hum, and sw stand for surface air temperature, precipitation, wind speed, cloud cover, relative humidity (or dew-point temperature) and short-wave radiation, respectively. Only significance level at 0.1 or below were considered in the interpretation.

Next, we use SA outputs to better describe the origin of the prediction skill, considering inertia, time integration as well as variable interactions. Assuming that climate signals in the ERA5 and SEAS5 input data over the warm-up, first lead month and target periods are additive sources of prediction skill, we can use Eqs 1–3 to partition the prediction skill originating from 465 those time periods, i.e., $P_{\text{warm-up}}$, P_{M0} and P_{season} , respectively. For Sau reservoir, this calculation yields $P_{\text{warm-up}}$ of 0.94 to 1.0 leaving only an unimportant fraction of prediction skill to the forcing data over the target season and the first lead month, as illustrated in Fig. 4. At this site, the output variables show in parallel very low sensitivity to input variables (Fig. 5 and 7) which supports a strong role of inertia or long-term time integration in hindcast predictive skill. The fact that 5 out of 470 the 6 windows of opportunity are for bottom water is also consistent with inertia as the main source of skill given the low circulation rate and inertia of hypolimnions. For Lake Vansjø, Eqs 1–3 yielded P_{season} of 0.003 (range: -0.19 to 0.18), P_{M0} of 0.19 (0.04 to 0.37) and $P_{\text{warm-up}}$ of 0.29 (0.09 to 0.60). Hence, a significant fraction of prediction skill is originating from the 475 SEAS5 boundary conditions although the largest source remains initial conditions through ERA5 data over the warm-up. Interestingly, the SEAS5 data over the first lead month is also a significant source of prediction skill. In fact, in decreasing order of importance, prediction skill originates from the warm-up, the first lead month and the target season. This progressive 480 decrease in prediction skill is only observed at Lake Vansjø and suggests that across-variable integration of climate signals persists through the first lead month and, in some cases, the target season, but is progressively deteriorating as we move into the target season. Indeed, there is additional consistency between the SEAS5 input variables showing some forecasting skill and the output variables. In fact, surface, and bottom temperature in spring at Vansjø are sensitive to surface air temperature and wind (Fig. 6b and c), and surface air temperature, wind u and v components are associated with some windows of 485 opportunity in spring (Table 6). Similarly, ice-off is sensitive to surface air temperature, as are snow quantities and melt intensities in the catchment (Fig. 6m–o). Hence, in contrast to Sau reservoir where most of the prediction skill seems to originate from inertia, at Lake Vansjø, across-variable integration contributes to predictive skills.

4 Discussion

4.1 Sources of skill

485 Our investigation into relationships between input and output variables and the sensitivity of predictive skill to meteorological data inputs over different time periods have yielded important insights into the sources of seasonal lake forecasting skill in our case study sites.

A key finding is that predictive skill is mostly sensitive to meteorological inputs over the warm-up and first lead months (Fig. 490 4, Section 3.4), although some specific windows of opportunity are also somewhat sensitive to the meteorological data over the target season. Hence, integration of the climate signal over time or across variables by catchment hydrologic and physical processes, e.g., snow accumulation (Harrigan et al., 2018) or heat accumulation in lakes, is likely a key source of predictive skill. In fact, Mercado-Bettin et al. (2021) already noted an increase in prediction skill when moving from weather to discharge

to lake temperature, i.e., in an increasing order of time and across variable integration of climate signals. Strong inertia is also a potential source of prediction skill.

495 After accounting for forecasting skill from the forcing data over various periods (Section 3.4), a large proportion of the skill still remains unexplained, especially for some selected windows of opportunity at Lake Vansjø in Norway. Bottom water temperature at Lake Vansjø in spring shows the highest residual skill after removal of skill from warm-up and first lead month (Fig. 4e–f). Surface and bottom temperature show a different degree of coupling with air temperature. In fact, while surface temperature responds tightly to changes in air temperature (Butcher et al., 2015; Schmid et al., 2014), bottom temperature 500 responds to a variety of complex interactions influenced by lake characteristics (e.g., fetch, surface area, depth, light penetration; Butcher et al., 2015). Indeed, bottom temperature in spring depends on preceding winter conditions but also on the intensity and length of the spring mixing event. To fully capture the intensity of this event, the model requires good initial water temperature inherited from previous winter but also skillful weather forcings, especially for surface air temperature and wind (Fig. 6c). In fact, for bottom temperature in spring to be higher than normal, it requires surface water to be heated up 505 more than normal, mainly through heat exchange with air temperature, but also the lake to remain mixed for a longer time period than normal. The interaction between skill from legacy and from weather forcing might thus be another source of predictive skill. The fact that the proportion of forecasting skill progressively decreases from warm-up, through the first lead month and the target season at Vansjø suggests that the interactions between input variables, which are incorporated in the process representation within the models, provide some skill but progressively deteriorates as we move forward in time. At 510 Sau reservoir in Spain, on the other hand, all skill is lost at the sharp boundary between the warm-up and the first lead month. This difference might be related to the presence of skill from the SEAS5 data at Vansjø (Table 6) and not at Sau. In other words, in the absence of skill in SEAS5 data, no additional skill can originate from interaction effects.

Literature on streamflow hindcasts broadly shows that beyond the first lead month, hindcasts forced with an ensemble of 515 boundary conditions resampled from historical meteorology are typically more skillful than hindcasts driven by seasonal meteorological predictions (Arnal et al., 2018; Bazile et al., 2017; Greuell et al., 2019). Hence, better lake forecasting skills could likely be achieved by simply forcing our models with climatology. Our results partly fit with these findings, as the skill of S-SA hindcasts for selected windows of opportunity were higher than the original hindcasts (Fig. 4a, h–k). These S-SA hindcasts are similar to climatology-driven hindcasts, although they are associated with higher uncertainty since they are driven by random SEAS5 data and should therefore be regarded as a minimum forecasting potential. For some windows of 520 opportunity, however, SEAS5 was a significant source of predictive skill (Fig. 4b, f, g and l). In those cases, only an improvement in SEAS5 forecasting skill is likely to improve lake forecasts. Improvement for only selected variables in SEAS5 would likely be enough to yield a significant increase in lake forecasting skill since most of the output variables presented here showed sensitivity to one or two input variables (Fig. 6).

4.2 Limitations and implications for seasonal lake forecasts

525 One apparent limitation of our study is the use of reanalysis weather data and pseudo-observations as inputs and benchmark output variables. Using pseudo-observations for skill assessment is a common methodology in streamflow forecasting studies (Alfieri et al., 2014; Wood et al., 2016) and it offers the opportunity to investigate the relationship between forecasting skills, initial and boundary conditions, while putting less emphasis on model errors and biases (Harrigan et al., 2018). Working with reanalysis weather data generates a less site-specific workflow and removes difficulties associated with dealing with temporal
530 and spatial heterogeneity in observed data. Nevertheless, here we also evaluate the forecasting skill against catchment and lake observations when possible (Table 7) and show that most of the windows of opportunity reported for water temperature held while those for discharge are no longer significant compared to observations. This discrepancy between discharge and water temperature can be related to the fact that discharge tends to be more variable than water temperature, with short-lived high peaks which are difficult to model. The catchment models therefore performed less well than the lake models. This further
535 suggests that evaluation against observations is likely more important for discharge than for water temperature.

Given that inertia and integration over time were the dominant sources of predictive skill at Sau reservoir and Lake Vansjø, useful hindcasts could already be issued without the use of SEAS5 data. In fact, our workflows show limited sensitivity to boundary conditions over the target season. Hence, future workflows should use selected climatology as forcing data over the target season, in addition to (or instead of) seasonal meteorological prediction. This benchmark forecasting workflow with
540 climatology will likely yield similar or more skillful forecasts, as well as being less time-consuming to set up. Indeed, even with randomly years from the selected SEAS5 data, which can be seen as a highly uncertain climatology, some windows of opportunity are more skillful than with the correct SEAS5 data (Fig. 4). Nevertheless, if seasonal meteorological prediction products become more skillful, they will likely be a real asset for lake seasonal forecasting enabling additional skills through interactions over time.

545 State-of-the-art modeling practices typically involve calibrating hydrologic and lake models against daily observations. Nevertheless, daily observations of water quality are often not available or only cover a fraction of the time of interest. Table 7 illustrates the challenges related to data coverage and model evaluation where many calibration and validation statistics could not be estimated because of the lack of observations. In addition, calibrating to daily data prioritizes model parameterizations which are able to capture daily variability, but not necessarily seasonal or interannual variability, which are more relevant for
550 seasonal forecasting. Calibrating the hydrologic and lake models using seasonal means or medians, in combination with daily data, could solve the observation coverage issue while improving seasonal predictive skill. Nevertheless, ones need to ensure that the seasonal averages are calculated from representative and well-distributed datasets. For Lake Vansjø, this would not have solved the lack of observations in Spring for example, because observations only cover April and May. For Sau and Wupper reservoirs, on the other hand, this would have been possible and potentially improve predictive skills. In any case,
555 having access to more complete, long-term and systematic observations on water temperature, inflow and outflow discharge, including abstraction and over- flows for reservoirs, would facilitate robust model calibration and validation, and likely model

predictive skills. The skill of water quality forecasting tools heavily depends on observation availability. Hence, continued efforts should be put on ensuring that observational programs are suited to providing the information needed by our models (Robson, 2014).

560 5 Conclusion

Lake seasonal forecasts could provide valuable knowledge for water managers to preserve drinking water reserves, as well as ecological and recreational services under increasing pressures from water demand and climate change. Nevertheless, their use is still limited in the water sector. Here we unravel the source of predictive skill of lake seasonal hindcasts at four case-studies across Europe and in Australia, including inflow discharge, surface and bottom water temperature as well as ice-off 565 dates. Through sensitivity analyses, we contribute to the demystification of lake forecasting tools with the long-term objective of facilitating their utilization in the water sector. In Spain, where the seasonal meteorological predictions have negligible skill, the source of predictive skill is mainly catchment and lake inertia. In Norway, where some seasonal meteorological predictions are skillful, the predictive skill is coming from, in decreasing order of importance, inertia, time- and across-variable integration of climate signals through catchment processes, and seasonal meteorological predictions over the target season (SEAS5). In 570 Norway, skillful SEAS5 meteorological hindcasts over specific seasons likely contribute to sustaining the predictive skill from antecedent conditions through to the target season.

Despite its central role in the probabilistic nature of the forecasting workflow, SEAS5 meteorological forcing data contributes to a limited extent to the predictive skill, and often reduces the performance of the hindcasts. Hence, our findings suggest that 575 using a probabilistic ensemble catchment-lake forecast without SEAS5 forcing data is currently likely to yield higher quality forecasts, as demonstrated by hindcasts driven with randomly selected SEAS5 data. Nevertheless, upon skill improvement of the seasonal meteorological forecasts, a small step would be needed to provide more skillful lake forecasts for better water management.

Index of abbreviations (in order of appearance)

SEAS5: seasonal meteorological forecast dataset from the European Centre for Medium Range Weather Forecasts.

580 ERA5: meteorological reanalysis dataset from the European Centre for Medium Range Weather Forecasts.

NSE: Nash–Sutcliffe efficiency coefficient

KGE: Kling-Gupta efficiency coefficient

RMSE: Root-mean squared error

Lake_PO: Lake pseudo-observations of water temperature, inflow discharge and ice-off produced with coupled catchment and 585 lake models forced with ERA5 meteorological data.

Lake_F: Seasonal lake hindcasts of water temperature, inflow discharge and ice-off produced with coupled catchment and lake models forced with SEAS5 meteorological data (25 members).

M0: First lead month

M1–M3: Month 1 to month 3 of the lake forecast, i.e., target season of the lake forecasts.

590 ROCSS: Relative Operating Characteristic Skill Score

RPSS: Ranked Probability Skill Score

FRPSS: Fair (or unbiased) RPSS

$ROCSS_{original}$: ROCSS for Lake_F as compared to reference forecast based on climatology from Lake_PO.

$ROCSS_{obs}$: ROCSS for Lake_F as compared to reference forecast based on local observations.

595 SA: Sensitivity analysis

S-SA: Sensitivity analysis of Lake_F to boundary conditions over the target season (M1–M3)

W-SA: Sensitivity analysis of Lake_F to boundary conditions over the warm-up period

W+M0-SA: Sensitivity analysis of Lake_F to boundary conditions over the period covering the warm-up and first lead month

$ROCSS_S$: ROCSS for Lake_F following S-SA as compared to reference forecast based on climatology from Lake_PO

600 $ROCSS_W$: ROCSS for Lake_F following W-SA as compared to reference forecast based on climatology from Lake_PO

$ROCSS_{W+M0}$: ROCSS for Lake_F following W+M0-SA as compared to reference forecast based on climatology from Lake_PO

OAT-SA: One at a time sensitivity analysis

PPCC: Partial correlation coefficient

605 airT: Surface air temperature

airP: Surface air pressure

cc: Cloud cover

hum: Relative humidity (or dew-point temperature)

sw: short-wave radiation

610 lw: downwelling long-wave radiation

U: U-component of wind speed

V: V-component of wind speed

p: Precipitation

615 **Computer code and models**

v

Acknowledgments

This study contributed to the WATExR project (https://nivanorge.github.io/seasonal_forecasting_watexr/), which is part of ERA4CS, an ERA-NET initiated by JPI Climate, and funded by MINECO-AEI (ES), FORMAS (SE), BMBF (DE), EPA (IE), 620 RCN (NO), and IFD (DK), with co-funding by the European Union (Grant 690462). MINECO-AEI funded this research through projects PCIN-2017-062 and PCIN-2017-092. We thank all water quality and quantity data providers: Ens d'Abastament d'Aigua Ter-Llobregat (ATL, <https://www.atl.cat/es>), SA Water (<https://www.sawater.com.au/>), Wupperverband (www.wupperverband.de), NIVA (www.niva.no) and NVE (<https://www.nve.no/english/>). We acknowledge ECMWF for providing the SEAS5 and ERA5 data.

625 Author contributions

RM, LJ-B, EdE, EJ, KR, LdvL, M-DF, and SH designed the study and provided guidance on modelling and forecasting approaches. FC, LJ-B, MNO, JS, DM, MS, AF, and TM contributed to the modelling, forcing data pre-processing and forecasting. FC, DM, MS, AF performed the sensitivity analyses. FC drafted the manuscript. All authors edited the manuscript.

References

630 Arnal, L., Cloke, H. L., Stephens, E., Wetterhall, F., Prudhomme, C., Neumann, J., Krzeminski, B., & Pappenberger, F. (2018). Skilful seasonal forecasts of streamflow over Europe? *Hydrology and Earth System Sciences*, 22(4), 2057–2072. <https://doi.org/10.5194/hess-22-2057-2018>

Baracchini, T., Wüest, A., & Bouffard, D. (2020). Meteolakes: An operational online three-dimensional forecasting platform for lake hydrodynamics. *Water Research*, 172, 115529. <https://doi.org/10.1016/j.watres.2020.115529>

635 Bazile, R., Boucher, M.-A., Perreault, L., & Leconte, R. (2017). Verification of ECMWF System 4 for seasonal hydrological forecasting in a northern climate. *Hydrology and Earth System Sciences*, 21(11), 5747–5762. <https://doi.org/10.5194/hess-21-5747-2017>

Blottiere, L. (2015). *The effects of wind-induced mixing on the structure and functioning of shallow freshwater lakes in a context of global change*. [Université Paris Saclay]. <https://tel.archives-ouvertes.fr/tel-01258843/document>

640 Butcher, J. B., Nover, D., Johnson, T. E., & Clark, C. M. (2015). Sensitivity of lake thermal and mixing dynamics to climate change. *Climatic Change*, 129(1), 295–305. <https://doi.org/10.1007/s10584-015-1326-1>

Ceglar, A., & Toreti, A. (2021). Seasonal climate forecast can inform the European agricultural sector well in advance of harvesting. *Npj Climate and Atmospheric Science*, 4(1), 1–8. <https://doi.org/10.1038/s41612-021-00198-3>

Coron, L., Thirel, G., Delaigue, O., Perrin, C., & Andréassian, V. (2017). The suite of lumped GR hydrological models in an 645 R package. *Environmental Modelling & Software*, 94, 166–171. <https://doi.org/10.1016/j.envsoft.2017.05.002>

Déqué, M. (2007). Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values. *Global and Planetary Change*, 57(1), 16–26. <https://doi.org/10.1016/j.gloplacha.2006.11.030>

Dokulil, M. T., de Etyo, E., Maberly, S. C., May, L., Weyhenmeyer, G. A., & Woolway, R. I. (2021). Increasing maximum lake surface temperature under climate change. *Climatic Change*, 165(3), 56. <https://doi.org/10.1007/s10584-021-03085-1>

Ducharne, A. (2008). Importance of stream temperature to climate change impact on water quality. *Hydrology and Earth System Sciences*, 12(3), 797–810. <https://doi.org/10.5194/hess-12-797-2008>

Ferro, C. a. T. (2014). Fair scores for ensemble forecasts. *Quarterly Journal of the Royal Meteorological Society*, 140(683), 1917–1923. <https://doi.org/10.1002/qj.2270>

Frías, M. D., Iturbide, M., Manzanas, R., Bedia, J., Fernández, J., Herrera, S., Cofiño, A. S., & Gutiérrez, J. M. (2018). An R package to visualize and communicate uncertainty in seasonal climate prediction. *Environmental Modelling & Software*, 99, 101–110. <https://doi.org/10.1016/j.envsoft.2017.09.008>

Giuliani, M., Crochemore, L., Pechlivanidis, I., & Castelletti, A. (2020). From skill to value: Isolating the influence of end user behavior on seasonal forecast assessment. *Hydrology and Earth System Sciences*, 24(12), 5891–5902. <https://doi.org/10.5194/hess-24-5891-2020>

Greuell, W., Franssen, W. H. P., & Hutjes, R. W. A. (2019). Seasonal streamflow forecasts for Europe – Part 2: Sources of skill. *Hydrology and Earth System Sciences*, 23(1), 371–391. <https://doi.org/10.5194/hess-23-371-2019>

Gutiérrez, J. M., Maraun, D., Widmann, M., Huth, R., Hertig, E., Benestad, R., Roessler, O., Wibig, J., Wilcke, R., Kotlarski, S., San Martín, D., Herrera, S., Bedia, J., Casanueva, A., Manzanas, R., Iturbide, M., Vrac, M., Dubrovsky, M., Ribalaygua, J., ... Pagé, C. (2019). An intercomparison of a large ensemble of statistical downscaling methods over Europe: Results from the VALUE perfect predictor cross-validation experiment. *International Journal of Climatology*, 39(9), 3750–3785. <https://doi.org/10.1002/joc.5462>

Harrigan, S., Prudhomme, C., Parry, S., Smith, K., & Tanguy, M. (2018). Benchmarking ensemble streamflow prediction skill in the UK. *Hydrology and Earth System Sciences*, 22(3), 2023–2039. <https://doi.org/10.5194/hess-22-2023-2018>

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., ... Thépaut, J.-N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>

Hipsey, M. R., Bruce, L. C., Boon, C., Busch, B., Carey, C. C., Hamilton, D. P., Hanson, P. C., Read, J. S., de Sousa, E., Weber, M., & Winslow, L. A. (2019). A General Lake Model (GLM 3.0) for linking with high-frequency sensor data from the Global Lake Ecological Observatory Network (GLEON). *Geoscientific Model Development*, 12(1), 473–523. <https://doi.org/10.5194/gmd-12-473-2019>

Iturbide, M., Bedia, J., Herrera, S., Baño-Medina, J., Fernández, J., Frías, M. D., Manzanas, R., San-Martín, D., Cimadevilla, E., Cofiño, A. S., & Gutiérrez, J. M. (2019). The R-based climate4R open framework for reproducible climate data access and
680 post-processing. *Environmental Modelling & Software*, 111, 42–54. <https://doi.org/10.1016/j.envsoft.2018.09.009>

Jackson-Blake, L. A., Clayer, F., de Eyto, E., French, A. S., Frías, M. D., Mercado-Bettín, D., Moore, T., Puertolas, L., Poole, R., Rinke, K., Shikhani, M., van der Linden, L., & Marcé, R. (2022). Opportunities for seasonal forecasting to support water management outside the tropics. *Hydrology and Earth System Sciences*, 26(5), 1389–1406. <https://doi.org/10.5194/hess-26-1389-2022>

685 Jackson-Blake, L. A., Sample, J. E., Wade, A. J., Helliwell, R. C., & Skeffington, R. A. (2017). Are our dynamic water quality models too complex? A comparison of a new parsimonious phosphorus model, SimplyP, and INCA-P. *Water Resources Research*, 53(7), 5382–5399. <https://doi.org/10.1002/2016WR020132>

Jeppesen, E., Pierson, D., & Jennings, E. (2021). Effect of Extreme Climate Events on Lake Ecosystems. *Water*, 13(3), 282. <https://doi.org/10.3390/w13030282>

690 Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche, S., Decremer, D., Weisheimer, A., Balsamo, G., Keeley, S. P. E., Mogensen, K., Zuo, H., & Monge-Sanz, B. M. (2019). SEAS5: The new ECMWF seasonal forecast system. *Geoscientific Model Development*, 12(3), 1087–1117. <https://doi.org/10.5194/gmd-12-1087-2019>

Jolliffe, I. T., & Stephenson, D. B. (2012). *Forecast Verification: A Practitioner's Guide in Atmospheric Science*. John Wiley & Sons.

695 Labrousse, C., Ludwig, W., Pinel, S., Sadaoui, M., & Lacquement, G. (2020). Unravelling Climate and Anthropogenic Forcings on the Evolution of Surface Water Resources in Southern France. *Water*, 12(12), 3581. <https://doi.org/10.3390/w12123581>

Lledó, Ll., Torralba, V., Soret, A., Ramon, J., & Doblas-Reyes, F. J. (2019). Seasonal forecasts of wind power generation. *700 Renewable Energy*, 143, 91–100. <https://doi.org/10.1016/j.renene.2019.04.135>

Lopez, A., & Haines, S. (2017). Exploring the Usability of Probabilistic Weather Forecasts for Water Resources Decision-Making in the United Kingdom. *Weather, Climate, and Society*, 9(4), 701–715. <https://doi.org/10.1175/WCAS-D-16-0072.1>

Manzanas, R., Frías, M. D., Cofiño, A. S., & Gutiérrez, J. M. (2014). Validation of 40 year multimodel seasonal precipitation forecasts: The role of ENSO on the global skill. *Journal of Geophysical Research: Atmospheres*, 119(4), 1708–1719. <https://doi.org/10.1002/2013JD020680>

705 Marcé, R., Rodríguez-Arias, M. À., García, J. C., & Armengol, J. (2010). El Niño Southern Oscillation and climate trends impact reservoir water quality. *Global Change Biology*, 16(10), 2857–2865. <https://doi.org/10.1111/j.1365-2486.2010.02163.x>

Marino, S., Hogue, I. B., Ray, C. J., & Kirschner, D. E. (2008). A Methodology For Performing Global Uncertainty And
710 Sensitivity Analysis In Systems Biology. *Journal of Theoretical Biology*, 254(1), 178–196. <https://doi.org/10.1016/j.jtbi.2008.04.011>

Mariotti, A., Baggett, C., Barnes, E. A., Becker, E., Butler, A., Collins, D. C., Dirmeyer, P. A., Ferranti, L., Johnson, N. C., Jones, J., Kirtman, B. P., Lang, A. L., Molod, A., Newman, M., Robertson, A. W., Schubert, S., Waliser, D. E., & Albers, J. (2020). Windows of Opportunity for Skillful Forecasts Subseasonal to Seasonal and Beyond. *Bulletin of the American Meteorological Society*, 101(5), E608–E625. <https://doi.org/10.1175/BAMS-D-18-0326.1>

Mercado-Bettin, D., Clayer, F., Shikhani, M., Moore, T. N., Frias, M. D., Jackson-Blake, L., Sample, J., Iturbide, M., Herrera, S., French, A. S., Norling, M. D., Rinke, K., & Marce, R. (2021). Forecasting water temperature in lakes and reservoirs using seasonal climate prediction. *Water Research*, 201, 117286. <https://doi.org/10.1016/j.watres.2021.117286>

Müller, W. A., Appenzeller, C., Doblas-Reyes, F. J., & Liniger, M. A. (2005). A Debiased Ranked Probability Skill Score to Evaluate Probabilistic Ensemble Forecasts with Small Ensemble Sizes. *Journal of Climate*, 18(10), 1513–1523. <https://doi.org/10.1175/JCLI3361.1>

Pagano, T. C., Wood, A. W., Ramos, M.-H., Cloke, H. L., Pappenberger, F., Clark, M. P., Cranston, M., Kavetski, D., Mathevet, T., Sorooshian, S., & Verkade, J. S. (2014). Challenges of Operational River Forecasting. *Journal of Hydrometeorology*, 15(4), 1692–1707. <https://doi.org/10.1175/JHM-D-13-0188.1>

Pechlivanidis, I. G., Crochemore, L., Rosberg, J., & Bosshard, T. (2020). What Are the Key Drivers Controlling the Quality of Seasonal Streamflow Forecasts? *Water Resources Research*, 56(6), e2019WR026987. <https://doi.org/10.1029/2019WR026987>

Pianosi, F., Beven, K., Freer, J., Hall, J. W., Rougier, J., Stephenson, D. B., & Wagener, T. (2016). Sensitivity analysis of environmental models: A systematic review with practical workflow. *Environmental Modelling & Software*, 79, 214–232. <https://doi.org/10.1016/j.envsoft.2016.02.008>

Piccolroaz, S., Healey, N. C., Lengers, J. D., Schladow, S. G., Hook, S. J., Sahoo, G. B., & Toffolon, M. (2018). On the predictability of lake surface temperature using air temperature in a changing climate: A case study for Lake Tahoe (U.S.A.). *Limnology and Oceanography*, 63(1), 243–261. <https://doi.org/10.1002/lno.10626>

Piccolroaz, S., Toffolon, M., & Majone, B. (2013). A simple lumped model to convert air temperature into surface water temperature in lakes. *Hydrology and Earth System Sciences*, 17(8), 3323–3338. <https://doi.org/10.5194/hess-17-3323-2013>

Portele, T. C., Lorenz, C., Dibrani, B., Laux, P., Bliefernicht, J., & Kunstmann, H. (2021). Seasonal forecasts offer economic benefit for hydrological decision making in semi-arid regions. *Scientific Reports*, 11(1), 10581. <https://doi.org/10.1038/s41598-021-89564-y>

Robson, B. J. (2014). State of the art in modelling of phosphorus in aquatic systems: Review, criticisms and commentary. *Environmental Modelling & Software*, 61, 339–359. <https://doi.org/10.1016/j.envsoft.2014.01.012>

Schmid, M., & Read, J. (2022). Heat Budget of Lakes. In T. Mehner & K. Tockner (Eds.), *Encyclopedia of Inland Waters* (Second Edition) (pp. 467–473). Elsevier. <https://doi.org/10.1016/B978-0-12-819166-8.00011-6>

Schmid, M., Hunziker, S., & Wüest, A. (2014). Lake surface temperatures in a changing climate: A global sensitivity analysis. *Climatic Change*, 124(1), 301–315. <https://doi.org/10.1007/s10584-014-1087-2>

745 Soares, M. B., Daly, M., & Dessai, S. (2018). Assessing the value of seasonal climate forecasts for decision-making. *WIREs Climate Change*, 9(4), e523. <https://doi.org/10.1002/wcc.523>

746 Staudinger, M., & Seibert, J. (2014). Predictability of low flow – An assessment with simulation experiments. *Journal of Hydrology*, 519, 1383–1393. <https://doi.org/10.1016/j.jhydrol.2014.08.061>

747 Stefan, H. G., & Preud'homme, E. B. (1993). Stream Temperature Estimation from Air Temperature1. *JAWRA Journal of the American Water Resources Association*, 29(1), 27–45. <https://doi.org/10.1111/j.1752-1688.1993.tb01502.x>

750 Themeßl, M., Gobiet, A., & Leuprecht, A. (2011). Empirical-statistical downscaling and error correction of daily precipitation from regional climate models. *International Journal of Climatology*, 31(10), 1530–1544. <https://doi.org/10.1002/joc.2168>

751 Toffolon, M., Piccolroaz, S., Majone, B., Soja, A.-M., Peeters, F., Schmid, M., & Wüest, A. (2014). Prediction of surface temperature in lakes with different morphology using air temperature. *Limnology and Oceanography*, 59(6), 2185–2202. <https://doi.org/10.4319/lo.2014.59.6.2185>

755 Troccoli, A. (2010). Seasonal climate forecasting. *Meteorological Applications*, 17(3), 251–268. <https://doi.org/10.1002/met.184>

756 Troin, M., Arsenault, R., Wood, A. W., Brissette, F., & Martel, J.-L. (2021). Generating Ensemble Streamflow Forecasts: A Review of Methods and Approaches Over the Past 40 Years. *Water Resources Research*, 57(7), e2020WR028392. <https://doi.org/10.1029/2020WR028392>

760 Werner, M., Cranston, M., Harrison, T., Whitfield, D., & Schellekens, J. (2009). Recent developments in operational flood forecasting in England, Wales and Scotland. *Meteorological Applications*, 16(1), 13–22. <https://doi.org/10.1002/met.124>

765 Wilcke, R. A. I., Mendlik, T., & Gobiet, A. (2013). Multi-variable error correction of regional climate models. *Climatic Change*, 120(4), 871–887. <https://doi.org/10.1007/s10584-013-0845-x>

770 Wood, A. W., Hopson, T., Newman, A., Brekke, L., Arnold, J., & Clark, M. (2016). Quantifying Streamflow Forecast Skill Elasticity to Initial Condition and Climate Prediction Skill. *Journal of Hydrometeorology*, 17(2), 651–668. <https://doi.org/10.1175/JHM-D-14-0213.1>

775 Wuijts, S., Claessens, J., Farrow, L., Doody, D. G., Klages, S., Christophoridis, C., Cvejić, R., Glavan, M., Nesheim, I., Platjouw, F., Wright, I., Rowbottom, J., Graversgaard, M., van den Brink, C., Leitão, I., Ferreira, A., & Boekhold, S. (2021). Protection of drinking water resources from agricultural pressures: Effectiveness of EU regulations in the context of local realities. *Journal of Environmental Management*, 287, 112270. <https://doi.org/10.1016/j.jenvman.2021.112270>

780 Yi, S., Sun, W., Feng, W., & Chen, J. (2016). Anthropogenic and climate-driven water depletion in Asia. *Geophysical Research Letters*, 43(17), 9061–9069. <https://doi.org/10.1002/2016GL069985>

785 Zhu, S., Piotrowski, A. P., Ptak, M., Napiorkowski, J. J., Dai, J., & Ji, Q. (2021). How does the calibration method impact the performance of the air2water model for the forecasting of lake surface water temperatures? *Journal of Hydrology*, 597, 126219. <https://doi.org/10.1016/j.jhydrol.2021.126219>

790 Zhu, S., Ptak, M., Yaseen, Z. M., Dai, J., & Sivakumar, B. (2020). Forecasting surface water temperature in lakes: A comparison of approaches. *Journal of Hydrology*, 585, 124809. <https://doi.org/10.1016/j.jhydrol.2020.124809>