

# Inter- and intra-event rainfall partitioning dynamics of two typical xerophytic shrubs in the Loess Plateau of China

Jinxia An<sup>1,2</sup>, Guangyao Gao<sup>1,2,4</sup>, Chuan Yuan<sup>3</sup>, Bojie Fu<sup>1,2,4</sup>

5

<sup>1</sup>State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

<sup>2</sup>University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup>State Key Laboratory of Subtropical Silviculture, Zhejiang A&F University, Hangzhou  
10 311300, China

<sup>4</sup>National Observation and Research Station of Earth Critical Zone on the Loess Plateau in Shaanxi, Xi'an 710061, China

*Correspondence to:* Guangyao Gao (gygao@rcees.ac.cn)

15

**Abstract.** Rainfall is known as the main water replenishment in dryland ecosystem, and rainfall partitioning by vegetation reshapes the spatial and temporal distribution patterns of rainwater entry into the soil. The dynamics of rainfall partitioning have been extensively studied at the inter-event scale, yet very few studies have explored its finer intra-event dynamics and the relating driving factors for shrubs. Here, we conducted a concurrent in-depth investigation of  
20 all rainfall partitioning components at inter- and intra-event scales for two typical xerophytic shrubs (*Caragana korshinskii* and *Salix psammophila*) in the Liudaogou catchment of the Loess Plateau, China. The event throughfall (TF), stemflow (SF), and interception loss (IC) and their temporal variations within the rainfall event as well as the meteorological factors and

25 vegetation characteristics were systematically measured during the 2014-2015 rainy seasons. The *C. korshinskii* had significantly higher SF percentage (9.2%) and lower IC percentage (21.4%) compared to *S. psammophila* (3.8% and 29.5%, respectively) ( $p < 0.05$ ), but their TF percentages were not significantly different (69.4% vs. 66.7%). At the intra-event scale, TF and SF of *S. psammophila* was initiated (0.1 vs. 0.3 h and 0.7 vs. 0.8 h) and peaked (1.8 vs. 2.0 h and 2.1 vs. 2.2 h) more quickly, and TF of *S. psammophila* lasted longer (5.2 vs. 4.8 h), delivered more intensely (4.3 vs. 3.8 mm·h<sup>-1</sup>), whereas SF of *C. korshinskii* lasted longer (4.6 vs. 4.1 h), delivered more intensely (753.8 vs. 471.2 mm·h<sup>-1</sup>). For both shrubs, rainfall amount was the most significant factor influencing inter-event rainfall partitioning, and rainfall intensity and duration controlled the intra-event TF and SF variables. The *C. korshinskii* with larger branch angle, more small branches and smaller canopy area, has an advantage to produce stemflow more efficiently over *S. psammophila*. The *S. psammophila* has lower canopy water storage capacity to generate and peak throughfall and stemflow earlier, and it has larger aboveground biomass and total canopy water storage of individual plant to produce higher interception loss compared to *C. korshinskii*. These findings contribute to the fine characterization of shrub-dominated eco-hydrological processes, and improve the accuracy of water balance estimation in dryland ecosystem.

## 1 Introduction

Rainfall is known as the main replenishment of water resources in arid and semi-arid areas, and water resource is the key factor limiting the function of arid ecosystems (Chesson et al.,

2004; Cayuela et al., 2018; Magliano et al., 2019a). Before entering into soil, rainfall is redistributed by plant canopies into throughfall (TF, diffuse water input), stemflow (SF, point water input), and interception loss (IC, evaporation). The sum of TF and SF is defined as "net rainfall". Differences in the distribution of net rainfall caused by plant canopy interception alter the spatial and temporal patterns of rainfall entry into the soil (Martinez-Meza and Whitford, 1996; Li et al., 2009; Van Stan II et al., 2020), and further profoundly affect the water use efficiency of vegetation and ecosystem sustainability (Xu and Li, 2006; Lacombe et al., 2018; Molina et al., 2019). In addition, net rainfall could regulate vegetation physiological metabolic processes through nutrient enrichment (Levia and Frost, 2003; Zhang et al., 2016; Van Stan II et al., 2017; Tonello et al., 2021), ultimately affecting the carbon balance of ecosystems (Chu et al., 2018; Jia et al., 2016). In light of the important role of rainfall partitioning in regulating soil water dynamics and vegetation patch pattern, investigations of the rainfall partitioning dynamics are imperative for a better understanding of the soil-water-vegetation relationships (Molina et al., 2019; Van Stan II et al., 2020; Zhang et al., 2021a).

Studies on rainfall partitioning have been broadly carried out in different climatic zones and various types of vegetation (Gordon et al. 2020; Rivera and Van Stan II, 2020; Zhang et al., 2021b; Yue et al., 2021). Based on a comprehensive global synthesis, Yue et al. (2021) concluded that most TF and SF observations were measured in forests ( $n = 718$  and  $n = 816$ , respectively), and that in shrublands was scarce ( $n = 43$  and  $n = 63$ , respectively), which was mainly due to that the shrubs have multiple branches and the rainfall partitioning of shrubs is difficult to be measured compared to forests. Shrubs are the dominant vegetation type in

drylands, providing important ecosystem goods and services (Levia and Frost, 2003; Llorens and Domingo, 2007; Soulsby et al., 2017). However, the lack of information on the detailed dynamics of rainfall partitioning processes induced by shrubs due to limited studies hinders us  
70 form a clear understanding of shrubs' eco-hydrological role in shaping and sustaining drylands.

Most of the existing studies on the rainfall partitioning by shrub are based on the inter-event scale (Garcia-Estringana et al., 2010; Magliano et al., 2019a). Magliano et al. (2019a) synthesized that for 27 shrub species in drylands, the mean event-based SF%, TF%, and IC% were 9.4%, 63.0% and 27.6%, respectively. Rainfall partitioning by shrubs has been reported  
75 to be determined by various meteorological factors, including rainfall amount, duration and intensity and others (Levia and Frost, 2003; Magliano et al., 2019b) and by vegetation structure characteristics (Martinez-Meza and Whitford, 1996; Garcia-Estringana et al., 2010; Yue et al., 2021). Take the later for example, trees/shrubs with smooth barks, more branches and vertical branching had advantages on SF generation (Honda et al., 2015; Magliano et al., 2019a;  
80 Whitworth-Hulse et al., 2020b), and a simple vegetation structure and low canopy density are generally corresponding to a relatively high TF rate and low IC rate (Soulsby et al., 2017; Yue et al., 2021). The complexity of shrub structure poses challenges to understand the causes of rainfall partitioning dynamics under different meteorological conditions, and it is necessary to substantially explore the differences of rainfall partitioning dynamics and main influencing  
85 factors among different shrub species (Levia et al., 2010; Sadeghi et al., 2020).

In addition to the inter-event studies, a detailed understanding of shrub rainfall partitioning dynamics at the intra-event scale is essential for better understanding of soil moisture

distribution and the hydrological cycle in arid regions. To understand the temporal fluctuations of shrub rainfall partitioning and its importance to hydrological processes, data with high temporal resolution are required (Levia et al., 2010; Levia and Germer, 2015; Cayuela et al., 2018; Zhang et al., 2021a). For instance, Zhang et al. (2018) investigated the spatial-temporal pattern of TF (at 10-min intervals) of *C. korshinskii* in arid area of northern China, highlighting the importance of recording high temporal resolution data. They found that temporal heterogeneity of rainfall clearly affected the spatiotemporal dynamics of TF beneath shrub canopies and wind directions were the main factor affecting TF in different radial directions. Yuan et al. (2019) described the branch SF variability of *C. korshinskii* and *S. psammophil*, and they showed that intra-event branch SF variability of xerophytic shrubs temporally depended on rainfall characteristics, and lag times were longer and more rainfall amount was required to initiate branch SF for *C. korshinskii* than *S. psammophila*. It can be found that those studies on temporal dynamics of shrub rainfall partitioning only explored the single-element process (throughfall or stemflow), and ignored interception loss. Concurrent investigation on all rainfall partitioning components and the associated influencing factors at the intra-event scale have rarely been reported.

This study was designed at the event and process scales to investigate inter-event and intra-event rainfall partitioning variability, based on field measurements on two dominant xerophytic shrubs (*C. korshinskii* and *S. psammophila*) during the rainy seasons of 2014-2015 in the Loess Plateau of China. This study integrated the inter-event and intra-event dynamics of rainfall partitioning by combining TF, SF and IC at the individual plant scale. We mainly

seek to (a) compare the dynamic processes of rainfall partitioning between the two shrubs at  
110 both inter-event and intra-event scales, and (b) elucidate the effects of rainfall characteristics  
and vegetation structure characteristics on rainfall partitioning at both scales. This work can  
help better understand and predict the role of shrubs in harvesting rainfall and recharging soil  
moisture, and thus enhance the understanding of eco-hydrological processes of shrubs.

## 2 Materials and methods

### 115 2.1 Site description and experimental design

This study was carried out in the Liudaogou catchment (110°21'-110°23' E, 38°46'-38°5' N) in  
Shenmu county, Shaanxi Province of China (Fig. 1a). The Liudaogou catchment (6.9 km<sup>2</sup>,  
altitude from 1094 to 1273 m) is located between the northern Loess Plateau and the south  
fringe of Mu Us sandy land in North China. This region is characterized by a moderate  
120 temperate continental climate with well-defined rainy and dry seasons. The mean annual  
rainfall is 437 mm ranging between 109-891 mm (dataset between 1971-2013), and the  
potential evaporation is 1337 mm yr<sup>-1</sup> (Jia et al., 2013). Approximately 70-80% of the rainfall  
events are concentrated in the warm months between July and September and most of them  
occur in the form of torrential rain (Yang et al., 2019). The Liudaogou catchment was  
125 characterized by the natural arid scrub steppe before it was artificially vegetated in the past 20  
or 30 years for soil and water conservation, windbreak and sand fixation. The main land use  
types include artificial grassland, artificial shrub and farmland, and the main vegetation species  
are *Stipa bungeana*, *C. korshinskii* and *S. psammophila*, which are widely distributed in the  
arid and semiarid areas of northwestern China (Yuan et al., 2019).

130 Two representative xerophytic shrubs, *C. korshinskii* and *S. psammophila* with 20 years  
old, were used for the study. Both species are multiple-stemmed deciduous perennial shrubs  
with inverted cone crowns and without trunks. They have minimal nutrient requirements,  
extensive adaptability and strong stress resistance, which makes them superior in adapting to  
resource-poor environments. According to the documentation of *Flora of China* (Chao and  
135 Gong, 1999; Liu et al., 2010) and field observations, the *S. psammophila* has an odd number  
of strip-shaped leaves with 2-4 mm in width and 40-80 mm in length, and the *C. korshinskii*  
has pinnate compound leaves arranged opposite or sub-opposite with 6-10 cm in length, and  
each pinna has 5 to 8 pairs of ovate leaflets (7-8 mm in length and 2-5 mm in width). We  
established two plots (one for *C. korshinskii* and the other for *S. psammophila*) at the  
140 southwestern catchment for field observation (Fig. 1a). The two plots share similar stand  
conditions, with the sizes of 3294 m<sup>2</sup> and 4056 m<sup>2</sup>, elevations of 1179 m and 1207 m, and  
slopes of 13° and 18°, respectively. The distance between the two plots do not exceed 1.5 km.

## 2.2 Field measurements

### 2.2.1 Measurements of rainfall and meteorological factors

145 This study focused on the individual shrub rainfall partitioning of *C. korshinskii* and *S.*  
*psammophila* during the 2014-2015 rainy seasons. Gross rainfall was measured using one  
tipping bucket rain gauge (TBRG, with 186.3 cm<sup>2</sup> collection area) (Onset® RG3-M, Onset  
Computer Corp., USA) in the open areas, which was recorded every 0.2 mm by a datalogger  
(Fig. 1b). The rainfall characteristics, e.g., rainfall amount (RA, mm), rainfall duration (RD, h),  
150 rainfall interval (RI, h), average rainfall intensity (I, mm·h<sup>-1</sup>), rainfall intensity at 10-min

interval ( $I_{10}$ ,  $\text{mm}\cdot\text{h}^{-1}$ ) were calculated accordingly. For  $I_{10}$ , the one after the onset of rainfall is defined as  $I_{10\_b}$  ( $\text{mm}\cdot\text{h}^{-1}$ ), and the maximum  $I_{10}$  during the rainfall process is defined as  $I_{10\_max}$  ( $\text{mm}\cdot\text{h}^{-1}$ ). As the TBRG has a resolution of 0.2 mm, we define a single rainfall event as one that is greater than 0.2 mm and not raining for at least 4 hours apart (Iida et al., 2012). A meteorological station was set up at the experimental plot to record wind speed (WS,  $\text{m}\cdot\text{s}^{-1}$ ) and wind direction (WD,  $^{\circ}$ ) (Model 03002, R. M. Young Company, Traverse City, Michigan, USA), air temperature (T,  $^{\circ}\text{C}$ ) and relative humidity (H, %) (Model HMP 155, Vaisala, Helsinki, Finland), which were logged at 10-min interval by a data logger (Model CR1000, Campbell Scientific, Inc., USA).

### 160 **2.2.2 Measurements of vegetation characteristics**

Three representative shrub plants with similar crown heights and crown areas were selected in each shrub species (Table 1). Based on plot investigation, the vegetation traits at the scale of single plant and branch were measured. For each plant, we measured shrub height (SH, m) with a graduated telescopic stick, counted the number of branches (NB), and calculated the projected canopy area (CA,  $\text{m}^2$ ) by measuring canopy diameter following the south-north and east-west direction. The total number of branches was 143 and 218 for selected *C. korshinskii* and *S. psammophila* plants, respectively. For each branch, we measured branch length (BL, cm) with a measuring tape, branch angle (BA,  $^{\circ}$ ) with a pocket geologic compass, and branch diameter (BD, mm) with a vernier caliper to calculate the total basal area of the shrub (TBA,  $\text{m}^2$ ). Thus, four BD categories (0-10, 10-15, 15-20 and  $> 20$  mm) were defined to ensure the appropriate branch amounts within each category. The measured vegetation traits of *C. korshinskii* and *S.*



*psammophila* plants are shown in Table 1.

Water storage capacity of the canopy is a key factor in determining the amount of interception loss (Levia and Herwitz, 2005; Garcia-Estringana et al., 2010) and SF yield (Van Stan II et al., 2020). We selected 10 representative branches for each shrub species outside the stands, to determine the canopy water storage capacity (C, mL/g) using water immersion method (Garcia-Estringana et al., 2010; Wang et al., 2012). The C was calculated as the difference between saturated weight and fresh weight divided by the dry biomass of the selected branch. The *C. korshinskii* and *S. psammophila* had a C of 0.85 mL/g and 0.38 mL/g, respectively. In addition, we estimated the total dry aboveground biomass of single plant (TB) for each species according to the allometric growth model developed by Yuan et al. (2017) in the same study area. The total canopy water storage of single plant ( $C_m = TB \text{ times } C$ ) was calculated to represent the amount of rainfall absorbed by the shrub canopy during the rainfall event (Table 1).

### 2.2.3 Measurements of inter-event rainfall partitioning

Manual rain gauges (314.12 cm<sup>2</sup> collection area) were used to measure event TF at eight radial directions (E, SE, S, SW, W, NW, N, NE) beneath each shrub canopy (Fig. 1b). For *C. korshinskii*, eight TF gauges were placed under each *C. korshinskii* plant with 50 cm distance from the base of stems in the eight directions. For *S. psammophila*, twenty TF gauges were placed under each plant, with twelve of them placed in 50 cm, 100 cm, and 150 cm distances from the base of stems in four directions (E, S, W, and N), and eight of them placed in 75 cm and 125 cm distances in the other four directions (SE, SW, NW, and NE). If the rainfall ended

during the daytime, we completed the collection of TF samples within two hours after the end of rainfall. If the rainfall ended at night, we completed the collection of samples as early as possible in the next day to avoid evaporation.

A total of 53 branches of *C. korshinskii* (17, 21, 7, 8 for BD categories of 0-10, 10-15, 15-20 and > 20 mm, respectively) and 98 branches of *S. psammophila* (20, 30, 20 and 28 for BD categories of 0-10, 10-15, 15-20, and > 20 mm, respectively) were used to determine SF yield. Funnels constructed of flexible aluminum foil plates were used to collect SF (Fig. 1b). The funnel was fixed to each branch near the base and sealed with neutral silicone caulk, and a 0.5 cm diameter PVC hose was attached vertically to transport SF from the funnel to a container with a lid (SF gauges) with minimum travel time.

#### **2.2.4 Measurements of intra-event rainfall partitioning**

Among the selected plants, one *C. korshinskii* and one *S. psammophila* plant were selected to record the volume and timing of TF and SF with TBRGs at intra-event scale. A TBRG was installed in each of four radial directions (E, S, W, N) beneath the shrub canopy of each species, to measure the temporal variations of TF within the rainfall event (Fig. 1b). And for each species, we selected six representative branches to characterize intra-event SF dynamics through strict selection conditions, including no crossover between the experimental branch and adjacent branches, no inflection point from the tip to the base of the branch, and easy of measurement. These branches were distributed across the four BD categories (0-10, 10-15, 15-20, and > 20 mm, respectively). Due to missing or incomplete measured data, four branches were finally identified for each species, located in each of the four BD categories to measure

intra-event SF (6.7, 13.5, 18.6, and 22.1 mm for *C. korshinskii* and 7.2, 14.4, 18.2, and 31.3  
215 mm for *S. psammophila*). SF TBRGs were covered with the polyethylene films to prevent the  
accessing of throughfall and splash (Fig. 1b).

## 2.3 Rainfall partitioning calculations

### 2.3.1 Inter-event rainfall partitioning calculations

For each individual shrub, we measured TF volume for each TF gauges, averaged them, and  
220 then converted the volume into TF depth (TF<sub>d</sub>, mm) at each rainfall event. And the percentage  
of TF (TF%, %) was calculated by dividing TF<sub>d</sub> by the RA, and the average TF intensity (TFI,  
mm·h<sup>-1</sup>) was calculated by dividing TF<sub>d</sub> by the TF duration (TFD, h). The TFD was recorded  
by TF TBRGs.

The inter-event SF yield was defined as the total SF volume of a single plant in a rainfall  
225 event. The SF volumes measured on the selected branches were averaged to obtain the average  
volume of SF on the branch scale, which multiply the number of branches to obtain the total  
SF volume from the plant. The shrub-scale SF equivalent water depth (SF<sub>d</sub>, mm) and the  
average SF intensity (SFI, mm·h<sup>-1</sup>) were calculated. The percentage of SF (SF%, %) was  
converted by dividing SF<sub>d</sub> by the RA. The SF<sub>d</sub> and SFI were calculated by the following  
230 equations (Hanchi and Rapp, 1997; Levia and Germer, 2015):

$$SF_d = (\overline{SF_b} \times n) / (1000 \times CA) \quad (1)$$

$$SFI = (\overline{SF_b} \times n) / (10 \times TBA \times SFD) \quad (2)$$

where  $\overline{SF_b}$  (ml) is the average volume of SF on the branch scale,  $n$  is the number of branches  
of individual plant,  $CA$  (m<sup>2</sup>) is the canopy area of individual plant,  $TBA$  (cm<sup>2</sup>) is the total basal

235 area of individual plant, and SFD is SF duration (h) recorded by SF TBRGs. The parameters  
1000 and 10 are the unit conversion factor.

The IC depth (IC<sub>d</sub>, mm) and percentage of IC (IC%, %) were estimated as:

$$IC_d = RA - TF_d - SF_d \quad (3)$$

$$IC\% = 100\% - TF\% - SF\% \quad (4)$$

240 The above inter-event rainfall partitioning variables and their explanations are  
summarized in Table 2.

### 2.3.2 Intra-event rainfall partitioning calculations

The TF and SF volume and timing within rainfall event were automatically recorded at dynamic  
intervals between neighboring TBRG tips (0.2 mm). To better reflect fluctuations in rainfall  
245 partitioning components, the intra-event TF and SF data were aggregated every 10 minutes.  
The four TF depths recorded by TBRGs were averaged to obtain the average TF depth at 10-  
min interval (TF<sub>d10</sub>, mm). The TF intensity at 10-min interval (TFI<sub>10</sub>, mm·h<sup>-1</sup>) was calculated  
by dividing the TF<sub>d10</sub> by the 10 min. Meanwhile, SF depth (SF<sub>d10</sub>, mm) and SF intensity at 10-  
min interval (SFI<sub>10</sub>, mm·h<sup>-1</sup>) were calculated as:

$$250 \quad SF_{d10} = \sum_{j=1}^4 (186.3 \times SF_{RG,j} \times n_j) / (100 \times CA) \quad (5)$$

$$SFI_{10} = \sum_{j=1}^4 (186.3 \times SF_{RG,j} \times n_j) / (TBA \times 1/6) \quad (6)$$

where SF<sub>RG,j</sub> (mm) is the SF depth of the selected jth category branch recorded by TBRG at  
10-min interval (1/6 h), n<sub>j</sub> is the number of branches in the jth category of single plant, 4 is the  
number of BD category (0-10, 10-15, 15-20, and > 20 mm), and 186.3 (cm<sup>2</sup>) is the collection

255 area of TBRG. The product of  $SF_{RG, j}$  and 186.3 is the SF volume from the branch. The parameter 100 is the unit conversion factor.

Based on the calculated  $TFI_{10}$  and  $SFI_{10}$ , the maximum TF and SF intensity at 10-min interval ( $TFI_{10\_max}$  and  $SFI_{10\_max}$ , respectively,  $mm \cdot h^{-1}$ ) of each rainfall event can be determined. The descriptive variables for the intra-event rainfall partitioning also include the lag times of TF or SF corresponding to the rainfall event. Based on the temporal data recorded by TBRGs (between neighboring tips), the following variables were calculated:  $LG_{TF}$  and  $LG_{SF}$  (h), the time lag of TF and SF generation after the start of rainfall, respectively;  $LM_{TF}$ ,  $LM_{SF}$  and  $LM_R$  (h), the time lag of  $TFI_{10\_max}$ ,  $SFI_{10\_max}$  and  $I_{10\_max}$  relative to the onset of rainfall, respectively; and  $LE_{TF}$  and  $LE_{SF}$  (h), the time lag of TF and SF ending after the end of rainfall. The intra-  
260 event rainfall partitioning variables and their explanations are summarized in Table 2.

## 2.4 Statistical analysis

Independent-samples T-tests were used to analyze differences in rainfall partitioning parameters between *C. korshinskii* and *S. psammophila* at both inter-event and intra-event scales. To detect the effects of meteorological factors on rainfall partitioning, Pearson  
270 correlation analysis was used to test the significance between rainfall partitioning parameters and meteorological factors at the two scales. The significant correlated factors were double-checked by partial correlation analysis to determine their individual effects on rainfall partitioning components. Stepwise regression of these indicators was performed by analytical tests at the 0.05 level of significance to select the most influential factors on rainfall partitioning  
275 variables at inter-event and intra-event scales, and the corresponding quantitative relationships

were established based on a qualifying level of significance ( $p < 0.05$ ) and the highest coefficient of determination ( $R^2$ ). Significance levels were set at 95% confidence intervals. Data analysis was performed using SPSS 21.0, Origin 2018, and Excel 2019.

### 3 Results

#### 280 3.1 Inter-event variations of rainfall partitioning

##### 3.1.1 Characteristics of inter-event rainfall partitioning variables

A total of 38 rainfall events were recorded for rainfall partitioning measurements, including 20 events (215.4 mm) in 2014 and 18 events (205.6 mm) in 2015, which accounted for 75.2% and 75.0% of total rainfall amount during the experimental period in 2014 and 2015, respectively (Fig. 2a). The RA ranged from 1.2-41.9 mm with an average of  $11.1 \pm 8.8$  mm (mean  $\pm$  standard). In general, rainfall events were unevenly distributed in terms of RA. Approximately 34.2% of rainfall events were smaller than 5 mm, 26.3% within 5-10 mm, 26.3% within 10-20 mm, and 13.2% larger than 20 mm, representing 8.8%, 17.5%, 36.3%, and 37.4% of the total rainfall amount, respectively (Fig. 2a). The average I varied from  $0.2 \text{ mm}\cdot\text{h}^{-1}$  to  $35.1 \text{ mm}\cdot\text{h}^{-1}$  with an average of  $6.0 \pm 1.3 \text{ mm}\cdot\text{h}^{-1}$ , and approximately 76.3% of the events was  $< 5 \text{ mm}\cdot\text{h}^{-1}$ , 13.2% was  $5\text{--}10 \text{ mm}\cdot\text{h}^{-1}$ , and 10.5% was  $> 10 \text{ mm}\cdot\text{h}^{-1}$ .  $I_{10\_max}$  ranged from  $1.2 \text{ mm}\cdot\text{h}^{-1}$  to  $68.4 \text{ mm}\cdot\text{h}^{-1}$  with an average of  $13.7 \pm 2.7 \text{ mm}\cdot\text{h}^{-1}$ , and approximately 42.1% of the events was  $< 5 \text{ mm}\cdot\text{h}^{-1}$ , 23.7% was  $5\text{--}10 \text{ mm}\cdot\text{h}^{-1}$ , and 34.2% was  $> 10 \text{ mm}\cdot\text{h}^{-1}$ . The RD ranged from 0.2 h to 28.9 h and averaged  $5.3 \pm 1.0$  h. The RD of most rainfall events was less than 5 h (68.4%), and only 5 rainfall events had RD greater than 10 h.

The  $TF_d$  for *C. korshinskii* ranged from 0.7 mm to 31.2 mm (coefficient of variation, CV

= 87.5%) with corresponding TF% ranging from 54.0 to 80.3% (CV = 10.6%) across the 38 events (Fig. 2b). The TF<sub>d</sub> values for *S. psammophila* were 0.4-33.4 mm (CV = 96%) and 28.5-82.7% (CV = 21.5%), respectively (Fig. 2c). The SF<sub>d</sub> for *C. korshinskii* ranged from 0.04 mm to 6.1 mm (CV = 106.6%), with corresponding SF % of 2.0-14.5% (CV = 34.2%) (Fig. 2b). The comparable SF<sub>d</sub> values for *S. psammophila* varied from 0.01 mm to 2.2 mm (CV = 98.6%) and 0.7-5.9% (CV = 38.9%), respectively (Fig. 2c). The IC<sub>d</sub> values for *C. korshinskii* varied from 0.5 mm to 2.9 mm (CV = 43.9%), with corresponding IC% of 5.7-40.8% (CV = 47.3%) (Fig. 2b), and the comparable values were 0.8-5.7 mm (CV = 44.8%) and 12.1-70.8% (CV = 53.3%) for *S. psammophila*, respectively (Fig. 2c). For *C. korshinskii*, TF represented the largest component of all rainfall events, while for *S. psammophila*, SF represented the smallest component of all rainfall events (Figs. 2 b and 2c).

The percentages of TF, SF, and IC in rainfall partitioning between two species are shown in Fig. 3. There was no significant difference ( $p > 0.05$ ) in average TF% between *C. korshinskii* ( $69.4 \pm 7.4\%$ ) and *S. psammophila* ( $66.7 \pm 14.6\%$ ). The SF% was significantly higher ( $p < 0.05$ ) for *C. korshinskii* ( $9.2 \pm 3.2\%$ ) than *S. psammophila* ( $3.8 \pm 1.5\%$ ) (Fig. 3b). The IC% was significantly lower ( $p < 0.05$ ) for *C. korshinskii* ( $21.4 \pm 10.2\%$ ) than *S. psammophila* ( $29.5 \pm 15.9\%$ ) (Fig. 3c). The variations of TF% and IC% among the rainfall events were greater for *S. psammophila*, but that of SF% was smaller compared to *C. korshinskii* (Fig. 3).

### 3.1.2 Relationships between inter-event rainfall partitioning variables and meteorological factors

Correlation analysis indicated that meteorological factors had a similar effect on rainfall

partitioning for the two species. Stepwise regression analysis identified that the SF parameters (SF<sub>d</sub> and SF%), TF parameters (TF<sub>d</sub> and TF%) and IC parameters (IC<sub>d</sub> and IC%) were all  
320 mainly controlled by RA. Following RA, the influences of rainfall intensity (I, I<sub>10\_max</sub>) were also significant ( $p < 0.05$ ). However, the other meteorological factors (RD, RI, WS, WD, T, H) had no significant effect on rainfall partitioning ( $p > 0.05$ ).

Significantly positive and linear relationships were found between TF<sub>d</sub> and RA for both  
*C. korshinskii* and *S. psammophila* (Fig. 4a). According to the regression equations, the  
325 threshold of rainfall amount for TF generation was 0.8 and 1.1 mm for *C. korshinskii* and *S. psammophila*, respectively. The TF% increased with increasing RA as an exponential function (Fig. 4b). When the RA reached 20 mm, the increasing of TF% became stabilized, and TF% of *C. korshinskii* and *S. psammophila* reached 79.2% and 80.0%, respectively. The SF<sub>d</sub> also had a significantly positive and linear relationship with RA for the two species (Fig. 4c). When RA  
330 was greater than 1.7 mm and 2.2 mm, *C. korshinskii* and *S. psammophila* began to produce SF, respectively. The SF% increased exponentially with increasing RA, and SF% of *C. korshinskii* was always higher than that of *S. psammophila*. The SF% approximately tended to be constant at 12.2% and 5.5% as  $RA \geq 20$  mm for *C. korshinskii* and *S. psammophila*, respectively (Fig. 4d). The IC<sub>d</sub> was also positively correlated with RA (Fig. 4e). However, IC% decreased  
335 exponentially with incremental RA, and IC% of *S. psammophila* was always higher than that of *C. korshinskii* (Fig. 4f). When RA reached 20 mm, IC% approximately tended to be constant at 9.0% and 14.5% for *C. korshinskii* and *S. psammophila*, respectively.



## 3.2 Intra-event variations of rainfall partitioning

### 3.2.1 Characteristics of intra-event rainfall partitioning variables

340 The intra-event TF and SF were well synchronized with rainfall process, in terms of the shape, number and location of the intensity peaks for both *C. korshinskii* and *S. psammophila*, which was vividly demonstrated at representative four rainfall events in Fig. 5. The SF intensity (SFI<sub>10</sub>) was much higher than TF intensity (TFI<sub>10</sub>) and rainfall intensity (I<sub>10</sub>) for both *C. korshinskii* and *S. psammophila*, whereas TFI<sub>10</sub> was less than or equal to I<sub>10</sub>. As expected, IC was the main  
345 component at the initial stage of rainfall, and then TF was the major component ( $\geq 50\%$ ) for rainfall partitioning (Fig. 5). As the continue of rainfall, the cumulative amount of TF, SF and IC increases, and finally the percentage of cumulative amounts (TF, SF, and IC) over the cumulative rainfall stabilized near a fixed value. The TF and SF generation thresholds measured using the TBRGs were  $0.4 \pm 0.2$  mm and  $1.0 \pm 0.7$  mm for *C. korshinskii*, and  $0.3 \pm$   
350  $0.1$  mm and  $0.7 \pm 0.3$  for *S. psammophila*, respectively. They were expected to be both smaller than the thresholds derived from the regression equation between TF<sub>d</sub> (or SF<sub>d</sub>) and RA aforementioned which assume that TF and SF start after the canopy is fully wet. This further demonstrates the importance of high-resolution data in rainfall partitioning studies.

Fig. 6 describes the difference in average intra-event TF and SF variables between *C*  
355 *korshinskii* and *S. psammophila*. Although there were no statistically significant differences between the two species in intensities, durations, or the lag time of TF and SF, some trends were observed. The TFI and TFI<sub>10\_max</sub> of both species were similar to I ( $6.0 \pm 1.3$  mm/h) and I<sub>10\_max</sub> ( $13.7 \pm 2.7$  mm/h), respectively. In contrast, SFI and SFI<sub>10\_max</sub> were significantly greater

than I and  $I_{10\_max}$ , respectively. Specifically, TFI and  $TFI_{10\_max}$  of *C. korshinskii* were  $3.8 \pm 1.2$   
360  $mm \cdot h^{-1}$  and  $13.3 \pm 4.9 mm \cdot h^{-1}$ , respectively, which were slighter lower than that of *S.*  
*psammophila* ( $4.3 \pm 1.5 mm \cdot h^{-1}$  and  $14.6 \pm 5.5 mm \cdot h^{-1}$ , respectively) (Fig. 6a). The SFI and  
 $SFI_{10\_max}$  of *C. korshinskii* ( $753.8 \pm 208.0 mm \cdot h^{-1}$  and  $3627.2 \pm 1424.7 mm \cdot h^{-1}$ , respectively)  
were higher than those of *S. psammophila* ( $471.2 \pm 170.2 mm \cdot h^{-1}$  and  $1424.8 \pm 538.3 mm \cdot h^{-1}$ ,  
respectively) (Fig. 6b).

365 Furthermore, a time lag was usually observed between the onset of rainfall and the  
generation of TF ( $LG_{TF}$ ) and SF ( $LG_{SF}$ ). Similarly, there is a time lag between rainfall and TF  
or SF in terms of the time to reach maximum intensity (LM) and the time to end (LE). The *S.*  
*psammophila* had a shorter lag time than *C. korshinskii* in terms of TF ( $LG_{TF}$ :  $0.1 \pm 0.04 h$  vs.  
 $0.3 \pm 0.1 h$ ) and SF production ( $LG_{SF}$ :  $0.7 \pm 0.3 h$  vs.  $0.8 \pm 0.3 h$ ), and their reaching maximum  
370 intensity ( $LM_{TF}$ :  $1.8 \pm 0.8 h$  vs.  $2.0 \pm 0.6 h$ ;  $LM_{SF}$ :  $2.1 \pm 0.7 h$  vs.  $2.2 \pm 0.5 h$ ) (Figs. 6c and 6d).  
However, the *S. psammophila* had longer TF duration ( $5.2 \pm 1.4 h$  vs.  $4.8 \pm 1.4 h$ ) and  $LE_{TF}$   
( $0.2 \pm 0.1 h$  vs.  $0.1 \pm 0.1 h$ ) than *C. korshinskii* (Fig. 6c). Conversely, the SF duration and  $LE_{SF}$   
in *C. korshinskii* ( $4.6 \pm 1.4 h$  and  $0.4 \pm 0.1 h$ , respectively) were longer than those in *S.*  
*psammophila* ( $4.1 \pm 1.3 h$  and  $0.2 \pm 0.2 h$ , respectively) (Fig. 6d). The above differences in TF  
375 and SF variables indicate that *S. psammophila* should be more conducive to generate TF than  
*C. korshinskii*, while *C. korshinskii* should be more conducive to produce SF than *S.*  
*psammophila*.

### 3.2.2 Relationships between intra-event rainfall partitioning variables and meteorological factors

380 Similar relationships existed between intra-event rainfall partitioning variables and meteorological factors for two species. For both shrubs, rainfall intensity (I, I<sub>10\_max</sub>, and I<sub>10\_b</sub>) and RD were the main influencing factors affecting intra-event TF variables (Fig. 7) and SF variables (Fig. 8). While the effects of other meteorological factors (RD, RI, WS, WD, T, H) on TF and SF variables within the event were not significant ( $p > 0.05$ ). The TFI, TFI<sub>10\_max</sub>, 385 LM<sub>TF</sub>, and TFD were linearly correlated with I, I<sub>10\_max</sub>, LM<sub>R</sub>, and RD, respectively, while LG<sub>TF</sub> was power functionally correlated with I<sub>10\_b</sub> ( $p < 0.05$ ). The TF intensities (TFI and TFI<sub>10\_max</sub>) of *S. psammophila* increased faster with rainfall intensities (I and I<sub>10\_max</sub>) than that of *C. korshinskii*. The SFI, SFI<sub>10\_max</sub>, LM<sub>SF</sub>, and SFD were also linearly correlated with I, I<sub>10\_max</sub>, LM<sub>R</sub>, and RD, respectively ( $p < 0.05$ ). The LG<sub>SF</sub> was power functionally correlated with I<sub>10\_b</sub> 390 ( $p = 0.14$  and  $p = 0.16$  for *C. korshinskii* and *S. psammophila*, respectively), which was weaker than the correlation between LG<sub>TF</sub> with I<sub>10\_b</sub>. The SF intensities (SFI and SFI<sub>10\_max</sub>) of *C. korshinskii* increased with rainfall intensities (I and I<sub>10\_max</sub>) more rapidly than that of *S. psammophila*. However, for both species, there was no significant relationship between LE<sub>TF</sub> or LE<sub>SF</sub> and RD (Figs. 7 and 8). The above results indicate that the intra-event rainfall 395 partitioning variables largely dependent on rainfall intensity and duration.

## 4 Discussion

### 4.1 Rainfall partitioning and influencing factors at inter-event scale

This study indicated that SF% of *C. korshinskii* (9.2%) was significantly higher than that of *S.*

*psammophila* (3.8%) (Fig. 3), which was comparable to the value of 10.4% and 6.3% reported  
400 by Yang et al. (2019) for the same species in similar semiarid regions of China. Under the same  
rainfall regimes, the difference in vegetation characteristics is the main reason for the difference  
in SF (Yuan et al., 2017; Whitworth-Hulse et al., 2020a; Yue et al., 2021). Comparing the  
structural properties of two shrubs with the same age (20 years), we found that CA, BD, BL,  
BA and NB values of *S. psammophila* were 4.51, 1.61, 1.94, 0.83 and 1.52 times of those of *C.*  
405 *korshinskii*, respectively (Table 1). On the branch scale, *C. korshinskii* had more small and short  
branches, but larger BA than that of *S. psammophila*, which was contributed to SF generation.  
Yuan et al. (2016) concluded that a beneficial branch architecture for SF production should  
include more relatively small branches and larger branch angles, and SF productivity decreased  
with BD size of branches. Furthermore, *C. korshinskii* with a smaller CA, and it had a  
410 larger SF<sub>d</sub> than *S. psammophila* under the same SF volume. Somewhat in line with Yuan  
et al. (2016) and Yue et al. (2021), our results suggest that a beneficial branch architecture for  
SF production of *C. korshinskii* should include relatively small CA, BD, BL and large BA  
(Table 1).

Leaf traits had been reported to exert a significant influence on rainfall partitioning  
415 (Garcia-Estringana et al., 2010; Magliano et al., 2019a). According to the documentation in  
*Flora of China* (Liu et al., 2010), *C. korshinskii* has pinnate compound leaves and each pinna  
has 5 to 8 pairs of ovate leaflets, and the leaves are lanceolate and concave, and the surface is  
densely sericeous. In comparison, *S. psammophila* has stripe or stripe oblanceolate leaves,  
margin revolute, and which upper surface of mature leaf blade is almost glabrous (Chao and

420 Gong, 1999). The branches of both shrubs are smooth, with a more developed cuticle layer on  
the surface of the *S. psammophila* branches, while the *C. korshinskii* branches contain oil and  
have waxy skin (Chao and Gong, 1999; Liu et al., 2010). The leaf morphology and epidermal  
characteristics of branches of *C. korshinskii* was more beneficial for SF generation than that of  
*S. psammophila* (Whitworth-Hulse et al., 2020b; Yuan et al. 2017). It was found that big  
425 biomass of leaves, concave leaf shape and leaf pubescence are beneficial to promote the  
generation of SF (Yuan et al., 2016). These factors together enable the leaves to function as a  
highly efficient natural water collecting system.

The mean IC% of *C. korshinskii* (21.4%) was significantly lower than that of *S.*  
*psammophila* (29.5%) in this study. The intercepts in the fitted formulas between interception  
430 loss and rainfall amount in Fig. 4e indicated that *C. korshinskii* (0.92 mm) had a lower canopy  
water storage than *S. psammophila* (1.15 mm), hence the potential interception loss of *C.*  
*korshinskii* was lower. Zhang et al. (2017) reported that IC% were higher in the *H. rhamnoides*  
stand (24.9%) than in the *S. pubescens* stand (19.2%), which was mainly attributed to the lower  
canopy water storage of *S. pubescens*. This study was done at the shrub-scale, so we compared  
435 the total canopy water storage of individual plant ( $C_m$ ), and we found that  $C_m$  of *S. psammophila*  
(6.0 L) was significantly higher than that of *C. korshinskii* (3.9 L) (Table 1). This was mainly  
due to the significantly higher average total dry aboveground biomass of *S. psammophila* (15.7  
kg per plant) than *C. korshinskii* (4.6 kg per plant). Consequently, individual *S. psammophila*  
absorbed more rain water to moisten the branches and leaves than that of individual *C.*  
440 *korshinskii*, which could explain higher IC% of *S. psammophila* than *C. korshinskii*. Thus, the

best predictors for interception loss were biomass-related parameters (i.e., woody biomass and total biomass) (Li et al., 2016).

#### **4.2 Rainfall partitioning and influencing factors at intra-event scale**

Temporal heterogeneity of rainfall clearly influences the amount and timing of TF and SF  
445 reaching the soil under the canopy, as explained by some previous intra-event rainfall  
partitioning studies from forested ecosystems (Owens et al., 2006; Levia et al., 2010; Molina  
et al., 2019). Our experiment investigated the intra-event dynamics of all rainfall partitioning  
components in xerophytic shrubs, which has rarely been reported before. Our results showed  
that the temporal dynamics of TF and SF under the shrub canopy almost matched the dynamics  
450 of rainfall (Fig. 5). It agreed with the reports of Zhang et al. (2018) and Yuan et al. (2019) who  
demonstrated the temporal synchronization of TF and SF with rainfall, respectively. The SF  
intensity is generally greater than rainfall intensity for different species (Fig. 6), which has eco-  
hydrological significance (Spencer and van Meerveld, 2016). The SF converges substantial  
rainwater to the shrub bases and then delivers it into the soil as a point input to recharge soil  
455 moisture (Germer et al., 2010; Cayuela et al., 2018; Jian et al., 2019). We assume that changes  
in SF inputs explain, at least in part, the temporal variation in subsurface moisture patterns.  
Spatial and temporal patterns in rainfall partitioning may influence various eco-hydrological  
processes under canopies, such as local soil moisture dynamics in arid and semi-arid regions  
(Li et al., 2009; Wang et al., 2011; Jian et al., 2019; Molina et al., 2019; Zhang et al., 2018).

460 The intensity variables and lag time of SF and TF relative to rainfall were the key to  
describe the intra-event rainfall partitioning (Fig. 6). The effects of meteorological factors on

SF and TF variables at the intra-event scale were derived from multiple regression analysis in this study. The SF and TF variables (intensity and temporal dynamics) were strongly influenced by rainfall intensity (e.g.,  $I$ ,  $I_{10\_max}$  and  $I_{10\_b}$ ) and duration (e.g.,  $RD$  and  $LM_R$ ). This is consistent with the results reported by Yuan et al. (2019) who indicated that there was a significant effect of rainfall intensity on the stemflow process of *C. korshinskii*. The main factors affecting intra-event SF and TF variables were the same, but the effects were still slightly different between the two shrubs. Under the same rainfall intensity, the average TF intensity under the canopy of *S. psammophila* was higher than *C. korshinskii* (Figs. 7a and 7b). But the average SF intensity of *C. korshinskii* was greater than *S. psammophila* at shrub scale (Figs. 8a and 8b), which was also found for the branch SF intensity reported by Yuan et al. (2019). In addition to the inter-shrub differences, the effects of  $I_{10\_b}$  on  $LG_{TF}$  and  $LG_{SF}$  were slightly different. The correlation between  $LG_{SF}$  and  $I_{10\_b}$  (Fig. 8c) was weaker than that between  $LG_{TF}$  and  $I_{10\_b}$  (Fig. 7c). This may be due to the fact that TF has two components, i.e., free TF and released TF (Staelens et al., 2008; Levia et al., 2017; Van Stan II et al., 2020), and that SF only starts to produce when a certain amount of rainfall is reached (Germer et al., 2010; Levia et al., 2010; Dunkerley, 2014; Yuan et al., 2019). Our results indicated that *S. psammophila* had dynamic characteristics (e.g., larger TFI,  $TFI_{10}$  and  $LE_{TF}$  as well as TFD, and shorter  $LG_{TF}$  and  $LM_{TF}$ ) producing larger TF depth ( $TFd = TFI * TFD$ ) (Figs. 6a and 6c), while *C. korshinskii* had dynamic characteristics (e.g., larger SFI,  $SFI_{10}$  and  $LE_{SF}$  as well as SFD) producing larger SF depth ( $SFd = SFI * SFD$ ) (Figs. 6b and 6d).

The vegetation characteristics have an important effect on the dynamics and the lag time

of TF and SF (Yuan et al., 2019; Zhang et al., 2018). Based on the temporal data recorded by TBRGs, we found that *C. korshinskii* produced TF and SF later than *S. psammophila* (Figs. 5 and 6), which was reported by Yuan et al. (2019) for branch SF of the same species. We inferred that this was due to the higher canopy water storage capacity of *C. korshinskii* (0.85 mL/g) compared to *S. psammophila* (0.38 mL/g). However, when the branches were moistened, SF production of *C. korshinskii* was greater than that of *S. psammophila* because of its branch and leaf characteristics as discussed in subsection 4.1 (Fig. 5). It was found that the great bark water storage capacity of forests could result in the further delay of TF and SF onset (Levia and Herwitz, 2005; Levia et al., 2010; Li et al., 2016; Pinos et al., 2021). The different intra-event TF and SF variables between species were attributed to a complex interaction of biotic and abiotic factors (Yuan et al., 2019; Zhang et al., 2018; Levia et al., 2010).

### **4.3 Implications and further scopes**

Most of previous rainfall partitioning investigations for shrubs were limited at inter-event scale, or only focused on TF or SF at intra-event scale. The intra-event rainfall partitioning dynamics, which could help have a better understanding of soil water replenishment and its distribution in soil and the key ecohydrological cycle in drylands, have been rarely explored. This study is the first time to investigate all the rainfall partitioning components (i.e., TF, SF and IC) for shrubs at both inter- and intra-event scales, which steps further and provides a full view of the reciprocal dynamics among interception loss, throughfall, and stemflow at the shrub-scale. This is the main novelty and a step forward compared with the previous related studies. We have also obtained the quantitative relationship between rainfall partitioning variables and rainfall



characteristics, and further elaborated the influence of vegetation structure characteristics (leaf,  
505 canopy structure, and biomass, etc..) on rainfall partitioning. The obtained new insights help to  
understand the fine characterization of shrub-dominated eco- hydrological processes, and  
improve the accuracy of water balance estimation in dryland ecosystem.

This study investigated the inter- and intra-event rainfall partitioning dynamics of two  
typical xerophytic shrubs in the Loess Plateau of China. There are several issues that need  
510 further investigation. Firstly, long-term observations of rainfall partitioning dynamics for more  
shrub plants and species are needed, and the rainfall partitioning models should be developed  
for shrubs. Secondly, the effects of rainfall partitioning on soil moisture dynamics, nutrient  
cycling, and plant transpiration should be substantially investigated to detect the interactions  
between water redistribution and vegetation physiological processes. Finally, the extension  
515 from the individual plant to stand and more larger scale remains a challenging topic for rainfall  
partitioning, which can help improve understanding the role of rainfall partitioning in the  
regional hydrologic cycle.

## **5 Conclusions**

In this study, we analyze the rainfall partitioning and the influences of bio-/abiotic factors of  
520 two typical shrubs at both inter- and intra-event scales in the Loess Plateau. To ensure a larger  
proportion of the rainfall is allocated under the canopy, two species can obtain more net rainfall  
through different mechanisms. At the event scale, there was no significant difference in TF  
percentage between the two shrubs, but *C. korshinskii* had significantly higher SF percentage  
and lower IC percentage compared to *S. psammophila*. At the intra-event scale, TF and SF of

525 two shrubs were well synchronized with the rainfall, but *C. korshinskii* had the advantage of stemflow production, while *S. psammophila* had the advantage of TF generation. For both shrubs, the inter-event rainfall partitioning amount and percentage depended more on rainfall amount, and rainfall intensity and duration controlled the intra-event TF and SF variables. The *C. korshinskii* has larger branch angles, more small branches and smaller canopy areas to  
530 produce SF more efficiently, and *S. psammophila* has larger biomass to intercept more rainfall amount. These findings could enhance our understanding of TF and SF dynamics and corresponding driving factors at inter- and intra-event scales, and help in modelling the critical eco-hydrological processes in arid and semi-arid regions.

535 **Data availability:** The data that support the findings of this study are available from the corresponding author upon request.

**Author contributions:** JXA: Formal analysis, Investigation, Methodology, Writing - original draft; GYG: Conceptualization, Methodology, Writing - review & editing; CY: Investigation,  
540 Writing - review & editing; BJF: Writing - review & editing.

**Competing interests:** The authors declare that they have no conflict of interest.

**Acknowledgements:** This research was supported by the National Natural Science Foundation  
545 of China (nos. 41991233 and 41822103), and the Youth Innovation Promotion Association  
CAS (no. Y202013). Special thanks are given to Shenmu Erosion and Environment Research  
Station for experimental support to this research. We thank David Dunkerley and two  
anonymous reviewers for their professional comments which greatly improved the quality of  
this paper.

550

## References

- Cayuela, C., Llorens, P., Sanchez-Costa, E., Levia, D. F., and Latron, J.: Effect of biotic and  
abiotic factors on inter- and intra-event variability in stemflow rates in oak and pine stands  
in a Mediterranean mountain area, *J. Hydrol.*, 560, 396-406, 2018.
- 555 Chao, P. N. and Gong, G. T.: *Salix (Salicaceae)*, in: Flora of China, edited by: Wu, Z. Y., Raven,  
P. H., and Hong, D. Y., Science Press, Beijing and Missouri Botanical Garden Press, St.  
Louis, 162–274, 1999.
- Chesson, P., Gebauer, R. L., Schwinning, S., Huntly, N., Wiegand, K., Ernest, M. S., Sher, A.,  
Novoplansky, A., and Weltzin, J. F.: Resource pulses, species interactions, and diversity  
560 maintenance in arid and semi-arid environments, *Oecologia*, 141, 236-253, 2004.
- Chu, X., Han, G., Xing, Q., Xia, J., Sun, B., Yu, J. B., and Li, D.: Dual effect of precipitation  
redistribution on net ecosystem CO<sub>2</sub> exchange of a coastal wetland in the Yellow River  
Delta, *Agr. Forest Meteorol.*, 249, 286-296, 2018.
- Dunkerley, D.: Stemflow production and intrastorm rainfall intensity variation: an

565 experimental analysis using laboratory rainfall simulation, *Earth. Surf. Proc. Land.*, 39,  
1741-1752, 2014.

Garcia-Estringana, P., Alonso-Blázquez, N., and Alegre, J.: Water storage capacity, stemflow  
and water funneling in Mediterranean shrubs, *J. Hydrol.*, 389, 363-372, 2010.

Germer, S., Werther, L., and Elsenbeer, H.: Have we underestimated stemflow? Lessons from  
570 an open tropical rainforest, *J. Hydrol.*, 395, 169-179, 2010.

Gordon, D. A., Coenders-Gerrits, M., Sellers, B. A., Sadeghi, S. M. M. and Van Stan II, J. T.:  
Rainfall interception and redistribution by a common North American understory and  
pasture forb, *Eupatorium capillifolium* (Lam. dogfennel), *Hydrol. Earth. Syst. Sci.*, 24,  
4587–4599, 2020.

575 Hanchi, A., and Rapp, M.: Stemflow determination in forest stands, *Forest Ecol. Manage.*, 97,  
231-235, 1997.

Honda, E. A., Mendonça, A. H., and Durigan, G.: Factors affecting the stemflow of trees in the  
Brazilian Cerrado, *Ecohydrology*, 8, 1351-1362, 2015.

Iida, S., Shimizu, T., Kabeya, N., Nobuhiro, T., Tamai, K., Shimizu, A., Ito, E., Ohnuki, Y., Abe,  
580 T., Tsuboyama, Y., Chann, S., and Keth, N.: Calibration of tipping-bucket flow meters and  
rain gauges to measure gross rainfall, throughfall, and stemflow applied to data from a  
Japanese temperate coniferous forest and a Cambodian tropical deciduous forest, *Hydrol.  
Process.*, 26, 2445-2454, 2012.

Jia, X. X., Shao, M. A., Wei, X. R., and Wang, Y. Q.: Hillslope scale temporal stability of soil  
585 water storage in diverse soil layers, *J. Hydrol.*, 498, 254-264, 2013.

- Jia, X., Zha, T. S., Gong, J. N., Wang, B., Zhang, Y. Q., Wu, B., Qin, S. G., and Peltola, H.: Carbon and water exchange over a temperate semi-arid shrubland during three years of contrasting precipitation and soil moisture patterns, *Agr. Forest Meteorol.*, 228, 120-129, 2016.
- 590 Jian, S. Q., Hu, C. H., Zhang, G. D., and Zhang, J. P.: Study on the throughfall, stemflow, and interception of two shrubs in the semiarid Loess region of China, *Agr. Forest Meteorol.*, 279, 107713, 2019.
- Lacombe, G., Valentin, C., Sounyafong, P., de Rouw, A., Soulileuth, B., Silvera, N., Pierret, A., Sengtaheuanghoung, O., and Ribolzi, O.: Linking crop structure, throughfall, soil surface  
595 conditions, runoff and soil detachment: 10 land uses analyzed in Northern Laos, *Sci. Total Environ.*, 616-617, 1330-1338, 2018.
- Levia, D. F. and Frost, E. E.: A review and evaluation of stemflow literature in the hydrologic and biogeochemical cycles of forested and agricultural ecosystems, *J. Hydrol.*, 274, 1-29, 2003.
- 600 Levia, D. F. and Germer, S.: A review of stemflow generation dynamics and stemflow-environment interactions in forests and shrublands, *Rev. Geophys.*, 53, 673-714, 2015.
- Levia, D.F. and Herwitz, S. R.: Interspecific variation of bark water storage capacity of three deciduous tree species in relation to stemflow yield and solute flux to forest soils, *Catena*, 64, 117-137, 2005.
- 605 Levia, D.F., Hudson, S. A., Llorens, P., and Nanko, K.: Throughfall drop size distributions: a review and prospectus for future research, *WIREs Water*, 4, 2017.

- Levia, D. F., Van Stan II, J. T., Mage, S. M., and Kelley-Hauske, P. W.: Temporal variability of stemflow volume in a beech-yellow poplar forest in relation to tree species and size, *J. Hydrol.*, 380, 112-120, 2010.
- 610 Li, X., Xiao, Q. F., Niu, J. Z., Dymond, S., van Doorn, N. S., Yu, X. X., Xie, B. Y., Lv, X. Z., Zhang, K. B., and Li, J.: Process-based rainfall interception by small trees in Northern China: The effect of rainfall traits and crown structure characteristics, *Agr. Forest Meteorol.*, 218-219, 65-73, 2016.
- Li, X. Y., Yang, Z. P., Li, Y. T., and Lin, H.: Connecting ecohydrology and hydrogeology in  
615 desert shrubs: stemflow as a source of preferential flow in soils, *Hydrol. Earth. Syst. Sci.*, 13, 1133-1144, 2009.
- Liu, Y. X., Chang, Z. Y. and Gennady, P. Y.: *Caragana (Fabaceae)*, in: *Flora of China*, edited by: Wu, Z. Y., Raven, P. H., and Hong, D. Y., Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, 528–545, 2010.
- 620 Llorens, P. and Domingo, F.: Rainfall partitioning by vegetation under Mediterranean conditions. A review of studies in Europe, *J. Hydrol.*, 335, 37-54, 2007.
- Magliano, P. N., Whitworth-Hulse, J. I., and Baldi, G.: Interception, throughfall and stemflow partition in drylands: Global synthesis and meta-analysis, *J. Hydrol.*, 568, 638-645, 2019a.
- Magliano, P. N., Whitworth-Hulse, J. I., Florio, E. L., Aguirre, E. C., and Blanco, L. J.:  
625 Interception loss, throughfall and stemflow by *Larrea divaricata*: The role of rainfall characteristics and plant morphological attributes, *Ecol. Res.*, 34, 753-764, 2019b.
- Martinez-Meza, E., Whitford, W. G.: Stemflow, throughfall and channelization of stemflow by

roots in three Chihuahuan desert shrubs, *J. Arid Environ.*, 32, 271-287, 1996.

Molina, A. J., Llorens, P., Garcia-Estringana, P., Moreno de Las Heras, M., Cayuela, C., Gallart,  
630 F., and Latron, J.: Contributions of throughfall, forest and soil characteristics to near-  
surface soil water-content variability at the plot scale in a mountainous Mediterranean  
area, *Sci. Total Environ.*, 647, 1421-1432, 2019.

Owens, M. K., Lyons, R. K., and Alejandro, C. L.: Rainfall partitioning within semiarid juniper  
communities: effects of event size and canopy cover, *Hydrol. Process.*, 20, 3179-3189,  
635 2006.

Pinos, J., Latron, J., Levia, D. F., and Llorens, P.: Drivers of the circumferential variation of  
stemflow inputs on the boles of *Pinus sylvestris* L. (Scots pine), *Ecohydrology*, 14, 2021.

Rivera, D. N. and Van Stan II, J. T.: Grand theft hydro? Stemflow interception and redirection  
by neighbouring *Tradescantia ohiensis* Raf. (spiderwort) plants, *Ecohydrology*, 13, 2020.

640 Sadeghi, S. M. M., Gordon, D. A., and Van Stan II, J. T.: A Global Synthesis of Throughfall  
and Stemflow Hydrometeorology. In: Van Stan II, J. T., Gutmann, E., Friesen, J. (Eds.),  
Precipitation Partitioning by Vegetation, Springer Nature Switzerland, Cham, pp. 49-70,  
2020.

Soulsby, C., Braun, H., Sprenger, M., Weiler, M., and Tetzlaff, D.: Influence of forest and shrub  
645 canopies on precipitation partitioning and isotopic signatures, *Hydrol. Process.*, 31, 4282-  
4296, 2017.

Spencer, S. A. and van Meerveld, H. J.: Double funnelling in a mature coastal British Columbia  
Forest: spatial patterns of stemflow after infiltration, *Hydrol. Process.*, 30, 4185-4201,

2016.

650 Staelens, J., De Schrijver, A., Verheyen, K., and Verhoest, N.E.C.: Rainfall partitioning into throughfall, stemflow, and interception within a single beech (*Fagus sylvatica* L.) canopy: influence of foliation, rain event characteristics, and meteorology, *Hydrol. Process.*, 22, 33-45, 2008.

Tonello, K. C., Rosa, A. G., Pereira, L. C., Matus, G. N., Guandique, M. E. G., and Navarrete, 655 A. A.: Rainfall partitioning in the Cerrado and its influence on net rainfall nutrient fluxes, *Agr. Forest Meteorol.*, 303, 108372, 2021.

Van Stan II, J. T., Gutmann, E., and Friesen, J. (Eds.): *Precipitation Partitioning by Vegetation — A Global Synthesis*, *Precipitation Partitioning by Vegetation*, Springer Nature Switzerland, 2020.

660 Van Stan II, J. T., Wagner, S., Guillemette, F., Whitetree, A., Lewis, J., Silva, L., and Stubbins, A.: Temporal Dynamics in the Concentration, Flux, and Optical Properties of Tree-Derived Dissolved Organic Matter in an Epiphyte-Laden Oak-Cedar Forest, *Journal of Geophysical Research*, 122, 2982-2997, 2017.

Wang, X. P., Zhang, Y. F., Hu, R., Pan, Y. X., and Berndtsson, R.: Canopy storage capacity of 665 xerophytic shrubs in Northwestern China, *J. Hydrol.*, 454, 152-159, 2012.

Wang, X., Wang, Z., Berndtsson, R., Zhang, Y., and Pan, Y.: Desert shrub stemflow and its significance in soil moisture replenishment, *Hydrol. Earth. Syst. Sci.*, 15, 561-567, 2011.

Whitworth-Hulse, J. I., Magliano, P. N., Zeballos, S. R., Gurvich, D. E., Spalazzi, F., and Kowaljow, E.: Advantages of rainfall partitioning by the global invader *Ligustrum*



- 670 lucidum over the dominant native *Lithraea molleoides* in a dry forest, *Agr. Forest Meteorol.*, 290, 108013, 2020a.
- Whitworth-Hulse, J. I., Magliano, P. N., Zeballos, S. R., Aguiar, S., and Baldi, G.: Global patterns of rainfall partitioning by invasive woody plants, *Global Ecol. Biogeogr.*, 30, 235-246, 2020b.
- 675 Xu, H. and Li, Y.: Water-use strategy of three central Asian desert shrubs and their responses to rain pulse events, *Plant and Soil*, 285, 5-17, 2006.
- Yang, X. L., Shao, M. A., and Wei, X. R.: Stemflow production differ significantly among tree and shrub species on the Chinese Loess Plateau, *J. Hydrol.*, 568, 427-436, 2019.
- Yuan, C., Gao, G. Y., and Fu, B. J.: Stemflow of a xerophytic shrub (*Salix psammophila*) in  
680 northern China: Implication for beneficial branch architecture to produce stemflow, *J. Hydrol.*, 539, 577-588, 2016.
- Yuan, C., Gao, G. Y., and Fu, B. J.: Comparisons of stemflow and its bio-/abiotic influential factors between two xerophytic shrub species, *Hydrol. Earth. Syst. Sci.*, 21, 1421-1438, 2017.
- 685 Yuan, C., Gao, G. Y., Fu, B. J., He, D. M., Duan, X. W., and Wei, X. H.: Temporally dependent effects of rainfall characteristics on inter- and intra-event branch-scale stemflow variability in two xerophytic shrubs, *Hydrol. Earth. Syst. Sci.*, 23, 4077-4095, 2019.
- Yue, K., De Frenne, P., Fornara, D. A., Van Meerbeek, K., Li, W., Peng, X., Ni, X., Peng, Y., Wu, F., Yang, Y., and Penuelas, J.: Global patterns and drivers of rainfall partitioning by  
690 trees and shrubs, *Glob Chang Biol*, 27, 3350-3357, 2021.

- Zhang, H., Fu, C., Liao, A., Zhang, C., Liu, J., Wang, N., and He, B.: Exploring the stemflow dynamics and driving factors at both inter- and intra-event scales in a typical subtropical deciduous forest, *Hydrol. Process.*, 35, 2021a.
- Zhang, Y., Li, X. Y., Li, W., Wu, X. C., Shi, F. Z., Fang, W. W., and Pei, T. T.: Modeling rainfall interception loss by two xerophytic shrubs in the Loess Plateau, *Hydrol. Process.*, 31, 1926-1937, 2017.
- Zhang, Y. F., Wang, X. P., Hu, R., and Pan, Y. X.: Meteorological influences on process-based spatial-temporal pattern of throughfall of a xerophytic shrub in arid lands of northern China, *Sci. Total Environ.*, 619-620, 1003-1013, 2018.
- 700 Zhang, Y. F., Wang, X. P., Pan, Y. X., and Hu, R.: Variations of Nutrients in Gross Rainfall, Stemflow, and Throughfall Within Revegetated Desert Ecosystems, *Water Air and Soil Pollution*, 227, ARTN 183, 2016.
- Zhang, Y. F., Wang, X. P., Pan, Y. X., Hu, R., Chen, N., and Sandel, R. O.: Global quantitative synthesis of effects of biotic and abiotic factors on stemflow production in woody ecosystems, *Global Ecol. Biogeogr.*, 30, 1713-1723, 2021b.
- 705

**Table 1.** Descriptive statistics (mean  $\pm$  standard error) of canopy morphology of *C. korshinskii* (CK1-CK3) and *S. psammophila* (SP1-SP3) plants. Values are mean  $\pm$  SD.


Plant ID	SH (m)	CA (m <sup>2</sup> )	NB	BL (cm)	BA (°)	BD (mm)	TBA (cm <sup>2</sup> )	TB (kg)	C <sub>m</sub> (L)
CK1	2.2	5.3	47	150.6 $\pm$ 5.1	60.26 $\pm$ 2.6	9.2 $\pm$ 0.5	34.9	4.0	3.4
CK2	2.3	5.2	47	123.3 $\pm$ 6.6	65.5 $\pm$ 2.1	8.5 $\pm$ 0.5	31.3	3.6	3.1
CK3	2.4	5.3	49	134.6 $\pm$ 6.7	65.4 $\pm$ 4.4	9.9 $\pm$ 0.7	45.8	6.2	5.2
Average	2.3a	5.27a	48a	136.2a	63.77a	9.2a	37.3a	4.6a	3.9a
SP1	3.5	23.9	85	262.2 $\pm$ 6.0	67.1 $\pm$ 1.4	13.8 $\pm$ 0.5	139.7	14.3	5.4
SP2	3.3	26.1	55	268.0 $\pm$ 7.7	56.0 $\pm$ 3.1	15.1 $\pm$ 0.6	124.3	15.9	6.0
SP3	3.6	21.4	78	262.0 $\pm$ 7.8	35.1 $\pm$ 2.9	15.3 $\pm$ 0.5	155.9	17.0	6.5
Average	3.5b	23.8b	73b	264.1b	52.7a	14.8b	140.0b	15.7b	6.0b

Note: SH: shrub height; CA: canopy area; NB: number of branches; BL: branch length; BA: branch angle; BD: basal diameter of branch; TBA: total basal area of the shrub; TB: total dry aboveground biomass; C<sub>m</sub>: total canopy storage per plant. Different letters indicate statistically significant differences between two species ( $p < 0.05$ ).


710

**Table 2.** Rainfall partitioning parameters at inter- and intra-event scales.


Scale	Parameter (unit)	Explanation
Inter-event	TF <sub>d</sub> (mm)	Throughfall depth per rainfall event
	SF <sub>d</sub> (mm)	Stemflow depth per rainfall event
	IC <sub>d</sub> (mm)	Interception loss depth per rainfall event
	TF%	Percentage of TF per rainfall event
	SF%	Percentage of SF per rainfall event
	IC%	Percentage of IC per rainfall event
	TFD (h)	Throughfall duration
	SFD (h)	Stemflow duration
	TFI (mm·h <sup>-1</sup> )	Average throughfall intensity
SFI (mm·h <sup>-1</sup> )	Average stemflow intensity	
Intra-event	I <sub>10</sub> (mm·h <sup>-1</sup> )	Rainfall intensity at 10-min interval
	I <sub>10_max</sub> (mm·h <sup>-1</sup> )	Maximum I <sub>10</sub> during the rainfall process
	TFI <sub>10</sub> (mm·h <sup>-1</sup> )	Throughfall intensity at 10-min interval
	TFI <sub>10_max</sub> (mm·h <sup>-1</sup> )	Maximum TFI <sub>10</sub> during the rainfall process
	SFI <sub>10</sub> (mm·h <sup>-1</sup> )	Stemflow intensity at 10-min interval
	SFI <sub>10_max</sub> (mm·h <sup>-1</sup> )	Maximum SFI <sub>10</sub> during the rainfall process
	LG <sub>TF</sub> (h)	Time lag of throughfall generation after the start of rainfall
	LG <sub>SF</sub> (h)	Time lag of stemflow generation after the start of rainfall
	LM <sub>R</sub> (h)	Time lag of I <sub>10_max</sub> occurrence relative to the onset of rainfall
	LM <sub>TF</sub> (h)	Time lag of TFI <sub>10_max</sub> occurrence relative to the onset of rainfall
	LM <sub>SF</sub> (h)	Time lag of SFI <sub>10_max</sub> occurrence relative to the onset of rainfall
	LE <sub>SF</sub> (h)	Time lag of throughfall ending after the end of rainfall
	LE <sub>SF</sub> (h)	Time lag of stemflow ending after the end of rainfall



**Figure 1.** The location and experimental settings in the plots of *C. korshinskii* and *S. psammophila*.




720 **Figure 2.** (a) individual rainfall amount (RA) ( $n = 38$ ), rainfall duration (RD), average rainfall intensity (I,  $\text{mm}\cdot\text{h}^{-1}$ ), maximum rainfall intensity in 10 minutes ( $I_{10\_max}$ ,  $\text{mm}\cdot\text{h}^{-1}$ ); and rainfall partitioning into TF %, SF %, and IC % of (b) *C. korshinskii* and (c) *S. psammophila*.




725 **Figure 3.** Box-plots of (a) TF%, (b) SF%, and (c) IC% for *C. korshinskii* (CK) and *S. psammophila* (SP). The horizontal thick black line indicates the median, boxes correspond to the 25th and 75th percentiles, and whiskers represent values that fall within 1.5 times the interquartile range. Mean values are represented with the black square. Different letters indicate significant differences between the two species ( $p < 0.05$ ).

730



**Figure 4.** Inter-event rainfall partitioning as a function of individual rainfall amount for *C. korshinskii* and *S. psammophila*.






735


**Figure 5.** Time series (10-min interval) of rainfall partitioning within four rainfall events for *C. korshinskii* (CK) and *S. psammophila* (SP). Events 1-4 occurred on August 3, September 17, September 28, and September 30 in 2015, respectively. The solid lines represent the rainfall, TF and SF intensity at 10-min interval. The dotted lines indicate the accumulated amount of

740 RA, TF, SF, and IC.



**Figure 6.** Intra-event TF (a, c) and SF (b, d) variables of *C. korshinskii* and *S. psammophila*.


All the variables are explained in Table 2.



745

**Figure 7.** Relationships of intra-event throughfall variables with meteorological characteristics

for *C. korshinskii* and *S. psammophila*. All the variables are explained in Table 2.



750 **Figure 8.** Relationships of intra-event stemflow variables with meteorological characteristics

for *C. korshinskii* and *S. psammophila*. All the variables are explained in Table 2.