

1 A generalised ecohydrological landscape classification for 2 assessing ecosystem risk in Australia due to an altering water 3 regime

4 Alexander Herr¹, Linda E. Merrin¹, Patrick J. Mitchell¹, Anthony P. O'Grady¹, Kate L. Holland¹,
5 Richard E. ~~Mount~~³Mount², David A. Post¹, Chris R. Pavey¹, Ashley D. Sparrow^{1,3}

6 ¹ Commonwealth Scientific and Industrial Research Organisation, Australia

7 ² formerly [at](#): Bureau of Meteorology, Australia

8 ³ Arthur Rylah Institute, Melbourne, Australia

9 *Correspondence to:* Dr Alexander Herr, email: alexander.herr@csiro.au

10 Abstract

11 Describing and classifying a landscape for environmental impact and risk assessment purposes is a
12 non-trivial challenge, ~~as standard~~ because this requires region specific landscape classifications that
13 cater for region specific impacts ~~do not exist~~. Assessing impacts on ecosystems from [the](#) extraction
14 of water resources across large regions requires linking ~~of~~ landscape features to their water
15 requirements. We present the rationale and implementation of an ecohydrological classification for
16 regions where coal mine and coal seam gas developments may impact on water. Our classification
17 provides the essential framework for modelling the potential impact of hydrological changes from
18 future coal resource developments at the landscape level.

19 We develop an attribute-based system that provides representations of the ecohydrological entities
20 and their connection to landscape features and make use of existing broad-level, classification
21 schemes into an attribute-based system. We incorporate a rule-set with prioritisation, which
22 underpin risk modelling and make the scheme resource efficient, where spatial landscape or
23 ecosystem classification schemes, developed for other purposes, already exist.

24 A consistent rule-set and conceptualised landscape processes and functions allow combining diverse
25 data with existing classification schemes. This makes the classification transparent, repeatable, and
26 adjustable, should new data become available. We apply the approach in three geographically
27 different regions, with widely disparate information sources, [for](#) the classification, [and](#) provide a
28 detailed example of its application. We propose that it is widely applicable around the world for
29 linking ecohydrology to environmental impacts.

30 **Keywords:** Typology, ecology, hydrology, causal pathway, bioregional assessments, environmental
31 impact, risk analysis

32 1 Introduction

33 The categorisation of the Earth's surface into geo-ecological landscape classes provides a way to
34 simplify the complexity of [the](#) form and function of the landscape and provides vital contextual
35 information to support land and water management, and policy initiatives. This includes identifying
36 geographical regions within which landscape-scale attributes, such as climate, topography, geology,
37 and land cover, that are homogeneous and distinctive compared to other regions, ~~and~~ [it](#) involves

38 identifying broad-scale, general patterns, processes, and functions. Landscape class units are
39 'ecologically equivalent', having the same dominant processes that sustain a similar suite of species,
40 and are likely to respond in similar ways to management initiatives or environmental changes. This
41 ecological equivalence enables the selection of assessment locations for monitoring, measurement
42 or experimentation, and it enables the extrapolation of results to all areas within the same
43 ecological class (Hawkins and Norris, 2000; MacMillan et al., 2003; Cullum et al., 2016a; Cullum et
44 al., 2016b).

45 Such a landscape classification also explains variation in ecological characteristics (e.g. assemblage
46 structure) and is predictive of the ecological attributes of those areas. This predictive quality is useful
47 for defining ecological criteria, identifying reference and degraded sites, defining conservation goals,
48 including the assessment of biodiversity, and the setting of restoration objectives (Hawkins et al.,
49 2000; McMahon et al., 2001; Snelder et al., 2004).

50 In summary, landscape classification is a way of dividing a landscape into components where the
51 characteristics within the components are more similar than the characteristics between the
52 components. That is, the components have their own distinct features that separate them from the
53 other components.

54 However, describing and classifying a landscape for environmental impact and risk assessment
55 purposes is a non-trivial challenge, where hydrological records are limited (see e.g. Wolfe et al.,
56 2019). This is the case for many regions in Australia, where low population densities, high
57 urbanisation and limits in (water) resource management information exist. For our purpose, which
58 was the assessment of risk to ecosystems within the regions of the Bioregional Assessments
59 Programme (Bioregional Assessments, 2018), we needed a landscape classification that reflected the
60 hydrological connectivity of surface and groundwater with ecosystems in the landscape. The
61 Bioregional Assessment Programme, an Australian regional scale impact assessment,- investigated
62 the impacts and risks of coal seam gas (CSG) and large coal mining developments on water resources
63 and water-dependent assets via a water pathway (Bioregional Assessments, 2018). [-This
64 investigation focussed on the landscape level, that is on areas within the regions where the
65 landscape is made up of different interacting land-uses and ecosystems.](#)

66 In our case, the broad scale assessments of impacts from resource developments on ecosystems
67 required an understanding of landscape composition and structure, and how these relate to the
68 ecosystems embedded in the landscape. The type and composition of the landscape components
69 are dependent on the focus of the assessment and therefore require careful consideration of the
70 questions the assessment seeks to answer (Wiens and Milne, 1989; Eigenbrot, 2016). For Australia,
71 there are several landscape level classifications available (see e.g. Thackway and Cresswell, 1995;
72 Pain et al., 2011; Hall et al., 2015; NVIS Technological Working Group, 2017; Gharari et al., 2011).
73 Unfortunately, these available classifications are not directly applicable for our assessment regions
74 because there is no alignment between the regions and existing classification boundaries, or the
75 classifications, even if they include ecohydrological elements, are limited to their locations or
76 domain of interest.

77 Identifying the water dependency of landscape components is a prerequisite when analysing the
78 potential impacts of proposed coal and gas resource developments on water resources at a regional
79 scale. For example, coal resource developments generally need to manage both groundwater and
80 surface water as part of their operations. With multiple developments within the one region,

81 impacts are likely to go beyond the local scale and affect ecosystems at the landscape level (see for
82 example Bioregional Assessments, 2018, 2019). In this context, there is a need for an ecological
83 classification of the landscape that identifies and causally connects the water dependency of its
84 components to activities of resource extraction, in a spatially explicit manner. Further, there is a
85 need to identify impact pathways between resource extraction sites and the ecosystems that show
86 causal connectivity between extraction activities and ecosystem impacts.

87 Land classification systems reveal patterns and underlying drivers of ecosystem structure and
88 function, or produce a tractable unit of assessment for evaluating environmental change (Hobbs and
89 McIntyre, 2005; Poff et al., 2010). Many different classification approaches and methodologies
90 currently exist to represent ecosystems in a landscape. This includes the interim bioregional
91 classification for Australia (IBRA), which provides the basis for defining and managing the national
92 reserve system and; the national vegetation information systems (NVIS), that describes the extent
93 and distribution of vegetation ecosystems for the Australian continent (Thackway and Cresswell,
94 1995; Department of Agriculture, 2021). Classifications addressing hydrology in Australia incorporate
95 a framework for river management that delineate boundaries between homogenous landscape
96 components, based on either their dependency on surface [water](#) or groundwater regimes (Poff et
97 al., 2010; Aquatic Ecosystems Task Group, 2012; Olden et al., 2012). However, none of these
98 classifications describe ecohydrological connections between waters and the wider landscape. For
99 example, IBRA and NVIS are based purely on vegetation classifications and so do not contain any
100 hydrological details, while the available hydrological classifications focus purely on the streams and
101 waterbodies within the landscape, as their focus is on aquatic organisms and environmental flows.
102 [While both these elements are part of the immediate landscape surrounding water bodies, they do
103 not in themselves provide conceptual and direct linkages between changes in water and ecosystem
104 responses in the wider landscape.](#) Therefore, a standardised approach to formulating classifications
105 that combine these two aspects, ecosystems and their water sources, is lacking. ~~The~~

106 [This](#) conundrum exists because different analysis contexts require classifications for different
107 purposes, ranging from conservation planning, habitat mapping, resource assessment and
108 vegetation modelling, and because there is contention between the generality of broad
109 classifications and their applicability at the local scale (Leathwick et al., 2003; Abella et al., 2003;
110 Poulter et al., 2011; Cullum et al., 2016b; Pyne et al., 2017). Hence, we needed a new classification
111 system, when evaluating water dependency in the context of regional scale for multiple [coal and
112 coal seam gas](#) resource developments. This new system must incorporate surface water and
113 groundwater regimes into a spatial demarcation of ecosystem boundaries in the landscape.
114 [Including surface water and groundwater regimes will provide the establishing of conceptual
115 connection between impacts from developments on surface water and groundwater within the
116 classification, and the classification must be spatially explicit, to enable a landscape wide analysis of
117 those impacts so that one can link changes in water at one part of the landscape to ecological
118 responses at another part of the landscape.](#)

119 With this context in mind, the objectives for this paper are to:

- 120 1. characterise a regional level landscape based on patterns in land use, ecology, geomorphology
121 and hydrology,
- 122 2. develop landscape classes of water-dependent, remnant and human-modified features, and
- 123 3. ensure landscape classes sit within a common framework that aids in formulating
124 conceptual models and patterns of water dependency across the landscape.

125 Here, we present the rationale, formulation, and implementation of an ecohydrological landscape
126 classification. Based on a generalised conceptual model of the typical hydrological connectivity
127 within landscape features in a region, the classification integrates pre-existing, broad-level
128 classification schemes into an attribute-based schema applied at the regional scale. It places the
129 landscape classification within a common framework ([i.e. a framework that is common to all](#)
130 [landscape elements in the region](#)) that aids in formulating conceptual models and patterns in water
131 dependency across the landscape. This makes our approach generally applicable for assessments
132 aimed at regional hydrological impacts [on](#), and risks to, ecosystems. Importantly, the classification
133 also provides the ability to [conceptually describe/develop a conceptual understanding of](#), and
134 causally connect, hydrological changes at the landscape level, with impacts on ecological entities
135 within the landscape. These causal pathways are the basis for spatially identifying the impacted
136 areas, and for developing an appropriate mitigation response, including for extractive resource
137 developments and water extraction.


138 We have applied this approach to several regions across eastern Australia with coal and CSG
139 resource developments. Here we will focus on its application in three regions; Namoi, Maranoa–
140 Balonne–Condamine and Galilee, and subsequently discuss why the approach is transferable to
141 other regional developments that may carry a hydrological [based](#) risk [to ecosystems](#), even those in a
142 different contextual setting with regards to data sources and existing landscape classifications.

143 **2 — Methods**

144 ~~In the following section, we show the development of a dataset-agnostic method to develop a~~
145 ~~regional-level landscape classification that is flexible in incorporating data sources at different scales,~~
146 ~~including region-specific datasets.~~

147 ~~The remainder of the paper is structured as follows: in the Methods section we describe the general~~
148 ~~approach for achieving the classification, including descriptive examples of existing data sources. It~~
149 ~~also provides a description of the three study regions in which we applied and tested the~~
150 ~~classification. The Results section provides evidence of the general applicability of our approach in~~
151 ~~that it shows the detailed ecological landscape classification for the three distinctively different~~
152 ~~region in terms of location, topography, and climate. In the Discussion section we provide an~~
153 ~~example on the use of the landscape classification. Here we describe an impact assessment in the~~
154 ~~Namoi region using modelling that includes a Bayesian expert assessment approach. We also discuss~~
155 ~~limitations and provide our conclusions.~~

156 ~~Figure 1 provides a visual outline of the paper, giving an overview of the workflow we applied. In this~~
157 ~~context the figure incorporates Methods and Results above the dashed line. Below the dashed line~~
158 ~~are the Discussion parts, which include applying our classification using quantitative and qualitative~~
159 ~~risk modelling in combination with surface water and groundwater modelling. Surface water and~~
160 ~~groundwater modelling establish a zone of hydrological change in which impacts are likely.~~



161

162

163

Figure 1: Visualisation of workflow for developing our ecological landscape classification (above the dashed line) and its application to develop an ecological risk assessment (below the dashed line).

164 **2 Methods**

165 [In the following section, we show the development of a dataset-agnostic method to develop a](#)
166 [regional-level landscape classification that is flexible in incorporating data sources at different scales,](#)
167 [including region-specific datasets. Ecological systems are complex and work at a range of scales](#)
168 [within regions/landscapes, and they exhibit interactions and feedbacks that work across scales.](#)
169 [Consequently, there is no one scale appropriate for a subsequent analysis of ecological impacts.](#)
170 [Here we use a variable scale range that is relevant for ecological impacts of water changes from coal](#)
171 [resource developments when using an expert assessment approach. Our classification focuses on a](#)
172 [scale range \(36,000 km² to 600,000 km²\) that is associated with eco-hydrological linkages \(and](#)
173 [associated causality\) between the response of ecological components to predicted hydrological](#)
174 [changes. This scale range is what most hydrologists would consider the “regional” scale range](#)
175 [\(Gleeson and Paszkowski, 2014\). It provides the basis and flexibility for experts to build their](#)
176 [conceptual understanding of causal pathways and use these to assess ecological impacts with the](#)
177 [landscape classes \(see also Figure 1\).](#)

178 **2.1 Study areas**

179 Our three study areas are the Namoi, the Maranoa–Balonne–Condamine and the Galilee regions in
180 eastern Australia. Each of these regions have coal resource developments within them and have
181 distinctly different landscape characteristics. They cover different state jurisdictions, or even cross
182 state jurisdictions, and range from approximately 36,000 km² to 600,000 km² in size. Consequently,
183 the classification is based on different state-based datasets. Each region’s classification relies on the
184 extent of [groundwater and](#) surface water [and groundwater](#) systems that existing and potential
185 future coal resource developments in the region may impact.

186 **2.1.1 Namoi region**

187 The Namoi region covers approximately 35,700 km² in eastern Australia, is located within New South
188 Wales and forms one catchment of the Murray-Darling Basin. The long-term mean annual rainfall
189 varies from 600 to 1100 mm and potential evapotranspiration (PET) varies from 1200 to 1400 mm. It
190 contains six operational coal mines (one underground mine and five open-cut mines), nine potential
191 future coal mines and one potential CSG development. The nine potential future coal mines consist
192 of two underground, one combined open cut and underground, and seven open cut mines. The
193 region covers most of the Namoi River catchment, with the Namoi River being the main river within
194 the region. It also contains two major aquifer systems – the Namoi Alluvial aquifer and the Pilliga
195 Sandstone aquifer (Figure 2a).

196 The main land use within the region is agriculture, both dryland and irrigated cropping, and
197 livestock grazing, as well as forestry. There is also a diverse range of landscapes and ecosystems
198 within the region, including the Liverpool and Kaputar ranges, the Liverpool Plains floodplains, and
199 Darling Riverine plains in the west of the region, open box woodlands on the slopes, and temperate
200 and sub-alpine forests in the east of the region. A range of aquatic habitats occur downstream of
201 Narrabri, with large areas of anabranches and billabong wetlands. The Pilliga Nature Reserve in the
202 upper catchment of Bohena Creek, together with The Pilliga State Forest, form the largest remaining
203 area of dry sclerophyll forest west of the Great Diving Range in New South Wales (Welsh et al.,
204 2014).

205 **2.1.2 Galilee region**

206 The Galilee region covers approximately 612,300 km² and is located mostly within Queensland,
207 Australia. PET far exceeds rainfall, particularly in the summer months. Yearly rainfall ranges from 300
208 to 700 mm and PET from 2200 to 2900 mm. There are 17 proposed coal resource developments in
209 the Galilee region. These include three open-cut coal mines, two underground coal mines, five
210 combined open-cut and underground coal mines, four coal mines of currently unknown type, and
211 three CSG projects (Figure 2b).

212 The Galilee region includes the headwaters of seven major drainage catchments. These catchments
213 are Bulloo, Burdekin, Cooper Creek, Diamantina, Flinders, Paroo and Warrego. The largest of these
214 catchments within the region are the Cooper Creek and Diamantina. Groundwater within the region
215 is a very important resource, as most of the streams are ephemeral. Groundwater is used for town
216 water, agriculture and industry. Most groundwater in the region is extracted from the Great Artesian
217 Basin (Figure 2b).

218 The region covers a range of environments, including mountains of the Great Dividing Range in the
219 east, through to semi-arid and arid areas in the central and western ~~part~~parts of the region. The
220 main land use in the region is livestock grazing on native vegetation. There is no intensive agriculture
221 in the region, and a low human population density, largely due to the low and unpredictable rainfall
222 (Evans et al., 2014).


223 **2.1.3 Maranoa–Balonne–Condamine region**

224 The Maranoa–Balonne–Condamine region covers approximately 130,000 km² and is located mostly
225 within south-east Queensland with about half the area within the Murray-Darling Basin. From east
226 to west, average annual rainfall decreases from 800 mm to 420 mm, as PET increases from 1500 to
227 2370 mm. The region overlies the Surat Basin and has five open-cut coal mines and five CSG projects,
228 as well as two proposed open-cut coal mines (Figure 2c).

229 The region contains the headwaters of the Condamine-Balonne, Moonie, Weir, Maranoa and
230 Dawson rivers. Most of the rivers within the region are ephemeral. Groundwater is therefore an
231 important water source and is used for stock and domestic purposes, and in some cases, town water
232 supply. The Great Artesian Basin is the main source of groundwater used within the region (Welsh et
233 al., 2015).


234 The main land use within the region is grazing on natural vegetation, with dryland cropping and
235 production forestry also major land uses. The main vegetation type within the region is grassy
236 woodlands, with river red gums, coolabah and river oak common riparian species. There are also six
237 wetlands of national significance within the region: Balonne River Floodplain, Boggomoss Springs,
238 Dalrymple and Blackfellow Creeks, Lake Broadwater, Palm Tree and Robinson Creeks, and The Gums
239 Lagoon (Welsh et al., 2015).

240




241


(a)



242




243



244

245 **Figure 2. Study areas for (a) the Namoi region (b) the Galilee region and (c) the Maranoa–Balonne–Condamine region,**
 246 **showing the potential coal resource development sites**



247

248
249

Figure 2 (cont). Study areas for (c) the Maranoa–Balonne–Condamine region, showing the potential coal resource development sites

250

2.2 Landscape classification development – overview and rationale

251

The purpose of this ecohydrological landscape classification is to characterise the landscape based on patterns in land use, ecology, geomorphology and hydrology, and from these, develop landscape classes of water-dependent, remnant and human-modified features. [ExistingWe chose these features because these three types represent a generally applicable delineation used in our spatial dataset. For example, in Australia the word remnant vegetation \(our remnant features\) describes all vegetation where there was no clearing or regrowth1 of \(semi-\) native vegetation, resulting in a vegetation community that resembles its predecessor’s structure. It represents areas with low to very minimal human interference. This is opposed to human-modified, where human activities are the defining features of the area, such as urban areas or other infrastructure. Water dependency is essential for establishing a conceptual linkage of water across landscape elements. Our classification employs a geographical information system to overlay existing spatial data for each region-forms. The spatial data are the basis for categorising the landscape features using a rule-set to prioritise the spatial data based on their attribute features-within the spatial-datasets. Depending on their origin and original purpose, the](#)

265

[The](#) datasets have a regional, state or national coverage. [ThisUsing a](#) feature-based classification helps to place the landscape classes within a common biophysical system that aids in formulating conceptual models and patterns in water dependency across the landscape of each region. This provides a classification that is aligned with the idiosyncrasies of each region. Maintaining regionality

267

268

269 is essential when developing conceptual models and quantitative models for assessing the risk to
270 ecological components from hydrological changes. For example, arid and semi-arid regions have very
271 different ecological environments, functions and processes than subtropical or temperate
272 woodlands.

273 Our approach uses a defined rule-set and priorities, which we apply to regionally available [data](#)
274 [sets/datasets](#) to achieve a landscape classification for each of our regions. [Tables 1 to 3 provide a list](#)
275 [of citations for example datasets used in this process.](#) This is different to most other landscape
276 classifications that may use climate, topography, hydrological assessment units and, remote sensing
277 data, and apply statistical dimensionality reduction and classifications such as proximity analysis ([see](#)
278 [e.g.](#) Gharari et al., 2011; Leibowitz et al., 2014; Sawicz et al., 2014; Zhang et al., 2016; Addicott et al.,
279 2021; Carlier et al., 2021; [see e.g.](#) Jones Jr et al., 2021).

280 When considering the characteristics of our regions, the following features form part of the [broad](#)
281 rule-set for defining landscape classes:

- 282 • broad habitat/land use type (remnant/human-modified).
283 [Note: In the Australian context, remnant vegetation are areas of natural vegetation that did](#)
284 [not experience significant human modification.](#)
285
- 286 • wetland (wetland/non-wetland)
- 287 • topography (upland/lowland, floodplain/non-floodplain)
- 288 • groundwater (groundwater-dependent/[non-groundwater dependent](#), Great Artesian Basin
289 (GAB)/non-GAB/[non-groundwater dependent](#).)
290 Note: identifies groundwater dependency and classifies this with [the presence/absence of](#)
291 Great Artesian basin groundwaters.
- 292 • vegetation type (riparian/woodland floodplain/grassy woodland/rainforest)
- 293 • water regime (permanent/ephemeral/null) of surface water

294 These features identify groups of [land forms/landforms](#) and use, streams and springs.

295

296 [For our work, where](#)The hydrological connectivity is the main reason for developing a new
297 classification, [as this allows us to assess the potential impact of coal resource developments on the](#)
298 [landscape via a water pathway. Therefore](#) the most important characteristics are the hydrological
299 features. [We](#)Describing the conceptual understanding of how water connects the landscape
300 [elements allows us to identify where in the landscape impacts are likely to occur. Therefore, we](#)
301 developed a hierarchical approach, where hydrological features have priority over other landscape
302 characteristics. This resulted in a spatially complete landscape classification, [where there are no](#)
303 [gaps in the mapping data.](#) The method of prioritisation depended on region-specific characteristics
304 and the data availability. [This resulted in a classification where the landscape classes have their](#)
305 [origin in the spatial datasets, and included the water dependency, which was a pre-requisite of the](#)
306 [prioritisation.](#) An example prioritisation assigned in order of highest to lowest is:

- 307 • aquatic ecosystems (e.g. wetlands, streams and lakes)
- 308 • remnant vegetation ~~—areas of vegetation that contain relatively intact plant communities~~
- 309 • other landscape components that are ‘non-remnant vegetation’ and are typically ‘human-
- 310 modified’.

311

312 Subsequent use of the landscape classification for risk identification with expert input also required
 313 combining landscape classes into broader landscape groups. [Landscape groups are sets of landscape](#)
 314 [classes that share hydrological properties.](#) These landscape groups provided efficiencies in the
 315 expert elicitation process [of the risk modelling](#), as they combined ~~more~~ similar ecological system
 316 components based on our landscape classes while also accounting for region specific differences. [For](#)
 317 [example, in the Namoi region there are two landscape groups where we do not expect any impact](#)
 318 [from coal resource developments. Firstly, the ‘Dryland remnant vegetation’ landscape group is ruled](#)
 319 [out from potential impacts because it comprises vegetation communities that are reliant on incident](#)
 320 [rainfall and local runoff and do not include features in the landscape that have potential hydrological](#)
 321 [connectivity to surface water or groundwater features. Secondly, the ‘Human-modified’ landscape](#)
 322 [group is excluded from the ecological impact assessment because it primarily comprises agricultural](#)
 323 [and urban landscapes that are highly modified by human activity. Here the impact assessment focus](#)
 324 [is on economic assets such as groundwater bores, and therefore beyond the scope of this](#)
 325 [publication.](#)

326 2.2.1 ~~Land form~~[Landform](#) classification

327 ~~Land form~~[Landform](#) classification relied on the dominant land type of either habitat or land use
 328 ~~(remnant/human-modified)~~ to determine landscapes that are relatively natural and those that have
 329 been ‘human-modified’. Relatively intact areas are more likely to contain ecological assets, such as
 330 species and ecological communities, than highly modified areas. Location within the region
 331 (topography—upland/lowland, floodplain/non—floodplain), groundwater dependency and water
 332 regime, were part of classifying the landscape. Determining areas that are subjected to flooding, or
 333 that have persistent water, assists in identifying landscapes that support water-dependent habitat
 334 and vegetation, and aquatic ecosystems (Table 1).

335 2.2.2 Stream classification

336 Stream classification in each of the study regions was based on stream position within the
 337 catchment, [\(e.g. upland/lowland\)](#), water regime (perennial/near permanent or
 338 ephemeral/temporary) and dependence and source of groundwater (Table 2). Catchment position is
 339 a potential indicator of stream morphology and flow patterns, while water regime is important when
 340 considering habitat suitability and physical processes within the channel and riparian zone. Streams
 341 can also gain and lose water to local and regional groundwater systems, interacting with
 342 groundwater-dependent ecosystems (Table 2).

343 2.2.3 Spring classification

344 The water source is the basis of spring classification. The source of groundwater is important when
 345 considering regional scale landscape classifications, due to the hydrological connectivity of aquifers
 346 and potential coal resource developments (Table 3).

347 **Table 1. Land form/Landform classification criteria and example datasets**

Characteristic	Classification	Example datasets
Habitat/land use	<ul style="list-style-type: none"> • Non-remnant • Remnant (and stream, Wetland) 	<p>Australian land use mapping (Australian Bureau of Agricultural and Resource Economics and Sciences, 2014) (national)</p> <p>NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)</p>
Topography	<ul style="list-style-type: none"> • Floodplain • Non-floodplain 	<p>NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)</p> <p>Namoi Valley Flood Plain Atlas 1979 (NSW Office of Environment and Heritage, 1979) (regional)</p> <p>Murray-Darling Basin aquatic ecosystem classification (Department of Sustainability, Environment, Water, Population and Communities, 2014) (regional)</p> <p>GEODATA TOPO 250K Series 3 (Geoscience Australia, 2006) (national)</p>
Groundwater	<ul style="list-style-type: none"> • Groundwater dependent (source) • Non-groundwater dependent 	<p>Namoi groundwater dependent ecosystems (NSW Office of Water, 2015) (regional)</p> <p>Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer (Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts, 2015) (state)</p> <p>Queensland Groundwater Dependent Ecosystems (Queensland Department of Science, Information Technology, Innovation and the Arts, 2013) (state)</p>
Water regime	<ul style="list-style-type: none"> • Temporary • Near-permanent • Fresh • Saline 	<p>Queensland wetland data version 3 - wetland areas (Queensland Department of Science, Information Technology, Innovation and the Arts, 2012) (state)</p> <p>Queensland wetland data version 3 - wetland areas (Queensland Department of Science, Information Technology, Innovation and the Arts, 2012) (state)</p> <p>Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer (Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts, 2015) (state)</p> <p>South Australian Wetlands – Groundwater Dependent Ecosystems (GDE) Classification (Sa Department for Water, 2010) (state)</p>
Vegetation	<ul style="list-style-type: none"> • Broad vegetation type 	<p>NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)</p>

348 NSW = New South Wales

349 **Table 2. Stream classification criteria and example datasets**

Characteristic	Classification	Example datasets
Topography	• Upland	Murray-Darling Basin aquatic ecosystem classification (Department of Sustainability, Environment, Water, Population and Communities, 2014) (regional)
	• Lowland	NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional) MrVBF (CSIRO, 2000) (national) MrVBF (CSIRO, 2000) (national)
Groundwater	• Groundwater dependent (source)	Asset database for the Namoi subregion (Bioregional Assessment Programme, 2016) (regional)
	• Non-groundwater dependent	Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer (Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts, 2015) (state) Queensland groundwater dependent ecosystems (Queensland Department of Science, Information Technology, Innovation and the Arts, 2013) (state)
Water regime	• Perennial	Murray-Darling Basin aquatic ecosystem classification (Department of Sustainability, Environment, Water, Population and Communities, 2014) (regional)
	• Ephemeral	Geofabric Surface Cartography (Bureau of Meteorology, 2012) (national)

350

351 **Table 3. Spring classification criteria and example datasets**

Characteristic	Classification	Example datasets
Groundwater	• Groundwater dependent (source)	Asset database for the Namoi subregion (Bioregional Assessment Programme, 2016) (regional)
	• Non-groundwater dependent	Asset database for the Maranoa–Balonne–Condamine subregion (Bioregional Assessment Programme, 2015) (regional) Spring vents assessed for the Surat Underground Water Impact Report 2012 (Office of Groundwater Impact Assessment, 2015) (regional)

352

353 **3 Results**

354 Below we present the resulting landscape classes for the three regions. For each region, we also
 355 combined the landscape classes into ~~groups (landscape groups) to gain efficiencies in a subsequent~~
 356 ~~expert elicitation process. These groups, which~~ were specific to ~~the each~~ region and were based on
 357 distinctions in ~~their~~ topography, ~~their~~ water dependency and association with GAB or non-GAB
 358 GDEs, floodplain/non-floodplain or upland/lowland environments and remnant/human-modified
 359 habitat types. ~~GDEs and remnant/human-modified habitat types.~~ The purpose of the landscape
 360 groups was to combine non-water dependent landscape classes and relate water dependent
 361 landscape classes to region specific aspects of their water dependency, ~~which.~~ This enabled ~~experts~~
 362 ~~to develop a~~ conceptualisation of the landscape for ~~modelling purposes, developing their ecological~~
 363 ~~impact models.~~ While the approach in defining the landscape classes is based on a consistent rule-

364 set and prioritisation, each of the regions has different landscape classes, which is a consequence of
365 the differences in location, jurisdictions and available ~~spatially explicit data~~[spatial datasets](#).

366 [The rule-set deriving from the landform classification \(Tables 1 to 3\) and prioritisation of](#)
367 [hydrological features is the main outcome of our approach and we present the rule-set as a decision](#)
368 [pathway visually below \(Figure 3\). For example, for the Namoi region, the rule-set includes: \(1\)](#)
369 [identify the habitat \(e.g. stream\) \(2\) select by topography \(e.g. upland\), \(3\) identify the groundwater](#)
370 [associations \(e.g. GDE\), and so on until one derives at the final landclass level \(see Figure 3\).](#)

371 **3.1 Landscape classes in the Namoi region**


372 There were 29 landscape classes within six landscape groups in the Namoi region ([Figure 2](#)).[Figure](#)
373 [3](#). Of these landscape groups, 'human-modified' (non-remnant) was the largest (59.3%; Table 4),
374 and included ~~land uses such as~~ urban, agriculture, plantations and other intensive land uses. The
375 dryland remnant vegetation was the second largest landscape group and consisted of the grassy
376 woodland landscape class (24.2%; Table 4). This landscape class was considered non-water
377 dependent as it did not intersect with floodplain, wetland or GDE features. The rainforest landscape
378 group was the smallest (0.5%; Table 4), with only a limited distribution (Figure 4a).

379 The stream network consisted of two landscape groups (floodplain or lowland riverine and non-
380 floodplain or upland riverine). The non-floodplain or upland riverine landscape group had a larger
381 proportion of stream network length (63.8%) compared to the floodplain or lowland riverine
382 landscape group (36.2%; Figure 4b). There were 22 springs identified within the Namoi region, with
383 seven of these associated with the GAB (Figure 4b).

Landscape classification

Topography	Groundwater	Water regime	Vegetation	Landscape class	Landscapes
Upland	GDE	Permanent	1. Permanent upland stream GDE	Non-floodplain
		Temporary	2. Temporary upland stream GDE	
	Non-GDE	Permanent	3. Permanent upland stream	
		Temporary	4. Temporary upland stream	
Lowland	GDE	Permanent	5. Permanent lowland stream GDE	Floodplain
		Temporary	6. Temporary lowland stream GDE	
	Non-GDE	Permanent	7. Permanent lowland stream	
		Temporary	8. Temporary lowland stream	
Floodplain	GDE	9. Floodplain wetland GDE	Non-floodplain	
	Non-GDE	10. Floodplain wetland		
Non-floodplain	GDE	11. Non-floodplain wetland GDE	Non-floodplain
	Non-GDE	12. Non-floodplain wetland	
	GAB GDE	22. GAB springs	
Springs	Non-GAB GDE	23. Non-GAB springs	Springs
	
Floodplain	GDE	Riparian forest	13. Floodplain riparian forest GDE	Floodplain
		Grassy woodlands	14. Floodplain grassy woodland GDE	
	Non-GDE	Riparian forest	15. Floodplain riparian forest	
		Grassy woodlands	16. Floodplain grassy woodland	
Non-floodplain	GDE	Rainforest	17. Rainforest GDE	Non-floodplain
		Riparian forest	18. Upland riparian forest GDE	
	Non-GDE	Grassy woodlands	19. Grassy woodland GDE	
		Rainforest	20. Rainforest	
.....	Grassy woodlands	21. Grassy woodland	Grassy woodlands
		
.....				24. Conservation and natural environments	Human-modified
.....				25. Production from relatively natural environments	
.....				26. Production from dryland agriculture and plantations	
.....				27. Production from irrigated agriculture and plantations	
.....				28. Intensive uses	
.....				29. Water	

385




386
387


Figure 3. Overview of the Namoi landscape classification schema, including criteria and attributes resulting in six landscape groups

388 **Table 4. Percentage of area of each landscape group for the Namoi region**

Landscape group	Percentage of region (%)	Number of classes	Landscape classification attributes
Human-modified	59.3%	6	Conservation of natural environments, production from relatively natural environments, production from dryland agriculture and plantations, production from irrigated agriculture and plantations, intensive uses or waters for production/consumption
Dryland remnant vegetation	24.2%	1	Non-floodplain, non-GDE, grassy woodland remnant vegetation
Non-floodplain or upland riverine	9.8%	8	Upland or non-floodplain streams, wetlands or remnant vegetation
Floodplain or lowland riverine	6.2%	10	Lowland or floodplain streams, wetlands or remnant vegetation
Rainforest	0.5%	2	Non-floodplain GDE or non-GDE rainforest remnant vegetation
Springs	<0.1%	2	GAB or non-GAB springs

389





391

392 Figure 4. (a) Landscape groups (excluding the 'Human-modified'), and (b) Stream network classified as 'upland' or
 393 'lowland' in the landscape classification and 'Springs' within the Namoi region. GAB = Great Artesian Basin
 394 Data: Bureau of Meteorology (2012); Bioregional Assessment Programme (2017)

395 **3.2 Landscape classes in the Galilee region**

396 The Galilee region has 31 landscape classes organised into 11 landscape groups (Figure 5). The
397 dryland landscape group was the largest group within the region and the only group to have no
398 water dependency (68.5%; Table 5). The landscape groups that covered the floodplain areas were
399 the next most dominant classes, with floodplain, terrestrial GDE (12.94%; Table 5) and floodplain,
400 non-wetland (11.8%; Table 5). The remaining three non-floodplain landscape groups consisted of
401 disconnected wetlands, and terrestrial and wetland GDEs (4.9% combined; Table 5).

402 The stream network was classified as groundwater dependent or non-groundwater dependent.
403 Most of the streams in the region were non-GDEs (87.7% compared to 12.3% for the streams, GDE
404 landscape group). There were also over 3000 springs in the region.

Landscape classification

		Landscape classification		
id	Groundwater	Water regime	Vegetation	Landscape class
	GDE	/	Remnant vegetation	Wetland GDE, remnant vegetation
			Non-remnant vegetation	Wetland GDE
	Disconnected	Saline	Remnant vegetation	Floodplain disconnected saline wetland, remnant vegetation
			Non-remnant vegetation	Floodplain disconnected saline wetland
		Non-saline	Remnant vegetation	Floodplain disconnected wetland, remnant vegetation
			Non-remnant vegetation	Floodplain disconnected wetland
GDE	/	Remnant vegetation	Terrestrial GDE, remnant vegetation	
		Non-remnant vegetation	Terrestrial GDE	
id	Disconnected	Non-saline	Remnant vegetation	Floodplain disconnected non-wetland, remnant vegetation
			Non-remnant vegetation	Floodplain disconnected non-wetland
GDE	/	/	Remnant vegetation	Non-floodplain wetland GDE, remnant vegetation
			Non-remnant vegetation	Non-floodplain wetland GDE
	Disconnected	Saline	Remnant vegetation	Non-floodplain disconnected saline wetland, remnant vegeta
			Non-remnant vegetation	Non-floodplain disconnected saline wetland
		Non-saline	Remnant vegetation	Non-floodplain disconnected wetland, remnant vegetation
			Non-remnant vegetation	Non-floodplain disconnected wetland
GDE	/	Remnant vegetation	Non-floodplain terrestrial GDE, remnant vegetation	
		Non-remnant vegetation	Non-floodplain terrestrial GDE	
id	Disconnected	Non-saline	Remnant vegetation	Dryland, remnant vegetation
			Non-remnant vegetation	Dryland
GDE	/	Temporary	Remnant vegetation	Temporary upland GDE stream
			Non-remnant vegetation	Near-permanent upland GDE stream
	Disconnected	Temporary	Remnant vegetation	Temporary upland stream
			Non-remnant vegetation	Near-permanent upland stream
GDE	/	Temporary	Remnant vegetation	Temporary lowland GDE stream
			Non-remnant vegetation	Near-permanent lowland GDE stream
Disconnected	/	Temporary	Remnant vegetation	Temporary lowland stream
			Non-remnant vegetation	Near-permanent lowland stream
	/	Temporary	Remnant vegetation	Near-permanent estuarine stream
			Non-remnant vegetation	Near-permanent estuarine stream
GDE	/	Near-permanent	Remnant vegetation	Spring
			Non-remnant vegetation	Spring

Landscape classification						
Topography	Wetland	Groundwater	Water regime	Vegetation	Landscape class	Landscape group
Vegetation				Remnant vegetation	... Wetland GDE, remnant vegetation	Floodplain, wetland GDE
	Wetland	GDE	<	Non-remnant vegetation	... Wetland GDE	
				Remnant vegetation	... Floodplain disconnected saline wetland, remnant vegetation	
	Wetland	Disconnected	< Saline	Non-remnant vegetation	... Floodplain disconnected saline wetland	Floodplain, disconnected wetland
				Remnant vegetation	... Floodplain disconnected wetland, remnant vegetation	
Floodplain	Non-wetland	GDE	< Non-saline	Non-remnant vegetation	... Floodplain disconnected wetland	
				Remnant vegetation	... Terrestrial GDE, remnant vegetation	Floodplain, terrestrial GDE
				Non-remnant vegetation	... Terrestrial GDE	
				Remnant vegetation	... Floodplain disconnected non-wetland, remnant vegetation	Floodplain, non-wetland
				Non-remnant vegetation	... Floodplain disconnected non-wetland	
				Remnant vegetation	... Non-floodplain wetland GDE, remnant vegetation	Non-floodplain, wetland GDE
	Wetland	GDE	<	Non-remnant vegetation	... Non-floodplain wetland GDE	
				Remnant vegetation	... Non-floodplain disconnected saline wetland, remnant vegetation	
	Wetland	Disconnected	< Saline	Non-remnant vegetation	... Non-floodplain disconnected saline wetland	Non-floodplain, disconnected wetland
				Remnant vegetation	... Non-floodplain disconnected wetland, remnant vegetation	
Non-floodplain	Non-wetland	GDE	< Non-saline	Non-remnant vegetation	... Non-floodplain disconnected wetland	
				Remnant vegetation	... Non-floodplain terrestrial GDE, remnant vegetation	Non-floodplain, terrestrial GDE
				Non-remnant vegetation	... Non-floodplain terrestrial GDE	
				Remnant vegetation	... Dryland, remnant vegetation	Dryland
				Non-remnant vegetation	... Dryland	
Stream network			 Temporary upland GDE stream	Streams, GDE
			 Near-permanent upland GDE stream	Streams, non-GDE
Upland		GDE	< Temporary Temporary upland stream	
			 Near-permanent upland stream	
			 Temporary lowland GDE stream	Streams, GDE
			 Near-permanent lowland GDE stream	
Lowland		GDE	< Temporary Temporary lowland stream	
			 Near-permanent lowland stream	
			 Near-permanent lowland stream	
			 Near-permanent lowland stream	
Estuarine			< Temporary Near-permanent estuarine stream	Streams, non-GDE
			 Near-permanent estuarine stream	
Springs			 Springs	Springs
			 Springs	

Figure 5. Landscape classification of the Galilee region

408

409 **Table 5. Percentage of area of each landscape group for the Galilee region**

Landscape group	Percentage of region (%)	Number of classes	Landscape classification attributes
Dryland	68.5%	2	Non-floodplain, non-wetland, disconnected and saline remnant or non-remnant vegetation
Floodplain, terrestrial GDE	12.9%	2	Floodplain, non-wetland and GDE remnant or non-remnant vegetation
Floodplain, non-wetland	11.8%	2	Floodplain, non-wetland, disconnected and non-saline remnant or non-remnant vegetation
Floodplain, disconnected wetland	1.1%	4	Floodplain, wetland, disconnected saline and non-saline remnant or non-remnant vegetation
Floodplain, wetland GDE	0.8%	2	Floodplain, wetland and GDE remnant or non-remnant vegetation
Non-floodplain, terrestrial GDE	3.4%	2	Non-floodplain, non-wetland and GDE remnant or non-remnant vegetation
Non-floodplain disconnected wetland	1.4%	4	Non-floodplain, wetland, disconnected, saline and non-saline remnant or non-remnant vegetation
Non-floodplain, wetland GDE	<0.1%	2	Non-floodplain, wetland and GDE remnant or non-remnant vegetation
Springs	<0.1%	1	GDE springs

410

411 **3.3 Landscape classes in the Maranoa–Balonne–Condamine region**

412 The landscape classification for the Maranoa–Balonne–Condamine resulted in 34 landscape classes
 413 within five landscape groups (Figure 6). The largest landscape group was the human-modified group
 414 (72.2%, Table 6), which included agricultural production, plantations and other intensive land uses.
 415 Of the remaining landscape groups, dryland remnant vegetation was the second most dominant
 416 (19.8%, Table 6). It was not considered water dependent, because it did not intersect with
 417 floodplain, wetland or GDE features.

418 There ~~are~~ were three landscape groups that cover the stream network. The most dominant
 419 landscape group ~~is~~ was floodplain or lowland riverine (including non-GAB GDEs) (47.8%), followed by
 420 non-floodplain or upland riverine (including non-GAB GDEs) (39.4%) and GAB GDEs (riverine, springs,
 421 floodplain or non-floodplain) (12.7%). There were 177 springs identified within the region. Most of
 422 the springs were GAB GDEs (riverine, springs, floodplain or non-floodplain) (86.4%, compared to
 423 13.6% for non-floodplain or upland riverine (including non-GAB GDEs)).

Landscape classification				
Groundwater	Wetland	Water regime	Landscape class	Landsc
getation				
GAB GDE	Wetland	Near-permanent	Floodplain GAB GDE, near-permanent wetland	GAB GDE
	Non-wetland	Temporary	Floodplain GAB GDE, temporary wetland	springs, f
			Floodplain GAB GDE	non-flood
Non-GAB GDE	Wetland	Near-permanent	Floodplain non-GAB GDE, near-permanent wetland	Floodplai
	Non-wetland	Temporary	Floodplain non-GAB GDE, temporary wetland	riverine (i
			Floodplain non-GAB GDE	GAB GDE
Non GDE	Wetland	Near-permanent	Floodplain, near-permanent wetland	
	Non-wetland	Temporary	Floodplain temporary wetland	
			Floodplain remnant vegetation	
GAB GDE	Wetland	Near-permanent	Non-floodplain GAB GDE, near-permanent wetland	GAB GDE
	Non-wetland	Temporary	Non-floodplain GAB GDE, temporary wetland	springs, f
			Non-floodplain GAB GDE	floodplair
Non-GAB GDE	Wetland	Near-permanent	Non-floodplain non-GAB GDE, near-permanent wetland	Non-flood
	Non-wetland	Temporary	Non-floodplain non-GAB GDE, temporary wetland	riverine (i
			Non-floodplain non-GAB GDE	GAB GDE
Non GDE	Wetland	Near-permanent	Non-floodplain, near-permanent wetland	
	Non-wetland	Temporary	Non-floodplain, temporary wetland	Dryland r
			Dryland remnant vegetation	vegetatio
it vegetation				
			Conservation and natural environments	
			Production from relatively natural environments	
			Production from dryland agriculture and plantations	Human-
			Production from irrigated agriculture and plantations	modified
			Intensive uses	
			Water	
ork				
GAB GDE		Temporary	Temporary upland GAB GDE stream	GAB GDE
Non-GAB GDE		Temporary	Temporary upland non-GAB GDE stream	Non-flood
Non GDE		Near-permanent	Near-permanent upland stream	riverine (i
		Temporary	Temporary upland stream	GAB GDE
GAB GDE		Temporary	Temporary lowland GAB GDE stream	GAB GDE
Non-GAB GDE		Temporary	Temporary lowland non-GAB GDE stream	Floodplai
Non GDE		Near-permanent	Near-permanent lowland stream	riverine (i
		Temporary	Temporary lowland stream	GAB GDE
GAB GDE			GAB springs	GAB GDE
Non-GAB GDE			Non-GAB springs	Non-flood

425
426
427
428

Landscape classification					
Topography	Groundwater	Wetland	Water regime	Landscape class	Landscape group
<u>Remnant vegetation</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary	Floodplain GAB GDE, near-permanent wetland Floodplain GAB GDE, temporary wetland Floodplain GAB GDE	GAB GDEs (riverine, springs, floodplain, non-floodplain)
		Non-wetland	Near-permanent Temporary	Floodplain non-GAB GDE, near-permanent wetland Floodplain non-GAB GDE, temporary wetland	Floodplain or lowland riverine (including non-GAB GDEs)
		Wetland	Near-permanent Temporary	Floodplain non-GAB GDE Floodplain, near-permanent wetland Floodplain, temporary wetland Floodplain remnant vegetation	
<u>Floodplain</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary	Non-floodplain GAB GDE, near-permanent wetland Non-floodplain GAB GDE, temporary wetland Non-floodplain GAB GDE	GAB GDEs (riverine, springs, floodplain, non-floodplain)
		Non-wetland	Near-permanent Temporary	Non-floodplain non-GAB GDE, near-permanent wetland Non-floodplain non-GAB GDE, temporary wetland	Non-floodplain or upland riverine (including non-GAB GDEs)
		Wetland	Near-permanent Temporary	Non-floodplain non-GAB GDE Non-floodplain, near-permanent wetland Non-floodplain, temporary wetland Dryland remnant vegetation	Dryland remnant vegetation
<u>Non-floodplain</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary	Conservation and natural environments Production from relatively natural environments Production from dryland agriculture and plantations Production from irrigated agriculture and plantations Intensive uses Water	Human-modified
		Non-wetland	Near-permanent Temporary	Temporary upland GAB GDE stream Temporary upland non-GAB GDE stream Near-permanent upland stream Temporary upland stream Temporary lowland GAB GDE stream Temporary lowland non-GAB GDE stream Near-permanent lowland stream Temporary lowland stream	GAB GDEs... Non-floodplain or upland riverine (including non-GAB GDEs) GAB GDEs... Floodplain or lowland riverine (including non-GAB GDEs)
		Wetland	Near-permanent Temporary	GAB springs Non-GAB springs	GAB GDEs... Non-floodplain...
<u>Non-remnant vegetation</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary		
		Non-wetland	Near-permanent Temporary		
		Wetland	Near-permanent Temporary		
<u>Stream network</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary		
		Non-wetland	Near-permanent Temporary		
		Wetland	Near-permanent Temporary		
<u>Upland</u>	GAB GDE Non-GAB GDE	Wetland	Near-permanent Temporary		
		Non-wetland	Near-permanent Temporary		
<u>Lowland</u>	GAB GDE Non-GAB GDE	Wetland	Near-permanent Temporary		
		Non-wetland	Near-permanent Temporary		
<u>Springs</u>	GAB GDE Non-GAB GDE	Wetland	Near-permanent Temporary		
		Non-wetland	Near-permanent Temporary		

Figure 6. Landscape classification of the Maranoa–Balonne–Condamine region
 GAB = Great Artesian Basin, GDE = groundwater-dependent ecosystem, GAB GDEs... = GAB GDEs (riverine, springs, floodplain, non-floodplain), Non-floodplain... = Non-floodplain or upland riverine (including or non-GAB GDEs)

429 **Table 6. Percentage of area of each landscape group for the Maranoa–Balonne–Condamine region**

Landscape group	Percentage of region (%)	Number of classes	Landscape classification attributes
Human-modified	72.2%	6	Conservation of natural environments, production from relatively natural environments, production from dryland agriculture and plantations, production from irrigated agriculture and plantations, intensive uses or waters for production/consumption
Dryland remnant vegetation	19.8%	1	Non-floodplain, non-GDE, non-wetland remnant vegetation
Floodplain or lowland riverine (including non-GAB GDEs)	4.5%	9	Floodplain or lowland, non-GAB GDE or non-GDE, temporary or near-permanent wetland, non-wetland or stream
Non-floodplain or upland (including non-GAB GDEs)	2.2%	9	Non-floodplain or upland, non-GAB GDE or non-GDE, temporary or near-permanent wetland, non-wetland, stream or spring
GAB GDEs (riverine, springs, floodplain or non-floodplain)	1.3%	9	Floodplain, non-floodplain or upland GAB-GDE, temporary or near-permanent wetland, non-wetland, stream or spring

430 GAB = Great Artesian Basin, GDE = groundwater-dependent ecosystem

431 **4 Discussion**

432 In Australia, there is no consistent national classification that links ecosystems at landscape level
 433 with their underlying hydrological system. While there are many different land classifications that
 434 incorporate hydrological aspects, they do not provide linkages between hydrology and landscape
 435 elements that enable a broad scale ecological assessment of impacts associated with changes in
 436 water flow and availability, and they are not sufficiently generic for the purpose of assessing
 437 landscape level water related impacts on ecosystems in a spatially explicit manner (Kilroy et al.,
 438 2008; Elmore et al., 2003; Brown et al., 2014; Liermann et al., 2012; Doody et al., 2017; Poff et al.,
 439 2010; Aquatic Ecosystems Task Group, 2012; Olden et al., 2012; Gharari et al., 2011). However, the
 440 bioregional assessment program needed to assess impacts of coal resource [extraction and developments](#)
 441 on ecological systems via a water pathway. Hence, we needed to develop an ecological landscape
 442 classification [for this purpose](#) that [could service would be applicable to](#) the different [regions of the](#)
 443 assessment [regions](#).

444 While our spatially explicit landscape classification provided experts with the ability to readily
 445 identify cause and effect relationships between landscape elements and landscape hydrology, there
 446 are obvious differences between the landscape classifications in the three regions [\(See Figure 2, reflecting their geographical differences \(see Figure 3, Figure 5 and Figure 6\) and this is a reflection of the locations and geographical differences of the regions.\)](#) It provides the specificity that is
 447 required in a regional impact assessment, where the boundaries are based on a combination of
 448 geology, water resources and administrative conditions. The regionality also means that there is
 449 need for different [data sets/datasets](#) describing the landscape features that would not be available
 450 from a classification covering the whole of Australia.
 451
 452

453 Nevertheless, each landscape classification provides a typology with an explicit connection of water
454 to the landscape class. This connection enables a causal [linkage](#) between hydrological change ~~in~~
455 ~~one part of the landscape~~ and impact to ecosystems represented by landscape classes. The causal
456 linkage is dependent on (i) a spatially explicit connection between water in the landscape and the
457 landscape classes, (ii) conceptual understanding how changes in water may result in a reaction of
458 specific ecosystem elements in the landscape class and/or landscape group and (iii) a way of
459 modelling quantitative changes in ecosystem elements related to changes in water. Our
460 ecohydrological classification approach for landscapes provides this spatially explicit connection and
461 has implicit ecohydrological elements that foster the conceptual understanding of the causal linkage.
462 For example, spatially modelling groundwater level drawdown enables a prediction on which
463 landscape elements classified as springs may be experiencing impacts from water extraction and,
464 with additional ecological modelling, by how much and when.

465 Subsequent ecological modelling using expert elicitation of potential impacts drew heavily on our
466 classification, which is based on a consistent rule-set and fosters conceptual understanding of
467 landscape processes and functions. It provides an essential framework for experts to understand and
468 conceptualise how modelled future hydrological changes from coal resource developments link to
469 potential ecological changes at the landscape level. It allows the incorporation of different data
470 sources and existing classification schemes. This consistency also makes the classification
471 development transparent, repeatable, and adjustable, should new data become available.

472 **4.1 In the remainder Application of this section the landscape classification based impact**
473 **assessment**

474 ~~Here~~ we show an application of ~~the how our classification~~ approach ~~in more detail can be used to~~
475 ~~substantiate our claim for~~ ~~assess the potential impact coal resource developments have on ecology~~
476 ~~using the general~~ ~~Namoi region as an example, demonstrating the~~ useability of our classification
477 approach ~~in water mitigated regional impact assessment of human developments.~~

478 **4.1 — Landscape classification based impact assessment**

479 The purpose of developing the landscape classification was to assess the risk of coal resource
480 development on the ecology of a region via a water pathway. ~~Our landscape classification provided~~
481 ~~the spatial framework on which experts can base their assessment of risk from coal resource~~
482 ~~development on the ecology of a region via a water pathway.~~ Details of the predicted changes in
483 ~~groundwater and~~ surface water ~~and groundwater~~ for the Namoi and Galilee regions are in Post et al.
484 (2020). Here, we demonstrate the assessment of potential ecological impacts using the Namoi
485 region. For full details of the analyses in each of the three regions see Holland et al. (2017); Herr et
486 al. (2018b); and Lewis et al. (2018). This work included expert assessment of ecological risk to
487 ecosystem components based on conceptual models. Hence, the models needed to identify water
488 mitigated linkages between hydrological changes, ecosystem components and the landscape classes.
489 ~~This occurred in a 3-step process. We briefly describe the expert assessment approach in a 3-step~~
490 ~~process below. For details we direct the reader to the above references and those listed below.~~

491 ~~In the following we briefly explain the 3-step process to illustrate~~ ~~The following describes an~~
492 ~~application of the landscape classification (see also Figure 1), and in doing so we demonstrate that it~~
493 ~~is a fit-for-purpose in the context of assessing potential ecological impact resulting from potential~~
494 ~~surface water and groundwater changes at different locations within the landscape. The 3-step~~

495 [process illustrates](#) the utility of our landscape classification approach for assessing the risk to
496 ecosystems in the landscape [classes and](#) groups. The process included experts identifying risk to
497 landscape classes using their knowledge on local ecosystems. Specifically, the experts used the
498 broad landscape groups and their underlying hydrogeological features to develop qualitative models
499 initially that then fed into building quantitative models. [Here the experts used outputs from surface](#)
500 [water and groundwater modelling to determine the potential changes in water and what this may](#)
501 [mean for ecological entities within the landscape classes and/or groups.](#) These models assessed the
502 future hydrological changes and risks to the ecosystems in the landscape groups- [\(see also Figure 1\).](#)

503 The [detailed](#) 3 step process included:

504 **Step 1:** Develop qualitative models to conceptualise and prioritise ecosystem components of
505 [theeach](#) landscape class and their linkage to hydrological variables.

506 Here we use the example of the upland riverine landscape class. A qualitative model for the upland
507 riverine landscape class agreed with the existing understanding that a reduction in overbank flows
508 and lowering of the water table resulted in a reduction in several ecosystem components, including
509 riparian habitat, amphibians and fish, and an increase in fine particulate matter, dissolved organic
510 matter and cyanobacteria (Holland et al., 2017; Herr et al., 2018b; Hosack et al., 2018). A qualitative
511 model has, at its basis, the conceptual understanding of ecosystem components and the direction of
512 their interactions, that is [a](#) positive, negative, or neutral influence of one component on another.
513 This understanding also incorporates feedback loops between the ecosystem components in [the](#)
514 form of [digraphs/sign directed graphs](#), and it enables [to direct](#) time intensive quantitative model
515 development to [be directed at](#) variables with the highest importance. The method is based on a
516 matrix level analysis of the component interactions (see for example Herr et al., 2016; Ickowicz et al.,
517 2018).

518 The hydrological variables, and relationships between ecosystem components that the qualitative
519 modelling process prioritised for upland riverine systems [in the Namoi region](#) were the
520 macroinvertebrate responses to riverine system change, presence of tadpoles and changes in
521 projected foliage cover in the riparian trees along the stream channel (Table 7).

522 **Step 2:** Use qualitative model priorities to develop quantitative models.

523 In this context, qualitative models highlighted critical relationships and variables that became the
524 focus of the quantitative models (see for example Herr et al., 2016; Hosack et al., 2018; Ickowicz et
525 al., 2018). [This process helped to focus on those critical ecosystem components that were important](#)
526 [quantitative models for an impact and risk assessment of landscape classes.](#) The focus of the
527 quantitative models was on [three](#) elements within the upland riverine landscape classes (Table 7):
528 (i) the response of upland riparian trees to changes in groundwater; (ii) macroinvertebrate
529 assemblage changes related to days with no consecutive water (zero-flow days) and the longest zero
530 flow event period; and (iii) the response of tadpoles to zero flow days and longest zero flow event
531 period. [Specific](#) Table 7 [provides a brief summary of these variables; specific](#) details of the variable
532 definitions are in Ickowicz et al. (2018).

533

534

535 **Table 7: Upland riverine ecosystem quantitative modelling variables that experts prioritised in the qualitative model and**
 536 **associated ecological and hydrological variables used in the development of the quantitative impact model (after**
 537 **Ickowicz et al., 2018)**

Expert prioritised relationship	Ecological variable (with associated sample units)	Hydrological variable
Response of the upland riparian forest to changes in hydrological regime and groundwater	Annual mean projected foliage cover of species group that includes: Casuarina, yellow box, Blakely's red gum, <i>Acacia salicina</i> , <i>Angophora floribunda</i> , grey box. Transect of 50 m length and 20 m width that extends from first bench ('toe') on both sides of stream	<ul style="list-style-type: none"> • The mean annual number of events with a peak daily flow exceeding the overbank flow events. • Maximum difference in drawdown under a baseline and under the expected drawdown • The year with the maximum difference in drawdown relative to the baseline
Response of fast-water macroinvertebrates to changes in number of zero-flow days and maximum zero-flow event	Average number of families of aquatic macroinvertebrates in riffle habitat sampled using the NSW AUSRIVAS method for riffles	<ul style="list-style-type: none"> • The number of zero-flow days per year, averaged over a 30-year period. • The maximum length of spells (in days per year) with zero flow, averaged over a 30-year period.
Response of tadpoles to changes in number of zero-flow days and maximum zero-flow event	Probability of presence of tadpoles from <i>Limnodynastes</i> genus (species <i>dumerilii</i> , <i>salmini</i> , <i>interioris</i> and <i>terraereginae</i>) sampled using standard 30 cm dip net	<ul style="list-style-type: none"> • The number of zero-flow days per year, averaged over a 30-year period. • The maximum length of spells (in days per year) with zero low, averaged over a 30-year period.

538


539 **Step 3:** Identify risk areas in the regions where quantitative modelling indicated significant changes
540 to landscape group components.

541 This quantitative modelling approach incorporated expert elicitation in a Bayesian framework to
542 predict changes in ecological system components because of expected changes in hydrology
543 conditions. The method dealt with complexity and limited knowledge that allows for updating with
544 new information, which is an important feature in evidence-based decision making (see for example
545 Hosack et al., 2017).

546 The modelling of risk to ecosystems at regional scale focuses on recognising which parts of the
547 region are potentially impacted and which parts are unlikely to experience harm. Using our
548 landscape classification as a crucial input, the modelling delineated impacted areas within each
549 region, based on a zone of potential hydrological change. This is the area in the landscape, where
550 hydrological modelling identified an expected change to surface [water](#) and groundwater from future
551 resource extraction. Risk levels across a landscape group are a result of aggregating individual risks
552 associated with each ecological variable and categorising the risks into three levels based on their
553 percentile spreads (for details see Herr et al., 2018b).

554 For the Namoi [subregion/region](#), for example, dryland remnant vegetation, human-modified
555 ecosystems, no-floodplain and upland riverine ecosystems and rainforests, will not experience
556 impacts, while floodplain and lowland ecosystems area and streams of floodplain and lowland
557 ecosystems will potentially experience impacts (Herr et al., 2018a). Figure 7 (a) shows the landscape
558 groups that are at risk of impact from hydrological changes as they are situated within the zone of
559 potential hydrological change, and Figure 7 (b) shows the risk level to these landscape groups from
560 the quantitative models. Note that there is a category “Remaining unquantified ‘floodplain and
561 lowland riverine’ classes”. The expert could not develop quantitative models for these classes,
562 because there was no surface water hydrological model available that could predict changes to
563 surface water flows. This was related to the lack of gauging data and groundwater interaction details
564 specific to the lowland drainage channels. Having lowland riverine classes whose risk remains
565 unquantified means there is additional work needed before an assessment and potential mitigation
566 of impacts from hydrological changes is possible (Herr et al., 2018b).

567




568

569

Figure 7a: Landscape classes overlaying areas of potential hydrological change (Herr et al., 2018b)

570

571



572

573

574

Figure 7b: Hydrological change risk level of lowland landscape class areas (Herr et al., 2018b)

575 **4.2 Limitations**

576 While the ecohydrological landscape classification approach provided the basis for the risk
577 assessment outlined above, there are some limitations that require consideration when attempting
578 to develop and apply this ecohydrological landscape classification approach.

579 An important issue for the landscape classification is formulating a typology that adequately reflects
580 both the functional and structural complexity of the ecosystem, while delivering a succinct and
581 consistent representation of the system that is 'fit for purpose' to assign hydrological connectivity
582 between the landscape classes, and within the general landscape. The systematic classification
583 imposes discrete boundaries among landscape components that may not adequately capture
584 gradients within and across landscape classes. This approach tends to simplify important
585 components of ecotones such as 'transition' zones or edges between landscape classes, where
586 ecosystem processes and/or biodiversity are likely to peak. If landscape classes are treated purely as
587 'closed' ecosystems, then the result may be a poor representation of the biotic interactions and
588 energy exchange between adjacent systems, and this could limit a conventional impact and risk
589 analyses. These conceptual challenges may be important considerations for subsequent impact
590 assessments, requiring special attention in assigning risk from human induced changes in hydrology.
591 However, conceptual modelling of impacts may be able to compensate for this shortfall, when for
592 example, incorporating riparian areas within riverine and wetland model development.

593 There are also spatial data issues that require additional consideration beyond just simply
594 incorporating existing data. There are several technical issues that constitute important gaps in the
595 landscape classification for the Namoi region, for example. Here, two different approaches to define
596 GDEs were required because one spatial dataset only included terrestrial vegetation and not riverine
597 systems mapped within the stream network (NSW Office of Water, 2015). A second GDE dataset
598 helped overcome this deficiency, and provided the basis to classify the stream network's
599 dependency on groundwater (Bioregional Assessment Programme, 2012).

600 Wetlands in large areas of Australia are not yet adequately mapped. The separation between
601 groundwater-dependent and surface water-dependent wetlands may not always be accurate. In
602 many areas there is little knowledge of groundwater – surface water interactions. There is also a
603 significant gap in the understanding of water thresholds for ecosystems associated with springs. In
604 part, this results from a lack of bores to provide meaningful groundwater data. Some examples of
605 these data gaps appear in the discussion of the functioning of springs in the Doongmabulla Springs
606 complex in the Galilee region, particularly in identifying the source aquifer (Fensham et al., 2016).

607 There is extensive work from Queensland that links regional ecosystems vegetation to their
608 groundwater needs, although the mapped areas are still small (Sattler and Williams, 1999;
609 Queensland Government, Queensland, 2016; Queensland Herbarium, 2021). However, in many parts
610 of Australia, GDE mapping and classification approaches are limited, and many areas lack systematic
611 ground-truthing. This is especially true in areas with extensive intact native vegetation remnants,
612 such as the Pilliga Forest of the Namoi region, where large areas of 'Grass woodland GDE' landscape
613 class exist, but the lack of published studies on vegetation–groundwater interactions limits a
614 definition of the nature of this interaction.

615 4.3 Conclusions

616 We showed that our [landscape classification](#) approach [worksworke](#)d in the three geographically
617 different regions, with widely disparate information sources that [feedfed](#) into [thea](#) landscape
618 classification. This also makes the approach resource efficient where existing spatial landscape or
619 ecosystem classification schemes, developed for other purposes, can be incorporated into the
620 classification.

621 The study was able to formulate and implement an attribute-based classification scheme to define
622 and delineate water-dependent features across three large regions. We conclude that this approach
623 allowed us to repurpose several existing schemas into an adaptable and practical typology of a
624 landscape classification. The conceptual framework of landscape ecohydrology forms the basis for
625 this classification, which is used to focus subsequent analysis of potential cumulative impacts on
626 water resources from multiple coal resource developments. The classification enabled the
627 development of specific conceptual and qualitative models that linked changes in hydrology to
628 potential impacts on ecosystems using the landscape classes. The classification provided crucial
629 inputs for a risk analysis of landscape components subjected to hydrological changes.

630 Applying our approach to different regions showed that it is sufficiently general and flexible to
631 enable the development of ecohydrological classifications in regions in Australia and potentially
632 [globally in other regions around the globe](#), given a sufficiently mature information base and data
633 availability.

634 5 References

635 Abella, S. R., Shelburne, V. B., and MacDonald, N. W.: Multifactor classification of forest landscape
636 ecosystems of Jocassee Gorges, southern Appalachian Mountains, South Carolina, Can J For Res, 33,
637 1933-1946, <https://doi.org/10.1139/x03-116>, 2003.

638 Addicott, E., Neldner, V. J., and Ryan, T.: Aligning quantitative vegetation classification and landscape
639 scale mapping: updating the classification approach of the Regional Ecosystem classification system
640 used in Queensland, Australian Journal of Botany, 69, 400-413, <https://doi.org/10.1071/BT20108>,
641 2021.

642 Aquatic Ecosystems Task Group: Aquatic Ecosystems Toolkit. Module 1: Aquatic Ecosystems Toolkit
643 Guidance Paper., Australian Government Department of Sustainability, Environment, Water,
644 Population and Communities, Canberra, [https://www.awe.gov.au/water/publications/aquatic-
645 ecosystems-toolkit-module-1-guidance-paper](https://www.awe.gov.au/water/publications/aquatic-ecosystems-toolkit-module-1-guidance-paper), 2012.

646 Australian Bureau of Agricultural and Resource Economics and Sciences: Catchment Scale Land Use
647 of Australia - 2014, Bioregional Assessment Source Dataset. [dataset],
648 <http://data.bioregionalassessments.gov.au/dataset/6f72f73c-8a61-4ae9-b8b5-3f67ec918826>, 2014.

649 Bioregional Assessment Programme: National Groundwater Dependent Ecosystems (GDE) Atlas,
650 Bioregional Assessment Derived Dataset [dataset],
651 <http://data.bioregionalassessments.gov.au/dataset/e358e0c8-7b83-4179-b321-3b4b70df857d>,
652 2012.

653 Bioregional Assessment Programme: Asset database for the Maranoa-Balonne-Condamine
654 subregion on 26 June 2015 [dataset], [http://data.bioregionalassessments.gov.au/dataset/35e95025-
655 f962-4425-83c7-767e2d6722e6](http://data.bioregionalassessments.gov.au/dataset/35e95025-f962-4425-83c7-767e2d6722e6), 2015.

656 Bioregional Assessment Programme: Asset database for the Namoi subregion on 18 February 2016,
657 Bioregional Assessment Derived Dataset [dataset],
658 <http://data.bioregionalassessments.gov.au/dataset/22061f2c-e86d-4ca8-9860-c349c2513fd8>, 2016.

659 Bioregional Assessment Programme: Landscape classification of the Namoi preliminary assessment
660 extent. Bioregional Assessment Derived Dataset. [dataset],
661 <http://data.bioregionalassessments.gov.au/dataset/360c39e5-1225-401d-930b-f5462fdb8005>,
662 2017.

663 Bioregional Assessments: Bioregional Assessment Programme,
664 <https://www.bioregionalassessments.gov.au/bioregional-assessment-program>, last access:
665 26/08/2019, 2018.

666 Bioregional Assessments: Geological and Bioregional Assessment Program,
667 <https://www.bioregionalassessments.gov.au/geological-and-bioregional-assessment-program>, last
668 access: 26/08/2019, 2019.

669 Brown, S. C., Lester, R. E., Versace, V. L., Fawcett, J., and Laurenson, L.: Hydrologic Landscape
670 Regionalisation Using Deductive Classification and Random Forests, PLoS ONE, 9,
671 <https://doi.org/10.1371/journal.pone.0112856>, 2014.

672 Bureau of Meteorology: Australian Hydrological Geospatial Fabric ('Geofabric'), version 2.1.1,
673 Canberra,
674 [http://www.bom.gov.au/water/geofabric/documents/v2_1/ahgf_dps_surface_cartography_V2_1_r](http://www.bom.gov.au/water/geofabric/documents/v2_1/ahgf_dps_surface_cartography_V2_1_release.pdf)
675 [elease.pdf](http://www.bom.gov.au/water/geofabric/documents/v2_1/ahgf_dps_surface_cartography_V2_1_release.pdf), 2012.

676 Carlier, J., Doyle, M., Finn, J. A., Ó hUallacháin, D., and Moran, J.: A landscape classification map of
677 Ireland and its potential use in national land use monitoring, Journal of Environmental Management,
678 289, 112498, <https://doi.org/10.1016/j.jenvman.2021.112498>, 2021.

679 CSIRO: Multi-resolution Valley Bottom Flatness MrVBF at three second resolution CSIRO 20000211
680 [dataset], [http://data.bioregionalassessments.gov.au/dataset/7dfc93bb-62f3-40a1-8d39-](http://data.bioregionalassessments.gov.au/dataset/7dfc93bb-62f3-40a1-8d39-0c0f27a83cb3)
681 [0c0f27a83cb3](http://data.bioregionalassessments.gov.au/dataset/7dfc93bb-62f3-40a1-8d39-0c0f27a83cb3), 2000.

682 Cullum, C., Brierley, G., Perry, G., and Witkowski, E.: Landscape archetypes for ecological
683 classification and mapping: The virtue of vagueness, Prog Phys Geogr, 41,
684 <http://doi.org/10.1177/0309133316671103>, 2016a.

685 Cullum, C., Rogers, K. H., Brierley, G., and Witkowski, E. T. F.: Ecological classification and mapping
686 for landscape management and science, Prog Phys Geogr, 40, 38-65,
687 <https://doi.org/10.1177/0309133315611573>, 2016b.

688 Department of Agriculture, W. a. t. E.: National Vegetation Information System (NVIS),
689 [https://www.awe.gov.au/agriculture-land/land/native-vegetation/national-vegetation-information-](https://www.awe.gov.au/agriculture-land/land/native-vegetation/national-vegetation-information-system)
690 [system](https://www.awe.gov.au/agriculture-land/land/native-vegetation/national-vegetation-information-system), last access: 22 Mar 2022, 2021.

691 Department of Sustainability, Environment, Water, Population and Communities: Murray-Darling
692 Basin aquatic ecosystem classification [dataset],
693 <http://data.bioregionalassessments.gov.au/dataset/a854a25c-8820-455c-9462-8bd39ca8b9d6>,
694 2014.

695 Doody, T. M., Barron, O. V., Dowsley, K., Emelyanova, I., Fawcett, J., Oyerton, I. C., Pritchard, J. L.,
696 Van Dijkf, A. I. J. M., and Warren, G.: Continental mapping of groundwater dependent ecosystems: A
697 methodological framework to integrate diverse data and expert opinion, *Journal of Hydrology-*
698 *Regional Studies*, 10, 61-81, <https://doi.org/10.1016/j.ejrh.2017.01.003>, 2017.

699 Eigenbrot, F.: Redefining Landscape Structure for Ecosystem Services, *Current Landscape Ecology*
700 *Reports*, 1, 80-86, <http://doi.org/10.1007/s40823-016-0010-0>, 2016.

701 Elmore, A. J., Mustard, J. F., and Manning, S. J.: Regional patterns of plant community response to
702 changes in water: Owens Valley, California, *Ecological Applications*, 13, 443-460,
703 [https://doi.org/10.1890/1051-0761\(2003\)013\[0443:RPOPCR\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)013[0443:RPOPCR]2.0.CO;2), 2003.

704 Evans, T., Tan, K., Magee, J., Karim, F., Sparrow, A., Lewis, S., Marshall, S., Kellett, J., and Galinec, V.:
705 Context statement for the Galilee subregion. Product 1.1 from the Lake Eyre Basin Bioregional
706 Assessment, Department of the Environment, Bureau of Meteorology, CSIRO and Geoscience
707 Australia, Australia, <http://data.bioregionalassessments.gov.au/product/LEB/GAL/1.1>, 2014.

708 Fensham, R., Silcock, J., Laffineur, B., and MacDermott, H.: Lake Eyre Basin Springs Assessment
709 Project: hydrogeology, cultural history and biological values of springs in the Barcardine, Springvale
710 and Flinders River supergroups, Galilee Basin springs and Tertiary springs of western Queensland.
711 Report to Office of Water Science, Department of Science, Information Technology and Innovation,
712 Brisbane., [https://publications.qld.gov.au/dataset/11c1af89-93b9-497a-b99f-
713 2ec6c7a8d339/resource/c5d1813b-73a4-4e05-aa86-39a8ed3045fb/download/lebsa-hchb-report-
714 springs-wst-qld.pdf](https://publications.qld.gov.au/dataset/11c1af89-93b9-497a-b99f-2ec6c7a8d339/resource/c5d1813b-73a4-4e05-aa86-39a8ed3045fb/download/lebsa-hchb-report-springs-wst-qld.pdf), 2016.

715 Geoscience Australia: GEODATA TOPO 250K Series 3 [dataset],
716 <http://data.bioregionalassessments.gov.au/dataset/a0650f18-518a-4b99-a553-44f82f28bb5f>, 2006.

717 Gharari, S., Hrachowitz, M., Fenicia, F., and Savenije, H. H. G.: Hydrological landscape classification:
718 investigating the performance of HAND based landscape classifications in a central European meso-
719 scale catchment, *Hydrol. Earth Syst. Sci.*, 15, 3275-3291, [https://doi.org/10.5194/hess-15-3275-
720 2011](https://doi.org/10.5194/hess-15-3275-2011), 2011.

721 [Gleeson, T. and Paszkowski, D.: Perceptions of scale in hydrology: what do you mean by regional](#)
722 [scale?](#), *Hydrological Sciences Journal*, 59, 99-107, 10.1080/02626667.2013.797581, 2014.

723 Hall, J., Storey, D., Piper, V., Bolton, E., Woodford, A., and Jolly, J.: Ecohydrological conceptualisation
724 for the eastern Pilbara region. A report prepared for BHP Billiton Iron Ore., Subiaco, 240,
725 [https://www.bhp.com/-/media/bhp/regulatory-information-media/iron-ore/western-australia-iron-
726 ore/0000/report-
727 appendices/160316_ironore_waio_pilbarastrategicassessment_state_appendix7_appendixd.pdf](https://www.bhp.com/-/media/bhp/regulatory-information-media/iron-ore/western-australia-iron-ore/0000/report-appendices/160316_ironore_waio_pilbarastrategicassessment_state_appendix7_appendixd.pdf),
728 2015.

729 Hawkins, C. P. and Norris, R. H.: Performance of different landscape classifications for aquatic
730 bioassessments: introduction to the series, *Journal of the North American Benthological Society*, 19,
731 367-369, <https://doi.org/10.1899/0887-3593-19.3.367>, 2000.

732 Hawkins, C. P., Norris, R. H., Gerritsen, J., Hughes, R. M., Jackson, S. K., Johnson, R. K., and Stevenson,
733 R. J.: Evaluation of the use of landscape classifications for the prediction of freshwater biota:
734 synthesis and recommendations, *Journal of the North American Benthological Society*, 19, 541-556,
735 <https://doi.org/10.2307/1468113> 2000.

736 Herr, A., Dambacher, J. M., Pinkard, E., Glen, M., Mohammed, C., and Wardlaw, T.: The uncertain
737 impact of climate change on forest ecosystems How qualitative modelling can guide future research
738 for quantitative model development, *Environ. Model. Software*, 76, 95-107,
739 <https://doi.org/10.1016/j.envsoft.2015.10.023>, 2016.

740 Herr, A., Brandon, C., Beringen, H., Merrin, L. E., Post, D. A., Mitchell, P. J., Crosbie, R., Aryal, S. K.,
741 Janarhanan, S., Schmidt, R. K., and Henderson, B. L.: Assessing impacts of coal resource development
742 on water resources in the Namoi subregion: key findings. Product 5: Outcome synthesis for the
743 Namoi subregion from the Northern Inland Catchments Bioregional Assessment,
744 <http://data.bioregionalassessments.gov.au/product/NIC/NAM/5>, 2018a.

745 Herr, A., Aryal, S. K., Brandon, C., Crawford, D., Crosbie, R., Davies, P., Dunne, R., Gonzalez, D., Hayes,
746 K. R., Henderson, B. L., Hosack, G., Ickowicz, A., Janarhanan, S., Marvanek, S., Mitchell, P. J., Merrin,
747 L. E., Herron, N. F., O'Grady, A. P., and Post, D. A.: Impact and risk analysis for the Namoi subregion.
748 Product 3-4 for the Namoi subregion from the Northern Inland Catchments Bioregional Assessment,
749 [https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-namoi-](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-namoi-subregion)
750 [subregion](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-namoi-subregion), 2018b.

751 Hobbs, R. J. and McIntyre, S.: Categorizing Australian landscapes as an aid to assessing the generality
752 of landscape management guidelines, *Global Ecology and Biogeography*, 14, 1-15,
753 <https://doi.org/10.1111/j.1466-822X.2004.00130.x>, 2005.

754 Holland, K., Beringen, H., Brandon, C., Crosbie, R., Davies, P., Gonzalez, D., Henderson, B.,
755 Janardhanan, S., Lewis, S., Merrin, L., Mitchell, P., Mount, R., O'Grady, A., Peeters, L., Post, D.,
756 Schmidt, R., Sudholz, C., and Turnadge, C.: Impact and risk analysis for the Maranoa-Balonne-
757 Condamine subregion. Product 3-4 for the Maranoa-Balonne-Condamine subregion from the
758 Northern Inland Catchments Bioregional Assessment, Department of the Environment and Energy,
759 Bureau of Meteorology, CSIRO and Geoscience Australia, Australia,
760 <http://data.bioregionalassessments.gov.au/product/NIC/MBC/3-4>, 2017.

761 Hosack, G., Ickowicz, A., Hayes, K. R., Dambacher, J. M., Barry, S. A., and Henderson, B. L.: Receptor
762 impact modelling. Submethodology M08 from the Bioregional Assessment Technical Programme,
763 <http://data.bioregionalassessments.gov.au/submethodology/M08>, 2018.

764 Hosack, G. R., Hayes, K. R., and Barry, S. C.: Prior elicitation for Bayesian generalised linear models
765 with application to risk control option assessment, *Reliab. Eng. Syst. Saf.*, 167, 351-361,
766 <https://doi.org/10.1016/j.res.2017.06.011>, 2017.

767 Ickowicz, A., Hosack, G., Mitchell, P. J., Dambacher, J. M., Hayes, K. R., O'Grady, A. P., Henderson, B.
768 L., and Herron, N. F.: Receptor impact modelling for the Namoi subregion. Product 2.7 for the Namoi
769 subregion from the Northern Inland Catchments Bioregional Assessment,
770 <http://data.bioregionalassessments.gov.au/product/NIC/NAM/2.7>, 2018.

771 Jones ~~Jr~~, C. E. J., Leibowitz, S. G., Sawicz, K. A., Comeleo, R. L., Stratton, L. E., Morefield, P. E., and
772 Weaver, C. P.: Using hydrologic landscape classification and climatic time series to assess hydrologic
773 vulnerability of the western U.S. to climate, *Hydrol. Earth Syst. Sci.*, 25, 3179-3206,
774 <https://doi.org/10.5194/hess-25-3179-2021>, 2021.

775 Kilroy, G., Ryan, J., Coxon, C., and Daly, D.: A Framework for the Assessment of Groundwater –
776 Dependent Terrestrial Ecosystems under the WaterFramework Directive,
777 <http://erc.epa.ie/safer/resource?id=b5799c70-224b-102c-b381-901ddd016b14>, 2008.

778 Leathwick, J. R., Overton, J. M., and McLeod, M.: An Environmental Domain Classification of New
779 Zealand and Its Use as a Tool for Biodiversity Management, *Conservation Biology*, 17, 1612-1623,
780 <http://dx.doi.org/10.1111/j.1523-1739.2003.00469.x>, 2003.

781 Leibowitz, S. G., Comeleo, R. L., Wigington Jr, P. J., Weaver, C. P., Morefield, P. E., Sproles, E. A., and
782 Ebersole, J. L.: Hydrologic landscape classification evaluates streamflow vulnerability to climate
783 change in Oregon, USA, *Hydrol. Earth Syst. Sci.*, 18, 3367-3392, <http://10.5194/hess-18-3367-2014>,
784 2014.

785 Lewis, S., Evans, T., Pavey, C., Holland, K., Henderson, B., Kilgour, P., Dehelean, A., Karim, F., Viney,
786 N., Post, D., Schmidt, R., Sudholz, C., Brandon, C., Zhang, Y., Lymburner, L., Dunn, B., Mount, R.,
787 Gonzalez, D., Peeters, L., O'Grady, A., Dunne, R., Ickowicz, A., Hosack, G., Hayes, K., Dambacher, J.,
788 and Barry, S.: Impact and risk analysis for the Galilee subregion. Product 3-4 for the Galilee subregion
789 from the Lake Eyre Basin Bioregional Assessment., Department of the Environment and Energy,
790 Bureau of Meteorology, CSIRO and Geoscience Australia, Australia, 348,
791 [https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-galilee-](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-galilee-subregion)
792 [subregion](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-galilee-subregion), 2018.

793 Liermann, C. A. R., Olden, J. D., Beechie, T. J., Kennard, M. J., Skidmore, P. B., Konrad, C. P., and
794 Imaki, H.: Hydrogeomorphic Classification of Washington State Rivers to Support Emerging
795 Environmental Flow Management Strategies, *River Research and Applications*, 28, 1340-1358,
796 <http://doi.org/10.1002/rra.1541>, 2012.

797 MacMillan, R., Martin, T., Earle, T., and McNabb, D.: Automated analysis and classification of
798 landforms using high-resolution digital elevation data: applications and issues, *Canadian Journal of*
799 *Remote Sensing*, 29, 592-606, <https://doi.org/10.5589/m03-031>, 2003.

800 McMahon, G., Gregonis, S. M., Waltman, S. W., Omernik, J. M., Thorson, T. D., Freeouf, J. A., Rorick,
801 A. H., and Keys, J. E.: Developing a spatial framework of common ecological regions for the
802 conterminous United States, *Environmental Management*, 28, 293-316,
803 <http://doi.org/10.1007/s0026702429> 2001.

804 NSW Office of Environment and Heritage: Namoi Valley Flood Plain Atlas 1979, Bioregional
805 Assessment Source Dataset [dataset], [http://data.bioregionalassessments.gov.au/dataset/a854a25c-](http://data.bioregionalassessments.gov.au/dataset/a854a25c-8820-455c-9462-8bd39ca8b9d6)
806 [8820-455c-9462-8bd39ca8b9d6](http://data.bioregionalassessments.gov.au/dataset/a854a25c-8820-455c-9462-8bd39ca8b9d6), 1979.

807 NSW Office of Environment and Heritage: Border Rivers Gwydir / Namoi Regional Native Vegetation
808 Map Version 2.0 VIS_ID_2004 [dataset],
809 <http://data.bioregionalassessments.gov.au/dataset/b3ca03dc-ed6e-4fdd-82ca-e9406a6ad74a>, 2015.

810 NSW Office of Water: Namoi CMA Groundwater Dependent Ecosystems [dataset],
811 <https://data.gov.au/data/dataset/a3e21ec4-ae53-4222-b06c-0dc2ad9838a8>, 2015.

812 NVIS Technological Working Group: Australian Vegetation Attribute Manual: National Vegetation
813 Information System, Version 7.0, Canberra,
814 [https://www.environment.gov.au/system/files/resources/292f10e2-8670-49b6-a72d-](https://www.environment.gov.au/system/files/resources/292f10e2-8670-49b6-a72d-25e892a92360/files/australian-vegetation-attribute-manual-v70.pdf)
815 [25e892a92360/files/australian-vegetation-attribute-manual-v70.pdf](https://www.environment.gov.au/system/files/resources/292f10e2-8670-49b6-a72d-25e892a92360/files/australian-vegetation-attribute-manual-v70.pdf), 2017.

816 Office of Groundwater Impact Assessment: Spring vents assessed for the Surat Underground Water
817 Impact Report 2012 [dataset], [http://data.bioregionalassessments.gov.au/dataset/6d2b59fc-e312-](http://data.bioregionalassessments.gov.au/dataset/6d2b59fc-e312-4c89-9f10-e1f1b20a7a6d)
818 [4c89-9f10-e1f1b20a7a6d](http://data.bioregionalassessments.gov.au/dataset/6d2b59fc-e312-4c89-9f10-e1f1b20a7a6d), 2015.

819 Olden, J. D., Kennard, M. J., and Pusey, B. J.: A framework for hydrologic classification with a review
820 of methodologies and applications in ecohydrology, *Ecohydrology*, 5, 503-518,
821 <http://dx.doi.org/10.1002/eco.251>, 2012.

822 Pain, C., Gregory, L., Wilson, P., and McKenzie, N.: The physiographic regions of Australia –
823 Explanatory notes 2011, CSIRO, Canberra,
824 <https://publications.csiro.au/rpr/download?pid=csiro:EP113843&dsid=DS4>, 2011.

825 Poff, N. L., Richter, B. D., Arthington, A. H., Bunn, S. E., Naiman, R. J., Kendy, E., Acreman, M., Apse,
826 C., Bledsoe, B. P., Freeman, M. C., Henriksen, J., Jacobson, R. B., Kennen, J. G., Merritt, D. M.,
827 O’Keeffe, J. H., Olden, J. D., Rogers, K., Tharme, R. E., and Warner, A.: The ecological limits of
828 hydrologic alteration (ELOHA): a new framework for developing regional environmental flow
829 standards, *Freshwater Biology*, 55, 147-170, <https://doi.org/10.1111/j.1365-2427.2009.02204.x>,
830 2010.

831 Post, D. A., Crosbie, R. S., Viney, N. R., Peeters, L. J., Zhang, Y., Herron, N. F., Janardhanan, S., Wilkins,
832 A., Karim, F., Aryal, S. K., Pena-Arancibia, J., Lewis, S., Evans, T., Vaze, J., Chiew, F. H. S., Marvanek, S.,
833 Henderson, B. L., Schmidt, B., and Herr, A.: Impacts of coal resource development in eastern
834 Australia on groundwater and surface water, *Journal of Hydrology*, 591,
835 <https://doi.org/10.1016/j.jhydrol.2020.125281>, 2020.

836 Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S., and Zimmermann, N. E.: Plant
837 functional type mapping for earth system models, *Geosci. Model Dev.*, 4, 993-1010,
838 <https://doi.org/10.5194/gmd-4-993-2011>, 2011.

839 Pyne, M. I., Carlisle, D. M., Konrad, C. P., and Stein, E. D.: Classification of California streams using
840 combined deductive and inductive approaches: Setting the foundation for analysis of hydrologic
841 alteration, *Ecohydrology*, 10, <https://doi.org/10.1002/eco.1802>, 2017.

842 Queensland Department of Science, Information Technology, Innovation and the Arts: Queensland
843 wetland data version 3 - wetland areas [dataset],
844 <http://data.bioregionalassessments.gov.au/dataset/2a187a00-b01e-4097-9ca4-c9683e7f4786>,
845 2012.

846 Queensland Department of Science, Information Technology, Innovation and the Arts: Queensland
847 groundwater dependent ecosystems [dataset],
848 <http://data.bioregionalassessments.gov.au/dataset/10940dfa-d7ef-44fb-8ac2-15d75068fff8>, 2013.

849 Queensland Government, Queensland: Groundwater dependent ecosystem mapping background,
850 <https://wetlandinfo.des.qld.gov.au/wetlands/facts-maps/gde-background/>, last access: 09 Sep 21,
851 2016.

852 Queensland Herbarium: Regional Ecosystem Description Database (REDD). Version 12 (March 2021),
853 Brisbane, [https://www.qld.gov.au/environment/plants-
854 animals/plants/ecosystems/descriptions/download](https://www.qld.gov.au/environment/plants-animals/plants/ecosystems/descriptions/download), 2021.

855 Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts:
856 Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer 20150714
857 [dataset], [http://data.bioregionalassessments.gov.au/dataset/3d36e3d4-b16b-43b3-b2eb-
858 c1aea7ef9193](http://data.bioregionalassessments.gov.au/dataset/3d36e3d4-b16b-43b3-b2eb-c1aea7ef9193), 2015.

- 859 SA Department for Water: South Australian Wetlands – Groundwater Dependent Ecosystems (GDE)
 860 Classification [dataset], [http://data.bioregionalassessments.gov.au/dataset/fc35d75a-f12e-494b-](http://data.bioregionalassessments.gov.au/dataset/fc35d75a-f12e-494b-a7d3-0f27e7159b05)
 861 [a7d3-0f27e7159b05](http://data.bioregionalassessments.gov.au/dataset/fc35d75a-f12e-494b-a7d3-0f27e7159b05), 2010.
- 862 Sattler, P. S. and Williams, R. D.: The conservation status of Queensland's bioregional ecosystems,
 863 Environmental Protection Agency, Brisbane 1999.
- 864 Sawicz, K. A., Kelleher, C., Wagener, T., Troch, P., Sivapalan, M., and Carrillo, G.: Characterizing
 865 hydrologic change through catchment classification, *Hydrol. Earth Syst. Sci.*, 18, 273-285,
 866 <https://doi.org/10.5194/hess-18-273-2014>, 2014.
- 867 Snelder, T. H., Cattaneo, F., Suren, A. M., and Biggs, B. J.: Is the River Environment Classification an
 868 improved landscape-scale classification of rivers?, *Journal of the North American Benthological*
 869 *Society*, 23, 580-598, [https://doi.org/10.1899/0887-3593\(2004\)023<0580:ITRECA>2.0.CO;2](https://doi.org/10.1899/0887-3593(2004)023<0580:ITRECA>2.0.CO;2), 2004.
- 870 Thackway, R. and Cresswell, I. D.: An Interim Biogeographic Regionalisation for Australia: A
 871 framework for setting priorities in the national reserves system cooperative program, Australian
 872 Nature Conservancy Agency, Canberra, 99,
 873 [https://www.environment.gov.au/system/files/resources/4263c26f-f2a7-4a07-9a29-](https://www.environment.gov.au/system/files/resources/4263c26f-f2a7-4a07-9a29-b1a81ac85acc/files/ibra-framework-setting-priorities-nrs-cooperative-program.pdf)
 874 [b1a81ac85acc/files/ibra-framework-setting-priorities-nrs-cooperative-program.pdf](https://www.environment.gov.au/system/files/resources/4263c26f-f2a7-4a07-9a29-b1a81ac85acc/files/ibra-framework-setting-priorities-nrs-cooperative-program.pdf), 1995.
- 875 Welsh, W., Herron, N., Rohead-O'Brien, H., Ransley, T., Aryal, S., Mitchell, P., Buckerfield, S., and
 876 Marshall, S.: Context statement for the Maranoa-Balonne-Condamine subregion. Product 1.1 for the
 877 Maranoa-Balonne-Condamine from the Northern Inland Catchments Bioregional Assessment,
 878 Department of the Environment, Bureau of Meteorology, CSIRO and Geoscience Australia, Australia,
 879 <http://data.bioregionalassessments.gov.au/product/LNIC/MBC/1.1>, 2015.
- 880 Welsh, W., Hodgkinson, J., Strand, J., Northey, J., Aryal, S., O'Grady, A., Slatter, E., Herron, N.,
 881 Pinetown, K., Carey, H., Yates, G., Raisbeck-Brown, N., and Lewis, S.: Context statement for the
 882 Cooper subregion. Product 1.1 for the Namoi subregion from the Northern Inland Catchments
 883 Bioregional Assessment. Department of the Environment, Bureau of Meteorology, Department of
 884 the Environment, Bureau of Meteorology, CSIRO and Geoscience Australia, Australia,
 885 <http://data.bioregionalassessments.gov.au/product/NIC/NAM/1.1>, 2014.
- 886 Wiens, J. A. and Milne, B. T.: Scaling of 'landscapes' in landscape ecology, or, landscape ecology from
 887 a beetle's perspective, *Landscape Ecology*, 3, 87-96, <https://doi.org/10.1007/BF00131172>, 1989.
- 888 Wolfe, J. D., Shook, K. R., Spence, C., and Whitfield, C. J.: A watershed classification approach that
 889 looks beyond hydrology: application to a semi-arid, agricultural region in Canada, *Hydrol. Earth Syst.*
 890 *Sci.*, 23, 3945-3967, <https://doi.org/10.5194/hess-23-3945-2019>, 2019.
- 891 Zhang, Z., Zang, R., Wang, G., and Huang, X.: Classification of Landscape Types Based on Land Cover,
 892 Successional Stages and Plant Functional Groups in a Species-Rich Forest in Hainan Island, China,
 893 *Tropical Conservation Science*, 9, 135-152, <https://doi.org/10.1177/194008291600900107>, 2016.

894

895

896 **6 Author contributions**

897 AH, LM undertook the original draft preparation. All authors contributed to review & editing,
 898 conceptualisation, methodology, and investigation.

899 **7 Competing interests**

900 The authors declare that they have no conflict of interest.

901 **8 Acknowledgements**

902 This research was carried out under the auspices of the Bioregional Assessment Programme, a
903 collaboration between the [Australian](#) Department of Environment and Energy, CSIRO, Geoscience
904 Australia and the Bureau of Meteorology.