

1 A generalised ecohydrological landscape classification for 2 assessing ecosystem risk in Australia due to an altering water 3 regime

4 Alexander Herr¹, Linda E. Merrin¹, Patrick J. Mitchell¹, Anthony P. O'Grady¹, Kate L. Holland¹,
5 Richard E. Mount², David A. Post¹, Chris R. Pavey¹, Ashley D. Sparrow^{1,3}

6 ¹ Commonwealth Scientific and Industrial Research Organisation, Australia

7 ² formerly at: Bureau of Meteorology, Australia

8 ³ Arthur Rylah Institute, Melbourne, Australia

9 *Correspondence to:* Dr Alexander Herr (alexander.herr@csiro.au, [email:-](#))

10 Abstract

11 Describing and classifying a landscape for environmental impact and risk assessment purposes is a
12 non-trivial challenge because this requires region specific landscape classifications that cater for
13 region specific impacts. Assessing impacts on ecosystems from the extraction of water resources
14 across large regions requires [linkinga causal link between](#) landscape features ~~to~~and their water
15 requirements. We present the rationale and implementation of an ecohydrological classification for
16 regions where coal mine and coal seam gas developments may impact on water. Our classification
17 provides the essential framework for modelling the potential impact of hydrological changes from
18 future coal resource developments at the landscape level.

19 We develop an attribute-based system that provides representations of the ecohydrological entities
20 and their connection to landscape features and make use of existing broad-level, classification
21 schemes into an attribute-based system. We incorporate a rule-set with prioritisation, which
22 underpin risk modelling and make the scheme resource efficient, where spatial landscape or
23 ecosystem classification schemes, developed for other purposes, already exist.

24 A consistent rule-set and conceptualised landscape processes and functions allow combining diverse
25 data with existing classification schemes. This makes the classification transparent, repeatable, and
26 adjustable, should new data become available. We apply the approach in three geographically
27 different regions, with widely disparate information sources, for the classification, and provide a
28 detailed example of its application. We propose that it is widely applicable around the world for
29 linking ecohydrology to environmental impacts.

30 **Keywords:** Typology, ecology, hydrology, causal pathway, bioregional assessments, environmental
31 impact, risk analysis

32 1 Introduction

33 The categorisation of the Earth's surface into geo-ecological landscape classes provides a way to
34 simplify the complexity of the form and function of the landscape and provides vital contextual
35 information to support land and water management, and policy initiatives. This includes identifying
36 geographical regions within which landscape-scale attributes, such as climate, topography, geology,
37 and land cover, that are homogeneous and distinctive compared to other regions. It involves

38 identifying broad-scale, general patterns, processes, and functions. Landscape class units are
39 'ecologically equivalent', having the same dominant processes that sustain a similar suite of species,
40 and are likely to respond in similar ways to management initiatives or environmental changes. This
41 ecological equivalence enables the selection of assessment locations for monitoring, measurement
42 or experimentation, and it enables the extrapolation of results to all areas within the same ecological
43 class (Hawkins and Norris, 2000; MacMillan et al., 2003; Cullum et al., 2016a; Cullum et al., 2016b).

44 Such a landscape classification also explains variation in ecological characteristics (e.g. assemblage
45 structure) and is predictive of the ecological attributes of those areas. This predictive quality is useful
46 for defining ecological criteria, identifying reference and degraded sites, defining conservation goals,
47 including the assessment of biodiversity, and the setting of restoration objectives (Hawkins et al.,
48 2000; McMahon et al., 2001; Snelder et al., 2004).

49 In summary, landscape classification is a way of dividing a landscape into components where the
50 characteristics within the components are more similar than the characteristics between the
51 components. That is, the components have their own distinct features that separate them from the
52 other components.

53 However, describing and classifying a landscape for environmental impact and risk assessment
54 purposes is a non-trivial challenge, where hydrological records are limited (see e.g. Wolfe et al.,
55 2019). This is the case for many regions in Australia, where low population densities, high
56 urbanisation and limits in (water) resource management information exist. For our purpose, which
57 was the assessment of risk to ecosystems within the regions of the Bioregional Assessments
58 Programme (Bioregional Assessments, 2018), we needed a landscape classification that reflected the
59 hydrological connectivity of surface and groundwater with ecosystems in the landscape. The
60 Bioregional Assessment Programme, an Australian regional scale impact assessment, investigated the
61 impacts and risks of coal seam gas (CSG) and large coal mining developments on water resources and
62 water-dependent assets via a water pathway (Bioregional Assessments, 2018). This investigation
63 focussed on the landscape level, that is on areas within the regions where the landscape is made up
64 of different interacting land-uses and ecosystems.

65 In our case, the broad scale assessments of impacts from resource developments on ecosystems
66 required an understanding of landscape composition and structure, and how these relate to the
67 ecosystems embedded in the landscape. The type and composition of the landscape components are
68 dependent on the focus of the assessment and therefore require careful consideration of the
69 questions the assessment seeks to answer (Wiens and Milne, 1989; Eigenbrot, 2016). For Australia,
70 there are several landscape level classifications available (see e.g. Thackway and Cresswell, 1995;
71 Pain et al., 2011; Hall et al., 2015; NVIS Technological Working Group, 2017; Gharari et al., 2011).
72 Unfortunately, these available classifications are not directly applicable for our assessment regions
73 because there is no alignment between the regions and existing classification boundaries, or the
74 classifications, even if they include ecohydrological elements, are limited to their locations or domain
75 of interest.

76 Identifying the water dependency of landscape components is a prerequisite when analysing the
77 potential impacts of proposed coal and gas resource developments on water resources at a regional
78 scale. For example, coal resource developments generally need to manage both groundwater and
79 surface water as part of their operations. With multiple developments within the one region, impacts
80 are likely to go beyond the local scale and affect ecosystems at the landscape level (see for example

81 Bioregional Assessments, 2018, 2019). In this context, there is a need for an ecological classification
82 of the landscape that identifies and causally connects the water dependency of its components to
83 activities of resource extraction, in a spatially explicit manner. Further, there is a need to identify
84 impact pathways between resource extraction sites and the ecosystems that show causal
85 connectivity between extraction activities and ecosystem impacts.

86 Land classification systems reveal patterns and underlying drivers of ecosystem structure and
87 function, or produce a tractable unit of assessment for evaluating environmental change (Hobbs and
88 McIntyre, 2005; Poff et al., 2010). Many different classification approaches and methodologies
89 currently exist to represent ecosystems in a landscape. This includes the interim bioregional
90 classification for Australia (IBRA), which provides the basis for defining and managing the national
91 reserve system and; the national vegetation information systems (NVIS), that describes the extent
92 and distribution of vegetation ecosystems for the Australian continent (Thackway and Cresswell,
93 1995; Department of Agriculture, 2021). Classifications addressing hydrology in Australia incorporate
94 a framework for river management that delineate boundaries between homogenous landscape
95 components, based on either their dependency on surface water or groundwater regimes (Poff et al.,
96 2010; Aquatic Ecosystems Task Group, 2012; Olden et al., 2012). However, none of these
97 classifications describe ecohydrological connections between waters and the wider landscape. For
98 example, IBRA and NVIS are based purely on vegetation classifications and so do not contain any
99 hydrological details, while the available hydrological classifications focus purely on the streams and
100 waterbodies within the landscape, as their focus is on aquatic organisms and environmental flows.
101 While both these elements are part of the immediate landscape surrounding water bodies, they do
102 not in themselves provide conceptual and direct linkages between changes in water and ecosystem
103 responses in the wider landscape. Therefore, a standardised approach to formulating classifications
104 that combine these two aspects, ecosystems and their water sources, is lacking.

105 This conundrum exists because different analysis contexts require classifications for different
106 purposes, ranging from conservation planning, habitat mapping, resource assessment and vegetation
107 modelling, and because there is contention between the generality of broad classifications and their
108 applicability at the local scale (Leathwick et al., 2003; Abella et al., 2003; Poulter et al., 2011; Cullum
109 et al., 2016b; Pyne et al., 2017). Hence, we needed a new classification system, when evaluating
110 water dependency in the context of regional scale for multiple coal and coal seam gas resource
111 developments. This new system must incorporate surface water and groundwater regimes into a
112 spatial demarcation of ecosystem boundaries in the landscape. Including surface water and
113 groundwater regimes will provide ~~the establishing of~~ conceptual connection between impacts from
114 developments on surface water and groundwater within the classification, ~~and~~. The classification
115 must also be spatially explicit, to enable a landscape wide analysis of those impacts, so that ~~one can~~
116 link changes in water at one part of the landscape can be linked to ecological responses at another
117 part of the landscape.

118 With this context in mind, the objectives for this paper are to:


- 119 1. characterise a regional level landscape based on patterns in land use, ecology, geomorphology
120 and hydrology,
- 121 2. develop landscape classes of water-dependent, remnant and human-modified features, and
- 122 3. ensure landscape classes sit within a common framework that aids in formulating
123 conceptual models and patterns of water dependency across the landscape.


124 Here, we present the rationale, formulation and implementation of an ecohydrological landscape
125 classification. Based on a generalised conceptual model of the typical hydrological connectivity
126 within landscape features in a region, the classification integrates pre-existing, broad-level
127 classification schemes into an attribute-based schema applied at the regional scale. It places the
128 landscape classification within a common framework (i.e. a framework that is common to all
129 landscape elements in the region) that aids in formulating conceptual models and patterns in water
130 dependency across the landscape. This makes our approach generally applicable for assessments
131 aimed at regional hydrological impacts on, and risks to, ecosystems. Importantly, the classification
132 also provides the ability to develop a conceptual understanding of, and causally connect, hydrological
133 changes at the landscape level, with impacts on ecological entities within the landscape. These
134 causal pathways are the basis for spatially identifying the impacted areas, and for developing an
135 appropriate mitigation response, including for extractive resource developments and water
136 extraction.

137 We have applied this approach to several regions across eastern Australia with coal and CSG resource
138 developments. Here we will focus on its application in three regions; Namoi, Maranoa–Balonne–
139 Condamine and Galilee, and subsequently discuss why the approach is transferable to other regional
140 developments that may carry a hydrological based risk to ecosystems, even those in a different
141 contextual setting with regards to data sources and existing landscape classifications.

142 The remainder of the paper is structured as follows: ~~in the Methods section we described~~ describes
143 the general approach for achieving the classification, including descriptive examples of existing data
144 sources. It also provides a description of the three study regions in which we applied and tested the
145 classification. The Results section provides evidence of the general applicability of our approach in
146 that it shows the detailed ecological landscape classification for the three distinctively different
147 region in terms of location, topography, and climate. In the Discussion ~~section~~ we provide an
148 example on the use of the landscape classification. Here we describe an impact assessment in the
149 Namoi region using modelling that includes ~~a Bayesian-expert assessment approach. We also~~
150 ~~discuss~~ assessments. ~~In the last section we provide a discussion of the landscape classification,~~
151 ~~including~~ limitations, and provide our conclusions.

152 Figure 1 provides a visual outline of the paper, ~~giving an overview of the~~ and workflow we applied.
153 ~~In this context the figure~~ It incorporates Methods ~~and~~, Results ~~above the dashed line. Below the~~
154 ~~dashed line are the~~ and Discussion (unshaded parts, ~~which include applying~~), and indicates where we
155 applied our classification using quantitative and qualitative risk modelling in combination with
156 surface water and groundwater modelling. (shaded parts; Section 4). Surface water and groundwater
157 modelling establish a zone of hydrological change in which impacts are likely. The red, more lightly
158 shaded circle shows the resulting risk assessment outcomes, where the landscape classification
159 provided the crucial details for experts to assign risks to landscape elements and classes.





161

162 **Figure 1: Visualisation of workflow for developing our ecological landscape classification (above the dashed line) and its**
 163 **application to develop an ecological risk assessment (below the dashed line), non-patterned, identifies focus of this**
 164 **paper) and its application in an ecological risk assessment, which we briefly summarise to show the classification's**
 165 **applicability (inside patterned rectangle, described in Section 4). The outcome of combining the landscape classification**
 166 **with hydrological modelling and risk modelling is the map of risk (identified in the lightly patterned red circle).**
 167 **Hydrological features are descriptors that have a hydrology component in their character. Ecohydrological elements are**
 168 **unique identifiable building blocks of the landscape that contain similar (hydrological) features.**

169 **2 Methods**

170 In the following section, we show the development of a dataset-agnostic method to develop a
 171 regional-level landscape classification that is flexible in incorporating data sources at different scales,
 172 including region-specific datasets. Ecological systems are complex and work at a range of scales

173 within regions/landscapes, and they exhibit interactions and feedbacks that work across scales.
174 Consequently, there is no one scale appropriate for a subsequent analysis of ecological impacts. Here
175 we use a variable scale range that is relevant for ecological impacts of water changes from coal
176 resource developments when using an expert assessment approach. Our classification focuses on a
177 scale range (36,000 km² to 600,000 km²) that is associated with eco-hydrological linkages (and
178 associated causality) between the response of ecological components to predicted hydrological
179 changes. This scale range is what most hydrologists would consider the “regional” scale range
180 (Gleeson and Paszkowski, 2014). It provides the basis and flexibility for experts to build their
181 conceptual understanding of causal pathways and use these to assess ecological impacts with the
182 landscape classes (see also Figure 1).

183 **2.1 Study areas**

184 Our three study areas are the Namoi, the Maranoa–Balonne–Condamine and the Galilee regions in
185 eastern Australia. Each of these regions have coal resource developments within them and have
186 distinctly different landscape characteristics. They cover different state jurisdictions, or even cross
187 state jurisdictions, and range from approximately 36,000 km² to 600,000 km² in size. Consequently,
188 the classification is based on different state-based datasets. Each region’s classification relies on the
189 extent of surface water and groundwater systems that existing and potential future coal resource
190 developments in the region may impact.

191 **2.1.1 Namoi region**

192 The Namoi region covers approximately 35,700 km² in eastern Australia, is located within New South
193 Wales and forms one catchment of the Murray-Darling Basin. The long-term mean annual rainfall
194 varies from 600 to 1100 mm and potential evapotranspiration (PET) varies from 1200 to 1400 mm. It
195 contains six operational coal mines (one underground mine and five open-cut mines), nine potential
196 future coal mines and one potential CSG development. The nine potential future coal mines consist
197 of two underground, one combined open cut and underground, and seven open cut mines. The
198 region covers most of the Namoi River catchment, with the Namoi River being the main river within
199 the region. It also contains two major aquifer systems – the Namoi Alluvial aquifer and the Pilliga
200 Sandstone aquifer (Figure 2a).

201 The main land use within the region is agriculture, both dryland and irrigated cropping, and livestock
202 grazing, as well as forestry. There is also a diverse range of landscapes and ecosystems within the
203 region, including the Liverpool and Kaputar ranges, the Liverpool Plains floodplains, and Darling
204 Riverine plains in the west of the region, open box woodlands on the slopes, and temperate and
205 sub-alpine forests in the east of the region. A range of aquatic habitats occur downstream of
206 Narrabri, with large areas of anabranches and billabong wetlands. The Pilliga Nature Reserve in the
207 upper catchment of Bohena Creek, together with The Pilliga State Forest, form the largest remaining
208 area of dry sclerophyll forest west of the Great Diving Range in New South Wales (Welsh et al., 2014).

209 **2.1.2 Galilee region**

210 The Galilee region covers approximately 612,300 km² and is located mostly within Queensland,
211 Australia. PET far exceeds rainfall, particularly in the summer months. Yearly rainfall ranges from 300
212 to 700 mm and PET from 2200 to 2900 mm. There are 17 proposed coal resource developments in
213 the Galilee region. These include three open-cut coal mines, two underground coal mines, five

214 combined open-cut and underground coal mines, four coal mines of currently unknown type, and
215 three CSG projects ([Figure 3b](#)).

216 The Galilee region includes the headwaters of seven major drainage catchments. These catchments
217 are Bulloo, Burdekin, Cooper Creek, Diamantina, Flinders, Paroo and Warrego. The largest of these
218 catchments within the region are the Cooper Creek and Diamantina. Groundwater within the region
219 is a very important resource, as most of the streams are ephemeral. Groundwater is used for town
220 water, agriculture and industry. Most groundwater in the region is extracted from the Great Artesian
221 Basin ([b](#)).

222 The region covers a range of environments, including mountains of the Great Dividing Range in the
223 east, through to semi-arid and arid areas in the central and western parts of the region. The main
224 land use in the region is livestock grazing on native vegetation. There is no intensive agriculture in the
225 region, and a low human population density, largely due to the low and unpredictable rainfall (Evans
226 et al., 2014).


227 **2.1.3 Maranoa–Balonne–Condamine region**

228 The Maranoa–Balonne–Condamine region covers approximately 130,000 km² and is located mostly
229 within south-east Queensland with about half the area within the Murray-Darling Basin. From east to
230 west, average annual rainfall decreases from 800 mm to 420 mm, as PET increases from 1500 to
231 2370 mm. The region overlies the Surat Basin and has five open-cut coal mines and five CSG projects,
232 as well as two proposed open-cut coal mines ([Figure 4c](#)).


233 The region contains the headwaters of the Condamine-Balonne, Moonie, Weir, Maranoa and Dawson
234 rivers. Most of the rivers within the region are ephemeral. Groundwater is therefore an important
235 water source and is used for stock and domestic purposes, and in some cases, town water supply.
236 The Great Artesian Basin is the main source of groundwater used within the region (Welsh et al.,
237 2015).

238 The main land use within the region is grazing on natural vegetation, with dryland cropping and
239 production forestry also major land uses. The main vegetation type within the region is grassy
240 woodlands, with river red gums, coolabah and river oak common riparian species. There are also six
241 wetlands of national significance within the region: Balonne River Floodplain, Boggomoss Springs,
242 Dalrymple and Blackfellow Creeks, Lake Broadwater, Palm Tree and Robinson Creeks, and The Gums
243 Lagoon (Welsh et al., 2015).


244



245 (a)



246 (b)




247


248

249

Figure 2. Study areas for (a) the Namoi region (b) the Galilee region and (c) the Maranoa-Balonne-Condamine region study area, showing the potential coal resource development sites



250




251

Figure 3

(e)

252 [\(cont.\) Study areas for \(c\) the Maranoa–Balonne–Condamine region. The Galilee region study area](#), showing the
253 potential coal resource development sites



255 **Figure 4. The Maranoa–Balonne–Condamine region study area, showing the potential coal resource development sites**

256 2.2 Landscape classification development – overview and rationale

257 The purpose of this ecohydrological landscape classification is to characterise the landscape based on
258 patterns in land use, ecology, geomorphology and hydrology, and from these, develop landscape
259 classes of water-dependent, remnant and human-modified features. We chose these features
260 because these three types represent a generally applicable delineation used in our spatial dataset.
261 For example, in Australia the word remnant vegetation (our remnant features) describes all
262 vegetation where there was no clearing or [regrowth1regrowth](#) of (semi-) native vegetation, resulting
263 in a vegetation community that resembles its predecessor’s structure. It represents areas with low to
264 very minimal human interference. This is opposed to human-modified, where human activities are
265 the defining features of the area, such as urban areas or other infrastructure. Water dependency is
266 essential for establishing a conceptual linkage of water across landscape elements. Our classification
267 employs a geographical information system to overlay existing spatial data for each region. The
268 spatial data are the basis for categorising the landscape features using a rule-set to prioritise the
269 spatial data based on their attribute features.

270 The datasets have a regional, state or national coverage. Using a feature-based classification helps to
271 place the landscape classes within a common biophysical system that aids in formulating
272 conceptual models and patterns in water dependency across the landscape of each region. This
273 provides a classification that is aligned with the idiosyncrasies of each region. Maintaining regionality

274 is essential when developing conceptual models and quantitative models for assessing the risk to
275 ecological components from hydrological changes. For example, arid and semi-arid regions have very
276 different ecological environments, functions and processes than subtropical or temperate
277 woodlands.

278 Our approach uses a defined rule-set and priorities, which we apply to regionally available datasets
279 to achieve a landscape classification for each of our regions. Tables 1 to 3 provide a list of citations
280 for example datasets used in this process. This is different to most other landscape classifications
281 that may use climate, topography, hydrological assessment units and remote sensing data, and apply
282 statistical dimensionality reduction and classifications such as proximity analysis (see e.g. Gharari et
283 al., 2011; Leibowitz et al., 2014; Sawicz et al., 2014; Zhang et al., 2016; Addicott et al., 2021; Carlier
284 et al., 2021; Jones et al., 2021).

285 When considering the characteristics of our regions, the following features form part of the rule-set
286 for defining landscape classes:

- 287 • broad habitat/land use type (remnant/human-modified).
- 288 • wetland (wetland/non-wetland)
- 289 • topography (upland/lowland, floodplain/non-floodplain)
- 290 • groundwater (groundwater dependent/non-groundwater dependent, Great Artesian Basin
291 (GAB)/non-GAB)
292 Note: identifies groundwater dependency and classifies this with the presence/absence of
293 Great Artesian basin groundwaters.
- 294 • vegetation type (riparian/woodland floodplain/grassy woodland/rainforest)
- 295 • water regime (permanent/ephemeral/null) of surface water

296 These features identify groups of landforms and use, streams and springs.

297

298 The hydrological connectivity is the main reason for developing a new classification as this allows us
299 to assess the potential impact of coal resource developments on the landscape via a water pathway.
300 Therefore, the most important characteristics are the hydrological features. Describing the
301 conceptual understanding of how water connects the landscape elements allows us to identify
302 where in the landscape impacts are likely to occur. [Therefore, In line with this](#) we developed a
303 hierarchical approach, where hydrological features have priority over other landscape characteristics.
304 This resulted in a spatially complete landscape classification, where there are no gaps in the mapping
305 data. The method of prioritisation depended on region-specific characteristics and the data
306 availability. This [resulted in yielded](#) a classification where the landscape classes have their origin in
307 the spatial datasets, and included the water dependency, which was a pre-requisite of the
308 prioritisation. An example prioritisation assigned in order of highest to lowest is:

- 309 • aquatic ecosystems (e.g. wetlands, streams and lakes)
- 310 • remnant vegetation
- 311 • other landscape components that are ‘non-remnant vegetation’ and are typically ‘human-
- 312 modified’.

313

314 Subsequent use of the landscape classification for risk identification with expert input also required
315 combining landscape classes into broader landscape groups. Landscape groups are sets of landscape
316 classes that share [hydrological](#)~~eco~~[hydrological](#) properties. These landscape groups provided
317 efficiencies in the expert elicitation process of the risk modelling, as they combined similar ecological
318 system components based on our landscape classes while also accounting for region specific
319 differences. For example, in the Namoi region there are two landscape groups where we do not
320 expect any impact from coal resource developments. Firstly, the ‘Dryland remnant vegetation’
321 landscape group is ruled out from potential impacts because it comprises vegetation communities
322 that are reliant on incident rainfall and local runoff and do not include features in the landscape that
323 have potential hydrological connectivity to surface water or groundwater features. Secondly, the
324 ‘Human-modified’ landscape group is excluded from the ecological impact assessment because it
325 primarily comprises agricultural and urban landscapes that are highly modified by human activity.
326 Here the impact assessment focus is on economic assets such as groundwater bores, and therefore
327 beyond the scope of this publication.

328 **2.2.1 Landform classification**

329 Landform classification relied on the dominant land type of either habitat or land use to determine
330 landscapes that are relatively natural and those that have been ‘human-modified’. Relatively intact
331 areas are more likely to contain ecological assets, such as species and ecological communities, than
332 highly modified areas. Location within the region (topography–upland/lowland, floodplain/non–
333 floodplain), groundwater dependency and water regime, were part of classifying the landscape.
334 Determining areas that are subjected to flooding, or that have persistent water, assists in identifying
335 landscapes that support water-dependent habitat and vegetation, and aquatic ecosystems (Table 1).

336 **2.2.2 Stream classification**

337 Stream classification in each of the study regions was based on stream position within the catchment
338 (e.g. upland/lowland), water regime (perennial/near permanent or ephemeral/temporary) and
339 dependence and source of groundwater (Table 2). Catchment position is a potential indicator of
340 stream morphology and flow patterns, while water regime is important when considering habitat
341 suitability and physical processes within the channel and riparian zone. Streams can also gain and
342 lose water to local and regional groundwater systems, interacting with groundwater-dependent
343 ecosystems (Table 2).

344 **2.2.3 Spring classification**

345 The water source is the basis of spring classification. The source of groundwater is important when
346 considering regional scale landscape classifications, due to the hydrological connectivity of aquifers
347 and potential coal resource developments (Table 3).

348 **Table 1. Landform classification criteria and example datasets**

Characteristic	Classification	Example datasets
Habitat/land use	<ul style="list-style-type: none"> • Non-remnant • Remnant (and stream, Wetland) 	<p>Australian land use mapping (Australian Bureau of Agricultural and Resource Economics and Sciences, 2014) (national)</p> <p>NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)</p>
Topography	<ul style="list-style-type: none"> • Floodplain • Non-floodplain 	<p>NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)</p> <p>Namoi Valley Flood Plain Atlas 1979 (NSW Office of Environment and Heritage, 1979) (regional)</p> <p>Murray-Darling Basin aquatic ecosystem classification (Department of Sustainability, Environment, Water, Population and Communities, 2014) (regional)</p> <p>GEODATA TOPO 250K Series 3 (Geoscience Australia, 2006) (national)</p>
Groundwater	<ul style="list-style-type: none"> • Groundwater dependent (source) • Non-groundwater dependent 	<p>Namoi groundwater dependent ecosystems (NSW Office of Water, 2015) (regional)</p> <p>Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer (Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts, 2015) (state)</p> <p>Queensland Groundwater Dependent Ecosystems (Queensland Department of Science, Information Technology, Innovation and the Arts, 2013) (state)</p>
Water regime	<ul style="list-style-type: none"> • Temporary • Near-permanent • Fresh • Saline 	<p>Queensland wetland data version 3 - wetland areas (Queensland Department of Science, Information Technology, Innovation and the Arts, 2012) (state)</p> <p>Queensland wetland data version 3 - wetland areas (Queensland Department of Science, Information Technology, Innovation and the Arts, 2012) (state)</p> <p>Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer (Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts, 2015) (state)</p> <p>South Australian Wetlands – Groundwater Dependent Ecosystems (GDE) Classification (Sa Department for Water, 2010) (state)</p>
Vegetation	<ul style="list-style-type: none"> • Broad vegetation type 	<p>NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)</p>

349 NSW = New South Wales

350 **Table 2. Stream classification criteria and example datasets**

Characteristic	Classification	Example datasets
Topography	• Upland	Murray-Darling Basin aquatic ecosystem classification (Department of Sustainability, Environment, Water, Population and Communities, 2014) (regional)
	• Lowland	NSW regional vegetation (NSW Office of Environment and Heritage, 2015) (regional)
		MrVBF (CSIRO, 2000) (national)
Groundwater	• Groundwater dependent (source)	Asset database for the Namoi subregion (Bioregional Assessment Programme, 2016) (regional)
	• Non-groundwater dependent	Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer (Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts, 2015) (state) Queensland groundwater dependent ecosystems (Queensland Department of Science, Information Technology, Innovation and the Arts, 2013) (state)
Water regime	• Perennial	Murray-Darling Basin aquatic ecosystem classification (Department of Sustainability, Environment, Water, Population and Communities, 2014) (regional)
	• Ephemeral	Geofabric Surface Cartography (Bureau of Meteorology, 2012) (national)

351

352 **Table 3. Spring classification criteria and example datasets**

Characteristic	Classification	Example datasets
Groundwater	• Groundwater dependent (source)	Asset database for the Namoi subregion (Bioregional Assessment Programme, 2016) (regional)
	• Non-groundwater dependent	Asset database for the Maranoa–Balonne–Condamine subregion (Bioregional Assessment Programme, 2015) (regional)
		Spring vents assessed for the Surat Underground Water Impact Report 2012 (Office of Groundwater Impact Assessment, 2015) (regional)

353

354 **3 Results**

355 Below we present the resulting landscape classes for the three regions. For each region, we also
 356 combined the landscape classes into landscape groups, which were specific to each region and were
 357 based on distinctions in topography, water dependency and association with GAB or non-GAB GDEs,
 358 floodplain/non-floodplain or upland/lowland environments and remnant/human-modified habitat
 359 types. The purpose of the landscape groups was to combine non-water dependent landscape classes
 360 and relate water dependent landscape classes to region specific aspects of their water dependency.
 361 This enabled experts to develop a conceptualisation of the landscape for developing their ecological
 362 impact models. While the approach in defining the landscape classes is based on a consistent rule-
 363 set and prioritisation, each of the regions has different landscape classes, which is a consequence of
 364 the differences in location, jurisdictions and available spatial datasets.


365 The rule-set deriving from the landform classification (Tables 1 to 3) and prioritisation of hydrological
366 features is the main outcome of our approach and we present the rule-set as a decision pathway
367 visually below (Figure 5). For example, for the Namoi region, the rule-set includes: (1) identify the
368 habitat (e.g. stream) (2) select by topography (e.g. upland), (3) identify the groundwater associations
369 (e.g. GDE), and so on until one derives at the final [landclasslandscape class](#) level (see Figure 5).

370 **3.1 Landscape classes in the Namoi region**

371 There were 29 landscape classes within six landscape groups in the Namoi region (Figure 5). Of these
372 landscape groups, 'human-modified' (non-remnant) was the largest (59.3%; Table 4), and included
373 urban, agriculture, plantations and other intensive land uses. The dryland remnant vegetation was
374 the second largest landscape group and consisted of the grassy woodland landscape class (24.2%;
375 Table 4). This landscape class was considered non-water dependent as it did not intersect with
376 floodplain, wetland or GDE features. The rainforest landscape group was the smallest (0.5%; Table 4),
377 with only a limited distribution (Figure 6a).

378 The stream network consisted of two landscape groups (floodplain or lowland riverine and non-
379 floodplain or upland riverine). The non-floodplain or upland riverine landscape group had a larger
380 proportion of stream network length (63.8%) compared to the floodplain or lowland riverine
381 landscape group (36.2%; Figure 6b). There were 22 springs identified within the Namoi region, with
382 seven of these associated with the GAB (Figure 6b).

383



384
385

Figure 5. Overview of the Namoi landscape classification schema, including criteria and attributes resulting in six landscape groups


386 **Table 4. Percentage of area of each landscape group for the Namoi region**

Landscape group	Percentage of region (%)	Number of classes	Landscape classification attributes
Human-modified	59.3%	6	Conservation of natural environments, production from relatively natural environments, production from dryland agriculture and plantations, production from irrigated agriculture and plantations, intensive uses or waters for production/consumption
Dryland remnant vegetation	24.2%	1	Non-floodplain, non-GDE, grassy woodland remnant vegetation
Non-floodplain or upland riverine	9.8%	8	Upland or non-floodplain streams, wetlands or remnant vegetation
Floodplain or lowland riverine	6.2%	10	Lowland or floodplain streams, wetlands or remnant vegetation
Rainforest	0.5%	2	Non-floodplain GDE or non-GDE rainforest remnant vegetation
Springs	<0.1%	2	GAB or non-GAB springs


387

388

(a)



(b)



389
390
391

Figure 6. (a) Landscape groups (excluding the 'Human-modified'), and (b) Stream network classified as 'upland' or 'lowland' in the landscape classification and 'Springs' within the Namoi region. GAB = Great Artesian Basin
Data: Bureau of Meteorology (2012); Bioregional Assessment Programme (2017)

392 **3.2 Landscape classes in the Galilee region**

393 The Galilee region has 31 landscape classes organised into 11 landscape groups (Figure 7). The
394 dryland landscape group was the largest group within the region and the only group to have no
395 water dependency (68.5%; Table 5). The landscape groups that covered the floodplain areas were
396 the next most dominant classes, with floodplain, terrestrial GDE (12.94%; Table 5) and floodplain,
397 non-wetland (11.8%; Table 5). The remaining three non-floodplain landscape groups consisted of
398 disconnected wetlands, and terrestrial and wetland GDEs (4.9% combined; Table 5).

399 The stream network was classified as groundwater dependent or non-groundwater dependent. Most
400 of the streams in the region were non-GDEs (87.7% compared to 12.3% for the streams, GDE
401 landscape group). There were also over 3000 springs in the region.

402

403

Landscape classification						
Topography	Wetland	Groundwater	Water regime	Vegetation	Landscape class	Landscape group
<u>Vegetation</u>				Remnant vegetation	Wetland GDE, remnant vegetation	Floodplain, wetland GDE
	Wetland	GDE	<	Non-remnant vegetation	Wetland GDE	
				Remnant vegetation	Floodplain disconnected saline wetland, remnant vegetation	
	Non-wetland	Disconnected	< Saline	Non-remnant vegetation	Floodplain disconnected saline wetland	Floodplain, disconnected wetland
Floodplain				Remnant vegetation	Floodplain disconnected wetland, remnant vegetation	
				Non-remnant vegetation	Floodplain disconnected wetland	
		GDE	<	Remnant vegetation	Terrestrial GDE, remnant vegetation	Floodplain, terrestrial GDE
		Disconnected	< Non-saline	Non-remnant vegetation	Terrestrial GDE	
				Remnant vegetation	Floodplain disconnected non-wetland, remnant vegetation	Floodplain, non-wetland
				Non-remnant vegetation	Floodplain disconnected non-wetland	
		GDE	<	Remnant vegetation	Non-floodplain wetland GDE, remnant vegetation	Non-floodplain, wetland GDE
	Wetland	Disconnected	< Saline	Non-remnant vegetation	Non-floodplain wetland GDE	
				Remnant vegetation	Non-floodplain disconnected saline wetland, remnant vegetation	
	Non-wetland	Disconnected	< Non-saline	Non-remnant vegetation	Non-floodplain disconnected saline wetland	Non-floodplain, disconnected wetland
Non-floodplain				Remnant vegetation	Non-floodplain disconnected wetland, remnant vegetation	
		GDE	<	Non-remnant vegetation	Non-floodplain disconnected wetland	
		Disconnected	< Non-saline	Remnant vegetation	Non-floodplain terrestrial GDE, remnant vegetation	Non-floodplain, terrestrial GDE
				Non-remnant vegetation	Non-floodplain terrestrial GDE	Dryland
<u>Stream network</u>				Remnant vegetation	Dryland, remnant vegetation	
		Disconnected	< Non-saline	Non-remnant vegetation	Dryland	
Upland		GDE	< Temporary	Temporary upland GDE stream	Streams, GDE
		Disconnected	< Near-permanent	Near-permanent upland GDE stream	Streams, non-GDE
				Temporary upland stream	
				Near-permanent upland stream	
Lowland		GDE	< Temporary	Temporary lowland GDE stream	Streams, GDE
		Disconnected	< Near-permanent	Near-permanent lowland GDE stream	
				Temporary lowland stream	
		Disconnected	< Near-permanent	Near-permanent lowland stream	Streams, non-GDE
Estuarine				Temporary estuarine stream	
<u>Springs</u>				Near-permanent estuarine stream	
		GDE	Springs

Figure 7. Landscape classification of the Galilee region

GAL-340-001

404

405 **Table 5. Percentage of area of each landscape group for the Galilee region**

Landscape group	Percentage of region (%)	Number of classes	Landscape classification attributes
Dryland	68.5%	2	Non-floodplain, non-wetland, disconnected and saline remnant or non-remnant vegetation
Floodplain, terrestrial GDE	12.9%	2	Floodplain, non-wetland and GDE remnant or non-remnant vegetation
Floodplain, non-wetland	11.8%	2	Floodplain, non-wetland, disconnected and non-saline remnant or non-remnant vegetation
Floodplain, disconnected wetland	1.1%	4	Floodplain, wetland, disconnected saline and non-saline remnant or non-remnant vegetation
Floodplain, wetland GDE	0.8%	2	Floodplain, wetland and GDE remnant or non-remnant vegetation
Non-floodplain, terrestrial GDE	3.4%	2	Non-floodplain, non-wetland and GDE remnant or non-remnant vegetation
Non-floodplain disconnected wetland	1.4%	4	Non-floodplain, wetland, disconnected, saline and non-saline remnant or non-remnant vegetation
Non-floodplain, wetland GDE	<0.1%	2	Non-floodplain, wetland and GDE remnant or non-remnant vegetation
Springs	<0.1%	1	GDE springs

406

407 **3.3 Landscape classes in the Maranoa–Balonne–Condamine region**

408 The landscape classification for the Maranoa–Balonne–Condamine resulted in 34 landscape classes
 409 within five landscape groups (Figure 8). The largest landscape group was the human-modified group
 410 (72.2%, Table 6), which included agricultural production, plantations and other intensive land uses.
 411 Of the remaining landscape groups, dryland remnant vegetation was the second most dominant
 412 (19.8%, Table 6). It was not considered water dependent, because it did not intersect with floodplain,
 413 wetland or GDE features.

414 There were three landscape groups that cover the stream network. The most dominant landscape
 415 group was floodplain or lowland riverine (including non-GAB GDEs) (47.8%), followed by non-
 416 floodplain or upland riverine (including non-GAB GDEs) (39.4%) and GAB GDEs (riverine, springs,
 417 floodplain or non-floodplain) (12.7%). There were 177 springs identified within the region. Most of
 418 the springs were GAB GDEs (riverine, springs, floodplain or non-floodplain) (86.4%, compared to
 419 13.6% for non-floodplain or upland riverine (including non-GAB GDEs)).

420
421
422
423

Landscape classification					
Topography	Groundwater	Wetland	Water regime	Landscape class	Landscape group
<u>Remnant vegetation</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary	Floodplain GAB GDE, near-permanent wetland Floodplain GAB GDE, temporary wetland Floodplain GAB GDE	GAB GDEs (riverine, springs, floodplain, non-floodplain)
		Non-wetland	Near-permanent Temporary	Floodplain non-GAB GDE, near-permanent wetland Floodplain non-GAB GDE, temporary wetland	Floodplain or lowland riverine (including non-GAB GDEs)
		Wetland	Near-permanent Temporary	Floodplain non-GAB GDE Floodplain, near-permanent wetland Floodplain, temporary wetland Floodplain remnant vegetation	
<u>Floodplain</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary	Non-floodplain GAB GDE, near-permanent wetland Non-floodplain GAB GDE, temporary wetland Non-floodplain GAB GDE	GAB GDEs (riverine, springs, floodplain, non-floodplain)
		Non-wetland	Near-permanent Temporary	Non-floodplain non-GAB GDE, near-permanent wetland Non-floodplain non-GAB GDE, temporary wetland	Non-floodplain or upland riverine (including non-GAB GDEs)
		Wetland	Near-permanent Temporary	Non-floodplain, near-permanent wetland Non-floodplain, temporary wetland Dryland remnant vegetation	Dryland remnant vegetation
<u>Non-floodplain</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary	Conservation and natural environments Production from relatively natural environments	Human-modified
		Non-wetland	Near-permanent Temporary	Production from irrigated agriculture and plantations Intensive uses Water	
		Wetland	Near-permanent Temporary		
<u>Non-remnant vegetation</u>	GAB GDE Non-GAB GDE Non GDE	Wetland	Near-permanent Temporary		
		Non-wetland	Near-permanent Temporary		
		Wetland	Near-permanent Temporary		
<u>Stream network</u>	GAB GDE Non-GAB GDE Non GDE GAB GDE Non-GAB GDE Non GDE	Wetland	Temporary Temporary Near-permanent Temporary Temporary Near-permanent Temporary	Temporary upland GAB GDE stream Temporary upland non-GAB GDE stream Near-permanent upland stream Temporary upland stream Temporary lowland GAB GDE stream Temporary lowland non-GAB GDE stream Near-permanent lowland stream Temporary lowland stream	GAB GDEs... Non-floodplain or upland riverine (including non-GAB GDEs) GAB GDEs... Floodplain or lowland riverine (including non-GAB GDEs)
		Non-wetland	Temporary		
		Wetland	Temporary		
<u>Springs</u>	GAB GDE Non-GAB GDE	Wetland		GAB springs Non-GAB springs	GAB GDEs... Non-floodplain...
		Non-wetland			

Figure 8. Landscape classification of the Maranoa–Balonne–Condamine region
 GAB = Great Artesian Basin, GDE = groundwater-dependent ecosystem, GAB GDEs... = GAB GDEs (riverine, springs, floodplain, non-floodplain), Non-floodplain... = Non-floodplain or upland riverine (including or non-GAB GDEs)

424 **Table 6. Percentage of area of each landscape group for the Maranoa–Balonne–Condamine region**

Landscape group	Percentage of region (%)	Number of classes	Landscape classification attributes
Human-modified	72.2%	6	Conservation of natural environments, production from relatively natural environments, production from dryland agriculture and plantations, production from irrigated agriculture and plantations, intensive uses or waters for production/consumption
Dryland remnant vegetation	19.8%	1	Non-floodplain, non-GDE, non-wetland remnant vegetation
Floodplain or lowland riverine (including non-GAB GDEs)	4.5%	9	Floodplain or lowland, non-GAB GDE or non-GDE, temporary or near-permanent wetland, non-wetland or stream
Non-floodplain or upland (including non-GAB GDEs)	2.2%	9	Non-floodplain or upland, non-GAB GDE or non-GDE, temporary or near-permanent wetland, non-wetland, stream or spring
GAB GDEs (riverine, springs, floodplain or non-floodplain)	1.3%	9	Floodplain, non-floodplain or upland GAB-GDE, temporary or near-permanent wetland, non-wetland, stream or spring

425 GAB = Great Artesian Basin, GDE = groundwater-dependent ecosystem

426 **4— Discussion**

427 In Australia, there is no consistent national classification that links ecosystems at landscape level with
 428 their underlying hydrological system. While there are many different land classifications that
 429 incorporate hydrological aspects, they do not provide linkages between hydrology and landscape
 430 elements that enable a broad-scale ecological assessment of impacts associated with changes in
 431 water flow and availability, and they are not sufficiently generic for the purpose of assessing
 432 landscape-level water-related impacts on ecosystems in a spatially explicit manner. However, the
 433 bioregional assessment program needed to assess impacts of coal resource developments on
 434 ecological systems via a water pathway. Hence, we needed to develop an ecological landscape
 435 classification that would be applicable to the different assessment regions.

436 While our spatially explicit landscape classification provided experts with the ability to readily
 437 identify cause and effect relationships between landscape elements and landscape hydrology, there
 438 are obvious differences between the landscape classifications in the three regions, reflecting their
 439 geographical differences (see, and). It provides the specificity that is required in a regional impact
 440 assessment, where the boundaries are based on a combination of geology, water resources and
 441 administrative conditions. The regionality also means that there is need for different datasets
 442 describing the landscape features that would not be available from a classification covering the
 443 whole of Australia.

444 Nevertheless, each landscape classification provides a typology with an explicit connection of water
 445 to the landscape class. This connection enables a causal link between hydrological change and
 446 impact to ecosystems represented by landscape classes. The causal linkage is dependent on (i) a
 447 spatially explicit connection between water in the landscape and the landscape classes, (ii)
 448 conceptual understanding how changes in water may result in a reaction of specific ecosystem

449 elements in the landscape class and/or landscape group and (iii) a way of modelling quantitative
450 changes in ecosystem elements related to changes in water. Our ecohydrological classification
451 approach for landscapes provides this spatially explicit connection and has implicit ecohydrological
452 elements that foster the conceptual understanding of the causal linkage. For example, spatially
453 modelling groundwater level drawdown enables a prediction on which landscape elements classified
454 as springs may be experiencing impacts from water extraction and, with additional ecological
455 modelling, by how much and when.

456 Subsequent ecological modelling using expert elicitation of potential impacts drew heavily on our
457 classification, which is based on a consistent rule set and fosters conceptual understanding of
458 landscape processes and functions. It provides an essential framework for experts to understand and
459 conceptualise how modelled future hydrological changes from coal resource developments link to
460 potential ecological changes at the landscape level. It allows the incorporation of different data
461 sources and existing classification schemes. This consistency also makes the classification
462 development transparent, repeatable, and adjustable, should new data become available.

463 **54** Application of the landscape classification based impact to the assessment of ecosystem risk

464 Here we show an application of how our classification approach can be used to assess. It shows the
465 potential impact coal resource developments can have on ecology using the Namoi region as an
466 example, thus, demonstrating the useability of our classification approach.

467 The purpose of developing the landscape classification was to assess the risk of coal resource
468 development on the ecology of a region via a water pathway. Our landscape classification provided
469 the spatial framework on which experts can could base their assessment of risk from coal resource
470 development on the ecology of a region via a water pathway. Details of the predicted changes in
471 surface water and groundwater for the Namoi and Galilee regions are in Post et al. (2020). Here, we
472 demonstrate the assessment of potential ecological impacts using the Namoi region. For full details
473 of the analyses in each of the three regions see Holland et al. (2017); Herr et al. (2018b); and Lewis
474 et al. (2018). This work included expert assessment of ecological risk to ecosystem components
475 based on conceptual models. Hence, The models needed to identify water mitigated linkages
476 between hydrological changes, ecosystem components and the landscape classes. We briefly
477 describe the expert assessment approach in a 3-step process below. For details we direct the reader
478 to the above references and those listed below.

479 The following describes an application of the landscape classification (see also Figure 1), and in doing
480 so we demonstrate that it is a fit-for-purpose in the context of for assessing potential ecological
481 impact resulting from potential predicted surface water and groundwater changes at different
482 locations within the landscape. The 3-step process illustrates the utility of our landscape
483 classification approach for assessing the risk to ecosystems in the landscape classes and groups. The
484 process included experts identifying risk to landscape classes using their knowledge on local
485 ecosystems within the landscape classes. Specifically, the experts used the broad landscape groups
486 and their underlying hydrogeological features to develop initial qualitative models initially that about
487 priority ecosystem components. These then fed into building quantitative models. Here the experts
488 used outputs from surface water and groundwater modelling to determine. This hydrological
489 modelling identified the potential changes in water and, which experts used to reach a consensus on
490 what this impact these changes may mean for have on ecological entities within the landscape classes

491 and/or groups. These [agreed impacts fed into quantitative](#) models ~~assessed~~[that outlined](#) the future
492 hydrological changes and risks to the ecosystems in the landscape groups (see also Figure 1).

493 [Here we use the example of the upland riverine landscape class in the Namoi region to outline](#) the 3-
494 step process ~~included~~:

495 **Step 1:** Develop qualitative models to conceptualise and prioritise ecosystem components of each
496 landscape class and their linkage to hydrological variables.

497 ~~Here we use the example of the upland riverine landscape class.~~ A qualitative model for the upland
498 riverine landscape class agreed with the existing understanding that a reduction in overbank flows
499 and lowering of the water table resulted in a reduction in several ecosystem components, including
500 riparian habitat, amphibians and fish, and an increase in fine particulate matter, dissolved organic
501 matter and cyanobacteria (Holland et al., 2017; Herr et al., 2018b; Hosack et al., 2018). A qualitative
502 model has, at its basis, the conceptual understanding of ecosystem components and the direction of
503 their interactions, that is a positive, negative, or neutral influence of one component on another. This
504 understanding also incorporates feedback loops between the ecosystem components in the form of
505 sign directed graphs, and it enables time intensive quantitative model development to be directed at
506 variables with the highest importance. The method is based on a matrix level analysis of the
507 component interactions (see for example Herr et al., 2016; Ickowicz et al., 2018).

508 [In the process of building a qualitative model, the expert developed a consensus on the overall scope](#)
509 [of the model, namely the model components and their interactions.](#) The hydrological variables, and
510 relationships between ecosystem components that the [experts prioritised in the](#) qualitative
511 modelling process ~~prioritised for upland riverine systems in the Namoi region~~ were the
512 macroinvertebrate responses to riverine system change, presence of tadpoles and changes in
513 projected foliage cover in the riparian trees along the stream channel (Table 7).

514 **Step 2:** Use qualitative model priorities to develop quantitative models.

515 In this context, qualitative models highlighted critical relationships and variables that became the
516 focus of the quantitative models (see for example Herr et al., 2016; Hosack et al., 2018; Ickowicz et
517 al., 2018). The focus of the quantitative models was on three elements within the upland riverine
518 landscape classes (Table 7): (i) the response of upland riparian trees to changes in groundwater; (ii)
519 macroinvertebrate assemblage changes related to days with no consecutive water (zero-flow days)
520 and the longest zero flow event period; and (iii) the response of tadpoles to zero flow days and
521 longest zero flow event period. Table 7 provides a brief summary of these variables; specific details
522 of the variable definitions are in Ickowicz et al. (2018).

523

524

525 **Table 7: Upland riverine ecosystem quantitative modelling variables that experts prioritised in the qualitative model and**
 526 **associated ecological and hydrological variables used in the development of the quantitative impact model (after**
 527 **Ickowicz et al., 2018)**

Expert prioritised relationship	Ecological variable (with associated sample units)	Hydrological variable
Response of the upland riparian forest to changes in hydrological regime and groundwater	Annual mean projected foliage cover of species group that includes: Casuarina, yellow box, Blakely's red gum, <i>Acacia salicina</i> , <i>Angophora floribunda</i> , grey box. Transect of 50 m length and 20 m width that extends from first bench ('toe') on both sides of stream	<ul style="list-style-type: none"> • The mean annual number of events with a peak daily flow exceeding the overbank flow events. • Maximum difference in drawdown under a baseline and under the expected drawdown • The year with the maximum difference in drawdown relative to the baseline
Response of fast-water macroinvertebrates to changes in number of zero-flow days and maximum zero-flow event	Average number of families of aquatic macroinvertebrates in riffle habitat sampled using the NSW AUSRIVAS method for riffles	<ul style="list-style-type: none"> • The number of zero-flow days per year, averaged over a 30-year period. • The maximum length of spells (in days per year) with zero flow, averaged over a 30-year period.
Response of tadpoles to changes in number of zero-flow days and maximum zero-flow event	Probability of presence of tadpoles from <i>Limnodynastes</i> genus (species <i>dumerilii</i> , <i>salmini</i> , <i>interioris</i> and <i>terraereginae</i>) sampled using standard 30 cm dip net	<ul style="list-style-type: none"> • The number of zero-flow days per year, averaged over a 30-year period. • The maximum length of spells (in days per year) with zero low, averaged over a 30-year period.

528


529 **Step 3:** Identify risk areas in the regions where quantitative modelling indicated significant changes
530 to landscape group components.

531 This quantitative modelling approach incorporated expert elicitation in a Bayesian framework to
532 predict changes in ecological system components because of expected changes in hydrology
533 conditions. The method dealt with complexity and limited knowledge that allows for updating with
534 new information, which is an important feature in evidence-based decision making (see for example
535 Hosack et al., 2017).

536 The modelling of risk to ecosystems at regional scale focuses on recognising which parts of the region
537 are potentially impacted and which parts are unlikely to experience harm. Using our landscape
538 classification as a crucial input, the modelling delineated impacted areas within each region, based
539 on a zone of potential hydrological change. This is the area in the landscape, where hydrological
540 modelling identified an expected change to surface water and groundwater from future resource
541 extraction. Risk levels across a landscape group are a result of aggregating individual risks associated
542 with each ecological variable and categorising the risks into three levels based on their percentile
543 spreads (for details see Herr et al., 2018b).

544 For the Namoi region, for example, dryland remnant vegetation, human-modified ecosystems, no-
545 floodplain and upland riverine ecosystems and rainforests, will not experience impacts, while
546 floodplain and lowland ecosystems area and streams of floodplain and lowland ecosystems will
547 potentially experience impacts (Herr et al., 2018a). Figure 9 (a) shows the landscape groups that are
548 at risk of impact from hydrological changes as they are situated within the zone of potential
549 hydrological change, and Figure 9 (b) shows the risk level to these landscape groups from the
550 quantitative models. Note that there is a category “Remaining unquantified ‘floodplain and lowland
551 riverine’ classes”. The expert could not develop quantitative models for these classes, because there
552 was no surface water hydrological model available that could predict changes to surface water flows.
553 This was related to the lack of gauging data and groundwater interaction details specific to the
554 lowland drainage channels. Having lowland riverine classes whose risk remains unquantified means
555 there is additional work needed before an assessment and potential mitigation of impacts from
556 hydrological changes is possible (Herr et al., 2018b).

557




Landscape group


- Floodplain or lowland riverine
- Non-floodplain or upland riverine
- Rainforest
- Springs

Zone of potential hydrological change

- Zone of potential hydrological change
- Zone of potential hydrological change - Pilliga region
- Assessment extent
- Subregion
- Watercourse

NAM-340-001







559

560


Figure 9a: Landscape classes overlaying areas of potential hydrological change (Herr et al., 2018b)



- | | | |
|---|---|--|
| Level of risk | Zone of potential hydrological change | Mines (underground and open-cut) |
| 'At minimal risk' | Remaining unquantified 'Floodplain or lowland riverine' classes | Narrabri Gas Project |
| 'At some risk' | Assessment extent | Watercourse |
| 'More at risk' | | |



NAM-340-013



564
565

566 **Figure 9b: Hydrological change risk level of lowland landscape class areas (Herr et al., 2018b)**

5 Discussion

In Australia, there is no consistent national classification that links ecosystems at landscape level with their underlying hydrological system. While there are many different land classifications that incorporate hydrological aspects, they do not provide linkages between hydrology and landscape elements. None of these enable a broad scale ecological assessment of impacts associated with changes in water flow and availability, and they are not sufficiently generic for the purpose of assessing landscape level water related impacts on ecosystems in a spatially explicit manner (Kilroy et al., 2008; Elmore et al., 2003; Brown et al., 2014; Liermann et al., 2012; Doody et al., 2017; Poff et al., 2010; Aquatic Ecosystems Task Group, 2012; Olden et al., 2012; Gharari et al., 2011). However, the bioregional assessment program needed to assess impacts of coal resource developments on ecological systems via a water pathway. Hence, we developed an ecological landscape classification that would be applicable to the markedly different assessment regions.

We developed this classification based on existing datasets that were readily available in the areas of interest. This is much more resource and time efficient than gathering new data, using for example, remote sensing and taking hydrological measurements (see e.g. Gharari et al., 2011; Leibowitz et al., 2014; Sawicz et al., 2014; Zhang et al., 2016; Addicott et al., 2021; Carlier et al., 2021; Jones et al., 2021). The latter would have also required intensive methodology development, and would, in our opinion, not have provided fit-for-purpose information for the expert elicitation process. The advantage of our approach was that it integrated the relationships between water in the landscape and the landscape classes from the multiple dimensions in the input datasets, which allowed experts to develop causal reasoning. This causal relationship would have been much less clear when using dimensionality reduction and classifications such as proximity analysis because such methods do not infer causality without external information.

Our classification identifies the causal pathways between the water dependency of its components and human activities that result in hydrological changes. Prioritising hydrological features ensures that there is a conceptual linkage between hydrology and landscape classes, as it identifies ecohydrological landscape elements. This was crucial for the experts' understanding of how hydrological changes impact the landscape. No currently existing ecohydrological classification was suitable to do this, either because these were not spatially explicit or they did not cover the landscape completely (Poff et al., 2010; Aquatic Ecosystems Task Group, 2012; Olden et al., 2012). A spatially complete coverage of the landscape is an important prerequisite of the risk analysis because it enables assigning risk levels to the whole landscape, and it allows to identify parts of the landscape where there is insufficient information from the other modelling components. In time critical environmental impact assessments, developing models of different environmental elements often occurs in parallel for those areas where data are available. Where data are unavailable, such modelling is left for future work to improve the risk assessment. In our case, as we had a complete spatial coverage of the landscape, it enables pinpointing which part of the risk modelling inputs needed to prioritise further work. It identified the areas where hydrological modelling needed further refinement because of the lack of gauging stations and knowledge of surface water - groundwater interactions in some of the lowland drainage channels (Figure 9b).

While our spatially explicit landscape classification provided experts with the ability to readily identify cause and effect relationships between landscape elements and landscape hydrology, there are obvious differences between the landscape classifications in the three regions, reflecting their geographical differences (see Figure 5, Figure 7 and Figure 8). It provides the specificity that is

611 [required in a regional impact assessment, where the boundaries are based on a combination of](#)
612 [geology, water resources and administrative conditions. The regionality also means that there is need](#)
613 [for different datasets describing the landscape features that would not be available from a single](#)
614 [classification covering the whole of Australia.](#)

615 [Nevertheless, each landscape classification provides a typology with an explicit connection of water](#)
616 [to the landscape class. This connection enables a causal link between hydrological change and](#)
617 [impact to ecosystems represented by landscape classes. The causal linkage is dependent on \(i\) a](#)
618 [spatially explicit connection between water in the landscape and the landscape classes, \(ii\)](#)
619 [conceptual understanding how changes in water may result in a reaction of specific ecosystem](#)
620 [elements in the landscape class and/or landscape group and \(iii\) a way of modelling quantitative](#)
621 [changes in ecosystem elements related to changes in water that incorporates causality. Our](#)
622 [ecohydrological classification approach for landscapes provides this spatially explicit connection and](#)
623 [has implicit ecohydrological elements that foster the conceptual understanding of the causal linkage.](#)
624 [For example, spatially modelling groundwater level drawdown enables a prediction of which springs](#)
625 [may be experiencing impacts from water extraction and, with additional modelling, by how much](#)
626 [and when. Linking this information with ecological expert inputs, will then allow the identification of](#)
627 [impacts on the spring communities and the risk to the communities.](#)

628 [Subsequent ecological modelling using expert elicitation of potential impacts drew heavily on our](#)
629 [classification, which is based on a consistent rule-set and fosters conceptual understanding of](#)
630 [landscape processes and functions. It provides an essential framework for experts to understand and](#)
631 [conceptualise how modelled future hydrological changes from coal resource developments link to](#)
632 [potential ecological changes at the landscape level. It is the basis for modelling the ecological risk to](#)
633 [the landscape from hydrological changes and it allows the incorporation of different data sources](#)
634 [and existing classification schemes. This consistency makes the classification development](#)
635 [transparent, repeatable, and adjustable, should new data become available.](#)

636 **5.1 Limitations**

637 While the ecohydrological landscape classification approach provided the basis for the risk
638 assessment outlined above, there are some limitations that require consideration when attempting
639 to develop and apply this ecohydrological landscape classification approach.

640 An important issue for the landscape classification is formulating a typology that adequately reflects
641 both the functional and structural complexity of the ecosystem, ~~while delivering. At the same time, it~~
642 [also needs](#) a succinct and consistent representation of the system that is 'fit for purpose' ~~to assign,~~
643 [which in our context means showing a](#) hydrological connectivity between the landscape classes, and
644 within the general landscape. The systematic classification imposes discrete boundaries among
645 landscape components that may not adequately capture gradients within and across landscape
646 classes. This approach tends to simplify important components of ecotones such as 'transition' zones
647 or edges between landscape classes, where ecosystem processes and/or biodiversity are likely to
648 peak [and tensions between human induced boundaries occur](#) (Ward et al., 1999; Ryberg et al.,
649 2021). If landscape classes are treated purely as 'closed' ecosystems, then the result may be a poor
650 representation of the biotic interactions and energy exchange between adjacent systems, and this
651 could limit a conventional impact and risk analyses. These conceptual challenges may be important
652 considerations for subsequent impact assessments, requiring special attention in assigning risk from
653 human induced changes in hydrology. However, [conceptual expert](#) modelling of impacts ~~may be able~~

654 ~~to can~~ compensate for this shortfall, when, for example, incorporating riparian areas ~~within~~ riverine
655 and wetland ~~impact~~ model development. [In our case, experts intrinsically applied the ecotone](#)
656 [concept to riparian areas when discussing and assigning impacts to stream ecosystem variables, thus](#)
657 [overcoming the tension of boundaries that the classification imposed](#) (see also Hosack et al., 2018;
658 Ickowicz et al., 2018).

659 There are also spatial data issues that require additional consideration beyond just simply
660 incorporating existing data. There are several technical issues that constitute important gaps in the
661 landscape classification for the Namoi region, for example. Here, two different approaches to define
662 GDEs were required because one spatial dataset only included terrestrial vegetation and not riverine
663 systems mapped within the stream network (NSW Office of Water, 2015). A second GDE dataset
664 helped overcome this deficiency, and provided the basis to classify the stream network's dependency
665 on groundwater (Bioregional Assessment Programme, 2012).

666 Wetlands in large areas of Australia are not yet adequately mapped. The separation between
667 groundwater-dependent and surface water-dependent wetlands may not always be accurate. In
668 many areas there is little knowledge of groundwater – surface water interactions. There is also a
669 significant gap in the understanding of water thresholds for ecosystems associated with springs. In
670 part, this results from a lack of bores to provide meaningful groundwater data. Some examples of
671 these data gaps appear in the discussion of the functioning of springs in the Doongmabulla Springs
672 complex in the Galilee region, particularly in identifying the source aquifer (Fensham et al., 2016).

673 There is extensive work from Queensland that links regional ecosystems vegetation to their
674 groundwater needs, although the mapped areas are still small (Sattler and Williams, 1999;
675 Queensland Government, Queensland, 2016; Queensland Herbarium, 2021). However, in many parts
676 of Australia, GDE mapping and classification approaches are limited, and many areas lack systematic
677 ground-truthing. This is especially true in areas with extensive intact native vegetation remnants,
678 such as the Pilliga Forest of the Namoi region, where large areas of 'Grass woodland GDE' landscape
679 class exist, but the lack of published studies on vegetation–groundwater interactions limits a
680 definition of the nature of this interaction. [This is where a risk approach can compensate for the lack](#)
681 [of knowledge because an elevated assigned risk can reflect the limits in understanding.](#)

682 **5.2 Conclusions**

683 We showed that our landscape classification approach worked in the three geographically different
684 regions, with widely disparate information sources that fed into a landscape classification. This also
685 makes the approach resource efficient where existing spatial landscape or ecosystem classification
686 schemes, developed for other purposes, can be incorporated into the classification.

687 The study was able to formulate and implement an attribute-based classification scheme to define
688 and delineate water-dependent features across three large regions. We conclude that this approach
689 allowed us to repurpose several existing schemas into an adaptable and practical typology of a
690 landscape classification. The conceptual framework of landscape ecohydrology forms the basis for
691 this classification, which is used to focus subsequent analysis of potential cumulative impacts on
692 water resources from multiple coal resource developments. The classification enabled the
693 development of specific conceptual and [qualitative/quantitative](#) models that linked changes in
694 hydrology to potential impacts on ecosystems using the landscape classes. The classification provided
695 crucial inputs for a risk analysis of landscape components subjected to hydrological changes.

696 Applying our approach to different regions showed that it is sufficiently general and flexible to enable
697 the development of ecohydrological classifications in regions in Australia and potentially in other
698 regions around the globe, given a sufficiently mature information base and data availability.

699 **6 References**

700 Abella, S. R., Shelburne, V. B., and MacDonald, N. W.: Multifactor classification of forest landscape
701 ecosystems of Jocassee Gorges, southern Appalachian Mountains, South Carolina, Canadian Journal
702 of Forest Research, 33, 1933-1946, <https://doi.org/10.1139/x03-116>, 2003.

703 Addicott, E., Neldner, V. J., and Ryan, T.: Aligning quantitative vegetation classification and landscape
704 scale mapping: updating the classification approach of the Regional Ecosystem classification system
705 used in Queensland, Australian Journal of Botany, 69, 400-413, <https://doi.org/10.1071/BT20108>,
706 2021.

707 Aquatic Ecosystems Task Group: Aquatic Ecosystems Toolkit. Module 1: Aquatic Ecosystems Toolkit
708 Guidance Paper., Australian Government Department of Sustainability, Environment, Water,
709 Population and Communities, Canberra, [https://www.awe.gov.au/water/publications/aquatic-
710 ecosystems-toolkit-module-1-guidance-paper](https://www.awe.gov.au/water/publications/aquatic-ecosystems-toolkit-module-1-guidance-paper), 2012.

711 Australian Bureau of Agricultural and Resource Economics and Sciences: Catchment Scale Land Use
712 of Australia - 2014, Bioregional Assessment Source Dataset. [dataset],
713 <http://data.bioregionalassessments.gov.au/dataset/6f72f73c-8a61-4ae9-b8b5-3f67ec918826>, 2014.

714 Bioregional Assessment Programme: National Groundwater Dependent Ecosystems (GDE) Atlas,
715 Bioregional Assessment Derived Dataset [dataset],
716 <http://data.bioregionalassessments.gov.au/dataset/e358e0c8-7b83-4179-b321-3b4b70df857d>,
717 2012.

718 Bioregional Assessment Programme: Asset database for the Maranoa-Balonne-Condamine subregion
719 on 26 June 2015 [dataset], [http://data.bioregionalassessments.gov.au/dataset/35e95025-f962-4425-
720 83c7-767e2d6722e6](http://data.bioregionalassessments.gov.au/dataset/35e95025-f962-4425-83c7-767e2d6722e6), 2015.

721 Bioregional Assessment Programme: Asset database for the Namoi subregion on 18 February 2016,
722 Bioregional Assessment Derived Dataset [dataset],
723 <http://data.bioregionalassessments.gov.au/dataset/22061f2c-e86d-4ca8-9860-c349c2513fd8>, 2016.

724 Bioregional Assessment Programme: Landscape classification of the Namoi preliminary assessment
725 extent. Bioregional Assessment Derived Dataset. [dataset],
726 <http://data.bioregionalassessments.gov.au/dataset/360c39e5-1225-401d-930b-f5462fdb8005>, 2017.

727 Bioregional Assessments: Bioregional Assessment Programme,
728 <https://www.bioregionalassessments.gov.au/bioregional-assessment-program>, last access:
729 26/08/2019, 2018.

730 Bioregional Assessments: Geological and Bioregional Assessment Program,
731 <https://www.bioregionalassessments.gov.au/geological-and-bioregional-assessment-program>, last
732 access: 26/08/2019, 2019.

733 Brown, S. C., Lester, R. E., Versace, V. L., Fawcett, J., and Laurenson, L.: Hydrologic Landscape
734 Regionalisation Using Deductive Classification and Random Forests, Plos One, 9,
735 <https://doi.org/10.1371/journal.pone.0112856>, 2014.

736 Bureau of Meteorology: Australian Hydrological Geospatial Fabric ('Geofabric'), version 2.1.1,
737 Canberra,
738 [http://www.bom.gov.au/water/geofabric/documents/v2_1/ahgf_dps_surface_cartography_V2_1_re-](http://www.bom.gov.au/water/geofabric/documents/v2_1/ahgf_dps_surface_cartography_V2_1_release.pdf)
739 [lease.pdf](http://www.bom.gov.au/water/geofabric/documents/v2_1/ahgf_dps_surface_cartography_V2_1_release.pdf), 2012.

740 Carlier, J., Doyle, M., Finn, J. A., Ó hUallacháin, D., and Moran, J.: A landscape classification map of
741 Ireland and its potential use in national land use monitoring, *Journal of Environmental Management*,
742 289, 112498, <https://doi.org/10.1016/j.jenvman.2021.112498>, 2021.

743 CSIRO: Multi-resolution Valley Bottom Flatness MrVBF at three second resolution CSIRO 20000211
744 [dataset], [http://data.bioregionalassessments.gov.au/dataset/7dfc93bb-62f3-40a1-8d39-](http://data.bioregionalassessments.gov.au/dataset/7dfc93bb-62f3-40a1-8d39-0c0f27a83cb3)
745 [0c0f27a83cb3](http://data.bioregionalassessments.gov.au/dataset/7dfc93bb-62f3-40a1-8d39-0c0f27a83cb3), 2000.

746 Cullum, C., Brierley, G., Perry, G., and Witkowski, E.: Landscape archetypes for ecological classification
747 and mapping: The virtue of vagueness, *Progress in Physical Geography*, 41,
748 <http://doi.org/10.1177/0309133316671103>, 2016a.

749 Cullum, C., Rogers, K. H., Brierley, G., and Witkowski, E. T. F.: Ecological classification and mapping for
750 landscape management and science, *Progress in Physical Geography*, 40, 38-65,
751 <https://doi.org/10.1177/0309133315611573>, 2016b.

752 Department of Agriculture, W. a. t. E.: National Vegetation Information System (NVIS),
753 [https://www.awe.gov.au/agriculture-land/land/native-vegetation/national-vegetation-information-](https://www.awe.gov.au/agriculture-land/land/native-vegetation/national-vegetation-information-system)
754 [system](https://www.awe.gov.au/agriculture-land/land/native-vegetation/national-vegetation-information-system), last access: 22 Mar 2022, 2021.

755 Department of Sustainability, Environment, Water, Population and Communities: Murray-Darling
756 Basin aquatic ecosystem classification [dataset],
757 <http://data.bioregionalassessments.gov.au/dataset/a854a25c-8820-455c-9462-8bd39ca8b9d6>, 2014.

758 Doody, T. M., Barron, O. V., Dowsley, K., Emelyanoya, I., Fawcett, J., Oyerton, I. C., Pritchard, J. L., Van
759 Dijkf, A. I. J. M., and Warren, G.: Continental mapping of groundwater dependent ecosystems: A
760 methodological framework to integrate diverse data and expert opinion, *Journal of Hydrology-*
761 *Regional Studies*, 10, 61-81, <https://doi.org/10.1016/j.ejrh.2017.01.003>, 2017.

762 Eigenbrot, F.: Redefining Landscape Structure for Ecosystem Services, *Current Landscape Ecology*
763 *Reports*, 1, 80-86, <http://doi.org/10.1007/s40823-016-0010-0>, 2016.

764 Elmore, A. J., Mustard, J. F., and Manning, S. J.: Regional patterns of plant community response to
765 changes in water: Owens Valley, California, *Ecol Appl*, 13, 443-460, [https://doi.org/10.1890/1051-](https://doi.org/10.1890/1051-0761(2003)013[0443:RPOPCR]2.0.CO;2)
766 [0761\(2003\)013\[0443:RPOPCR\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)013[0443:RPOPCR]2.0.CO;2), 2003.

767 Evans, T., Tan, K., Magee, J., Karim, F., Sparrow, A., Lewis, S., Marshall, S., Kellett, J., and Galinec, V.:
768 Context statement for the Galilee subregion. Product 1.1 from the Lake Eyre Basin Bioregional
769 Assessment, Department of the Environment, Bureau of Meteorology, CSIRO and Geoscience
770 Australia, Australia, <http://data.bioregionalassessments.gov.au/product/LEB/GAL/1.1>, 2014.

771 Fensham, R., Silcock, J., Laffineur, B., and MacDermott, H.: Lake Eyre Basin Springs Assessment
772 Project: hydrogeology, cultural history and biological values of springs in the Barcardine, Springvale
773 and Flinders River supergroups, Galilee Basin springs and Tertiary springs of western Queensland.
774 Report to Office of Water Science, Department of Science, Information Technology and Innovation,
775 Brisbane., <https://publications.qld.gov.au/dataset/11c1af89-93b9-497a-b99f->

776 [2ec6c7a8d339/resource/c5d1813b-73a4-4e05-aa86-39a8ed3045fb/download/lebsa-hchb-report-](https://data.bioregionalassessments.gov.au/dataset/a0650f18-518a-4b99-a553-44f82f28bb5f)
777 [springs-wst-qld.pdf](https://data.bioregionalassessments.gov.au/dataset/a0650f18-518a-4b99-a553-44f82f28bb5f), 2016.

778 Geoscience Australia: GEODATA TOPO 250K Series 3 [dataset],
779 <http://data.bioregionalassessments.gov.au/dataset/a0650f18-518a-4b99-a553-44f82f28bb5f>, 2006.

780 Gharari, S., Hrachowitz, M., Fenicia, F., and Savenije, H. H. G.: Hydrological landscape classification:
781 investigating the performance of HAND based landscape classifications in a central European meso-
782 scale catchment, *Hydrol. Earth Syst. Sci.*, 15, 3275-3291, <https://doi.org/10.5194/hess-15-3275-2011>,
783 2011.

784 Gleeson, T. and Paszkowski, D.: Perceptions of scale in hydrology: what do you mean by regional
785 scale?, *Hydrological Sciences Journal*, 59, 99-107, <https://doi.org/10.1080/02626667.2013.797581>,
786 2014.

787 Hall, J., Storey, D., Piper, V., Bolton, E., Woodford, A., and Jolly, J.: Ecohydrological conceptualisation
788 for the eastern Pilbara region. A report prepared for BHP Billiton Iron Ore., Subiaco, 240,
789 [https://www.bhp.com/-/media/bhp/regulatory-information-media/iron-ore/western-australia-iron-](https://www.bhp.com/-/media/bhp/regulatory-information-media/iron-ore/western-australia-iron-ore/0000/report-appendices/160316_ironore_waio_pilbarastrategicassessment_state_appendix7_appendixd.pdf)
790 [ore/0000/report-](https://www.bhp.com/-/media/bhp/regulatory-information-media/iron-ore/western-australia-iron-ore/0000/report-appendices/160316_ironore_waio_pilbarastrategicassessment_state_appendix7_appendixd.pdf)
791 [appendices/160316_ironore_waio_pilbarastrategicassessment_state_appendix7_appendixd.pdf](https://www.bhp.com/-/media/bhp/regulatory-information-media/iron-ore/western-australia-iron-ore/0000/report-appendices/160316_ironore_waio_pilbarastrategicassessment_state_appendix7_appendixd.pdf),
792 2015.

793 Hawkins, C. P. and Norris, R. H.: Performance of different landscape classifications for aquatic
794 bioassessments: introduction to the series, *Journal of the North American Benthological Society*, 19,
795 367-369, <https://www.journals.uchicago.edu/doi/pdfplus/10.1899/0887-3593-19.3.367>, 2000.

796 Hawkins, C. P., Norris, R. H., Gerritsen, J., Hughes, R. M., Jackson, S. K., Johnson, R. K., and Stevenson,
797 R. J.: Evaluation of the use of landscape classifications for the prediction of freshwater biota:
798 synthesis and recommendations, *Journal of the North American Benthological Society*, 19, 541-556,
799 <https://doi.org/10.2307/1468113> 2000.

800 Herr, A., Dambacher, J. M., Pinkard, E., Glen, M., Mohammed, C., and Wardlaw, T.: The uncertain
801 impact of climate change on forest ecosystems How qualitative modelling can guide future research
802 for quantitative model development, *Environ Modell Softw*, 76, 95-107,
803 <https://doi.org/10.1016/j.envsoft.2015.10.023>, 2016.

804 Herr, A., Brandon, C., Beringen, H., Merrin, L. E., Post, D. A., Mitchell, P. J., Crosbie, R., Aryal, S. K.,
805 Janarhanan, S., Schmidt, R. K., and Henderson, B. L.: Assessing impacts of coal resource development
806 on water resources in the Namoi subregion: key findings. Product 5: Outcome synthesis for the
807 Namoi subregion from the Northern Inland Catchments Bioregional Assessment,
808 <http://data.bioregionalassessments.gov.au/product/NIC/NAM/5>, 2018a.

809 Herr, A., Aryal, S. K., Brandon, C., Crawford, D., Crosbie, R., Davies, P., Dunne, R., Gonzalez, D., Hayes,
810 K. R., Henderson, B. L., Hosack, G., Ickowicz, A., Janarhanan, S., Marvanek, S., Mitchell, P. J., Merrin, L.
811 E., Herron, N. F., O'Grady, A. P., and Post, D. A.: Impact and risk analysis for the Namoi subregion.
812 Product 3-4 for the Namoi subregion from the Northern Inland Catchments Bioregional Assessment,
813 [https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-namoi-](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-namoi-subregion)
814 [subregion](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-namoi-subregion), 2018b.

815 Hobbs, R. J. and McIntyre, S.: Categorizing Australian landscapes as an aid to assessing the generality
816 of landscape management guidelines, *Global Ecology and Biogeography*, 14, 1-15,
817 <http://dx.doi.org/10.1111/j.1466-822X.2004.00130.x>, 2005.

818 Holland, K., Beringen, H., Brandon, C., Crosbie, R., Davies, P., Gonzalez, D., Henderson, B.,
819 Janardhanan, S., Lewis, S., Merrin, L., Mitchell, P., Mount, R., O'Grady, A., Peeters, L., Post, D.,
820 Schmidt, R., Sudholz, C., and Turnadge, C.: Impact and risk analysis for the Maranoa-Balonne-
821 Condamine subregion. Product 3-4 for the Maranoa-Balonne-Condamine subregion from the
822 Northern Inland Catchments Bioregional Assessment, Department of the Environment and Energy,
823 Bureau of Meteorology, CSIRO and Geoscience Australia, Australia,
824 <http://data.bioregionalassessments.gov.au/product/NIC/MBC/3-4>, 2017.

825 Hosack, G., Ickowicz, A., Hayes, K. R., Dambacher, J. M., Barry, S. A., and Henderson, B. L.: Receptor
826 impact modelling. Submethodology M08 from the Bioregional Assessment Technical Programme,
827 <http://data.bioregionalassessments.gov.au/submethodology/M08>, 2018.

828 Hosack, G. R., Hayes, K. R., and Barry, S. C.: Prior elicitation for Bayesian generalised linear models
829 with application to risk control option assessment, *Reliab Eng Syst Safe*, 167, 351-361,
830 <https://doi.org/10.1016/j.ress.2017.06.011>, 2017.

831 Ickowicz, A., Hosack, G., Mitchell, P. J., Dambacher, J. M., Hayes, K. R., O'Grady, A. P., Henderson, B. L.,
832 and Herron, N. F.: Receptor impact modelling for the Namoi subregion. Product 2.7 for the Namoi
833 subregion from the Northern Inland Catchments Bioregional Assessment,
834 <http://data.bioregionalassessments.gov.au/product/NIC/NAM/2.7>, 2018.

835 Jones, C. E. J., Leibowitz, S. G., Sawicz, K. A., Comeleo, R. L., Stratton, L. E., Morefield, P. E., and
836 Weaver, C. P.: Using hydrologic landscape classification and climatic time series to assess hydrologic
837 vulnerability of the western U.S. to climate, *Hydrol. Earth Syst. Sci.*, 25, 3179-3206,
838 <https://doi.org/10.5194/hess-25-3179-2021>, 2021.

839 Kilroy, G., Ryan, J., Coxon, C., and Daly, D.: A Framework for the Assessment of Groundwater –
840 Dependent Terrestrial Ecosystems under the WaterFramework Directive,
841 <http://erc.epa.ie/safer/resource?id=b5799c70-224b-102c-b381-901ddd016b14>, 2008.

842 Leathwick, J. R., Overton, J. M., and McLeod, M.: An Environmental Domain Classification of New
843 Zealand and Its Use as a Tool for Biodiversity Management, *Conservation Biology*, 17, 1612-1623,
844 <http://dx.doi.org/10.1111/j.1523-1739.2003.00469.x>, 2003.

845 Leibowitz, S. G., Comeleo, R. L., Wigington Jr, P. J., Weaver, C. P., Morefield, P. E., Sproles, E. A., and
846 Ebersole, J. L.: Hydrologic landscape classification evaluates streamflow vulnerability to climate
847 change in Oregon, USA, *Hydrol. Earth Syst. Sci.*, 18, 3367-3392, <http://10.5194/hess-18-3367-2014>,
848 2014.

849 Lewis, S., Evans, T., Pavey, C., Holland, K., Henderson, B., Kilgour, P., Dehelean, A., Karim, F., Viney, N.,
850 Post, D., Schmidt, R., Sudholz, C., Brandon, C., Zhang, Y., Lymburner, L., Dunn, B., Mount, R.,
851 Gonzalez, D., Peeters, L., O'Grady, A., Dunne, R., Ickowicz, A., Hosack, G., Hayes, K., Dambacher, J.,
852 and Barry, S.: Impact and risk analysis for the Galilee subregion. Product 3-4 for the Galilee subregion
853 from the Lake Eyre Basin Bioregional Assessment., Department of the Environment and Energy,
854 Bureau of Meteorology, CSIRO and Geoscience Australia, Australia, 348,
855 [https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-galilee-](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-galilee-subregion)
856 [subregion](https://www.bioregionalassessments.gov.au/assessments/3-4-impact-and-risk-analysis-galilee-subregion), 2018.

857 Liermann, C. A. R., Olden, J. D., Beechie, T. J., Kennard, M. J., Skidmore, P. B., Konrad, C. P., and Imaki,
858 H.: Hydrogeomorphic Classification of Washington State Rivers to Support Emerging Environmental
859 Flow Management Strategies, *River Res Appl*, 28, 1340-1358, <http://doi.org/10.1002/rra.1541>, 2012.

860 MacMillan, R., Martin, T., Earle, T., and McNabb, D.: Automated analysis and classification of
861 landforms using high-resolution digital elevation data: applications and issues, *Canadian Journal of*
862 *Remote Sensing*, 29, 592-606, <https://doi.org/10.5589/m03-031>, 2003.

863 McMahon, G., Gregonis, S. M., Waltman, S. W., Omernik, J. M., Thorson, T. D., Freeouf, J. A., Rorick,
864 A. H., and Keys, J. E.: Developing a spatial framework of common ecological regions for the
865 conterminous United States, *Environmental Management*, 28, 293-316,
866 <http://doi.org/10.1007/s0026702429> 2001.

867 NSW Office of Environment and Heritage: Namoi Valley Flood Plain Atlas 1979, Bioregional
868 Assessment Source Dataset [dataset], [http://data.bioregionalassessments.gov.au/dataset/a854a25c-
869 8820-455c-9462-8bd39ca8b9d6](http://data.bioregionalassessments.gov.au/dataset/a854a25c-8820-455c-9462-8bd39ca8b9d6), 1979.

870 NSW Office of Environment and Heritage: Border Rivers Gwydir / Namoi Regional Native Vegetation
871 Map Version 2.0 VIS_ID_2004 [dataset],
872 <http://data.bioregionalassessments.gov.au/dataset/b3ca03dc-ed6e-4fdd-82ca-e9406a6ad74a>, 2015.

873 NSW Office of Water: Namoi CMA Groundwater Dependent Ecosystems [dataset],
874 <https://data.gov.au/data/dataset/a3e21ec4-ae53-4222-b06c-0dc2ad9838a8>, 2015.

875 NVIS Technological Working Group: Australian Vegetation Attribute Manual: National Vegetation
876 Information System, Version 7.0, Canberra,
877 [https://www.environment.gov.au/system/files/resources/292f10e2-8670-49b6-a72d-
878 25e892a92360/files/australian-vegetation-attribute-manual-v70.pdf](https://www.environment.gov.au/system/files/resources/292f10e2-8670-49b6-a72d-25e892a92360/files/australian-vegetation-attribute-manual-v70.pdf), 2017.

879 Office of Groundwater Impact Assessment: Spring vents assessed for the Surat Underground Water
880 Impact Report 2012 [dataset], [http://data.bioregionalassessments.gov.au/dataset/6d2b59fc-e312-
881 4c89-9f10-e1f1b20a7a6d](http://data.bioregionalassessments.gov.au/dataset/6d2b59fc-e312-4c89-9f10-e1f1b20a7a6d), 2015.

882 Olden, J. D., Kennard, M. J., and Pusey, B. J.: A framework for hydrologic classification with a review
883 of methodologies and applications in ecohydrology, *Ecohydrology*, 5, 503-518,
884 <http://dx.doi.org/10.1002/eco.251>, 2012.

885 Pain, C., Gregory, L., Wilson, P., and McKenzie, N.: The physiographic regions of Australia –
886 Explanatory notes 2011, CSIRO, Canberra,
887 <https://publications.csiro.au/rpr/download?pid=csiro:EP113843&dsid=DS4>, 2011.

888 Poff, N. L., Richter, B. D., Arthington, A. H., Bunn, S. E., Naiman, R. J., Kendy, E., Acreman, M., Apse, C.,
889 Bledsoe, B. P., Freeman, M. C., Henriksen, J., Jacobson, R. B., Kennen, J. G., Merritt, D. M., O’Keeffe, J.
890 H., Olden, J. D., Rogers, K., Tharme, R. E., and Warner, A.: The ecological limits of hydrologic
891 alteration (ELOHA): a new framework for developing regional environmental flow standards,
892 *Freshwater Biology*, 55, 147-170, <https://doi.org/10.1111/j.1365-2427.2009.02204.x>, 2010.

893 Post, D. A., Crosbie, R. S., Viney, N. R., Peeters, L. J., Zhang, Y., Herron, N. F., Janardhanan, S., Wilkins,
894 A., Karim, F., Aryal, S. K., Pena-Arancibia, J., Lewis, S., Evans, T., Vaze, J., Chiew, F. H. S., Marvanek, S.,
895 Henderson, B. L., Schmidt, B., and Herr, A.: Impacts of coal resource development in eastern Australia
896 on groundwater and surface water, *Journal of Hydrology*, 591,
897 <https://doi.org/10.1016/j.jhydrol.2020.125281>, 2020.

898 Poulter, B., Ciais, P., Hodson, E., Lischke, H., Maignan, F., Plummer, S., and Zimmermann, N. E.: Plant
899 functional type mapping for earth system models, *Geosci. Model Dev.*, 4, 993-1010,
900 <https://doi.org/10.5194/gmd-4-993-2011>, 2011.

901 Pyne, M. I., Carlisle, D. M., Konrad, C. P., and Stein, E. D.: Classification of California streams using
902 combined deductive and inductive approaches: Setting the foundation for analysis of hydrologic
903 alteration, *Ecohydrology*, 10, <https://doi.org/10.1002/eco.1802>, 2017.

904 Queensland Department of Science, Information Technology, Innovation and the Arts: Queensland
905 wetland data version 3 - wetland areas [dataset],
906 <http://data.bioregionalassessments.gov.au/dataset/2a187a00-b01e-4097-9ca4-c9683e7f4786>, 2012.

907 Queensland Department of Science, Information Technology, Innovation and the Arts: Queensland
908 groundwater dependent ecosystems [dataset],
909 <http://data.bioregionalassessments.gov.au/dataset/10940dfa-d7ef-44fb-8ac2-15d75068fff8>, 2013.

910 Queensland Government, Queensland: Groundwater dependent ecosystem mapping background,
911 <https://wetlandinfo.des.qld.gov.au/wetlands/facts-maps/gde-background/>, last access: 09 Sep 21,
912 2016.

913 Queensland Herbarium: Regional Ecosystem Description Database (REDD). Version 12 (March 2021),
914 Brisbane, <https://www.qld.gov.au/environment/plants-animals/plants/ecosystems>, 2021.

915 Queensland Herbarium, Department of Science, Information Technology, Innovation and the Arts:
916 Queensland Groundwater Dependent Ecosystems and Shallowest Watertable Aquifer 20150714
917 [dataset], [http://data.bioregionalassessments.gov.au/dataset/3d36e3d4-b16b-43b3-b2eb-](http://data.bioregionalassessments.gov.au/dataset/3d36e3d4-b16b-43b3-b2eb-c1aea7ef9193)
918 [c1aea7ef9193](http://data.bioregionalassessments.gov.au/dataset/3d36e3d4-b16b-43b3-b2eb-c1aea7ef9193), 2015.

919 Ryberg, T., Davidsen, J., Bernhard, J., and Larsen, M. C.: Ecotones: a Conceptual Contribution to
920 Postdigital Thinking, *Postdigital Science and Education*, 3, 407-424, [https://doi.org/10.1007/s42438-](https://doi.org/10.1007/s42438-020-00213-5)
921 [020-00213-5](https://doi.org/10.1007/s42438-020-00213-5), 2021.

922 SA Department for Water: South Australian Wetlands – Groundwater Dependent Ecosystems (GDE)
923 Classification [dataset], [http://data.bioregionalassessments.gov.au/dataset/fc35d75a-f12e-494b-](http://data.bioregionalassessments.gov.au/dataset/fc35d75a-f12e-494b-a7d3-0f27e7159b05)
924 [a7d3-0f27e7159b05](http://data.bioregionalassessments.gov.au/dataset/fc35d75a-f12e-494b-a7d3-0f27e7159b05), 2010.

925 Sattler, P. S. and Williams, R. D.: The conservation status of Queensland's bioregional ecosystems,
926 Environmental Protection Agency, Brisbane 1999.

927 Sawicz, K. A., Kelleher, C., Wagener, T., Troch, P., Sivapalan, M., and Carrillo, G.: Characterizing
928 hydrologic change through catchment classification, *Hydrol. Earth Syst. Sci.*, 18, 273-285,
929 <https://doi.org/10.5194/hess-18-273-2014>, 2014.

930 Snelder, T. H., Cattaneo, F., Suren, A. M., and Biggs, B. J.: Is the River Environment Classification an
931 improved landscape-scale classification of rivers?, *Journal of the North American Benthological*
932 *Society*, 23, 580-598, [https://doi.org/10.1899/0887-3593\(2004\)023<0580:ITRECA>2.0.CO;2](https://doi.org/10.1899/0887-3593(2004)023<0580:ITRECA>2.0.CO;2), 2004.

933 Thackway, R. and Cresswell, I. D.: An Interim Biogeographic Regionalisation for Australia: A
934 framework for setting priorities in the national reserves system cooperative program, Australian
935 Nature Conservancy Agency, Canberra, 99,
936 [https://www.environment.gov.au/system/files/resources/4263c26f-f2a7-4a07-9a29-](https://www.environment.gov.au/system/files/resources/4263c26f-f2a7-4a07-9a29-b1a81ac85acc/files/ibra-framework-setting-priorities-nrs-cooperative-program.pdf)
937 [b1a81ac85acc/files/ibra-framework-setting-priorities-nrs-cooperative-program.pdf](https://www.environment.gov.au/system/files/resources/4263c26f-f2a7-4a07-9a29-b1a81ac85acc/files/ibra-framework-setting-priorities-nrs-cooperative-program.pdf), 1995.

938 Ward, J. V., Tockner, K., and Schiemer, F.: Biodiversity of floodplain river ecosystems: ecotones and
939 connectivity1, *Regulated Rivers: Research & Management*, 15, 125-139, 10.1002/(sici)1099-
940 1646(199901/06)15:1/3<125::Aid-rrr523>3.0.Co;2-e, 1999.

941 Welsh, W., Herron, N., Rohead-O'Brien, H., Ransley, T., Aryal, S., Mitchell, P., Buckerfield, S., and
942 Marshall, S.: Context statement for the Maranoa-Balonne-Condamine subregion. Product 1.1 for the
943 Maranoa-Balonne-Condamine from the Northern Inland Catchments Bioregional Assessment,
944 Department of the Environment, Bureau of Meteorology, CSIRO and Geoscience Australia, Australia,
945 <http://data.bioregionalassessments.gov.au/product/LNIC/MBC/1.1>, 2015.

946 Welsh, W., Hodgkinson, J., Strand, J., Northey, J., Aryal, S., O'Grady, A., Slatter, E., Herron, N.,
947 Pinetown, K., Carey, H., Yates, G., Raisbeck-Brown, N., and Lewis, S.: Context statement for the
948 Cooper subregion. Product 1.1 for the Namoi subregion from the Northern Inland Catchments
949 Bioregional Assessment. Department of the Environment, Bureau of Meteorology, Department of the
950 Environment, Bureau of Meteorology, CSIRO and Geoscience Australia, Australia,
951 <http://data.bioregionalassessments.gov.au/product/NIC/NAM/1.1>, 2014.

952 Wiens, J. A. and Milne, B. T.: Scaling of 'landscapes' in landscape ecology, or, landscape ecology from
953 a beetle's perspective, *Landscape Ecology*, 3, 87-96, <https://doi.org/10.1007/BF00131172>, 1989.

954 Wolfe, J. D., Shook, K. R., Spence, C., and Whitfield, C. J.: A watershed classification approach that
955 looks beyond hydrology: application to a semi-arid, agricultural region in Canada, *Hydrol. Earth Syst.*
956 *Sci.*, 23, 3945-3967, <https://10.5194/hess-23-3945-2019>, 2019.

957 Zhang, Z., Zang, R., Wang, G., and Huang, X.: Classification of Landscape Types Based on Land Cover,
958 Successional Stages and Plant Functional Groups in a Species-Rich Forest in Hainan Island, China,
959 *Tropical Conservation Science*, 9, 135-152, <https://doi.org/10.1177/194008291600900107>, 2016.

960

961 **7 Author contributions**

962 AH, LM undertook the original draft preparation. All authors contributed to review and editing,
963 conceptualisation, methodology and investigation.

964 **8 Competing interests**

965 The authors declare that they have no conflict of interest.

966 **9 Acknowledgements**

967 This research was carried out under the auspices of the Bioregional Assessment Programme, a
968 collaboration between the Australian Department of Environment and Energy, CSIRO, Geoscience
969 Australia and the Bureau of Meteorology. [The Authors wish to thank Willem Vervoort and two](#)
970 [anonymous reviewers for valuable suggestions that provided valuable improvements on the clarity of](#)
971 [the manuscript.](#)