

# Monitoring the combined effects of drought and salinity stress on crops using remote sensing in the Netherlands

Wen Wen<sup>1,\*</sup>, Joris Timmermans<sup>1,2,3</sup>, Qi Chen<sup>1</sup> and Peter M. van Bodegom<sup>1</sup>

<sup>1</sup>Institute of Environmental Sciences (CML), Leiden University, Box 9518, 2300 RA Leiden, The Netherlands

<sup>2</sup>Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 1090 GE Amsterdam, The Netherlands

<sup>3</sup>Lifewatch ERIC, vLab & Innovation Centre, 1090 GE Amsterdam, The Netherlands

Correspondence: Wen Wen ([w.wen@cml.leidenuniv.nl](mailto:w.wen@cml.leidenuniv.nl))

**Abstract.** Global sustainable agricultural systems are under threat, due to ~~projected~~ increasing and ~~of~~ co-occurring drought and salinity ~~stresses with climate change~~. Combined effects of these stresses drought and salinity on agricultural crops have traditionally been evaluated in small-scale experimental studies. ~~Consequently, As such the need exists for large~~ large-scale studies need to be performed that to increase our understanding and assessment of the combined impacts in agricultural practice in ~~real-real~~ real-life scenarios. This study aims to provide a new monitoring approach using remote sensing observations to evaluate estimate and compare the joint impacts of drought, ~~and~~ salinity ~~and their combination~~ on crop traits. In our tests over the Netherlands at large spatial (138.74 km<sup>2</sup>) and temporal extents in the Netherlands using remote sensing observations. ~~Specifically, for both maize and potato,~~ we calculated five functional traits for both maize and potato from Sentinel-2 observations, namely: leaf area index (LAI), the fraction of absorbed photosynthetically active radiation (FAPAR), the fraction of vegetation cover (FVC), leaf chlorophyll content (Cab) and leaf water content (Cw). Individual and combined effects of the stresses on the seasonal dynamics in crop traits were determined using both one-way and two-way ANOVAs. We found that both stresses (individual and co-occurring) affected the functional traits of both crops significantly (with R<sup>2</sup> ranging from 0.326 to 0.796), though with stronger sensitivities to drought than to salinity. While we found exacerbating effects within co-occurrent stresses, the impact-level depended strongly on the moment in the growing season. For both crops, LAI, FAPAR and FVC dropped the most under severe drought stress conditions. The patterns for Cab and Cw were more inhibited by co-occurring drought and salinity. Consequently, our study constitutes a way towards evaluating drought and salinity impacts in agriculture with the possibility of potential large-scale application for a sustainable food security.

**Keywords:** Drought; Salinity; Agriculture; Remote sensing; Functional traits

## 1 Introduction

Food production is required to increase by 70% to satisfy the growing population demand by the year 2050 (Godfray et al., 2010). Meanwhile, food security is becoming increasingly threatened due to the increasing abiotic stresses under the influence of global climate change. ~~Currently, abiotic stresses, including drought, soil salinity, nutrient stress and heavy metals, are estimated to constrain crop productivity by 50% ~ 80% (Shinozaki et al., 2015). Of these stresses, drought and salinity have been identified as the two main factors to limit crop growth, affecting respectively 40% and 11% of the global irrigated areas (FAO, 2020; Dunn et al., 2020). With drought and salinity forecasted to increase spatially and in severity (Schwalm et al., 2017; Trenberth et al., 2013; Rozema and Flowers, 2008), and with predictions of higher co-occurrence around the world (Wang et al., 2013; Corwin, 2020; Jones and van Vliet, 2018), food production will be more deeply deeper~~ challenged by both stresses.

Field Code Changed

Field Code Changed

Field Code Changed

37 Numerous small-scale experimental studies for a large variety of crops have shown that the impact of co-occurring drought  
38 and salinity stress is ~~exacerbated~~~~additive~~. ~~It was found that e~~Co-occurrence of drought and salinity stress ~~is found to~~  
39 ~~decreased~~ the yield of spinach (Ors and Suarez, 2017) and ~~of~~ the forage grass *Panicum antidotale* (Hussain et al., 2020)  
40 ~~more than~~ compared with the occurrence of one of these stresses only. Likewise, cotton root growth ~~tends was observed~~ to  
41 be more inhibited under the co-occurrence of drought and ~~salinity~~~~water stress~~ than by isolated occurrences (Zhang et al.,  
42 2013). Similarly, the exacerbating effect of co-occurring stresses ~~has been shown to limits~~ both maize reproductive growth  
43 and grain formation (Liao et al., 2022). While these ~~small-scale experimental~~ studies demonstrate the exacerbating effects  
44 of ~~co-occurring~~ drought and salinity ~~stress~~, they have limitations in projecting the impact towards real farmers' conditions  
45 due to their small-scale experimental nature. Thus, ~~there is still a significant knowledge gap concerning the large scale~~  
46 ~~evaluation of research focusing on~~ the combined impacts of drought and salinity ~~with respect to large-scale evaluation is~~  
47 ~~still a knowledge gap~~.

48 Remote sensing (RS) provides a huge potential to close this knowledge gap due to its capability ~~of to~~ monitoring continuous  
49 large areas at a frequent interval. ~~For this~~~~Traditionally,~~ remote sensing has ~~traditionally~~ used vegetation indices, such as  
50 Normalized Difference Vegetation Index (NDVI) (Tucker, 1979), ~~to monitor the impact on crop growth~~.  
51 ~~Nevertheless~~~~However,~~ such indices provide limited information on how ~~this the~~ impact is achieved (e.g. Wen et al., 2020)  
52 and how it can be mitigated. With the launch of better multispectral and high-resolution satellite sensors (such as Sentinel-  
53 2), new RS methods (e.g., hyperspectral, thermal infrared, microwave) have been identified to detect stress in both natural  
54 vegetation (Gerhards et al., 2019; Vereecken et al., 2012) as well as for agricultural applications (Homolova et al., 2013;  
55 Weiss et al., 2020). Specifically, these new RS methods allow for the retrieval of plant traits that directly link to plant  
56 processes, such as leaf biochemistry and photosynthetic processes, and thereby provide high potential for agricultural  
57 applications. RS plant traits of specific interest to monitor crop health include: leaf area index (LAI) (Wengert et al., 2021),  
58 canopy chlorophyll content (Cab\*LAI) (Gitelson et al., 2005), canopy water content (Cw\*LAI) (Kriston-Vizi et al., 2008),  
59 the fraction of absorbed photosynthetically active radiation (FAPAR) (Zhang et al., 2015), and the fraction of vegetation  
60 cover (FVC) (Yang et al., 2018). However, while there have been several attempts to monitor the response of crop health  
61 based on a multi-trait, multi-crop, and either drought or salinity focus, not much research has taken these ~~three~~ factors into  
62 account simultaneously (Wen et al., 2020).

63 In this study, we propose a novel approach to estimate, compare and evaluate the impacts of drought, salinity, and their  
64 combination on crop traits using remote sensing. ~~To allow for a detailed evaluation of this approach we applied it to analyse~~  
65 ~~the impacts of the the 2018 summer drought in the Netherlands on agricultural crops~~. ~~In this, A~~ a stress co-occurrence map  
66 was created by overlaying a high-resolution drought map of 2018 with a groundwater salinity map. Then, we characterized  
67 the response of maize and potato to different stress conditions based on five plant's traits (LAI, FAPAR, FVC, Cab and  
68 Cw). Two-way ANOVAs were adopted to test the main effects and the interactive effect between stress combinations and  
69 time on crop traits. Moreover, the effect of drought and salinity on crop traits was determined across the growing season  
70 with one-way ANOVAs. Consequently, this approach facilitates simultaneously monitoring crop health at various scales  
71 (regional, national, continental) across multiple stresses (drought, salinity) and multiple species.

## 72 2 Methodology


73 To achieve our aim of monitoring the impacts of (co-occurring) drought and salinity on agricultural production, we  
74 developed a new approach to estimate crop traits from remote sensing observations. Specifically, we developed an approach

Field Code Changed

Field Code Changed

75 that integrates image-processing techniques, such as image classification, co-registration, land surface parameter retrieval,  
 76 and time-series analysis. Using these techniques, we were able to estimate the drought, salinity, and **crop vegetation**-growth.

77



78

79 **Figure 1.** Technical workflow of the maps and data framework.

80 To allow for a detailed evaluation, we focused on the 2018 summer drought in the Netherlands. This period was selected  
 81 because of the extreme drought that affected a large part of Europe (Masante D., 2018). Within parts of the selected area  
 82 salinity was reported to increase during that same period (Broekhuizen, 2018). Hence this study area provides us with the  
 83 opportunity to investigate the combined impacts of these stresses on crops. In the following paragraphs, we provide more  
 84 information on the specific processing steps.

## 85 2.1 Study area and data

### 86 2.1.1 Drought map

87 A drought map of the Netherlands in 2018 was created based on the standardized precipitation evapotranspiration index  
 88 (SPEI) drought index, which was calculated from long-term precipitation data and potential evapotranspiration, from 2004  
 89 to 2018 (Chen et al., 2022) (Chen et al., 2021). Specifically, SPEI was estimated using a 3-month sliding time window, as  
 90 this [was](#) found best to investigate the impacts on the local ecosystems. We have extracted SPEI-3 data from April 1st to  
 91 October 30th, totally 214 days, as this coincided with the crop growth period of both maize and potato. Then, the drought  
 92 map was resampled to 250m resolution using the nearest neighbor interpolation and reprojected to RD\_new projection. [The RD\\_new projection \(EPSG:28992\) is a projected coordinate reference system of the Netherlands. All maps were](#)  
 93 [projected to RD\\_new projection to create consistent data layers. -We defined -1 and -1.5 as daily thresholds for different](#)  
 94

95 ~~drought severity classes according to previous classifications~~ (McKee et al., 1993; Tao et al., 2014). ~~Thus, (cumulative)~~  
96 ~~SPEI for no drought should be between -214 to 0. SPEI for moderate drought should be between -321 to -214 and for~~  
97 ~~severe drought, SPEI should be lower than -321 when calculated for the whole growing period Finally, the drought map~~  
98 ~~was classified into three classes namely no drought (SPEI from -214 to 0), moderate drought (SPEI from -321 to -214), and~~  
99 ~~severe drought (SPEI <= -321)~~ (McKee et al., 1993; Tao et al., 2014) (Fig. 2a).

#### 100 **2.1.1 Salinity map**

101 A ~~topsoil~~top-soil salinity map of the Netherlands was created based on a nationwide fresh-salt groundwater dataset, which  
102 derived chloride concentrations as a salinity indicator (<https://data.nhi.nu/>). To obtain the topsoil-salinity map of the top-  
103 soil ~~(~~, 15 layers of the groundwater salinity were extracted from the 3D groundwater salinity map. For each location, the  
104 layer closest to the corresponding to location's elevation (according to the Digital Elevation Model), i.e. closest to the soil  
105 surface, was selected. The salinity map was resampled to 250 m resolution and reprojected to RD\_new projection.  
106 Ultimately, the salinity map was classified into three levels namely no-salinity (0.1 g·L<sup>-1</sup> to 0.8 g·L<sup>-1</sup>), moderate salinity (0.8  
107 g·L<sup>-1</sup> to 2.5 g·L<sup>-1</sup>), severe salinity (>= 2.5 g·L<sup>-1</sup>) according to the salt-resistant capacity of various crops cultivated in the  
108 Netherlands (Mulder et al., 2018; Stuyt, 2016) (Fig. 2b).

#### 109 **2.1.3 Crop map**

110 The crop map of the Netherlands in 2018 was collected from the Key Register of Parcels (BRP) of the Netherlands  
111 Enterprise Agency (<https://www.pdok.nl/introductie/-/article/basisregistratie-gewaspercelen-brp->). The crop map was  
112 resampled to 250m resolution and reprojected to RD\_new projection (Fig. 2d).

#### 113 **2.1.4 Co-occurrence map of drought and salinity**


114 The drought map and the salinity map were overlain to evaluate co-occurrences of drought and salinity of the Netherlands  
115 in 2018 (Fig. 2c). By classifying the three stress levels for the individual occurrences, we obtained nine stress classes of  
116 co-occurring drought and salinity, namely no stress, moderate drought only (MD), severe drought only (SD), moderate  
117 salinity only (MS), severe salinity only (SS), moderate drought and moderate salinity (MD+MS), moderate drought and  
118 severe salinity (MD+SS), severe drought and moderate salinity (SD+MS), and severe drought and severe salinity (SD+SS).

Field Code Changed

Formatted: Font: 10 pt


Field Code Changed

Formatted: Font: 10 pt



Formatted: Centered

119



120

121

122 **Figure 2.** The overlap map of the Netherlands overlaying a) drought and b) salinity in the Netherlands to show c) the co-occurrence  
 123 of drought and salinity. The selected study area is indicated by black yellow-lines in panel c. d) The associated crop map of the study  
 124 area.

125 **2.1.5 Study area selection**

126 Based on the national map of the Netherlands (Fig. 2c), a single region with similar soil type, climate, tillage systems, and  
127 irrigation methods was chosen to minimize the interference of these factors on the observed trait expressions. The province  
128 of North-Holland was selected because it contained the most (7 out of 9) combinations of drought and salt stress (Fig. 2c),  
129 namely: no stress, MD, SD, MS, SS, MD+MS, and SD+SS. Moreover, both maize and potato were cultivated across all  
130 stress combinations in this province. For further analysis, MS and SS were grouped into a new class of salinity stress since  
131 the area of MS and SS was quite limited. Therefore, six classes of stress combinations namely no stress, MD, SD, salinity  
132 (MS+SS), MD+MS, and MD+SS were analyzed for the study area.

## 133 2.2 Traits retrieval

### 134 2.2.1 Satellite data

135 ~~The Sentinel-2 mission consists of two satellites equipped with the high-resolution Multispectral Instrument (MSI) in the~~  
136 ~~same orbit. This sensor The satellites acquires 13 spectral bands (with varying spatial resolutions) from in the visible~~  
137 ~~spectrum to and near the short-wavelength infrared spectrum in at 5 days of revisit times at a spatial resolution of 10m,~~  
138 ~~20m, and 60m (ESA, 2015). In our study, we used both the 10m and 20m Level 2A resolution observations, downloaded~~  
139 ~~from the The Copernicus Open Access Hub (<https://scihub.copernicus.eu/>), to facilitate the requirement as of the Sentinel~~  
140 ~~Application Platform (SNAP) toolbox requires for both optical and near-infrared observations to be available for~~  
141 ~~determining the functional traits. To create consistency across the bands, those Bands in with a 20m resolution (including~~  
142 ~~B5, B6, B7, B8A, B11 and B12) were resampled to the 10m resolution of to match consistency with B3 and B4. In total,~~  
143 ~~Eight eight cloud-free scenes were found (21/04/2018, 06/05/2018, 26/05/2018, 30/06/2018, 15/07/2018, 13/09/2018,~~  
144 ~~13/10/2018, and 28/10/2018) to cover the crop growth cycle. Although additional cloud-free scenes were found in prior~~  
145 ~~analyses, we found that none of the scenes in August (04/08/2018, 09/08/2018, 14/08/2018, 19/08/2018, 24/08/2018, and~~  
146 ~~29/08/2018), none was were of high quality, i.e. with low cloud cover, and we therefore choose to omit August from our~~  
147 ~~analysis. After this prior analysis, we downloaded Level 2A (L2A) data from The Copernicus Open Access Hub~~  
148 ~~(<https://scihub.copernicus.eu/>). Then, bands in 20 m and 60 m resolution were resampled to 10 m resolution to match~~  
149 ~~consistency for traits retrieval.~~

### 150 2.2.2 Traits selection

151 Plant traits (e.g. LAI, FAPAR, FVC, Cab and Cw) were selected in consideration of their corresponding impacts on crop  
152 functioning and their potential for assessment by remote sensing. LAI is a critical vegetation structural trait related to  
153 various plant functioning processes such as primary productivity, photosynthesis, and transpiration (Jarlan et al., 2008;  
154 Asner et al., 2003; Boussetta et al., 2012; Fang et al., 2019). FAPAR depends on vegetation structure, energy exchange,  
155 and illumination conditions while FAPAR is also an important parameter to assess primary productivity (Liang, 2020;  
156 Weiss and Baret, 2016). FVC is a promising parameter corresponding to the energy balance process such as temperature  
157 and evapotranspiration (Weiss and Baret, 2016). Cab is an effective indicator of stress and is strongly related to  
158 photosynthesis and resource strategy (Croft et al., 2017). Cw plays an important role in transpiration, stomatal conductance,  
159 photosynthesis, and respiration (Bowman, 1989; Zhu et al., 2017), as well as in drought assessment (Steidle Neto et al.,  
160 2017).

## 161 2.3 dataset processing

162 ~~The biophysical processor of within the Sentinel Application Platform (SNAP) toolbox was used to derives the five traits,~~  
163 ~~namely LAI, FAPAR, FVC, canopy chlorophyll content (CCC), and canopy water content (CWC), for each pixel from the~~

Formatted: Font: (Asian) +Body Asian (DengXian),  
(Asian) Chinese (China)

Field Code Changed

Field Code Changed

Field Code Changed

164 ~~Sentinel-2 top of canopy reflectance data. This biophysical processor utilizes is driven by an artificial neural~~  
165 ~~network (ANN) approach, trained using the PROSAIL simulated database (Weiss and Baret, 2016). This training utilized~~  
166 ~~canopy traits rather than leaf traits (estimated by multiplication with LAI) to improve their neural network performance.~~  
167 ~~For CCC and CWC, these two canopy traits are internally (in SNAP) calculated as (Cab\*LAI) and (Cw\*LAI), i.e. based~~  
168 ~~on their leaf equivalents. Cab and Cw use in our approach To obtain their leaf counterparts (Cw and Cab), to create fully~~  
169 ~~independent variables, CCC and CWC thus were need to be divided by LAI to obtain since both CCC and CWC are~~  
170 ~~calculated from LAI. Thus, Cab (=CCC / LAI) and Cw (=CWC / LAI). The biophysical processor of Sentinel Application~~  
171 ~~Platform (SNAP) was used to compute the selected canopy traits (LAI, FAPAR, FVC, Cab\*LAI, and Cw\*LAI) for each~~  
172 ~~pixel from the Sentinel 2 top of canopy reflectance data. This biophysical processor is driven by an artificial neural~~  
173 ~~network (ANN) approach, trained using the PROSAIL simulated database (Weiss and Baret, 2016). To eliminate the effects~~  
174 ~~of crop biomass on canopy levels of chlorophyll and water, they were expressed as their leaf content counterparts, namely~~  
175 ~~Cab (=Cab\*LAI / LAI) and Cw (=Cw\*LAI / LAI).~~

176 Pixels with quality flags were eliminated from the dataset. It was observed that in April no crop had yet been planted.  
177 Instead, we observed that only along the edge of the plots, e.g. in ditches, vegetation was found. This feature was used to  
178 generate a ditch map and to mask out pixels in trait maps for the other months. For each variable and each date, only data  
179 within the 95% confidence interval were taken to increase data robustness.

## 180 2.4 Analysis

181 ~~Due to the unbalance in the occurrence of stress conditions, Since the pixel counts of the six classes of stress combinations~~  
182 ~~namely no stress, MD, SD, salinity, MD+MS, and MD+SS were (highly) different, drought and salinity were not considered~~  
183 ~~as two independent factors. drought and salinity were not considered as two independent factors. Instead, two-way~~  
184 ANOVAs were adopted to test the main effects and the interactive effect between stress combinations (consisting of 6  
185 levels) and time (5 months) on crop traits. Significant effects of the main stress condition were investigated through post  
186 hoc tests to test whether interaction effects between drought and salinity had occurred. Two-way ANOVAs were run  
187 separately for each trait and each crop type (maize and potato) as we expected different patterns.  
188 In the Netherlands, potato and maize are planted between mid-April to early May. Crops are surfacing in May and harvested  
189 in October. Therefore, to evaluate the response of crops to stresses across the growing season, the effect of drought and  
190 salinity on crop traits was determined for May, June, July, and September with a one-way ANOVA. Tukey HSD post hoc  
191 tests were performed to identify the differences among the six stress combinations. All statistical analyses were performed  
192 with SPSS 27.0 (SPSS Inc., USA).

## 193 3 Results

### 194 3.1 Stress impacts depend on the moment in the growing season

195 The two-way ANOVAs revealed strong effects of date and stress level on the five traits with effect sizes of the response  
196 ( $R^2$ ) ranging from 0.326 to 0.796 for the five traits, which was similar for maize and potato. For both maize and potato,  $R^2$   
197 values were lowest for Cab and highest for LAI, FAPAR, and FVC. For maize, we found a significant main effect of both  
198 date and stress ( $p < 0.05$ ) for Cab, Cw, FAPAR, and FVC. In contrast, LAI was not significantly different across the  
199 different stress conditions. For potato, all main effects of date and stress were significant for all five crop traits (Table 1).

200 For all traits and both crops, the interaction between the effects of time and stress conditions was significant ( $p < 0.05$ )  
 201 (Table 1), indicating that the impact of stress depended on the moment in the growing season. Despite the significant  
 202 interaction terms, the partial Eta squared values (Table 1) showed that the effects of time in the growing season were much  
 203 stronger than those of stress or the interaction of date and stress. The effects of date for maize were stronger than for potato.  
 204 Interestingly, the effects of the interaction between date and stress were stronger than those of the main effects of stress,  
 205 suggesting strongly time-specific impacts of stress on the crop traits investigated. The interaction terms were strongest for  
 206 FVC.

207 **Table 1.** Two-way ANOVA for different crop traits by time series and stress interactions.

Crops	Traits	Factors	F	$p$	Partial Eta Squared	$R^2$
Maize	LAI	date	2144.5	0.000	0.636	0.766
		stress	1.4	0.226	0.001	
		date*stress	8.5	0.000	0.033	
	Cab	date	333.9	0.000	0.222	0.326
		stress	10.7	0.000	0.008	
		date*stress	3.6	0.000	0.015	
	Cw	date	952.1	0.000	0.449	0.590
		stress	9.9	0.000	0.007	
		date*stress	4.0	0.000	0.017	
	FAPAR	date	1865.9	0.005	0.603	0.738
		stress	3.3	0.000	0.002	
		date*stress	8.5	0.000	0.033	
FVC	date	2022.5	0.000	0.622	0.761	
	stress	22.1	0.000	0.015		
	date*stress	28.7	0.000	0.105		
Potato	LAI	date	752.1	0.000	0.273	0.782
		stress	13.7	0.000	0.006	
		date*stress	8.1	0.000	0.020	
	Cab	date	96.4	0.000	0.050	0.329
		stress	54.2	0.000	0.024	
		date*stress	8.7	0.000	0.023	
	Cw	date	347.4	0.000	0.158	0.571
		stress	68.1	0.000	0.030	
		date*stress	10.3	0.000	0.027	
	FAPAR	date	612.7	0.000	0.234	0.744
		stress	25.8	0.000	0.011	
		date*stress	14.0	0.000	0.034	
FVC	date	844.0	0.000	0.297	0.796	
	stress	18.8	0.000	0.008		
	date*stress	13.6	0.000	0.033		

208 **Note:**  $F$  indicates the test statistic of the  $F$ -test;  $p$  indicates whether the effect is statistically significant in comparison to the significance  
 209 level ( $p < 0.05$ ); Partial Eta Squared indicates the effect size of different factors;  $R^2$  indicates the percentage that the model coincides  
 210 with the data.

### 211 3.2 Response of LAI, FAPAR, FVC to drought and salinity

212 Given the significance of both date and stress and their interactions, subsequent one-way ANOVAs were performed to  
 213 compare the effects of drought and salinity on LAI, FAPAR, and FVC for maize and potato in May, June, July, and  
 214 September separately (Fig. 3). The patterns for LAI, FAPAR, and FVC were very similar, although they differ in details  
 215 and were therefore treated together.

Formatted: Font: Times New Roman, 9 pt

Formatted: Font: Times New Roman, 9 pt

Formatted: Normal, Line spacing: Multiple 1.25 li


Formatted: Font: Times New Roman, 9 pt, Italic

Formatted: Font: 9 pt

216 For maize, all of LAI, FAPAR, and FVC ~~LAI had the lowest ( $p < 0.05$ ) value under severe drought (SD) conditions while~~  
217 ~~both FAPAR and FVC~~ obtained their lowest value under MD+SS stress conditions in May. In June, both LAI and FVC  
218 dropped the most under salinity stress and it was significantly ( $p < 0.05$ ) different from MD, MD+MS, and MD+SS  
219 conditions, but not significantly different from no stress conditions. In contrast, FAPAR also reached its ~~the~~-lowest value  
220 (under MD+MS stress conditions) in June but had a significant difference ( $p < 0.05$ ) compared with no stress conditions.  
221 Both in July and September, LAI, FAPAR, and FVC all had the lowest value under SD conditions, and the difference was  
222 significant compared with no stress conditions.

223 For potato, LAI, FAPAR, and FVC had the lowest ( $p < 0.05$ ) value under MD+MS and MD+SS stress conditions in May.  
224 In June, LAI, FAPAR as well as FVC reached the lowest value under SD conditions and were significantly lower than in  
225 most other stress conditions even though the difference was not significant from no stress conditions. In July, there was a  
226 tendency for LAI, FAPAR, and FVC to be lower at under stress conditions, although none of the effects were significant.  
227 In September, however, LAI, FAPAR, and FVC significantly decreased under MD, MD+MS, and MD+SS conditions, and  
228 the difference was significant compared with no stress conditions. In addition, the difference was not significant among  
229 these three stress conditions.

230 Therefore, both for maize and potato, LAI, FAPAR, and FVC dropped the most under SD stress conditions when they  
231 reached their respective maximum value, compared with other stress conditions. At the same time, maize and potato were  
232 more sensitive to drought than salinity since no significant change was observed between drought conditions and conditions  
233 with a combination of drought and salinity stress.



235 **Figure 3.** Expressions of LAI, FAPAR, and FVC under various stress conditions in May, June, July, and September. Different letters in  
 236 each panel indicate significant differences ( $p < 0.05$ ). MD, moderate drought only; Salinity, salinity only; MD+MS, moderate drought  
 237 and moderate salinity; MD+SS, moderate drought and severe salinity (MD+SS); SD, severe drought only.


**Formatted:** Font: (Default) Times New Roman, 9 pt, Bold  
**Formatted:** Normal, Line spacing: Multiple 1.25 li  
**Formatted:** Font: (Default) Times New Roman, 9 pt  
**Formatted:** Font: Bold

### 3.3 Response of leaf chlorophyll and water content to drought and salinity

239 The one-way ANOVAs revealed that there were significant ( $p < 0.05$ ) impacts of the various stress conditions on Cab and  
 240 Cw (Fig. 4). For maize, Cab obtained its lowest value under salinity stress in May and June while it was not significantly  
 241 different from no stress conditions. However, in July, Cab reached the lowest value under MD+MS conditions although  
 242 the difference was not significant from other stress conditions. There were no significant changes observed for Cab in  
 243 September. For potato, Cab dropped the most under salinity conditions in May although the difference was not significant  
 244 from no stress conditions. Furthermore, Cab significantly decreased under MD+SS conditions in June and July, compared  
 245 with other conditions. Although Cab dropped the most under salinity conditions in September, the difference was not  
 246 significantly different from other conditions. In addition, compared with no stress, potato had the lowest Cab under MD+SS  
 247 conditions while there was no significant difference between MD+SS and salinity conditions only in most growing periods.  
 248 Cw decreased under all stress conditions in May, June, and July for both maize and potato, except for SD conditions in  
 249 May, compared with no stress conditions. At the same time, Cw reached its lowest value under MD+SS co-occurring  
 250 conditions and it was significantly different from under no stress conditions. Nonetheless, there were different changes for  
 251 both maize and potato in September. Cw was not significantly different among any conditions for maize while it was the  
 252 lowest under salinity conditions for potato.

253 Therefore, this analysis illustrates that salinity affected maize less than drought since crop responses were more obvious to  
 254 drought than salinity for Cab and Cw. In contrast, salinity showed a more severe effect on maize and potato at the early  
 255 growth stages for Cab. Meanwhile, Cab was affected by co-occurring drought and salinity in June and July for potato. It  
 256 seems that there was a non-additive effect of drought and salinity for Cw, since the changes were not significant between  
 257 MD+MS, MD+SS, MD, and salinity conditions. MD conditions, salinity, and MD+MS compared to no stress conditions.

**Formatted:** Font color: Auto  
**Formatted:** Font color: Auto  
**Formatted:** Font color: Auto



258 **Figure 4.** Trait expressions of Cab and Cw under various stress conditions in May, June, July, and September. Different letters in each  
 259 panel indicate significant differences ( $p < 0.05$ ). MD, moderate drought only; Salinity, salinity only; MD+MS, moderate drought and  
 260 moderate salinity; MD+SS, moderate drought and severe salinity (MD+SS); SD, severe drought only.

**Formatted:** Space After: 0 pt, Line spacing: Multiple 1.25 li  
**Formatted:** Font: (Asian) +Body Asian (DengXian), (Asian) Chinese (China), (Other) English (United States)  
**Formatted:** Font: (Asian) +Body Asian (DengXian), (Asian) Chinese (China), (Other) English (United States)  
**Formatted:** Font: (Asian) +Body Asian (DengXian), Not Italic, (Asian) Chinese (China), (Other) English (United States)  
**Formatted:** Font: (Asian) +Body Asian (DengXian), (Asian) Chinese (China), (Other) English (United States)  
**Formatted:** Font: (Asian) +Body Asian (DengXian), 9 pt, (Asian) Chinese (China), (Other) English (United States)  
**Formatted:** Font: Bold  
**Formatted:** Normal

263 **4 Discussion**

264 In this study, we quantified the large-scale impacts of co-occurring drought and salinity on a variety of crop traits using  
265 satellite remote sensing. We observed that –in contrast to our expectations – the impacts of salinity were not highly  
266 pronounced at this scale, with most strong impacts originating due to drought stress during the 2018 drought. Although  
267 irrigation may modify the severity of drought impacts on crops, we have evidence that irrigation did not play a major role  
268 in the patterns found in this case since all croplands included within our research area have been identified as rainfed  
269 cropland according to the ESA/CCI land cover map in 2018 (<https://maps.elie.ucl.ac.be/CCI/viewer/>). In addition, while -  
270 in the area- farmers are known to irrigate their cropland, the Dutch government announced a temporary national irrigation  
271 ban in various areas including our research area in 2018 (Perry de Louw, 2020) to spare water. Therefore, we assumed that  
272 irrigation management was absent during our study period. However, At specific moments in the growing season, salinity  
273 and/or the combined effects of salinity and drought pronouncedly affected individual crop traits. In this way, with  
274 increasing salinity driven by more intensive droughts, water allocation should not only be governed by the amount of water  
275 shortage, but also the salinity of the remaining water. In this paper, we provide the first evidence that those impacts can be  
276 monitored through remote sensing. This might provide a basis towards a monitoring system for multiple crops with multiple  
277 stresses as well as better governance policies to release this problem.

278 **4.1 Drought stress is more important than salinity stress in farmers' conditions**

279 The exacerbating effects of co-occurrent drought and salinity (Fig. 3 and Fig. 4) that we found are consistent with findings  
280 of small-scale experiments (e.g. greenhouses). Consistent with our results, synergistic effects of co-occurring water stress  
281 and salinity stress have been found on maize reproductive growth and grain formation in a field study (Liao et al., 2022).  
282 Spinach (*Spinaciaoleracea* L., cv. Ragoon) yield decreased more under co-occurring water-salinity stress in comparison  
283 with separate water stress and salinity (Ors and Suarez, 2017). The co-occurring drought and salinity stress was more  
284 harmful to cotton root growth compared to their individual effects- (Zhang et al., 2013). Moreover, the combined negative  
285 effect of drought and salinity stress on *Panicum antidotale* was stronger than that of single stress (Hussain et al., 2020).  
286 Our research showed that the outcomes of these small-scale experimental studies also apply to real large-scale  
287 environments, where different sources of variance are present. Specifically, we show that in real farmers' conditions, the  
288 co-occurrence of drought and salinity indeed can constitute a severe threat due to its interactive effects for-on crop growth.  
289 In addition, we evaluated whether drought or salinity stress has more impact on crop performance. We observed that maize  
290 and potato were generally more sensitive to drought than salinity in this study (Fig. 3 and Fig. 4). This is consistent with  
291 results of previous studies that highlight that drought impacts are generally more detrimental than salinity stress for crops,  
292 e.g. for sesame (*Sesamum indicum*) (Harfi et al., 2016), *Mentha pulegium* L. (Azad et al., 2021), durum wheat (Sayar et  
293 al., 2010), grass pea (Tokarz et al., 2020), and sweet sorghum (Patane et al., 2013). However, given that the threshold of  
294 salinity at which crop damage occurs (according to the FAO guidelines (Ayers and Westcot, 1985)) was surpassed in all  
295 situations in which salinity stress was imposed (including in our study), we initially expected salinity to be a stronger  
296 explanatory variable than drought. As such, salinity impacts on crop performance (by the FAO) may have been  
297 overestimated. Indeed, in an experimental field situation in which drought stress was carefully avoided, higher thresholds  
298 of salinity-induced damage were observed for potato (van Straten et al., 2021).

299 In combination, the results from our study (supported by results from other studies) suggest that salinity particularly induces  
300 adverse effects when co-occurring with drought stress. Water stress impacts on photosynthesis and biomass of plants were  
301 extenuated by salinity since salinity enhances the synthesis of ATP and NADPH by promoting photosynthetic pigments

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, 10 pt, Bold, Font color: Black, Kern at 16 pt

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Field Code Changed

Field Code Changed

Formatted: English (United States)

Formatted: English (United States)

Formatted: Font color: Red

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, 10 pt

Formatted: Normal, Line spacing: 1.5 lines

302 ~~and photosystem II efficiency. The impacts of combined drought and salinity stress on plant growth, chlorophyll content,~~  
303 ~~water use efficiency, and photosynthesis were less severe compared to drought alone. This indicates compensating effects~~  
304 ~~on carbon assimilation due to osmotic adjustments induced by Na<sup>+</sup> and Cl<sup>-</sup> (Hussain et al., 2020). Thus, the detrimental~~  
305 ~~effect of single drought stress on crop growth is considered to be mitigated by salinity. Thus, the detrimental effect of single~~  
306 ~~drought stress on crop growth is considered to be mitigated by salinity, which might be associated with the synergetic effect~~  
307 ~~in carbon assimilation and osmotic adjustment by Na<sup>+</sup> and Cl<sup>-</sup> (Hussain et al., 2020).~~

#### 309 4.2 Drought and salinity stress differ between growth stages

310 The responses to drought and salinity stress were different at different growth stages of the crops. This was expressed by  
311 the significant interactions between the effects of time and stress conditions for all of our crop responses (Table 1). We  
312 found that during the grain filling (maize) and tuber bulking phase (potato), the sensitivities of these crops are expressed  
313 distinctly in the non-harvested aboveground tissues (Fig. 3 and Fig.4), with clear differences in the remote sensing plant  
314 traits.

315 Given that we were not able to monitor the harvestable products, multiple mechanisms may explain these patterns. The  
316 relatively high leaf coverage (as related to LAI, FAPAR, and FVC) at salinity and severe drought conditions at the end of  
317 the growing season may be an expression of a compensation process. Specifically, early and prolonged drought could have  
318 led to more assimilates allocated to non-harvestable potato parts for drought resistance since the number of tubers reduced  
319 (Jefferies, 1995; Schittenhelm et al., 2006). In that case, we should consider their higher leaf coverage at the end of the  
320 season as a survival mechanism, rather than true drought tolerance, leading to reduced tuber yields (Daryanto et al., 2016b).

321 Future studies that combine remote sensing with harvesting data may be able to evaluate this mechanism in more detail.  
322 In our study, different response patterns of maize and potato occurred to the different stresses over the growing season.  
323 This is consistent with previous studies focusing on the impact of drought and/or salinity onsets. For potato, it has been  
324 suggested that tuber yields particularly decreased when drought stress occurs during the vegetative and tuber initiation  
325 stages than during the tuber bulking stage (Wagg et al., 2021), although another study observed the reverse pattern  
326 (Daryanto et al., 2016b). For maize, on the other hand, drought seems to have the most detrimental impact during the  
327 maturation stage (Mi et al., 2018; Zhang et al., 2019), and the reproductive phase (Daryanto et al., 2017; Daryanto et al.,  
328 2016a). Considering the additional co-varying factors within our 'real-life' study, it is very ~~probable~~promising that we  
329 were able to detect similar effects. This suggests that we may use satellite remote sensing –albeit less spatially precise than  
330 e.g. sensing through drones- as a cost-effective early warning signal for detecting drought and salinity stress at moments  
331 during the growing season when differences in crop performance are still subtle.

#### 332 4.3 A multi-trait approach to understanding crop responses to stress

333 In addition to ~~facilitating the evaluation of being able to evaluate~~ crop performance during multiple stages of the growing  
334 season (in contrast to most destructive methods), remote sensing also allows a multi-trait approach to better understand the  
335 mechanisms involved in crop responses. ~~Each of the five traits is associated with different functions of plants that might~~  
336 ~~be individually impacted by the different stresses. Therefore, focusing on only one individual metric (as commonly done;~~  
337 ~~see Wen et al. (2020) for a review) limits our capacity to gain full insight into drought and salinity responses. Hence, given~~  
338 ~~that individual crop traits may respond differently to drought and salinity reflecting its stress resistance and tolerance~~  
339 ~~strategy, the evaluation of these distinct responses may help to understand this strategy. In our study, Cab and Cw had a~~

Field Code Changed

Field Code Changed

Field Code Changed

340 ~~response to drought and salinity distinct from LAI, FAPAR, and FVC, which showed a similar pattern (Fig. 3 and Fig. 4).~~  
341 ~~Given that individual crop traits may differently respond to drought and salinity to reflect their stress-resistance and~~  
342 ~~tolerance strategies, the evaluation of these distinct responses may help understanding these strategies.~~

343 In this study, Cw was consistently lower in all drought and salinity treatments ~~as compared to~~ ~~compared to~~ no stress  
344 conditions in May, June, and July. Indeed, this is a common response of plants in response to drought and salinity (e.g.  
345 Wen et al., 2020). In ~~this~~ ~~that~~ respect, it is interesting that no decrease in Cw was observed at the end of the growing season,  
346 in ~~September~~ ~~October~~. Whether the phenomenon is related to the survival mechanism mentioned above or to the lower  
347 transpiration demands at the end of the season because of lower aboveground biomass, cannot be concluded from these  
348 data. Some evidence pointing to the survival mechanism is the finding (Ghosh et al., 2001; Levy, 1992) that the leaf dry  
349 matter increased for potato under drought/salinity stress (like in our study) while the dry matter of the tubers appeared to  
350 have a greater decline.

351 With respect to chlorophyll contents, we observed a decline in Cab ~~under~~ ~~at~~ ~~salinity~~ ~~conditions~~ ~~,~~ ~~at~~ ~~the~~ ~~salinity~~ ~~treatment~~  
352 in May and the MS+SS treatment in June and July, while no decrease was observed in any of the treatments exposed to  
353 drought only. This indicates that while total leaf area was not (much) affected by salinity, the salinity did negatively affect  
354 crop performance. It has been reported that chlorophyll content in maize was significantly reduced upon salinity, along  
355 with other plant traits including plant height, shoot/root biomass, and leaf numbers (Fatima et al., 2021; Mahmood et al.,  
356 2021). Likewise, similar patterns were obtained in potato plants in saline soil (Efimova et al., 2018). Hence, this implies  
357 that soil salinity tends to negatively affect crop growth and restrict nutrient uptake.

358 ~~Cab and Cw responses to drought and salinity were distinct from responses of LAI, FAPAR, and FVC (Fig. 3 and Fig. 4).~~  
359 ~~LAI, FAPAR, and FVC showed similar patterns to stress due to their highly physical correlation (Hu et al., 2020). The~~  
360 ~~different patterns of Cw and Cab point to different drought and salinity resistance strategy components associated with~~  
361 ~~these traits: LAI (and FAPAR/FVC) reflect the decrease in biomass due to stress, partly because stress directly and~~  
362 ~~negatively impacts growth and partly because having lower biomass decreases the evapotranspiration demands of the crop,~~  
363 ~~which increases the resilience of the crop to deal with drought. Cw represents another pathway to reduce evapotranspiration~~  
364 ~~demands, i.e. by reducing the amount of water per gram of leaves. Also, this response may be a direct effect of the more~~  
365 ~~negative pressure heads due to drought or due to increased osmotic pressures (due to salinity). It may also be part of the~~  
366 ~~adaptive strategy of the crop to increase its resilience. Cab also responds to drought and salinity, but in its own way, i.e. by~~  
367 ~~adapting its photosynthetic capacity while being affected by a lower stomatal conductance (due to drought and/or salinity).~~  
368 ~~See e.g. Wright et al. (2003) for a framework explaining these nitrogen-water interactions.~~

369 ~~In addition, our approach gives the insight to analyze the effect of stresses on yield based on the five traits, even though~~  
370 ~~yield cannot directly be derived from remote sensing. Traits including Cab, LAI, and FAPAR, have been used (either~~  
371 ~~separately or in combination) as a proxy for final yield estimates from remote sensing in many studies. For instance, NDVI~~  
372 ~~-which is based on the combination of LAI and Cab- is extensively used to estimate crop yield (Huang et al., 2014;~~  
373 ~~Mkhabela et al., 2011; Vannoppen et al., 2020). Also, LAI itself has been used for predicting the final yield (Sun et al.,~~  
374 ~~2017; Dente et al., 2008; Doraiswamy et al., 2005). Meanwhile, Cab and FAPAR were also proven to be highly correlated~~  
375 ~~with crop yield (López-Lozano et al., 2015; Ghimire et al., 2015). Thus, while yield cannot be estimated directly from~~  
376 ~~remote sensing or ground truth data at the desired high spatial resolution, our indicators do relate to yield and can be used~~  
377 ~~in more application-based contexts to inform on yield impacts.~~

Field Code Changed

## 378 5 Conclusions

379 In this study, we represent the first attempt to evaluate the [real-life](#) effects of drought, salinity, and their combination on  
380 crop [health traits in real-life conditions based using on multiple traits from](#) remote sensing [information monitoring](#). Our  
381 approach gives new insights for monitoring [multi-crop](#) growth under co-occurring stresses at a large scale with high-  
382 resolution data. We found that while in general temporal patterns –reflecting crop growth dynamics- were stronger than  
383 effects of stress conditions, stress impacts depended on the time of the growing season. Furthermore, we also found that  
384 the temporal dynamics in crop responses to drought and salinity were different for maize vs. potato. In general, the five  
385 investigated traits were more negatively affected by a combination of drought and salinity stress compared to individual  
386 stress. Meanwhile, both maize and potato responded more prominently to drought, thus demonstrating a stronger  
387 sensitivity, than to salinity. Specifically, LAI, FAPAR, and FVC dropped the most under severe drought stress conditions.  
388 Consequently, the proposed new approach poses a facilitated way for simultaneously monitoring the effect of drought and  
389 salinity on crops in large-scale agricultural applications.

390  
391 *Data availability.* The drought map of the Netherlands in 2018 is retrieved from Chen et al. (2022)–Chen et al. (2021);  
392 (Chen et al., 2022). The [topsoil](#) salinity map of the Netherlands is retrieved from The Netherlands Hydrological  
393 Instrumentarium (NHI) (<https://data.nhi.nu/>). The crop map of the Netherlands in 2018 is retrieved from the Key Register  
394 of Parcels (BRP) of the Netherlands Enterprise Agency ([https://www.pdok.nl/introductie/-/article/basisregistratie-  
395 gewaspercelen-brp-](https://www.pdok.nl/introductie/-/article/basisregistratie-gewaspercelen-brp-)). All satellite scenes are downloaded from The Copernicus Open Access Hub  
396 (<https://scihub.copernicus.eu/>). The dataset relevant to this study is available upon request from the corresponding author.

397  
398 *Author contributions.* Conceptualization, JT, PVB, and WW; methodology, JT, QC, WW, and PVB.; investigation, WW  
399 and QC; writing—original draft preparation, WW; writing—review and editing, PVB. and JT; supervision, PVB, and JT  
400 All authors have read and agreed to the published version of the manuscript.

401  
402 *Competing interests.* The authors declare no conflict of interest.

403  
404 *Financial support.* This work was supported by the China Scholarship Council (CSC).

## 405 References

- 406 Asner, G. P., Scurlock, J. M. O., and Hicke, J. A.: Global synthesis of leaf area index observations: Implications for  
407 ecological and remote sensing studies, *Glob. Ecol. Biogeogr.*, 12, 191-205, [https://doi.org/10.1046/j.1466-  
408 822X.2003.00026.x](https://doi.org/10.1046/j.1466-822X.2003.00026.x), 2003.
- 409 Ayers, R. S., and Westcot, D. W.: Water quality for agriculture, Food and Agriculture Organization of the United Nations  
410 Rome, 1985.
- 411 Azad, N., Rezayian, M., Hassanpour, H., Niknam, V., and Ebrahimzadeh, H.: Physiological mechanism of salicylic acid  
412 in mentha pulegium l. Under salinity and drought stress, *Braz. J. Bot.*, 44, 359-369, [https://doi.org/10.1007/s40415-021-  
413 00706-y](https://doi.org/10.1007/s40415-021-00706-y), 2021.
- 414 Boussetta, S., Balsamo, G., Beljaars, A., Kral, T., and Jarlan, L.: Impact of a satellite-derived leaf area index monthly  
415 climatology in a global numerical weather prediction model, *Int. J. Remote Sens.*, 34, 3520-3542,  
416 <https://doi.org/10.1080/01431161.2012.716543>, 2012.
- 417 Bowman, W. D.: The relationship between leaf water status, gas-exchange, and spectral reflectance in cotton leaves,  
418 *Remote Sens. Environ.*, 30, 249-255, [https://doi.org/10.1016/0034-4257\(89\)90066-7](https://doi.org/10.1016/0034-4257(89)90066-7), 1989.
- 419 ‘Storm duurt dagen, droogte duurt maanden’: [https://www.rijkswaterstaat.nl/nieuws/archief/2018/08/storm-duurt-dagen-  
420 droogte-duurt-maanden](https://www.rijkswaterstaat.nl/nieuws/archief/2018/08/storm-duurt-dagen-droogte-duurt-maanden), 2018.
- 421 Chen, Q., Timmermans, J., Wen, W., and van Bodegom, P. M.: A multi-metric assessment of drought vulnerability across

422 different vegetation types using high resolution remote sensing, *Sci. Total Environ.*, 2021.

423 Chen, Q., Timmermans, J., Wen, W., and van Bodegom, P. M.: A multi-metric assessment of drought vulnerability across  
424 different vegetation types using high-resolution remote sensing, *Sci. Total Environ.*, 154970,  
425 <https://doi.org/10.1016/j.scitotenv.2022.154970>, 2022.

426 Corwin, D. L.: Climate change impacts on soil salinity in agricultural areas, *Eur. J. Soil Sci.*, 72, 842-862,  
427 <https://doi.org/10.1111/ejss.13010>, 2020.

428 Croft, H., Chen, J. M., Luo, X., Bartlett, P., Chen, B., and Staebler, R. M.: Leaf chlorophyll content as a proxy for leaf  
429 photosynthetic capacity, *Glob. Change Biol.*, 23, 3513-3524, <https://doi.org/10.1111/gcb.13599>, 2017.

430 Daryanto, S., Wang, L., and Jacinthe, P. A.: Global synthesis of drought effects on maize and wheat production, *PLoS One*,  
431 11, e0156362, <https://doi.org/10.1371/journal.pone.0156362>, 2016a.

432 Daryanto, S., Wang, L. X., and Jacinthe, P. A.: Drought effects on root and tuber production: A meta-analysis, *Agric.*  
433 *Water Manag.*, 176, 122-131, <https://doi.org/10.1016/j.agwat.2016.05.019>, 2016b.

434 Daryanto, S., Wang, L. X., and Jacinthe, P. A.: Global synthesis of drought effects on cereal, legume, tuber and root crops  
435 production: A review, *Agric. Water Manag.*, 179, 18-33, <https://doi.org/10.1016/j.agwat.2016.04.022>, 2017.

436 Dente, L., Satalino, G., Mattia, F., and Rinaldi, M.: Assimilation of leaf area index derived from asar and meris data into  
437 cereals-wheat model to map wheat yield, *Remote Sens. Environ.*, 112, 1395-1407, <https://doi.org/10.1016/j.rse.2007.05.023>,  
438 2008.

439 Doraiswamy, P. C., Sinclair, T. R., Hollinger, S., Akhmedov, B., Stern, A., and Prueger, J.: Application of modis derived  
440 parameters for regional crop yield assessment, *Remote Sens. Environ.*, 97, 192-202,  
441 <https://doi.org/10.1016/j.rse.2005.03.015>, 2005.

442 Dunn, R. J. H., Stanitski, D. M., Gobron, N., Willett, K. M., Ades, M., Adler, R., Allan, R., Allan, R. P., Anderson, J.,  
443 Argüez, A., Arosio, C., Augustine, J. A., Azorin-Molina, C., Barichivich, J., Barnes, J., Beck, H. E., Becker, A., Bellouin,  
444 N., Benedetti, A., Berry, D. I., Blenkinsop, S., Bock, O., Bosilovich, M. G., Boucher, O., Buehler, S. A., Carrea, L.,  
445 Christiansen, H. H., Chouza, F., Christy, J. R., Chung, E. S., Coldewey-Egbers, M., Compo, G. P., Cooper, O. R., Covey,  
446 C., Crotwell, A., Davis, S. M., de Eyto, E., de Jeu, R. A. M., VanderSat, B. V., DeGasperi, C. L., Degenstein, D., Di  
447 Girolamo, L., Dokulil, M. T., Donat, M. G., Dorigo, W. A., Durre, I., Dutton, G. S., Duveiller, G., Elkins, J. W., Fioletov,  
448 V. E., Flemming, J., Foster, M. J., Frey, R. A., Frith, S. M., Froidevaux, L., Garforth, J., Gupta, S. K., Haimberger, L., Hall,  
449 B. D., Harris, I., Heidinger, A. K., Hemming, D. L., Ho, S.-p., Hubert, D., Hurst, D. F., Hüser, I., Inness, A., Isaksen, K.,  
450 John, V., Jones, P. D., Kaiser, J. W., Kelly, S., Khaykin, S., Kidd, R., Kim, H., Kipling, Z., Kraemer, B. M., Kratz, D. P.,  
451 La Fuente, R. S., Lan, X., Lantz, K. O., Leblanc, T., Li, B., Loeb, N. G., Long, C. S., Loyola, D., Marszelewski, W.,  
452 Martens, B., May, L., Mayer, M., McCabe, M. F., McVicar, T. R., Mears, C. A., Menzel, W. P., Merchant, C. J., Miller,  
453 B. R., Miralles, D. G., Montzka, S. A., Morice, C., Mühle, J., Myneni, R., Nicolas, J. P., Noetzli, J., Osborn, T. J., Park, T.,  
454 Pasik, A., Paterson, A. M., Pelto, M. S., Perkins-Kirkpatrick, S., Pétron, G., Phillips, C., Pinty, B., Po-Chedley, S., Polvani,  
455 L., Preimesberger, W., Pulkkanen, M., Randel, W. J., Rémy, S., Ricciardulli, L., Richardson, A. D., Rieger, L., Robinson,  
456 D. A., Rodell, M., Rosenlof, K. H., Roth, C., Rozanov, A., Rusak, J. A., Rusanovskaya, O., Rutishäuser, T., Sánchez-Lugo,  
457 A., Sawaengphokhai, P., Scanlon, T., Schenzinger, V., Schladow, S. G., Schlegel, R. W., Schmid, M. E., Selkirk, H. B.,  
458 Sharma, S., Shi, L., Shimaraeva, S. V., Silow, E. A., Simmons, A. J., Smith, C. A., Smith, S. L., Soden, B. J., Sofieva, V.,  
459 Sparks, T. H., Stackhouse, P. W., Steinbrecht, W., Streletskiy, D. A., Taha, G., Telg, H., Thackeray, S. J., Timofeyev, M.  
460 A., Tourpali, K., Tye, M. R., van der A, R. J., van der Schalie, R. V. B. V., van der Schrier, W. Paul, G., van der Werf, G.  
461 R., Verburg, P., Vernier, J.-P., Vömel, H., Vose, R. S., Wang, R., Watanabe, S. G., Weber, M., Weyhenmeyer, G. A.,  
462 Wiese, D., Wilber, A. C., Wild, J. D., Wong, T., Woolway, R. I., Yin, X., Zhao, L., Zhao, G., Zhou, X., Ziemke, J. R., and  
463 Ziese, M.: Global climate-state of the climate in 2019, *Bull. Amer. Meteor. Soc.*, 0003-0007 1520-0477, S9-S128, 2020.

464 Efimova, M. V., Kolomeichuk, L. V., Boyko, E. V., Malofii, M. K., Vidershpan, A. N., Plyusnin, I. N., Golovatskaya, I.  
465 F., Murgan, O. K., and Kuznetsov, V. V.: Physiological mechanisms of solanum tuberosum l. Plants' tolerance to chloride  
466 salinity, *Russ. J. Plant Physiol.*, 65, 394-403, <https://doi.org/10.1134/S1021443718030020>, 2018.

467 ESA: Sentinel-2 user handbook, [https://sentinel.esa.int/documents/247904/685211/sentinel-2\\_user\\_handbook](https://sentinel.esa.int/documents/247904/685211/sentinel-2_user_handbook), 2015.

468 Fang, H., Baret, F., Plummer, S., and Schaepman-Strub, G.: An overview of global leaf area index (lai): Methods, products,  
469 validation, and applications, *Rev. Geophys.*, 57, 739-799, <https://doi.org/10.1029/2018RG000608>, 2019.

470 FAO, I., UNICEF, WFP and WHO: The state of food security and nutrition in the world 2020, FAO, Rome978-92-5-  
471 132901-6, 2020.

472 Fatima, A., Hussain, S., Hussain, S., Ali, B., Ashraf, U., Zulfikar, U., Aslam, Z., Al-Robai, S. A., Alzahrani, F. O., Hano,  
473 C., and El-Esawi, M. A.: Differential morphophysiological, biochemical, and molecular responses of maize hybrids to  
474 salinity and alkalinity stresses, *Agronomy-Basel*, 11, 1150, <https://doi.org/10.3390/agronomy11061150>, 2021.

475 Gerhards, M., Schlerf, M., Mallick, K., and Udelhoven, T.: Challenges and future perspectives of multi-/hyperspectral  
476 thermal infrared remote sensing for crop water-stress detection: A review, *Remote Sens.*, 11, 1240-1264,  
477 <https://doi.org/10.3390/rs11101240>, 2019.

478 Ghimire, B., Timsina, D., and Nepal, J.: Analysis of chlorophyll content and its correlation with yield attributing traits on  
479 early varieties of maize (zea mays l.), *J. Maize Res. Dev.*, 1, 134-145, <https://doi.org/10.3126/jmrd.v1i1.14251>, 2015.

480 Gitelson, A. A., Vina, A., Ciganda, V., Rundquist, D. C., and Arkebauer, T. J.: Remote estimation of canopy chlorophyll  
481 content in crops, *Geophys. Res. Lett.*, 32, L08403, <https://doi.org/10.1029/2005GL022688>, 2005.

482 Godfray, H. C., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S.  
483 M., and Toulmin, C.: Food security: The challenge of feeding 9 billion people, *Science*, 327, 812-818,  
484 <https://doi.org/10.1126/science.1185383>, 2010.

485 Harfi, M. E., Hanine, H., Rizki, H., Latrache, H., and Nabloussi, A.: Effect of drought and salt stresses on germination and  
486 early seedling growth of different color-seeds of sesame (*sesamum indicum*), *Int. J. Agric. Biol.*, 18, 1088-1094,  
487 <https://doi.org/10.17957/ijab/15.0145>, 2016.

488 Homolova, L., Maenovsky, Z., Clevers, J. G. P. W., Garcia-Santos, G., and Schaepnran, M. E.: Review of optical-based  
489 remote sensing for plant trait mapping, *Ecol. Complex.*, 15, 1-16, <https://doi.org/10.1016/j.ecocom.2013.06.003>, 2013.

490 Hu, Q., Yang, J., Xu, B., Huang, J., Memon, M. S., Yin, G., Zeng, Y., Zhao, J., and Liu, K.: Evaluation of global decametric-  
491 resolution lai, fapar and fvc estimates derived from sentinel-2 imagery, *Remote Sens.*, 12,  
492 <https://doi.org/10.3390/rs12060912>, 2020.

493 Huang, J., Wang, H., Dai, Q., and Han, D.: Analysis of ndvi data for crop identification and yield estimation, *IEEE J. Sel.*  
494 *Top. Appl. Earth Obs. Remote Sens.*, 7, 4374-4384, <https://doi.org/10.1109/JSTARS.2014.2334332>, 2014.

495 Hussain, T., Koyro, H. W., Zhang, W., Liu, X., Gul, B., and Liu, X.: Low salinity improves photosynthetic performance  
496 in panicum antidotale under drought stress, *Front. Plant Sci.*, 11, 481, <https://doi.org/10.3389/fpls.2020.00481>, 2020.

497 Jarlan, L., Balsamo, G., Lafont, S., Beljaars, A., Calvet, J. C., and Mougou, E.: Analysis of leaf area index in the ecmwf  
498 land surface model and impact on latent heat and carbon fluxes: Application to west africa, *J. Geophys. Res. Atmos.*, 113,  
499 D24117, <https://doi.org/10.1029/2007jd009370>, 2008.

500 Jefferies, R.: Physiology of crop response to drought, in: *Potato ecology and modelling of crops under conditions limiting*  
501 *growth*, Springer, 61-74, 1995.

502 Jones, E., and van Vliet, M. T. H.: Drought impacts on river salinity in the southern us: Implications for water scarcity, *Sci.*  
503 *Total Environ.*, 644, 844-853, <https://doi.org/10.1016/j.scitotenv.2018.06.373>, 2018.

504 Kriston-Vizi, J., Umeda, M., and Miyamoto, K.: Assessment of the water status of mandarin and peach canopies using  
505 visible multispectral imagery, *Biosyst. Eng.*, 100, 338-345, <https://doi.org/10.1016/j.biosystemseng.2008.04.001>, 2008.

506 Liang, S. L. W., J. D.: Chapter 11 - fraction of absorbed photosynthetically active radiation, in: *Advanced remote sensing*  
507 *(second edition)*, edited by: Liang, S., and Wang, J., Academic Press, 447-476, 2020.

508 Liao, Q., Gu, S. J., Kang, S. Z., Du, T. S., Tong, L., Wood, J. D., and Ding, R. S.: Mild water and salt stress improve water  
509 use efficiency by decreasing stomatal conductance via osmotic adjustment in field maize, *Sci. Total Environ.*, 805,  
510 <https://doi.org/10.1016/j.scitotenv.2021.150364>, 2022.

511 López-Lozano, R., Duveiller, G., Seguini, L., Meroni, M., García-Condado, S., Hooker, J., Leo, O., and Baruth, B.:  
512 Towards regional grain yield forecasting with 1km-resolution eo biophysical products: Strengths and limitations at pan-  
513 european level, *Agric. For. Meteorol.*, 206, 12-32, <https://doi.org/10.1016/j.agrformet.2015.02.021>, 2015.

514 Mahmood, U., Hussain, S., Hussain, S., Ali, B., Ashraf, U., Zamir, S., Al-Robai, S. A., Alzahrani, F. O., Hano, C., and El-  
515 Esawi, M. A.: Morpho-physio-biochemical and molecular responses of maize hybrids to salinity and waterlogging during  
516 stress and recovery phase, *Plants (Basel)*, 10, <https://doi.org/10.3390/plants10071345>, 2021.

517 Masante D., B. P., McCormick N.: Drought in central-northern europe – august 2018, Report of the Copernicus European  
518 Drought Observatory (EDO) and Emergency Response Coordination Center (ERCC) Analytical Team 1-13, 2018.

519 McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales, *Proceedings*  
520 *of the 8th Conference on Applied Climatology*, 1993, 179-183,

521 Mi, N., Cai, F., Zhang, Y. S., Ji, R. P., Zhang, S. J., and Wang, Y.: Differential responses of maize yield to drought at  
522 vegetative and reproductive stages, *Plant Soil Environ.*, 64, 260-267, <https://doi.org/10.17221/141/2018-Pse>, 2018.

523 Mkhabela, M. S., Bullock, P., Raj, S., Wang, S., and Yang, Y.: Crop yield forecasting on the canadian prairies using modis  
524 ndvi data, *Agric. For. Meteorol.*, 151, 385-393, <https://doi.org/10.1016/j.agrformet.2010.11.012>, 2011.

525 Mulder, M., Hack-ten Broeke, M., Bartholomeus, R., van Dam, J., Heinen, M., van Bakel, J., Walvoort, D., Kroes, J.,  
526 Hoving, I., and Holshof, G.: *Waterwijzer landbouw: Instrumentarium voor kwantificeren van effecten van waterbeheer en*  
527 *klimaat op landbouwproductie*, 2018-48, Stowa, 2018.

528 Ors, S., and Suarez, D. L.: Spinach biomass yield and physiological response to interactive salinity and water stress, *Agric.*  
529 *Water Manag.*, 190, 31-41, <https://doi.org/10.1016/j.agwat.2017.05.003>, 2017.

530 Patane, C., Saita, A., and Sortino, O.: Comparative effects of salt and water stress on seed germination and early embryo  
531 growth in two cultivars of sweet sorghum, *J. Agron. Crop Sci.*, 199, 30-37, <https://doi.org/10.1111/j.1439-037X.2012.00531.x>, 2013.

532 Perry de Louw, V. K., Harry Massop, Ab Veldhuizen Beregening: Deltafact, Alterra - Soil, water and land use, Amersfoort  
533 2020.

534 Rozema, J., and Flowers, T.: Ecology. Crops for a salinized world, *Science*, 322, 1478-1480,  
535 <https://doi.org/10.1126/science.1168572>, 2008.

536 Sayar, R., Behini, H., Mosbahi, M., and Khemira, H.: Response of durum wheat (*triticum durum* desf.) growth to salt and  
537 drought stresses, *Czech J. Genet. Plant. Breed.*, 46, 54-63, <https://doi.org/10.17221/85/2009-CJGPB>, 2010.

538 Schittenhelm, S., Sourell, H., and Lopmeier, F. J.: Drought resistance of potato cultivars with contrasting canopy  
539 architecture, *Eur. J. Agron.*, 24, 193-202, <https://doi.org/10.1016/j.eja.2005.05.004>, 2006.

540 Schwalm, C. R., Anderegg, W. R. L., Michalak, A. M., Fisher, J. B., Biondi, F., Koch, G., Litvak, M., Ogle, K., Shaw, J.

542 D., Wolf, A., Huntzinger, D. N., Schaefer, K., Cook, R., Wei, Y., Fang, Y., Hayes, D., Huang, M., Jain, A., and Tian, H.:  
543 Global patterns of drought recovery, *Nature*, 548, 202-205, <https://doi.org/10.1038/nature23021>, 2017.  
544 Shinozaki, K., Uemura, M., Bailey-Serres, J., Bray, E., and Weretilnyk, E.: Responses to abiotic stress. *Biochemistry and*  
545 *molecular biology of plants*, Wiley Blackwell, 1051-1100 pp., 2015.  
546 Steidle Neto, A. J., Lopes, D. d. C., Silva, T. G. F. d., Ferreira, S. O., and Grossi, J. A. S.: Estimation of leaf water content  
547 in sunflower under drought conditions by means of spectral reflectance, *Eng. Agric. Environ. Food*, 10, 104-108,  
548 <https://doi.org/10.1016/j.eaef.2016.11.006>, 2017.  
549 Stuyt, L. C. P. M., Blom-Zandstra, M., & Kselik, R. A. L.: Inventarisatie en analyse zouttolerantie van landbouwgewassen  
550 op basis van bestaande gegevens, Wageningen environmental research rapport, Wageningen Environmental Research,  
551 2016.  
552 Sun, L., Gao, F., Anderson, M. C., Kustas, W. P., Alsina, M. M., Sanchez, L., Sams, B., McKee, L., Dulaney, W., White,  
553 W. A., Alfieri, J. G., Prueger, J. H., Melton, F., and Post, K.: Daily mapping of 30 m lai and ndvi for grape yield prediction  
554 in california vineyards, *Remote Sens.*, 9, <https://doi.org/10.3390/rs9040317>, 2017.  
555 Tao, H., Borth, H., Fraedrich, K., Su, B., and Zhu, X.: Drought and wetness variability in the tarim river basin and  
556 connection to large-scale atmospheric circulation, *Int. J. Climatol.*, 34, 2678-2684, <https://doi.org/10.1002/joc.3867>, 2014.  
557 Tokarz, B., Wójtowicz, T., Makowski, W., Jędrzejczyk, R. J., and Tokarz, K. M.: What is the difference between the  
558 response of grass pea (*lathyrus sativus* L.) to salinity and drought stress?—a physiological study, *Agronomy*, 10, 833,  
559 <https://doi.org/10.3390/agronomy10060833>, 2020.  
560 Trenberth, K. E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., and Sheffield, J.: Global warming  
561 and changes in drought, *Nat. Clim. Chang.*, 4, 17-22, <https://doi.org/10.1038/nclimate2067>, 2013.  
562 Tucker, C. J.: Red and photographic infrared linear combinations for monitoring vegetation, *Remote Sens. Environ.*, 8,  
563 127-150, [https://doi.org/10.1016/0034-4257\(79\)90013-0](https://doi.org/10.1016/0034-4257(79)90013-0), 1979.  
564 van Straten, G., Bruning, B., de Vos, A. C., González, A. P., Rozema, J., and van Bodegom, P. M.: Estimating cultivar-  
565 specific salt tolerance model parameters from multi-annual field tests for identification of salt tolerant potato cultivars,  
566 *Agric. Water Manag.*, 252, <https://doi.org/10.1016/j.agwat.2021.106902>, 2021.  
567 Vannoppen, A., Gobin, A., Kotova, L., Top, S., De Cruz, L., Viksna, A., Aniskevich, S., Bobylev, L., Buntmeyer, L.,  
568 Caluwaerts, S., De Troch, R., Gnatiuk, N., Hamdi, R., Reca Remedio, A., Sakalli, A., Van De Vyver, H., Van Schaeuybroeck,  
569 B., and Termonia, P.: Wheat yield estimation from ndvi and regional climate models in latvia, *Remote Sens.*, 12,  
570 <https://doi.org/10.3390/rs12142206>, 2020.  
571 Vereecken, H., Weihermuller, L., Jonard, F., and Montzka, C.: Characterization of crop canopies and water stress related  
572 phenomena using microwave remote sensing methods: A review, *Vadose Zone J.*, 11, vzj2011.0138ra,  
573 <https://doi.org/10.2136/vzj2011.0138ra>, 2012.  
574 Wagg, C., Hann, S., Kupriyanovich, Y., and Li, S.: Timing of short period water stress determines potato plant growth,  
575 yield and tuber quality, *Agric. Water Manag.*, 247, <https://doi.org/10.1016/j.agwat.2020.106731>, 2021.  
576 Wang, J. L., Huang, X. J., Zhong, T. Y., and Chen, Z. G.: Climate change impacts and adaptation for saline agriculture in  
577 north jianguo province, china, *Environ. Sci. Policy*, 25, 83-93, <https://doi.org/10.1016/j.envsci.2012.07.011>, 2013.  
578 Weiss, M., and Baret, F.: S2toolbox level 2 products: Lai, fapar, fcover, version 1.1, ESA Contract nr 4000110612/14/I-  
579 BG, 52, 2016.  
580 Weiss, M., Jacob, F., and Duveiller, G.: Remote sensing for agricultural applications: A meta-review, *Remote Sens.*  
581 *Environ.*, 236, 111402, <https://doi.org/10.1016/j.rse.2019.111402>, 2020.  
582 Wen, W., Timmermans, J., Chen, Q., and van Bodegom, P. M.: A review of remote sensing challenges for food security  
583 with respect to salinity and drought threats, *Remote Sens.*, 13, <https://doi.org/10.3390/rs13010006>, 2020.  
584 Wengert, M., Piepho, H. P., Astor, T., Grass, R., Wijesingha, J., and Wachendorf, M.: Assessing spatial variability of  
585 barley whole crop biomass yield and leaf area index in silvoarable agroforestry systems using uav-borne remote sensing,  
586 *Remote Sens.*, 13, 2751, <https://doi.org/10.3390/rs13142751>, 2021.  
587 Wright, I. J., Reich, P. B., and Westoby, M.: Least-cost input mixtures of water and nitrogen for photosynthesis, *Am. Nat.*,  
588 161, 98-111, <https://doi.org/10.1086/344920>, 2003.  
589 Yang, L., Jia, K., Liang, S., Liu, M., Wei, X., Yao, Y., Zhang, X., and Liu, D.: Spatio-temporal analysis and uncertainty of  
590 fractional vegetation cover change over northern china during 2001–2012 based on multiple vegetation data sets, *Remote*  
591 *Sens.*, 10, <https://doi.org/10.3390/rs10040549>, 2018.  
592 Zhang, F., Zhou, G. S., and Nilsson, C.: Remote estimation of the fraction of absorbed photosynthetically active radiation  
593 for a maize canopy in northeast china, *J. Plant Ecol.*, 8, 429-435, <https://doi.org/10.1093/jpe/rtu027>, 2015.  
594 Zhang, H., Han, M., Comas, L. H., DeJonge, K. C., Gleason, S. M., Trout, T. J., and Ma, L.: Response of maize yield  
595 components to growth stage - based deficit irrigation, *Agron. J.*, 111, 3244-3252,  
596 <https://doi.org/10.2134/agronj2019.03.0214>, 2019.  
597 Zhang, L., Chen, B., Zhang, G., Li, J., Wang, Y., Meng, Y., and Zhou, Z.: Effect of soil salinity, soil drought, and their  
598 combined action on the biochemical characteristics of cotton roots, *Acta Physiol. Plant*, 35, 3167-3179,  
599 <https://doi.org/10.1007/s11738-013-1350-6>, 2013.  
600 Zhu, X., Wang, T. J., Skidmore, A. K., Darvishzadeh, R., Niemann, K. O., and Liu, J.: Canopy leaf water content estimated  
601 using terrestrial lidar, *Agric. For. Meteorol.*, 232, 152-162, <https://doi.org/10.1016/j.agrformet.2016.08.016>, 2017.

