

1 Monitoring the combined effects of drought and salinity stress on 2 crops using remote sensing in the Netherlands

3 Wen Wen^{1,*}, Joris Timmermans^{1,2,3}, Qi Chen¹ and Peter M. van Bodegom¹

4¹Institute of Environmental Sciences (CML), Leiden University, Box 9518, 2300 RA Leiden, The Netherlands

5²Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, 1090 GE Amsterdam, The Netherlands

6³Lifewatch ERIC, vLab & Innovation Centre, 1090 GE Amsterdam, The Netherlands

7 *Correspondence:* Wen Wen (w.wen@cml.leidenuniv.nl)

8 **Abstract.** Global sustainable agricultural systems are under threat, due to increasing and co-occurring drought and salinity
9 stresses. Combined effects of these stresses on agricultural crops have traditionally been evaluated in small-scale
10 experimental studies. Consequently, large-scale studies need to be performed to increase our understanding and assessment
11 of the combined impacts in agricultural practice in real-life scenarios. This study aims to provide a new monitoring
12 approach using remote sensing observations to evaluate the joint impacts of drought and salinity on crop traits. In our tests
13 over the Netherlands at large spatial (138.74 km²), we calculated five functional traits for both maize and potato from
14 Sentinel-2 observations, namely: leaf area index (LAI), the fraction of absorbed photosynthetically active radiation
15 (FAPAR), the fraction of vegetation cover (FVC), leaf chlorophyll content (Cab) and leaf water content (Cw). Individual
16 and combined effects of the stresses on the seasonal dynamics in crop traits were determined using both one-way and two-
17 way ANOVAs. We found that both stresses (individual and co-occurring) affected the functional traits of both crops
18 significantly (with R² ranging from 0.326 to 0.796), though with stronger sensitivities to drought than to salinity. While we
19 found exacerbating effects within co-occurring stresses, the impact-level depended strongly on the moment in the growing
20 season. For both crops, LAI, FAPAR and FVC dropped the most under severe drought stress conditions. The patterns for
21 Cab and Cw were more inhibited by co-occurring drought and salinity. Consequently, our study constitutes a way towards
22 evaluating drought and salinity impacts in agriculture with the possibility of potential large-scale application for sustainable
23 food security.

24 **Keywords:** Drought; Salinity; Agriculture; Remote sensing; Functional traits

25 1 Introduction

26 Food production is required to increase by 70% to satisfy the growing population demand by the year 2050 (Godfray et al.,
27 2010). Meanwhile, food security is becoming increasingly threatened due to the increasing abiotic stresses under the
28 influence of global climate change; abiotic stresses including drought, soil salinity, nutrient stress and heavy metals are
29 estimated to constrain crop productivity by 50% ~ 80% (Shinozaki et al., 2015). Of these stresses, drought and salinity
30 stress have been identified as the two main factors to limit crop growth, affecting respectively 40% and 11% of the global
31 irrigated areas (Dunn et al., 2020; FAO, 2020). With drought and salinity forecasted to increase spatially and in severity
32 (Rozema and Flowers, 2008; Schwalm et al., 2017; Trenberth et al., 2013), and with predictions of higher co-occurrence
33 around the world (Corwin, 2020; Jones and van Vliet, 2018; Wang et al., 2013), food production will be more deeply
34 challenged by both stresses.

35 Numerous small-scale experimental studies for a large variety of crops have shown that the impact of co-occurring drought
36 and salinity stress is exacerbated. Co-occurrence of drought and salinity stress is found to decrease the yield of spinach
37 (Ors and Suarez, 2017) and the forage grass *Panicum antidotale* (Hussain et al., 2020) more compared with the occurrence


38 of one of these stresses only. Likewise, cotton root growth tends to be more inhibited under the co-occurrence of drought
39 and salinity than by isolated occurrences (Zhang et al., 2013). Similarly, the exacerbating effect of co-occurring stresses
40 limits both maize reproductive growth and grain formation (Liao et al., 2022). While these studies demonstrate the
41 exacerbating effects of co-occurring drought and salinity stress, they have limitations in projecting the impact towards real
42 farmers' conditions due to their small-scale experimental nature. Thus, there is still a significant knowledge gap concerning
43 the large scale evaluation of the combined impacts of drought and salinity.

44 Remote sensing (RS) provides a huge potential to close this knowledge gap due to its capability to monitor continuous
45 large areas at a frequent interval. For this, remote sensing has traditionally used vegetation indices, such as Normalized
46 Difference Vegetation Index (NDVI) (Tucker, 1979). However, such indices provide limited information on how the
47 impact is achieved (e.g. Wen et al., 2020) and how it can be mitigated. With the launch of better multispectral and high-
48 resolution satellite sensors (such as Sentinel-2), new RS methods (e.g., hyperspectral, thermal infrared, microwave) have
49 been identified to detect stress in both natural vegetation (Gerhards et al., 2019; Vereecken et al., 2012) as well as for
50 agricultural applications (Homolova et al., 2013; Weiss et al., 2020). Specifically, these new RS methods allow for the
51 retrieval of plant traits that directly link to plant processes, such as leaf biochemistry and photosynthetic processes, and
52 thereby provide high potential for agricultural applications. RS plant traits of specific interest to monitor crop health include
53 leaf area index (LAI) (Wengert et al., 2021), canopy chlorophyll content (Cab*LAI) (Gitelson et al., 2005), canopy water
54 content (Cw*LAI) (Kriston-Vizi et al., 2008), the fraction of absorbed photosynthetically active radiation (FAPAR) (Zhang
55 et al., 2015) and the fraction of vegetation cover (FVC) (Yang et al., 2018). Canopy chlorophyll content and mean leaf
56 equivalent water thickness (EWT) of maize differed remarkably under drought stress using hyperspectral remote sensing
57 data (Zhang and Zhou, 2015). Using a look-up-table approach, LAI and chlorophyll content of wheat obtained from a
58 radiative transfer model showed potential to assess drought levels (Richter et al., 2008). However, while there have been
59 several attempts to monitor the response of crop health with either a drought or salinity focus, not much research has taken
60 these factors into account simultaneously (Wen et al., 2020).

61 In this study, we propose a novel approach to estimate, compare and evaluate the impacts of drought, salinity, and their
62 combination on crop traits using remote sensing. To allow for a detailed evaluation of this approach we applied it to analyze
63 the impacts of the 2018 summer drought in the Netherlands on agricultural crops. In this, a stress co-occurrence map was
64 created by overlaying a high-resolution drought map of 2018 with a groundwater salinity map. Then, we characterized the
65 response of maize and potato to different stress conditions based on five plant traits (LAI, FAPAR, FVC, Cab, and Cw).
66 Two-way ANOVAs were adopted to test the main effects and the interactive effect between stress combinations and time
67 on crop traits. Moreover, the effect of drought and salinity on crop traits was determined across the growing season with
68 one-way ANOVAs. Consequently, this approach facilitates simultaneously monitoring crop health at various scales
69 (regional, national, continental) across multiple stresses (drought, salinity) and multiple species.

70 **2 Methodology**

71 To achieve our aim of monitoring the impacts of (co-occurring) drought and salinity on agricultural production, we
72 developed a new approach to estimate crop traits from remote sensing observations. Specifically, we developed an approach
73 that integrates image-processing techniques, such as image classification, co-registration, land surface parameter retrieval,
74 and time-series analysis. Using these techniques, we were able to estimate the drought, salinity, and crop growth.



76

77 **Figure 1.** Technical workflow of the maps and data framework.

78 To allow for a detailed evaluation, we focused on the 2018 summer drought in the Netherlands. This period was selected
 79 because of the extreme drought that affected a large part of Europe (Masante D., 2018). Within parts of the selected area
 80 salinity was reported to increase during that same period (Broekhuizen, 2018). Hence this study area provides us with the
 81 opportunity to investigate the combined impacts of these stresses on crops. In the following paragraphs, we provide more
 82 information on the specific processing steps.

83 **2.1 Study area and data**84 **2.1.1 Drought map**

85 A drought map of the Netherlands in 2018 was created based on the standardized precipitation evapotranspiration index
 86 (SPEI) drought index, which was calculated from long-term precipitation data and potential evapotranspiration, from 2004
 87 to 2018 (Chen et al., 2022). Specifically, SPEI was estimated using a 3-month sliding time window, as this was found best
 88 to investigate the impacts on the local ecosystems. We have extracted SPEI-3 data from April 1st to October 30th, totally
 89 214 days, as this coincided with the crop growth period of both maize and potato. Then, the drought map was resampled
 90 to 250m resolution using the nearest neighbor interpolation and reprojected to RD_new projection. The RD_new projection
 91 (EPSG:28992) is a projected coordinate reference system of the Netherlands. All maps were projected to RD_new
 92 projection to create consistent data layers. We defined -1 and -1.5 as daily thresholds for different drought severity classes
 93 according to previous classifications (McKee et al., 1993; Tao et al., 2014). Thus, (cumulative) SPEI for no drought should
 94 be between -214 to 0, SPEI for moderate drought should be between -321 to -214 and for severe drought, SPEI should be
 95 lower than -321 when calculated for the whole growing period (Fig. 2a).

96 **2.1.1 Salinity map**


97 A top-soil salinity map of the Netherlands was created based on a nationwide fresh-salt groundwater dataset, which derived
98 chloride concentrations as a salinity indicator (<https://data.nhi.nu/>). To obtain the salinity map of the top-soil, 15 layers of
99 the groundwater salinity were extracted from the 3D groundwater salinity map. For each location, the layer closest to the
100 corresponding to location's elevation (according to the Digital Elevation Model), i.e. closest to the soil surface, was
101 selected. The salinity map was resampled to 250 m resolution and reprojected to RD_new projection. Ultimately, the
102 salinity map was classified into three levels namely no-salinity ($0.1 \text{ g}\cdot\text{L}^{-1}$ to $0.8 \text{ g}\cdot\text{L}^{-1}$), moderate salinity ($0.8 \text{ g}\cdot\text{L}^{-1}$ to $2.5 \text{ g}\cdot\text{L}^{-1}$),
103 severe salinity ($\geq 2.5 \text{ g}\cdot\text{L}^{-1}$) according to the salt-resistant capacity of various crops cultivated in the Netherlands
104 (Mulder et al., 2018; Stuyt, 2016) (Fig. 2b).

105 **2.1.3 Crop map**

106 The crop map of the Netherlands in 2018 was collected from the Key Register of Parcels (BRP) of the Netherlands
107 Enterprise Agency (<https://www.pdok.nl/introductie/-/article/basisregistratie-gewaspercelen-brp->). The crop map was
108 resampled to 250m resolution and reprojected to RD_new projection (Fig. 2d).

109 **2.1.4 Co-occurrence map of drought and salinity**

110 The drought map and the salinity map were overlain to evaluate co-occurrences of drought and salinity of the Netherlands
111 in 2018 (Fig. 2c). By classifying the three stress levels for the individual occurrences, we obtained nine stress classes of
112 co-occurring drought and salinity, namely no stress, moderate drought only (MD), severe drought only (SD), moderate
113 salinity only (MS), severe salinity only (SS), moderate drought and moderate salinity (MD+MS), moderate drought and
114 severe salinity (MD+SS), severe drought and moderate salinity (SD+MS), and severe drought and severe salinity (SD+SS).



115

116

117 **Figure 2.** Map of the Netherlands overlaying a) drought and b) salinity to show c) the co-occurrence of drought and salinity in 2018.
 118 The selected study area is indicated by black lines in panel c. d) The associated crop map of the study area in 2018.

119 **2.1.5 Study area selection**

120 Based on the national map of the Netherlands (Fig. 2c), a single region with similar soil type, climate, tillage systems, and
 121 irrigation methods was chosen to minimize the interference of these factors on the observed trait expressions. The province
 122 of North-Holland was selected because it contained the most (7 out of 9) combinations of drought and salt stress (Fig. 2c),
 123 namely: no stress, MD, SD, MS, SS, MD+MS, and SD+SS. Moreover, both maize and potato were cultivated across all
 124 stress combinations in this province. For further analysis, MS and SS were grouped into a new class of salinity stress since
 125 the area of MS and SS was quite limited. Therefore, six classes of stress combinations namely no stress, MD, SD, salinity
 126 (MS+SS), MD+MS, and MD+SS were analyzed for the study area.

127 **2.2 Traits retrieval**

128 **2.2.1 Satellite data**

129 The Sentinel-2 mission consists of two satellites equipped with the high-resolution Multispectral Instrument (MSI) in the
130 same orbit. This sensor acquires 13 spectral bands (with varying spatial resolutions) in the visible and near-infrared
131 spectrum at 5 days of revisit times (ESA, 2015). In our study, we used both the 10m and 20m Level 2A observations,
132 downloaded from The Copernicus Open Access Hub (<https://scihub.copernicus.eu/>), to facilitate the requirement of the
133 Sentinel Application Platform (SNAP) toolbox for both optical and near-infrared observations to be available for
134 determining the functional traits. To create consistency across the bands, those with a 20m resolution (B5, B6, B7, B8A,
135 B11, and B12) were resampled to the 10m resolution of B3 and B4. In total, eight cloud-free scenes were found
136 (21/04/2018, 06/05/2018, 26/05/2018, 30/06/2018, 15/07/2018, 13/09/2018, 13/10/2018, and 28/10/2018) to cover the crop
137 growth cycle. Although additional cloud-free scenes were found in August (04/08/2018, 09/08/2018, 14/08/2018,
138 19/08/2018, 24/08/2018, and 29/08/2018), none were of high quality, and we therefore choose to omit August from our
139 analysis.

140 **2.2.2 Traits selection**

141 Plant traits (e.g. LAI, FAPAR, FVC, Cab and Cw) were selected in consideration of their corresponding impacts on crop
142 functioning and their potential for assessment by remote sensing. LAI is a critical vegetation structural trait related to
143 various plant functioning processes such as primary productivity, photosynthesis, and transpiration (Asner et al., 2003;
144 Boussetta et al., 2012; Fang et al., 2019; Jarlan et al., 2008). FAPAR depends on vegetation structure, energy exchange,
145 and illumination conditions while FAPAR is also an important parameter to assess primary productivity (Liang, 2020;
146 Weiss and Baret, 2016). FVC is a promising parameter corresponding to the energy balance process such as temperature
147 and evapotranspiration (Weiss and Baret, 2016). Cab is an effective indicator of stress and is strongly related to
148 photosynthesis and resource strategy (Croft et al., 2017). Cw plays an important role in transpiration, stomatal conductance,
149 photosynthesis, and respiration (Bowman, 1989; Zhu et al., 2017), as well as in drought assessment (Steidle Neto et al.,
150 2017).

151 **2.3 Dataset processing**

152 The biophysical processor within the SNAP toolbox derives the five traits, namely LAI, FAPAR, FVC, canopy chlorophyll
153 content (CCC), and canopy water content (CWC), for each pixel from the Sentinel-2 top of canopy reflectance data at a
154 10m-resolution for each month. This processor utilizes an artificial neural network (ANN) approach, trained using the
155 PROSAIL simulated database (Weiss and Baret, 2016). This training utilized canopy traits rather than leaf traits (estimated
156 by multiplication with LAI) to improve their neural network performance. To obtain their leaf counterparts (Cw and Cab),
157 to create fully independent variables, CCC and CWC thus need to be divided by LAI to obtain Cab (=CCC / LAI) and Cw
158 (=CWC / LAI). Pixels with quality flags were eliminated from the dataset. It was observed that in April no crop had yet
159 been planted. Instead, we observed that only along the edge of the plots, e.g. in ditches, vegetation was found. This feature
160 was used to generate a ditch map and to mask out pixels in trait maps for the other months. For each variable and each
161 date, only data within the 95% confidence interval were taken to increase data robustness.

162 **2.4 Analysis**

163 Since the pixel counts of the six classes of stress combinations namely no stress, MD, SD, salinity, MD+MS, and MD+SS
164 were (highly) different, drought and salinity were not considered as two independent factors. Instead, two-way analysis of
165 variance (ANOVA) was applied to test the main effects and the interactive effect between stress combinations (consisting
166 of 6 levels) and time (5 months) on each individual crop trait. Significant effects of the main stress condition were

167 investigated through post hoc tests to test whether interaction effects between drought and salinity had occurred. Two-way
 168 ANOVAs were run separately for each trait and each crop type (maize and potato) as we expected different patterns. In the
 169 Netherlands, potato and maize are planted between mid-April to early May. Crops are surfacing in May and harvested in
 170 October. Therefore, to evaluate the response of crops to stresses across the growing season, the effect of drought and
 171 salinity on crop traits was determined for May, June, July, and September with a one-way ANOVA. Tukey HSD post hoc
 172 tests were performed to identify the differences among the six stress combinations. All statistical analyses were performed
 173 with SPSS 27.0 (SPSS Inc., USA).

174 **3 Results**

175 **3.1 Stress impacts depend on the moment in the growing season**

176 The two-way ANOVAs revealed strong effects of date and stress level on the five traits with effect sizes of the response
 177 (R^2) ranging from 0.326 to 0.796 for the five traits, which was similar for maize and potato. For both maize and potato, R^2
 178 values were lowest for Cab and highest for LAI, FAPAR, and FVC. For maize, we found a significant main effect of both
 179 date and stress ($p < 0.05$) for Cab, Cw, FAPAR, and FVC. In contrast, LAI was not significantly different across the
 180 different stress conditions. For potato, all main effects of date and stress were significant for all five crop traits (Table 1).
 181 For all traits and both crops, the interaction between the effects of time and stress conditions was significant ($p < 0.05$)
 182 (Table 1), indicating that the impact of stress depended on the moment in the growing season. Despite the significant
 183 interaction terms, the partial Eta squared values (Table 1) showed that the effects of time in the growing season were much
 184 stronger than those of stress or the interaction of date and stress. The effects of date for maize were stronger than for potato.
 185 Interestingly, the effects of the interaction between date and stress were stronger than those of the main effects of stress,
 186 suggesting strongly time-specific impacts of stress on the crop traits investigated. The interaction terms were strongest for
 187 FVC.

188 **Table 1.** Two-way ANOVA for different crop traits by time series and stress interactions.

Crops	Traits	Factors	F	p	Partial Eta Squared	R^2
Maize	LAI	date	2144.5	0.000	0.636	0.766
		stress	1.4	0.226	0.001	
		date*stress	8.5	0.000	0.033	
	Cab	date	333.9	0.000	0.222	
		stress	10.7	0.000	0.008	
		date*stress	3.6	0.000	0.015	
Potato	Cw	date	952.1	0.000	0.449	0.590
		stress	9.9	0.000	0.007	
		date*stress	4.0	0.000	0.017	
	FAPAR	date	1865.9	0.005	0.603	
		stress	3.3	0.000	0.002	
		date*stress	8.5	0.000	0.033	
	FVC	date	2022.5	0.000	0.622	
		stress	22.1	0.000	0.015	
		date*stress	28.7	0.000	0.105	
	LAI	date	752.1	0.000	0.273	0.782
		stress	13.7	0.000	0.006	
		date*stress	8.1	0.000	0.020	
	Cab	date	96.4	0.000	0.050	0.329

	stress	54.2	0.000	0.024
	date*stress	8.7	0.000	0.023
Cw	date	347.4	0.000	0.158
	stress	68.1	0.000	0.030
	date*stress	10.3	0.000	0.027
FAPAR	date	612.7	0.000	0.234
	stress	25.8	0.000	0.011
	date*stress	14.0	0.000	0.034
FVC	date	844.0	0.000	0.297
	stress	18.8	0.000	0.008
	date*stress	13.6	0.000	0.033

189 Note: F indicates the test statistic of the F -test; p indicates whether the effect is statistically significant in comparison to the significance
 190 level ($p < 0.05$); Partial Eta Squared indicates the effect size of different factors; R^2 indicates the percentage that the model coincides
 191 with the data.


192 3.2 Response of LAI, FAPAR, FVC to drought and salinity

193 Given the significance of both date and stress and their interactions, subsequent one-way ANOVAs were performed to
 194 compare the effects of drought and salinity on LAI, FAPAR, and FVC for maize and potato in May, June, July, and
 195 September separately (Fig. 3). The patterns for LAI, FAPAR, and FVC were very similar, although they differ in details
 196 and were therefore treated together.

197 For maize, all of LAI, FAPAR, and FVC obtained their lowest value under MD+SS stress conditions in May. In June, both
 198 LAI and FVC dropped the most under salinity stress and it was significantly ($p < 0.05$) different from MD, MD+MS, and
 199 MD+SS conditions, but not significantly different from no stress conditions. In contrast, FAPAR also reached its lowest
 200 value (under MD+MS stress conditions) in June but had a significant difference ($p < 0.05$) compared with no stress
 201 conditions. Both in July and September, LAI, FAPAR, and FVC all had the lowest value under SD conditions, and the
 202 difference was significant compared with no stress conditions.

203 For potato, LAI, FAPAR, and FVC had the lowest ($p < 0.05$) value under MD+MS and MD+SS stress conditions in May.
 204 In June, LAI, FAPAR as well as FVC reached the lowest value under SD conditions and were significantly lower than in
 205 most other stress conditions even though the difference was not significant from no stress conditions. In July, there was a
 206 tendency for LAI, FAPAR, and FVC to be lower under stress conditions, although none of the effects were significant. In
 207 September, however, LAI, FAPAR, and FVC significantly decreased under MD, MD+MS, and MD+SS conditions, and
 208 the difference was significant compared with no stress conditions. In addition, the difference was not significant among
 209 these three stress conditions.

210 Therefore, both for maize and potato, LAI, FAPAR, and FVC dropped the most under SD stress conditions when they
 211 reached their respective maximum value, compared with other stress conditions. At the same time, maize and potato were
 212 more sensitive to drought than salinity since no significant change was observed between drought conditions and conditions
 213 with a combination of drought and salinity stress.




215 **Figure 3.** Expressions of LAI, FAPAR, and FVC under various stress conditions in May, June, July, and September 2018. Different
 216 letters in each panel indicate significant differences ($p < 0.05$). MD, moderate drought only; Salinity, salinity only; MD+MS, moderate
 217 drought and moderate salinity; MD+SS, moderate drought and severe salinity (MD+SS); SD, severe drought only.

218 3.3 Response of leaf chlorophyll and water content to drought and salinity

219 The one-way ANOVAs revealed that there were significant ($p < 0.05$) impacts of the various stress conditions on Cab and
 220 Cw (Fig. 4). For maize, Cab obtained its lowest value under salinity stress in May and June while it was not significantly
 221 different from no stress conditions. However, in July, Cab reached the lowest value under MD+MS conditions although
 222 the difference was not significant from other stress conditions. There were no significant changes observed for Cab in
 223 September. For potato, Cab dropped the most under salinity conditions in May although the difference was not significant
 224 from no stress conditions. Furthermore, Cab significantly decreased under MD+SS conditions in June and July, compared
 225 with other conditions. Although Cab dropped the most under salinity conditions in September, the difference was not
 226 significantly different from other conditions. In addition, compared with no stress, potato had the lowest Cab under MD+SS
 227 conditions while there was no significant difference between MD+SS and salinity conditions in most growing periods.
 228 Cw decreased under all stress conditions in May, June, and July for both maize and potato, except for SD conditions in
 229 May, compared with no stress conditions. At the same time, Cw reached its lowest value under MD+SS conditions and it
 230 was significantly different from under no stress conditions. Nonetheless, there were different changes for maize and potato
 231 in September. Cw was not significantly different among any conditions for maize while it was the lowest under salinity
 232 conditions for potato.

233 Therefore, this analysis illustrates that salinity affected maize less than drought since crop responses were more obvious to
 234 drought than salinity for Cw. In contrast, salinity showed a more severe effect on maize and potato at the early growth
 235 stages for Cab. Meanwhile, Cab was affected by co-occurring drought and salinity in June and July for potato. It seems
 236 that there was a non-additive effect of drought and salinity for Cw since the changes were not significant between MD+MS,
 237 MD+SS, MD, and salinity conditions.



238 **Figure 4.** Expressions of Cab and Cw under various stress conditions in May, June, July, and September 2018. Different letters in each
 239 panel indicate significant differences ($p < 0.05$). MD, moderate drought only; Salinity, salinity only; MD+MS, moderate
 240 drought and moderate salinity; MD+SS, moderate drought and severe salinity (MD+SS); SD, severe drought only.

242 **4 Discussion**

243 In this study, we quantified the large-scale impacts of co-occurring drought and salinity on a variety of crop traits using
244 satellite remote sensing. We observed that –in contrast to our expectations – the impacts of salinity were not highly
245 pronounced at this scale, with most strong impacts originating due to drought stress during the 2018 drought. At specific
246 moments in the growing season, salinity and/or the combined effects of salinity and drought pronouncedly affected
247 individual crop traits. In this way, with increasing salinity driven by more intensive droughts, water allocation should not
248 only be governed by the amount of water shortage, but also the salinity of the remaining water. In this paper, we provide
249 the first evidence that those impacts can be monitored through remote sensing. This might provide a basis towards a
250 monitoring system for multiple crops with multiple stresses as well as better governance policies to release this problem.

251 **4.1 Drought stress is more important than salinity stress in farmers' conditions**

252 The exacerbating effects of co-occurrent drought and salinity (Fig. 3 and Fig. 4) that we found are consistent with findings
253 of small-scale experiments (e.g. greenhouses). Consistent with our results, synergistic effects of co-occurring water stress
254 and salinity stress have been found on maize reproductive growth and grain formation in a field study (Liao et al., 2022).
255 Spinach (*Spinacia oleracea* L., cv. Racoon) yield decreased more under co-occurring water-salinity stress in comparison
256 with separate water stress and salinity (Ors and Suarez, 2017). The co-occurring drought and salinity stress was more
257 harmful to cotton root growth compared to their individual effects (Zhang et al., 2013). Moreover, the combined negative
258 effect of drought and salinity stress on *Panicum antidotale* was stronger than that of single stress (Hussain et al., 2020).
259 Our research showed that the outcomes of these small-scale experimental studies also apply to real large-scale
260 environments, where different sources of variance are present. Specifically, we show that in real farmers' conditions, the
261 co-occurrence of drought and salinity indeed can constitute a severe threat due to its interactive effects on crop growth.
262 In addition, we evaluated whether drought or salinity stress has more impact on crop performance. We observed that maize
263 and potato were generally more sensitive to drought than salinity in this study (Fig. 3 and Fig. 4). This is consistent with
264 results of previous studies that highlight that drought impacts are generally more detrimental than salinity stress for crops,
265 e.g. for sesame (*Sesamum indicum*) (Harfi et al., 2016), *Mentha pulegium* L. (Azad et al., 2021), durum wheat (Sayar et
266 al., 2010), grass pea (Tokarz et al., 2020), and sweet sorghum (Patane et al., 2013). However, given that the threshold of
267 salinity at which crop damage occurs (according to the FAO guidelines (Ayers and Westcot, 1985)) was surpassed in all
268 situations in which salinity stress was imposed (including in our study), we initially expected salinity to be a stronger
269 explanatory variable than drought. As such, salinity impacts on crop performance (by the FAO) may have been
270 overestimated. Indeed, in an experimental field situation in which drought stress was carefully avoided, higher thresholds
271 of salinity-induced damage were observed for potato (van Straten et al., 2021).
272 In combination, the results from our study (supported by results from other studies) suggest that salinity particularly induces
273 adverse effects when co-occurring with drought stress. Water stress impacts on photosynthesis and biomass of plants were
274 extenuated by salinity since salinity enhances the synthesis of ATP and NADPH by promoting photosynthetic pigments
275 and photosystem II efficiency. The impacts of combined drought and salinity stress on plant growth, chlorophyll content,
276 water use efficiency, and photosynthesis were less severe compared to drought alone. This indicates compensating effects
277 on carbon assimilation due to osmotic adjustments induced by Na^+ and Cl^- (Hussain et al., 2020). Thus, the detrimental
278 effect of single drought stress on crop growth is considered to be mitigated by salinity.

279 **4.2 Drought and salinity stress differ between growth stages**

280 The responses to drought and salinity stress were different at different growth stages of the crops. This was expressed by
281 the significant interactions between the effects of time and stress conditions for all of our crop responses (Table 1). We
282 found that during the grain filling (maize) and tuber bulking phase (potato), the sensitivities of these crops are expressed
283 distinctly in the non-harvested aboveground tissues (Fig. 3 and Fig.4), with clear differences in the remote sensing plant
284 traits.

285 Given that we were not able to monitor the harvestable products, multiple mechanisms may explain these patterns. The
286 relatively high leaf coverage (as related to LAI, FAPAR, and FVC) at salinity and severe drought conditions at the end of
287 the growing season may be an expression of a compensation process. Specifically, early and prolonged drought could have
288 led to more assimilates allocated to non-harvestable potato parts for drought resistance since the number of tubers reduced
289 (Jefferies, 1995; Schittenhelm et al., 2006). In that case, we should consider their higher leaf coverage at the end of the
290 season as a survival mechanism, rather than true drought tolerance, leading to reduced tuber yields (Daryanto et al., 2016b).

291 Future studies that combine remote sensing with harvesting data may be able to evaluate this mechanism in more detail.
292 In our study, different response patterns of maize and potato occurred to the different stresses over the growing season.
293 This is consistent with previous studies focusing on the impact of drought and/or salinity onsets. For potato, it has been
294 suggested that tuber yields particularly decreased when drought stress occurs during the vegetative and tuber initiation
295 stages than during the tuber bulking stage (Wagg et al., 2021), although another study observed the reverse pattern
296 (Daryanto et al., 2016b). For maize, on the other hand, drought seems to have the most detrimental impact during the
297 maturation stage (Mi et al., 2018; Zhang et al., 2019), and the reproductive phase (Daryanto et al., 2016a; Daryanto et al.,
298 2017). Considering the additional co-varying factors within our ‘real-life’ study, it is very probable that we were able to
299 detect similar effects. This suggests that we may use satellite remote sensing –albeit less spatially precise than e.g. sensing
300 through drones- as a cost-effective early warning signal for detecting drought and salinity stress at moments during the
301 growing season when differences in crop performance are still subtle.

302 **4.3 Crop responses to stress can be better understood with a multi-trait approach**

303 In addition to facilitating the evaluation of crop performance during multiple stages of the growing season (in contrast to
304 most destructive methods), remote sensing also allows a multi-trait approach to better understand the mechanisms involved
305 in crop responses. Each of the five traits is associated with different functions of plants that might be individually impacted
306 by the different stresses. Therefore, focusing on only one individual metric (as commonly done; see Wen et al. (2020) for
307 a review) limits our capacity to gain full insight into drought and salinity responses. Hence, given that individual crop traits
308 may respond differently to drought and salinity reflecting its stress resistance and tolerance strategy, the evaluation of these
309 distinct responses may help to understand this strategy.

310 In this study, Cw was consistently lower in all drought and salinity treatments as compared to no stress conditions in May,
311 June, and July. Indeed, this is a common response of plants in response to drought and salinity (e.g. Wen et al., 2020). In
312 this respect, it is interesting that no decrease in Cw was observed at the end of the growing season, in September. Whether
313 the phenomenon is related to the survival mechanism mentioned above or to the lower transpiration demands at the end of
314 the season because of lower aboveground biomass, cannot be concluded from these data. Some evidence pointing to the
315 survival mechanism is the finding (Ghosh et al., 2001; Levy, 1992) that the leaf dry matter increased for potato under
316 drought/salinity stress (like in our study) while the dry matter of the tubers appeared to have a greater decline.

317 With respect to chlorophyll contents, we observed a decline in Cab under salinity conditions in May and the MS+SS
318 treatment in June and July, while no decrease was observed in any of the treatments exposed to drought only. This indicates
319 that while total leaf area was not (much) affected by salinity, the salinity did negatively affect crop performance. It has
320 been reported that chlorophyll content in maize was significantly reduced upon salinity, along with other plant traits
321 including plant height, shoot/root biomass, and leaf numbers (Fatima et al., 2021; Mahmood et al., 2021). Likewise, similar
322 patterns were obtained in potato plants in saline soil (Efimova et al., 2018). Hence, this implies that soil salinity tends to
323 negatively affect crop growth and restrict nutrient uptake.

324 Cab and Cw responses to drought and salinity were distinct from responses of LAI, FAPAR, and FVC (Fig. 3 and Fig. 4).
325 LAI, FAPAR, and FVC showed similar patterns to stress due to their highly physical correlation (Hu et al., 2020). The
326 different patterns of Cw and Cab point to different drought and salinity resistance strategy components associated with
327 these traits: LAI (and FAPAR/FVC) reflect the decrease in biomass due to stress, partly because stress directly and
328 negatively impacts growth and partly because having lower biomass decreases the evapotranspiration demands of the crop,
329 which increases the resilience of the crop to deal with drought. Cw represents another pathway to reduce evapotranspiration
330 demands, i.e. by reducing the amount of water per gram of leaves. Also, this response may be a direct effect of the more
331 negative pressure heads due to drought or due to increased osmotic pressures (due to salinity). It may also be part of the
332 adaptive strategy of the crop to increase its resilience. Cab also responds to drought and salinity, but in its own way, i.e. by
333 adapting its photosynthetic capacity while being affected by a lower stomatal conductance (due to drought and/or salinity).
334 See e.g. Wright et al. (2003) for a framework explaining these nitrogen-water interactions.

335 In addition, our approach gives the insight to analyze the effect of stresses on yield based on the five traits, even though
336 yield cannot directly be derived from remote sensing. Traits including Cab, LAI, and FAPAR, have been used (either
337 separately or in combination) as a proxy for final yield estimates from remote sensing in many studies. For instance, NDVI
338 -which is based on the combination of LAI and Cab- is extensively used to estimate crop yield (Huang et al., 2014;
339 Mkhabela et al., 2011; Vannoppen et al., 2020). Also, LAI itself has been used for predicting the final yield (Dente et al.,
340 2008; Doraiswamy et al., 2005; Sun et al., 2017). Meanwhile, Cab and FAPAR were also proven to be highly correlated
341 with crop yield (Ghimire et al., 2015; López-Lozano et al., 2015). Thus, while yield cannot be estimated directly from
342 remote sensing or ground truth data at the desired high spatial resolution, our indicators do relate to yield and can be used
343 in more application-based contexts to inform on yield impacts.

344 **4.4 Implications for future research and management.**

345 The number of studies that evaluate the effects of drought and salinity stress on crops is limited (Wen et al., 2020). In
346 general, studies focus on small-scale experimental studies under strictly control of all variables with only a limited number
347 of crops (Hussain et al., 2020; Ors and Suarez, 2017). To our knowledge, this is the first study that uses satellite remote
348 sensing to investigate drought and salinity impacts for a large area under real-life conditions necessary for constructing
349 stress management policies.

350 In such real-life conditions, as investigated here, irrigation of crops is commonly applied as management practice during
351 drought events to reduce the severity of drought impacts (Deb et al., 2022; Lu et al., 2020). In this study, however, we have
352 evidence that irrigation did not play a major role in the patterns found since all croplands included in our research area
353 were identified as rainfed cropland (according to the ESA/CCI land cover map in 2018;
354 <https://maps.elie.ucl.ac.be/CCI/viewer/>). In addition, while farmers in the area are known to irrigate their cropland, the
355 Dutch government announced a temporary national irrigation ban in 2018 (for various areas including our research area)

356 to spare water (Perry de Louw, 2020). As a consequence, we could not analyze the impacts of irrigation management on
357 the combined effects of drought and salinity. This might potentially be solved by investigating other drought historic events
358 with moderate severity in Europe, such as the year of 2003 (Ciais et al., 2005) or 2015 (Ionita et al., 2017) in Europe, when
359 such a ban was not executed. Unfortunately, satellite remote sensing observations with the required 20-30m resolutions of
360 these events are limited, as Sentinel-2 was only launched in 2015 and the Landsat satellites provide a too coarse temporal
361 resolution.

362 Likewise, impacts of salinity and drought are moderated by crop selection. Traditionally, farmers do not plant highly
363 vulnerable crops in moderate/high salinity areas. In fact, we found crops sensitive to salinity such as apple (Ivanov, 1970)
364 and broccoli (Bernstein and Ayers, 1949) to be abundant in non-saline areas but only little in saline areas. To ensure an
365 accurate evaluation of salinity impacts, we only investigated those crops with a significant abundance in all available stress
366 conditions. More sensitive crops might even respond more strongly.

367 5 Conclusions

368 In this study, we present the first attempt to evaluate the real-life effects of drought, salinity, and their combination on crop
369 health using multiple traits from remote sensing monitoring during 2018 over the Netherlands. Our approach gives new
370 insights for monitoring crop growth under co-occurring stresses at a large scale with high-resolution data. We found that
371 while in general temporal patterns –reflecting crop growth dynamics- were stronger than effects of stress conditions, stress
372 impacts depended on the time of the growing season. Furthermore, we also found that the temporal dynamics in crop
373 responses to drought and salinity were different for maize vs. potato. In general, the five investigated traits were more
374 negatively affected by a combination of drought and salinity stress compared to individual stress. Meanwhile, both maize
375 and potato responded more prominently to drought, thus demonstrating a stronger sensitivity, than to salinity. Specifically,
376 LAI, FAPAR, and FVC dropped the most under severe drought stress conditions. Consequently, the proposed new
377 approach poses a facilitated way for simultaneously monitoring the effect of drought and salinity on crops in large-scale
378 agricultural applications.

379
380 *Data availability.* The drought map of the Netherlands in 2018 is retrieved from Chen et al. (2022). The top-soil salinity
381 map of the Netherlands is retrieved from The Netherlands Hydrological Instrumentarium (NHI) (<https://data.nhi.nu/>). The
382 crop map of the Netherlands in 2018 is retrieved from the Key Register of Parcels (BRP) of the Netherlands Enterprise
383 Agency (<https://www.pdok.nl/introductie/-/article/basisregistratie-gewaspercelen-brp->). All satellite scenes are
384 downloaded from The Copernicus Open Access Hub (<https://scihub.copernicus.eu/>). The dataset relevant to this study is
385 available upon request from the corresponding author.

386
387 *Author contributions.* Conceptualization, JT, PVB, and WW; methodology, JT, QC, WW, and PVB.; investigation, WW
388 and QC; writing—original draft preparation, WW; writing—review and editing, PVB. and JT; supervision, PVB, and JT
389 All authors have read and agreed to the published version of the manuscript.

390
391 *Competing interests.* The authors declare no conflict of interest.

392
393 *Financial support.* This work was supported by the China Scholarship Council (CSC).

References

395 Asner, G. P., Scurlock, J. M. O., and Hicke, J. A.: Global synthesis of leaf area index observations: Implications for
 396 ecological and remote sensing studies, *Glob. Ecol. Biogeogr.*, 12, 191-205, <https://doi.org/10.1046/j.1466-822X.2003.00026.x>, 2003.

398 Ayers, R. S., and Westcot, D. W.: Water quality for agriculture, Food and Agriculture Organization of the United Nations
 399 Rome, 1985.

400 Azad, N., Rezayian, M., Hassanpour, H., Niknam, V., and Ebrahimzadeh, H.: Physiological mechanism of salicylic acid
 401 in mentha pulegium l. Under salinity and drought stress, *Braz. J. Bot.*, 44, 359-369, <https://doi.org/10.1007/s40415-021-00706-y>, 2021.

403 Bernstein, L., and Ayers, A.: Salt tolerance of cabbage and broccoli, United States Salinity Laboratory Report to
 404 Collaborators, Riverside, CA, 39, 1949.

405 Boussetta, S., Balsamo, G., Beljaars, A., Kral, T., and Jarlan, L.: Impact of a satellite-derived leaf area index monthly
 406 climatology in a global numerical weather prediction model, *Int. J. Remote Sens.*, 34, 3520-3542,
 407 <https://doi.org/10.1080/01431161.2012.716543>, 2012.

408 Bowman, W. D.: The relationship between leaf water status, gas-exchange, and spectral reflectance in cotton leaves,
 409 *Remote Sens. Environ.*, 30, 249-255, [https://doi.org/10.1016/0034-4257\(89\)90066-7](https://doi.org/10.1016/0034-4257(89)90066-7), 1989.

410 'Storm duurt dagen, droogte duurt maanden': <https://www.rijkswaterstaat.nl/nieuws/archief/2018/08/storm-duurt-dagen-droogte-duurt-maanden>, 2018.

412 Chen, Q., Timmermans, J., Wen, W., and van Bodegom, P. M.: A multi-metric assessment of drought vulnerability across
 413 different vegetation types using high-resolution remote sensing, *Sci. Total Environ.*, 154970,
 414 <https://doi.org/10.1016/j.scitotenv.2022.154970>, 2022.

415 Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara,
 416 A., Chevallier, F., De Noblet, N., Friend, A. D., Friedlingstein, P., Grünwald, T., Heinesch, B., Kersten, P., Knohl, A.,
 417 Krinner, G., Loustau, D., Manca, G., Matteucci, G., Miglietta, F., Ourcival, J. M., Papale, D., Pilegaard, K., Rambal, S.,
 418 Seufert, G., Soussana, J. F., Sanz, M. J., Schulze, E. D., Vesala, T., and Valentini, R.: Europe-wide reduction in primary
 419 productivity caused by the heat and drought in 2003, *Nature*, 437, 529-533, <https://doi.org/10.1038/nature03972>, 2005.

420 Corwin, D. L.: Climate change impacts on soil salinity in agricultural areas, *Eur. J. Soil Sci.*, 72, 842-862,
 421 <https://doi.org/10.1111/ejss.13010>, 2020.

422 Croft, H., Chen, J. M., Luo, X., Bartlett, P., Chen, B., and Staebler, R. M.: Leaf chlorophyll content as a proxy for leaf
 423 photosynthetic capacity, *Glob. Change Biol.*, 23, 3513-3524, <https://doi.org/10.1111/gcb.13599>, 2017.

424 Daryanto, S., Wang, L., and Jacinthe, P. A.: Global synthesis of drought effects on maize and wheat production, *PLoS One*,
 425 11, e0156362, <https://doi.org/10.1371/journal.pone.0156362>, 2016a.

426 Daryanto, S., Wang, L. X., and Jacinthe, P. A.: Drought effects on root and tuber production: A meta-analysis, *Agric.
 427 Water Manag.*, 176, 122-131, <https://doi.org/10.1016/j.agwat.2016.05.019>, 2016b.

428 Daryanto, S., Wang, L. X., and Jacinthe, P. A.: Global synthesis of drought effects on cereal, legume, tuber and root crops
 429 production: A review, *Agric. Water Manag.*, 179, 18-33, <https://doi.org/10.1016/j.agwat.2016.04.022>, 2017.

430 Deb, P., Moradkhani, H., Han, X., Abbaszadeh, P., and Xu, L.: Assessing irrigation mitigating drought impacts on crop
 431 yields with an integrated modeling framework, *J. Hydrol.*, 609, 127760, <https://doi.org/10.1016/j.jhydrol.2022.127760>,
 432 2022.

433 Dente, L., Satalino, G., Mattia, F., and Rinaldi, M.: Assimilation of leaf area index derived from asar and meris data into
 434 ceres-wheat model to map wheat yield, *Remote Sens. Environ.*, 112, 1395-1407, <https://doi.org/10.1016/j.rse.2007.05.023>,
 435 2008.

436 Doraiswamy, P. C., Sinclair, T. R., Hollinger, S., Akhmedov, B., Stern, A., and Prueger, J.: Application of modis derived
 437 parameters for regional crop yield assessment, *Remote Sens. Environ.*, 97, 192-202,
 438 <https://doi.org/10.1016/j.rse.2005.03.015>, 2005.

439 Dunn, R. J. H., Stanitski, D. M., Gobron, N., Willett, K. M., Ades, M., Adler, R., Allan, R., Allan, R. P., Anderson, J.,
 440 Argüez, A., Arosio, C., Augustine, J. A., Azorin-Molina, C., Barichivich, J., Barnes, J., Beck, H. E., Becker, A., Bellouin,
 441 N., Benedetti, A., Berry, D. I., Blenkinsop, S., Bock, O., Bosilovich, M. G., Boucher, O., Buehler, S. A., Carrea, L.,
 442 Christiansen, H. H., Chouza, F., Christy, J. R., Chung, E. S., Coldewey-Egbers, M., Compo, G. P., Cooper, O. R., Covey,
 443 C., Crotwell, A., Davis, S. M., de Eyto, E., de Jeu, R. A. M., VanderSat, B. V., DeGasperi, C. L., Degenstein, D., Di
 444 Girolamo, L., Dokulil, M. T., Donat, M. G., Dorigo, W. A., Durre, I., Dutton, G. S., Duveiller, G., Elkins, J. W., Fioletov,
 445 V. E., Flemming, J., Foster, M. J., Frey, R. A., Frith, S. M., Froidevaux, L., Garforth, J., Gupta, S. K., Haimberger, L., Hall,
 446 B. D., Harris, I., Heidinger, A. K., Hemming, D. L., Ho, S.-p., Hubert, D., Hurst, D. F., Hüser, I., Inness, A., Isaksen, K.,
 447 John, V., Jones, P. D., Kaiser, J. W., Kelly, S., Khaykin, S., Kidd, R., Kim, H., Kipling, Z., Kraemer, B. M., Kratz, D. P.,
 448 La Fuente, R. S., Lan, X., Lantz, K. O., Leblanc, T., Li, B., Loeb, N. G., Long, C. S., Loyola, D., Marszelewski, W.,
 449 Martens, B., May, L., Mayer, M., McCabe, M. F., McVicar, T. R., Mears, C. A., Menzel, W. P., Merchant, C. J., Miller,
 450 B. R., Miralles, D. G., Montzka, S. A., Morice, C., Mühle, J., Myneni, R., Nicolas, J. P., Noetzli, J., Osborn, T. J., Park, T.,
 451 Pasik, A., Paterson, A. M., Pelto, M. S., Perkins-Kirkpatrick, S., Pétron, G., Phillips, C., Pinty, B., Po-Chedley, S., Polvani,
 452 L., Preimesberger, W., Pulkkanen, M., Randel, W. J., Rémy, S., Ricciardulli, L., Richardson, A. D., Rieger, L., Robinson,

453 D. A., Rodell, M., Rosenlof, K. H., Roth, C., Rozanov, A., Rusak, J. A., Rusanovskaya, O., Rutishauser, T., Sánchez-Lugo,
 454 A., Sawaengphokhai, P., Scanlon, T., Schenzinger, V., Schladow, S. G., Schlegel, R. W., Schmid, M. E., Selkirk, H. B.,
 455 Sharma, S., Shi, L., Shimaraeva, S. V., Silow, E. A., Simmons, A. J., Smith, C. A., Smith, S. L., Soden, B. J., Sofieva, V.,
 456 Sparks, T. H., Stackhouse, P. W., Steinbrecht, W., Streletsckiy, D. A., Taha, G., Telg, H., Thackeray, S. J., Timofeyev, M.
 457 A., Tourpali, K., Tye, M. R., van der A, R. J., van der Schalie, R. V. B. V., van der Schrier, W. Paul, G., van der Werf, G.
 458 R., Verburg, P., Vernier, J.-P., Vömel, H., Vose, R. S., Wang, R., Watanabe, S. G., Weber, M., Weyhenmeyer, G. A.,
 459 Wiese, D., Wilber, A. C., Wild, J. D., Wong, T., Woolway, R. I., Yin, X., Zhao, L., Zhao, G., Zhou, X., Ziemke, J. R., and
 460 Ziese, M.: Global climate-state of the climate in 2019, *Bull. Amer. Meteor.* 0003-0007 1520-0477, S9-S128, 2020.

461 Efimova, M. V., Kolomeichuk, L. V., Boyko, E. V., Malofii, M. K., Vidershpan, A. N., Plyusnin, I. N., Golovatskaya, I.
 462 F., Murgan, O. K., and Kuznetsov, V. V.: Physiological mechanisms of *solanum tuberosum* l. Plants' tolerance to chloride
 463 salinity, *Russ. J. Plant Physiol.*, 65, 394-403, <https://doi.org/10.1134/S1021443718030020>, 2018.

464 ESA: Sentinel-2 user handbook, https://sentinel.esa.int/documents/247904/685211/sentinel-2_user_handbook, 2015.

465 Fang, H., Baret, F., Plummer, S., and Schaepman-Strub, G.: An overview of global leaf area index (lai): Methods, products,
 466 validation, and applications, *Rev. Geophys.*, 57, 739-799, <https://doi.org/10.1029/2018RG000608>, 2019.

467 FAO, I., UNICEF, WFP and WHO: The state of food security and nutrition in the world 2020, FAO, Rome978-92-5-
 468 132901-6, 2020.

469 Fatima, A., Hussain, S., Hussain, S., Ali, B., Ashraf, U., Zulfiqar, U., Aslam, Z., Al-Robai, S. A., Alzahrani, F. O., Hano,
 470 C., and El-Esawi, M. A.: Differential morphophysiological, biochemical, and molecular responses of maize hybrids to
 471 salinity and alkalinity stresses, *Agronomy*, 11, 1150, <https://doi.org/10.3390/agronomy11061150>, 2021.

472 Gerhards, M., Schlerf, M., Mallick, K., and Udelhoven, T.: Challenges and future perspectives of multi-/hyperspectral
 473 thermal infrared remote sensing for crop water-stress detection: A review, *Remote Sens.*, 11, 1240-1264,
 474 <https://doi.org/10.3390/rs11101240>, 2019.

475 Ghimire, B., Timinsina, D., and Nepal, J.: Analysis of chlorophyll content and its correlation with yield attributing traits on
 476 early varieties of maize (*zea mays* l.), *J. Maize Res. Dev.*, 1, 134-145, <https://doi.org/10.3126/jmrd.v1i1.14251>, 2015.

477 Gitelson, A. A., Vina, A., Ciganda, V., Rundquist, D. C., and Arkebauer, T. J.: Remote estimation of canopy chlorophyll
 478 content in crops, *Geophys. Res. Lett.*, 32, L08403, <https://doi.org/10.1029/2005GL022688>, 2005.

479 Godfray, H. C., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S.
 480 M., and Toulmin, C.: Food security: The challenge of feeding 9 billion people, *Science*, 327, 812-818,
 481 <https://doi.org/10.1126/science.1185383>, 2010.

482 Harfi, M. E., Hanine, H., Rizki, H., Latrache, H., and Naboussi, A.: Effect of drought and salt stresses on germination and
 483 early seedling growth of different color-seeds of sesame (*sesamum indicum*), *Int. J. Agric. Biol.*, 18, 1088-1094,
 484 <https://doi.org/10.17957/ijab/15.0145>, 2016.

485 Homolova, L., Maenovsky, Z., Clevers, J. G. P. W., Garcia-Santos, G., and Schaeprnan, M. E.: Review of optical-based
 486 remote sensing for plant trait mapping, *Ecol. Complex.*, 15, 1-16, <https://doi.org/10.1016/j.ecocom.2013.06.003>, 2013.

487 Hu, Q., Yang, J., Xu, B., Huang, J., Memon, M. S., Yin, G., Zeng, Y., Zhao, J., and Liu, K.: Evaluation of global decametric-
 488 resolution lai, fapar and fvc estimates derived from sentinel-2 imagery, *Remote Sens.*, 12,
 489 <https://doi.org/10.3390/rs12060912>, 2020.

490 Huang, J., Wang, H., Dai, Q., and Han, D.: Analysis of ndvi data for crop identification and yield estimation, *IEEE J. Sel.
 491 Top. Appl. Earth Obs. Remote Sens.*, 7, 4374-4384, <https://doi.org/10.1109/JSTARS.2014.2334332>, 2014.

492 Hussain, T., Koyro, H. W., Zhang, W., Liu, X., Gul, B., and Liu, X.: Low salinity improves photosynthetic performance
 493 in *panicum antidotale* under drought stress, *Front. Plant Sci.*, 11, 481, <https://doi.org/10.3389/fpls.2020.00481>, 2020.

494 Ionita, M., Tallaksen, L. M., Kingston, D. G., Stagge, J. H., Laaha, G., Van Lanen, H. A. J., Scholz, P., Chelcea, S. M.,
 495 and Haslinger, K.: The european 2015 drought from a climatological perspective, *Hydrol. Earth Syst. Sci.*, 21, 1397-1419,
 496 <https://doi.org/10.5194/hess-21-1397-2017>, 2017.

497 Ivanov, V.: Main principles of fruit crop salt resistance determination, *Pochvovedenie*, 4, 78-85, 1970.

498 Jarlan, L., Balsamo, G., Lafont, S., Beljaars, A., Calvet, J. C., and Mougin, E.: Analysis of leaf area index in the ecmwf
 499 land surface model and impact on latent heat and carbon fluxes: Application to west africa, *J. Geophys. Res. Atmos.*, 113,
 500 D24117, <https://doi.org/10.1029/2007jd009370>, 2008.

501 Jefferies, R.: Physiology of crop response to drought, in: Potato ecology and modelling of crops under conditions limiting
 502 growth, Springer, 61-74, 1995.

503 Jones, E., and van Vliet, M. T. H.: Drought impacts on river salinity in the southern us: Implications for water scarcity, *Sci.
 504 Total Environ.*, 644, 844-853, <https://doi.org/10.1016/j.scitotenv.2018.06.373>, 2018.

505 Kriston-Vizi, J., Umeda, M., and Miyamoto, K.: Assessment of the water status of mandarin and peach canopies using
 506 visible multispectral imagery, *Biosyst. Eng.*, 100, 338-345, <https://doi.org/10.1016/j.biosystemseng.2008.04.001>, 2008.

507 Liang, S. L. W., J. D.: Chapter 11 - fraction of absorbed photosynthetically active radiation, in: Advanced remote sensing
 508 (second edition), edited by: Liang, S., and Wang, J., Academic Press, 447-476, 2020.

509 Liao, Q., Gu, S. J., Kang, S. Z., Du, T. S., Tong, L., Wood, J. D., and Ding, R. S.: Mild water and salt stress improve water
 510 use efficiency by decreasing stomatal conductance via osmotic adjustment in field maize, *Sci. Total Environ.*, 805,
 511 <https://doi.org/10.1016/j.scitotenv.2021.150364>, 2022.

512 López-Lozano, R., Duveiller, G., Seguini, L., Meroni, M., García-Condado, S., Hooker, J., Leo, O., and Baruth, B.:

513 Towards regional grain yield forecasting with 1km-resolution eo biophysical products: Strengths and limitations at pan-
 514 european level, *Agric. For. Meteorol.*, 206, 12-32, <https://doi.org/10.1016/j.agrformet.2015.02.021>, 2015.

515 Lu, J., Carbone, G. J., Huang, X., Lackstrom, K., and Gao, P.: Mapping the sensitivity of agriculture to drought and
 516 estimating the effect of irrigation in the united states, 1950–2016, *Agric. For. Meteorol.*, 292-293, 108124,
 517 <https://doi.org/10.1016/j.agrformet.2020.108124>, 2020.

518 Mahmood, U., Hussain, S., Hussain, S., Ali, B., Ashraf, U., Zamir, S., Al-Robai, S. A., Alzahrani, F. O., Hano, C., and El-
 519 Esawi, M. A.: Morpho-physio-biochemical and molecular responses of maize hybrids to salinity and waterlogging during
 520 stress and recovery phase, *Plants (Basel)*, 10, <https://doi.org/10.3390/plants10071345>, 2021.

521 Masante D., B. P., McCormick N.: Drought in central-northern europe – august 2018, Report of the Copernicus European
 522 Drought Observatory (EDO) and Emergency Response Coordination Center (ERCC) Analytical Team 1-13, 2018.

523 McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales, *Proceedings*
 524 of the 8th Conference on Applied Climatology

525 1993, 179-183,

526 Mi, N., Cai, F., Zhang, Y. S., Ji, R. P., Zhang, S. J., and Wang, Y.: Differential responses of maize yield to drought at
 527 vegetative and reproductive stages, *Plant Soil Environ.*, 64, 260-267, <https://doi.org/10.17221/141/2018-Pse>, 2018.

527 Mkhabela, M. S., Bullock, P., Raj, S., Wang, S., and Yang, Y.: Crop yield forecasting on the canadian prairies using modis
 528 ndvi data, *Agric. For. Meteorol.*, 151, 385-393, <https://doi.org/10.1016/j.agrformet.2010.11.012>, 2011.

529 Mulder, M., Hack-ten Broeke, M., Bartholomeus, R., van Dam, J., Heinen, M., van Bakel, J., Walvoort, D., Kroes, J.,
 530 Hoving, I., and Holshof, G.: Waterwijzer landbouw: Instrumentarium voor kwantificeren van effecten van waterbeheer en
 531 klimaat op landbouwproductie, 2018-48, Stowa, 2018.

532 Ors, S., and Suarez, D. L.: Spinach biomass yield and physiological response to interactive salinity and water stress, *Agric.*
 533 *Water Manag.*, 190, 31-41, <https://doi.org/10.1016/j.agwat.2017.05.003>, 2017.

534 Patane, C., Saita, A., and Sortino, O.: Comparative effects of salt and water stress on seed germination and early embryo
 535 growth in two cultivars of sweet sorghum, *J. Agron. Crop Sci.*, 199, 30-37, <https://doi.org/10.1111/j.1439-037X.2012.00531.x>, 2013.

536 Perry de Louw, V. K., Harry Massop, Ab Veldhuizen Beregening: Deltafact, Alterra - Soil, water and land use, Amersfoort
 537 2020.

538 Richter, K., Rischbeck, P., Eitzinger, J., Schneider, W., Suppan, F., and Weihs, P.: Plant growth monitoring and potential
 539 drought risk assessment by means of earth observation data, *Int. J. Remote Sens.*, 29, 4943-4960,
 540 <https://doi.org/10.1080/01431160802036268>, 2008.

541 Rozema, J., and Flowers, T.: Ecology. Crops for a salinized world, *Science*, 322, 1478-1480,
 542 <https://doi.org/10.1126/science.1168572>, 2008.

543 Sayar, R., Bchini, H., Mosbahi, M., and Khemira, H.: Response of durum wheat (triticum durum desf.) growth to salt and
 544 drought stresses, *Czech J. Genet. Plant. Breed.*, 46, 54-63, <https://doi.org/10.17221/85/2009-CJGPB>, 2010.

545 Schittenhelm, S., Sourell, H., and Lopmeier, F. J.: Drought resistance of potato cultivars with contrasting canopy
 546 architecture, *Eur. J. Agron.*, 24, 193-202, <https://doi.org/10.1016/j.eja.2005.05.004>, 2006.

547 Schwalm, C. R., Anderegg, W. R. L., Michalak, A. M., Fisher, J. B., Biondi, F., Koch, G., Litvak, M., Ogle, K., Shaw, J.
 548 D., Wolf, A., Huntzinger, D. N., Schaefer, K., Cook, R., Wei, Y., Fang, Y., Hayes, D., Huang, M., Jain, A., and Tian, H.:
 549 Global patterns of drought recovery, *Nature*, 548, 202-205, <https://doi.org/10.1038/nature23021>, 2017.

550 Shinozaki, K., Uemura, M., Bailey-Serres, J., Bray, E., and Weretilnyk, E.: Responses to abiotic stress, *Biochemistry and*
 551 *molecular biology of plants*, Wiley Blackwell, 1051-1100 pp., 2015.

552 Steidle Neto, A. J., Lopes, D. d. C., Silva, T. G. F. d., Ferreira, S. O., and Grossi, J. A. S.: Estimation of leaf water content
 553 in sunflower under drought conditions by means of spectral reflectance, *Eng. Agric. Environ. Food*, 10, 104-108,
 554 <https://doi.org/10.1016/j.eaef.2016.11.006>, 2017.

555 Stuyt, L. C. P. M., Blom-Zandstra, M., & Kselik, R. A. L. : Inventarisatie en analyse zouttolerantie van landbouwgewassen
 556 op basis van bestaande gegevens, Wageningen environmental research rapport, Wageningen Environmental Research,
 557 2016.

558 Sun, L., Gao, F., Anderson, M. C., Kustas, W. P., Alsina, M. M., Sanchez, L., Sams, B., McKee, L., Dulaney, W., White,
 559 W. A., Alfieri, J. G., Prueger, J. H., Melton, F., and Post, K.: Daily mapping of 30 m lai and ndvi for grape yield prediction
 560 in california vineyards, *Remote Sens.*, 9, <https://doi.org/10.3390/rs9040317>, 2017.

561 Tao, H., Borth, H., Fraedrich, K., Su, B., and Zhu, X.: Drought and wetness variability in the tarim river basin and
 562 connection to large-scale atmospheric circulation, *Int. J. Climatol.*, 34, 2678-2684, <https://doi.org/10.1002/joc.3867>, 2014.

563 Tokarz, B., Wójtowicz, T., Makowski, W., Jędrzejczyk, R. J., and Tokarz, K. M.: What is the difference between the
 564 response of grass pea (*lathyrus sativus* l.) to salinity and drought stress?—a physiological study, *Agronomy*, 10, 833,
 565 <https://doi.org/10.3390/agronomy10060833>, 2020.

566 Trenberth, K. E., Dai, A., van der Schrier, G., Jones, P. D., Barichivich, J., Briffa, K. R., and Sheffield, J.: Global warming
 567 and changes in drought, *Nat. Clim. Chang.*, 4, 17-22, <https://doi.org/10.1038/nclimate2067>, 2013.

568 Tucker, C. J.: Red and photographic infrared linear combinations for monitoring vegetation, *Remote Sens. Environ.*, 8,
 569 127-150, [https://doi.org/10.1016/0034-4257\(79\)90013-0](https://doi.org/10.1016/0034-4257(79)90013-0), 1979.

570 van Straten, G., Bruning, B., de Vos, A. C., González, A. P., Rozema, J., and van Bodegom, P. M.: Estimating cultivar-
 571 specific salt tolerance model parameters from multi-annual field tests for identification of salt tolerant potato cultivars,
 572

573 Agric. Water Manag., 252, <https://doi.org/10.1016/j.agwat.2021.106902>, 2021.
574 Vannoppen, A., Gobin, A., Kotova, L., Top, S., De Cruz, L., Vīksna, A., Aniskevich, S., Bobylev, L., Bunteleyer, L.,
575 Caluwaerts, S., De Troch, R., Gnatiuk, N., Hamdi, R., Reca Remedio, A., Sakalli, A., Van De Vyver, H., Van Schaeybroeck,
576 B., and Termonia, P.: Wheat yield estimation from ndvi and regional climate models in latvia, Remote Sens., 12,
577 <https://doi.org/10.3390/rs12142206>, 2020.
578 Vereecken, H., Weihermuller, L., Jonard, F., and Montzka, C.: Characterization of crop canopies and water stress related
579 phenomena using microwave remote sensing methods: A review, Vadose Zone J., 11, vzb2011.0138ra,
580 <https://doi.org/10.2136/vzj2011.0138ra>, 2012.
581 Wagg, C., Hann, S., Kupriyanovich, Y., and Li, S.: Timing of short period water stress determines potato plant growth,
582 yield and tuber quality, Agric. Water Manag., 247, <https://doi.org/10.1016/j.agwat.2020.106731>, 2021.
583 Wang, J. L., Huang, X. J., Zhong, T. Y., and Chen, Z. G.: Climate change impacts and adaptation for saline agriculture in
584 north jiangsu province, china, Environ. Sci. Policy, 25, 83-93, <https://doi.org/10.1016/j.envsci.2012.07.011>, 2013.
585 Weiss, M., and Baret, F.: S2toolbox level 2 products: Lai, fapar, fcover, version 1.1, ESA Contract nr 4000110612/14/I-
586 BG, 52, 2016.
587 Weiss, M., Jacob, F., and Duveiller, G.: Remote sensing for agricultural applications: A meta-review, Remote Sens.
588 Environ., 236, 111402, <https://doi.org/10.1016/j.rse.2019.111402>, 2020.
589 Wen, W., Timmermans, J., Chen, Q., and van Bodegom, P. M.: A review of remote sensing challenges for food security
590 with respect to salinity and drought threats, Remote Sens., 13, <https://doi.org/10.3390/rs13010006>, 2020.
591 Wengert, M., Piepho, H. P., Astor, T., Grass, R., Wijesingha, J., and Wachendorf, M.: Assessing spatial variability of
592 barley whole crop biomass yield and leaf area index in silvoarable agroforestry systems using uav-borne remote sensing,
593 Remote Sens., 13, 2751, <https://doi.org/10.3390/rs13142751>, 2021.
594 Wright, I. J., Reich, P. B., and Westoby, M.: Least-cost input mixtures of water and nitrogen for photosynthesis, Am. Nat.,
595 161, 98-111, <https://doi.org/10.1086/344920>, 2003.
596 Yang, L., Jia, K., Liang, S., Liu, M., Wei, X., Yao, Y., Zhang, X., and Liu, D.: Spatio-temporal analysis and uncertainty of
597 fractional vegetation cover change over northern china during 2001–2012 based on multiple vegetation data sets, Remote
598 Sens., 10, <https://doi.org/10.3390/rs10040549>, 2018.
599 Zhang, F., and Zhou, G.: Estimation of canopy water content by means of hyperspectral indices based on drought stress
600 gradient experiments of maize in the north plain china, Remote Sens., 7, 15203-15223, <https://doi.org/10.3390/rs71115203>,
601 2015.
602 Zhang, F., Zhou, G. S., and Nilsson, C.: Remote estimation of the fraction of absorbed photosynthetically active radiation
603 for a maize canopy in northeast china, J. Plant Ecol., 8, 429-435, <https://doi.org/10.1093/jpe/rtu027>, 2015.
604 Zhang, H., Han, M., Comas, L. H., DeJonge, K. C., Gleason, S. M., Trout, T. J., and Ma, L.: Response of maize yield
605 components to growth stage - based deficit irrigation, Agron. J., 111, 3244-3252,
606 <https://doi.org/10.2134/agronj2019.03.0214>, 2019.
607 Zhang, L., Chen, B., Zhang, G., Li, J., Wang, Y., Meng, Y., and Zhou, Z.: Effect of soil salinity, soil drought, and their
608 combined action on the biochemical characteristics of cotton roots, Acta Physiol. Plant., 35, 3167-3179,
609 <https://doi.org/10.1007/s11738-013-1350-6>, 2013.
610 Zhu, X., Wang, T. J., Skidmore, A. K., Darvishzadeh, R., Niemann, K. O., and Liu, J.: Canopy leaf water content estimated
611 using terrestrial lidar, Agric. For. Meteorol., 232, 152-162, <https://doi.org/10.1016/j.agrformet.2016.08.016>, 2017.