

Quantifying overlapping and differing information of global precipitation for GCM forecasts and El Niño–Southern Oscillation

Tongtiegang Zhao¹, Haoling Chen¹, Yu Tian², Denghua Yan², Weixin Xu³, Huayang Cai⁴, Jiabiao Wang¹, and Xiaohong Chen¹

5 ¹ Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), School of Civil Engineering, Sun Yat-Sen University, Guangzhou, China

² Department of Water Resources, Institute of Water Resources and Hydropower Research of China, Beijing, China

³ School of Atmospheric Sciences, Sun Yat-Sen University, Zhuhai, China

⁴ School of Marine Engineering and Technology, Sun Yat-Sen University, Zhuhai, China

10 *Correspondence to:* Tongtiegang Zhao (zhaottg@mail.sysu.edu.cn), Haoling Chen (chenhling@mail2.sysu.edu.cn), Denghua Yan (yandh@iwhr.com) and Xiaohong Chen (eescxh@mail.sysu.edu.cn)

Abstract. While El Niño–Southern Oscillation (ENSO) teleconnection has long been used in statistical precipitation forecasting, global climate models (GCMs) provide increasingly available dynamical precipitation forecasts for hydrological modelling and water resources management. It is not yet known to what extent dynamical GCM forecasts provide new information compared to statistical teleconnection. This paper develops a novel Set Operations of Coefficients of Determination (SOCD) method to explicitly quantify the overlapping and differing information for GCM forecasts and ENSO teleconnection. Specifically, the intersection operation of the coefficient of determination derives the overlapping information for GCM forecasts and Niño3.4 index; and then the difference operation determines the differing information in GCM forecasts (Niño3.4 index) from Niño3.4 index (GCM forecasts). A case study is devised for the Climate Forecast System version 2 (CFSv2) seasonal forecasts of global precipitation in December-January-February. The results show that the overlapping information for GCM forecasts and Niño3.4 index is significant for 34.94% of global land grid cells, the differing information in GCM forecasts from Niño3.4 index is significant for 31.18% of grid cells and the differing information in Niño3.4 index from GCM forecasts is significant for 11.37% of grid cells. These results confirm the effectiveness of GCMs in capturing the ENSO-related variability of global precipitation and illustrate where there is room for improvements of GCM forecasts. Overall, the bootstrapping significance tests of the three types of information facilitate in total eight patterns to disentangle the close but divergent association of GCM forecast correlation skill with ENSO teleconnection.

1 Introduction

Seasonal precipitation forecasts are important for agricultural scheduling, water management and drought mitigation (Sheffield et al., 2014; Anghileri et al., 2016; Peng et al., 2018; He et al., 2019; Zhao et al., 2019). Performing hydrological forecasting into the future, the uncertainty generally arises from catchment initial conditions and future climate forcings (Wood and Lettenmaier, 2006; Yuan et al., 2014; Huang et al., 2020). In a short lead time up to about one month, initial conditions tend

to outweigh climate forcings; at longer lead times, climate forcings become a more important contributor (Li et al., 2009; Yossef et al., 2013). Therefore, besides remote sensing-based estimations of initial conditions of snow cover, soil moisture and groundwater storage (Mei et al., 2020; Xu et al., 2020b; Sheffield et al., 2014), efforts have been devoted to developing
35 sub-seasonal to seasonal forecasts of temperature and precipitation (Schepen et al., 2020; Strazzo et al., 2019; Bennett et al., 2016; Cash et al., 2019; Li et al., 2017). While temperature forecasts have been improved substantially in the past decades, skilful precipitation forecasts remain challenging (Becker et al., 2022).

Climate indices, in particular El Niño–Southern Oscillation (ENSO) (Mason and Goddard, 2001), have been conventionally used in precipitation forecasting (Hamlet and Lettenmaier, 1999; Hidalgo and Dracup, 2003; Peel et al., 2004).
40 Teleconnections with climate indices generally reflect slowly varying and recurrent components, such as sea surface temperature (SST), of atmospheric circulations that link climate anomalies over large distances in both the tropics and extratropics (Webster and Yang, 1992; Mason and Goddard, 2001; Lim et al., 2021). As one of the most remarkable teleconnections, ENSO affects the global climate through eastward propagating Kelvin waves, westward propagating Rossby waves and Walker circulations that span the tropical Pacific, Indian and Atlantic Oceans (Yang et al., 2018; Webster and Yang,
45 1992). For regions exhibiting teleconnection patterns, various forecasting models have been developed, including historical resampling methods (Hamlet and Lettenmaier, 1999; Wood and Lettenmaier, 2006; Lim et al., 2021), statistical and Bayesian methods (Hidalgo and Dracup, 2003; Strazzo et al., 2019; Emerton et al., 2017) and machine learning methods (Xu et al., 2020a; Li et al., 2021).

Major climate centers develop global climate models (GCMs) to generate operational forecasts of global climate (Bauer et al.,
50 2015; Saha et al., 2014; Khan et al., 2017; Johnson et al., 2019a; Kirtman et al., 2014). For example, the United States National Centers for Environmental Prediction (NCEP) runs the Climate Forecast System version 2 (CFSv2) (Saha et al., 2014) and the European Centre for Medium-Range Weather Forecasts operates the fifth-generation seasonal forecast system (SEAS5) (Johnson et al., 2019b). In contrast to teleconnections that are generally “statistical”, GCM forecasts are “dynamical” in that GCMs assimilate observational information to reduce initial state uncertainty and couple atmosphere, land, ocean and sea ice
55 modules to formulate complex interactions among different components of the earth system (Bauer et al., 2015; Corti et al., 2015; Becker et al., 2022). Previous studies found that GCM forecasts tend to be skilful in regions subject to prominent ENSO teleconnection and also highlighted that GCM forecasts can be skilful in some extratropical regions where there is limited ENSO teleconnection (Johnson et al., 2019b; Kirtman et al., 2014; Delworth et al., 2020).

Conventional ENSO-based statistical forecasts and emerging GCM dynamical forecasts generally represent two different
60 sources of information (Wood and Lettenmaier, 2006; Bauer et al., 2015; Emerton et al., 2017; Delworth et al., 2020; He et al., 2021). Despite the facts that both of them are valuable and that they can be combined to generate improved forecasts (Madadgar et al., 2016; Wanders et al., 2017; Strazzo et al., 2019), it is not yet known to what extent their information overlaps or differs. Small overlap and large difference highlight that GCM forecasts do offer new information comparing to ENSO teleconnection, while large overlap and small difference imply that GCM forecasts might provide little additional information.
65 Zhao et al. (2021) investigated the overlapping information to attribute GCM forecast correlation skill to ENSO teleconnection.

In this paper, we build a Set Operations of Coefficients of Determination (SOCD) method upon Zhao et al. (2021) to furthermore account for the differing information. As will be demonstrated through the methods and results, besides the overlapping information, there exist two types of differing information, i.e., the differing information in GCM forecasts from ENSO and the differing information in ENSO from GCM forecasts. The three types of information facilitate eight patterns to
70 disentangle the close but divergent association of GCM correlation skill with ENSO teleconnection.

2 Data description

GCM precipitation forecasts are generally five-dimensional data (Kirtman et al., 2014; Saha et al., 2014; Delworth et al., 2020; Zhao et al., 2021; Becker et al., 2022). Taking the NCEP-CFSv2 forecasts as an example, the five dimensions are: 1) forecast
75 initial time s , which represents the time at which forecasts are generated, is measured by the number of months since January 1960; 2) lead time l , which represents the months ahead the initial time, ranges from 0 to 9; 3) ensemble member n , which is meant to explicitly account for forecast uncertainty, ranges from 1 to 24, i.e., 24 ensemble members in total; 4) latitude y ; and 5) longitude x . GCM forecasts are therefore formulated as:

$$F = [f_{s,l,n,y,x}], \quad (1)$$

where f represents individual forecast value under the five dimensions and all the forecast values form a dataset F .

80 The observed precipitation corresponding to the forecasts has three dimensions:

$$O = [o_{t,y,x}] \quad (t = s + l), \quad (2)$$

in which o represents individual observation value and O the dataset of observations. The three dimensions are target time t , latitude y and longitude x . It is important to note that target time t is mathematically the sum of initial time s and lead time l in aligning observations with forecasts.

Niño3.4 index that indicates the SST of the East Central Tropical Pacific (5°N–5°S, 170°–120°W) is one of the most popular
85 indicators of the status of ENSO (Hamlet and Lettenmaier, 1999; Emerton et al., 2017; Lin et al., 2020):

$$Niño3.4 = [niño3.4_t], \quad (3)$$

in which there is only one dimension, i.e., time t , for $Niño3.4$.

F , O and $Niño3.4$ shown in Eqs. (1) to (3) lay the basis for the analysis of overlapping and differing information in this paper. In the North American Multi-Model Ensemble (NMME) experiment (Kirtman et al., 2014), CFSv2 retrospective forecasts that range from 1982 to 2010 have been temporally aggregated to monthly and spatially regridded to a $1.0^\circ \times 1.0^\circ$ resolution. In the
90 meantime, the daily Unified Rain-gauge Database (Chen et al., 2008) of the Climate Prediction Center (CPC-URD) precipitation observations over land have also been aggregated and regridded by the NMME. In the analysis, both CFSv2 forecasts and CPC-URD observations are obtained from the International Research Institute of the Columbia University

(<https://iridl.ldeo.columbia.edu/SOURCES/.Models/.NMME/>). Monthly Niño3.4 index is obtained from the CPC (<https://www.cpc.ncep.noaa.gov/data/indices/>).

95

3 Methods

3.1 Consideration of seasonality

Precipitation worldwide exhibits seasonality, e.g., wet and dry seasons of monsoonal precipitation (Webster and Yang, 1992; Zhao et al., 2017; Liu et al., 2022). As a result, the predictive performance of GCM forecasts varies across different seasons
 100 (Kirtman et al., 2014; Bauer et al., 2015; Strazzo et al., 2019) and ENSO teleconnection also exhibits seasonal variabilities (Mason and Goddard, 2001; Peel et al., 2004; Emerton et al., 2017). By fixing the target season, lead time l would be determined by initial time s . Taking December-January-February (DJF) for an example, forecasts generated at the start of December are at 0-month lead time, forecasts at the start of November are at 1-month lead time and so on.

Considering seasonality, the initial time s in Eq. (1) is re-formulated by month m and year k , e.g., December 1982, December
 105 1983, ..., and December 2010. By fixing the target season and specifying the start month, GCM forecasts are then extracted from F :

$$F_{Dec \rightarrow DJF, y, x} = [\bar{f}_k]. \quad (4)$$

The five dimensions of F (Eq. 1) are handled as follows: 1) initial time s and lead time l are replaced by $Dec \rightarrow DJF$ and then represented by k , i.e., aggregating monthly forecasts into seasonal and pooling forecasts across different years; 2) ensemble member n is eliminated by taking the mean value (\bar{f}), i.e., the ensemble mean, of all ensemble members (Saha et al., 2014;
 110 Yuan et al., 2016; Khan et al., 2017); and 3) latitude y and longitude x are pre-specified for the extraction of forecasts.

The observations corresponding to the forecasts (Eq. 4) are extracted from O :

$$O_{DJF, y, x} = [o_k]. \quad (5)$$

In Eq. (5) is observed precipitation in the target season (DJF) across multiple years at the selected grid cell (y, x) . Similar to forecasts, monthly observations are aggregated into seasonal.

Furthermore, the Niño3.4 index in the same season as observed precipitation is obtained:

$$Niño3.4_{DJF} = [niño3.4_k]. \quad (6)$$

115 In Eq. (6) is the concurrent $Niño3.4$ of the target season (DJF) across multiple years.

3.2 Quantification of information in forecasts and Niño3.4

The coefficient of determination (R^2) is effective in quantifying the proportion of the variance of dependent variable explained by a regression model that is built upon some independent variable(s) (Pham, 2006). In this paper, the dependent variable is observed seasonal precipitation (Eq. 5). The candidate independent variables are GCM precipitation forecasts (Eq. 4) and Niño3.4 index (Eq. 6). Three classic simple linear regression models are set up to account for the information of observations in forecast ensemble mean and Niño3.4 index.

The first model regresses observed seasonal precipitation o against ensemble mean \bar{f} of GCM precipitation forecasts:

$$o_k = \alpha_1 + \beta_1 \bar{f}_k + \varepsilon_{1,k} \Rightarrow R^2(o \sim \bar{f}) = 1 - \frac{\sum_k \varepsilon_{1,k}^2}{\sum_k (o_k - \bar{o})^2}, \quad (7)$$

in which α_1 and β_1 are respectively the intercept and slope parameters. The unexplained variance indicated by the sum of squared residuals, i.e., $\sum_k \varepsilon_{1,k}^2$, is compared to the variance of observed precipitation $\sum_k (o_k - \bar{o})^2$. In this way, the proportion of variance explained by ensemble mean is quantified.

The second model regresses observed seasonal precipitation o against *niño*3.4:

$$o_k = \alpha_2 + \beta_2 \tilde{niño}3.4_k + \varepsilon_{2,k} \Rightarrow R^2(o \sim \tilde{niño}3.4) = 1 - \frac{\sum_k \varepsilon_{2,k}^2}{\sum_k (o_k - \bar{o})^2}, \quad (8)$$

in which α_2 , β_2 and $\varepsilon_{2,k}$ are respectively the intercept parameter, slope parameter and residual of regression. This regression quantifies the proportion of variance of observed precipitation explained by *Niño*3.4 index.


The third model regresses observed seasonal precipitation o against both ensemble mean \bar{f} and *niño*3.4:

$$o_k = \alpha_3 + \beta_{3,1} \bar{f}_k + \beta_{3,2} \tilde{niño}3.4_k + \varepsilon_{3,k} \\ \Rightarrow R^2(o \sim \bar{f} \cup \tilde{niño}3.4) = 1 - \frac{\sum_k \varepsilon_{3,k}^2}{\sum_k (o_k - \bar{o})^2}, \quad (9)$$

in which α_3 , $\beta_{3,1}$, $\beta_{3,2}$ and $\varepsilon_{3,k}$ are respectively the intercept parameter, slope parameter of ensemble mean, slope parameter of *Niño*3.4 and residual of regression. The proportion of the variance of observed precipitation explained by the union of ensemble mean and *Niño*3.4 is therefore measured by this bi-variate regression.

3.3 Quantification of overlapping and differing information

As shown by Venn diagrams in Figure 1, the information of observed precipitation contained in forecast ensemble mean, Niño3.4 index and their union are respectively quantified by $R^2(o \sim \bar{f})$, $R^2(o \sim \tilde{niño}3.4)$ and $R^2(o \sim \bar{f} \cup \tilde{niño}3.4)$.



140 **Figure 1: Venn diagram representation of the set operations of union, intersection and difference to quantify the overlapping information and the two types of differing information. The different terms of information are measured by the classic coefficient of determination**

Following the classic set theory, the SOCD method performs the set operations of intersection and difference to quantify the overlapping and differing information:

1) The proportion of variance explained by ensemble mean but not by Niño3.4 index is derived by the difference operation:

$$R^2(o \sim \bar{f} / \text{Niño}3.4) = R^2(o \sim \bar{f} \cup \text{Niño}3.4) - R^2(o \sim \text{Niño}3.4). \quad (10)$$

In Eq. (10), $R^2(o \sim \bar{f} / \text{Niño}3.4)$ measures the differing information of GCM forecasts on observed precipitation from Niño3.4 index.

2) The intersection operation derives the proportion of variance of seasonal precipitation explained by both ensemble mean and Niño3.4 index:

$$\begin{aligned}
& R^2(o \sim \bar{f} \cap \tilde{ni\~{n}o3.4}) \\
& = R^2(o \sim \bar{f}) + R^2(o \sim \tilde{ni\~{n}o3.4}) - R^2(o \sim \bar{f} \cup \tilde{ni\~{n}o3.4})
\end{aligned} \tag{11}$$

In Eq. (11), $R^2(o \sim \bar{f} \cap \tilde{ni\~{n}o3.4})$ represents the overlapping information.

3) The proportion of variance explained by Niño3.4 index but not by ensemble mean is derived by the difference operation:

$$R^2(o \sim \tilde{ni\~{n}o3.4}/\bar{f}) = R^2(o \sim \bar{f} \cup \tilde{ni\~{n}o3.4}) - R^2(o \sim \bar{f}). \tag{12}$$

In Eq. (12), $R^2(o \sim \tilde{ni\~{n}o3.4}/\bar{f})$ represents the differing information of Niño3.4 index from GCM forecasts.

155 3.4 Eight patterns for overlapping and differing information

The significance of overlapping and differing information is tested by bootstrapping (Efron and Tibshirani, 1986). The null hypothesis is that the three variables under investigation, i.e., o , \bar{f} and $\tilde{ni\~{n}o3.4}$, were fully independent from one another.

Under the null hypothesis, the samples in Eqs. (4), (5) and (6) are randomly selected with replacement to calculate the overlapping and differing information; one thousand such recalculations formulate the respective reference distributions for these R^2 values. Comparing the R^2 values for the original samples respectively to their reference distributions, the p-values are obtained to tell how extreme the R^2 values for the original samples are. In this way, the significance is tested (Efron and Tibshirani, 1986; Pham, 2006). As the null hypothesis is full independence, the R^2 values, which indicate the amount of information of the dependent variable contained in independent variable(s), are expected to be rather small. From this perspective, the larger the R^2 values for the original samples are, the more extreme they are and the less likely the null hypothesis holds. Therefore, the one-tailed test is implemented for the significance of the R^2 values (Pham, 2006). Specifically, under the significance level of 0.10, the SOCD method pays attention to whether the R^2 value falls into the top 10% of the corresponding bootstrapping-derived reference distribution.

The one type of overlapping information and the two types of differing information each have two cases of significance, i.e., significant or non-significant. Therefore, in Table 1, a three-digit number is devised to represent the results of significance test.

The first digit indicates the significance of $R^2(o \sim \bar{f}/\tilde{ni\~{n}o3.4})$, the second digit the significance of $R^2(o \sim \bar{f} \cap \tilde{ni\~{n}o3.4})$ and the third digit the significance of $R^2(o \sim \tilde{ni\~{n}o3.4}/\bar{f})$. As is shown in Table 1, there are in total 8 ($2*2*2$) patterns, with 1 representing the significant case and 0 indicating the non-significant case. The meanings of the eight patterns are illustrated in the last column of Table 1.

175 **Table 1: Three-digit representations of the eight patterns of overlapping and differing information**

$R^2(o \sim \bar{f}/\tilde{nino3.4})$	$R^2(o \sim \bar{f} \cap \tilde{nino3.4})$	$R^2(o \sim \tilde{nino3.4}/\bar{f})$	Meaning
0	0	0	Neither overlapping information nor differing information is significant
0	0	1	Only the differing information in Niño3.4 index from GCM forecasts is significant
0	1	0	Only the overlapping information is significant
0	1	1	Both overlapping information and differing information in Niño3.4 index from GCM forecasts are significant
1	0	0	Only the differing information in GCM forecasts from Niño3.4 index is significant
1	0	1	Both differing information in GCM forecasts from Niño3.4 index and differing information in Niño3.4 index from GCM forecasts are significant, but the overlapping information is not significant
1	1	0	Both differing information in GCM forecasts from Niño3.4 index and overlapping information are significant
1	1	1	Differing information in GCM forecasts from Niño3.4 index, overlapping information and differing information in Niño3.4 index from GCM forecasts are all significant

4 Results

4.1 Spatial plots of correlation skill and ENSO teleconnection

180 GCM forecast correlation skill and ENSO teleconnection for DJF are shown in the left-hand side of Figure 2. The correlation skill is mathematically the Pearson's correlation coefficient between GCM forecast ensemble mean and observed precipitation. In the upper left part of Figure 2, it is observed that the correlation skill is higher than 0.3 in a substantial number of grid cells around the world. This result indicates that ensemble mean is generally indicative of observed precipitation, i.e., high values of ensemble mean coincide with high values of observed precipitation and vice versa (Saha et al., 2014; Yuan et al., 2014;

Cash et al., 2019). In the lower left part is ENSO teleconnection that mathematically represents the Pearson's correlation coefficient between Niño3.4 index and observed precipitation. Both positive and negative ENSO teleconnections are observed. For example, the teleconnection tends to be positive in southern North America, south-eastern South America, southern China and Eastern Africa, implying above-average precipitation in El Niño years but below-average precipitation in La Niña years; and it turns out to be negative in the northern part of South America, southern Africa as well as Southeast Asia, i.e., there can be below-average precipitation in El Niño years and above-average precipitation in La Niña years (Mason and Goddard, 2001; Emerton et al., 2017; Yang et al., 2018).

The SOCD method facilitates in total eight patterns to characterize the overlapping and differing information for GCM forecast ensemble mean and Niño3.4 index. As the Venn diagram in Figure 1 is presented for the purpose of conceptual illustration, the right-hand side of Figure 2 showcases the Venn diagrams generated from real-world data. The eight patterns in Table 1 are illustrated for eight grid cells selected from the left-hand side of Figure 2. Grid cell A (36° N, 115° E) is under the pattern 000: the areas of the circles that indicate the ratios of explained variance are rather small, implying little information of observed precipitation in GCM forecasts and Niño3.4 index. By contrast, the areas of the circles are larger for the other seven selected grid cells, suggesting the existence of significant overlapping or differing information. For example, the significant overlapping information is highlighted in grid cell D (36°N, 115°E); the significant differing information in GCM forecasts from Niño3.4 index is shown in grid cell E (46°N, 0°); and the significant differing information in Niño3.4 index from GCM forecasts is highlighted in grid cell B (35°N, 117°E).




Figure 2: Spatial plots of correlation skill (upper left part) and ENSO teleconnection (lower left part) for global precipitation in DJF, and Venn diagrams (right part) of overlapping and differing information for eight selected grid cells under the eight patterns.





Figure 3: Spatial distribution of the eight patterns of overlapping and differing information.

The spatial distribution of the eight patterns is shown in Figure 3 by applying the SOCD method and the bootstrapping
 210 significance test to all the land grid cells. Grid cells under the pattern 000, which indicates poor GCM correlation skill and
 limited ENSO teleconnection, are in grey. In the meantime, it is noted that a considerable amount of grid cells around the
 world are colored. That is, for the overlapping information and the two types of differing information, at least one of them is
 significant. From the left-hand side of Figure 2, it can be found that positive correlation skill corresponds to positive ENSO
 teleconnection in southern North America and Eastern Africa and that positive correlation skill corresponds to negative
 215 teleconnection over the northern part of South America, southern Africa and Southeast Asia. In the meantime, from Figure 3
 it can be observed that in these regions a considerable number of grid cells fall under the patterns 010, 110 and 011, indicating
 significant overlapping information.

4.2 Patterns of overlapping and differing information

220 The eight patterns serve as a link between correlation skill and ENSO teleconnection. The pattern 010 that is concentrated on
 the overlapping information is shown in Figure 4. At the left-hand side of the figure are the results for grid cells under the
 pattern 010 (the results for the other grid cells are masked). The overlapping information is significant in southern North
 America where positive correlation skill (upper left part of Figure 4) coincides with positive ENSO teleconnection (lower left
 part of Figure 4). It is also significant in southern Africa and northern South America where positive correlation skill and
 225 negative ENSO teleconnection coexist. As both correlation skill and ENSO teleconnection are mathematically the Pearson's


230 correlation coefficient, they each can be classified into three cases, i.e., significantly positive (P), non-significant (ns) and significantly negative (N) (Kirtman et al., 2014; Emerton et al., 2017; Huang and Zhao, 2022). At the right-hand side of Figure 4, the Sankey diagram shows that 18.95% of the global land grid cells exhibit the pattern 010. For this pattern, 8.98% of grid cells exhibit significantly positive correlation skill, 9.85% non-significant correlation skill and 0.12% significantly negative correlation skill; in the meantime, 3.77% exhibiting significantly positive ENSO teleconnection, 10.92% non-significant ENSO teleconnection and 4.25% significantly negative ENSO teleconnection.



235 **Figure 4: Illustrations of correlation skill (upper left part) and ENSO teleconnection (lower left part) under the pattern 010 (middle left part) and the Sankey diagram showing the percentage of grid cells under the pattern 010 and the percentages of grid cells exhibiting significantly positive (P), non-significant (ns) and significantly negative (N) correlation skill/ENSO teleconnection (right part). Grid cells under the other patterns are masked and therefore not shown in the spatial plots and the Sankey diagram.**

240 The pattern 100 that focuses on the significant differing information of global precipitation in GCM forecasts from Niño3.4 index is shown in Figure 5. From the left-hand side of the figure, it can be observed that this pattern (middle left part) tends to cover grid cells where correlation skill is around or above 0.3 (upper left part) but ENSO teleconnection is nearly zero (lower left part). This observation is confirmed by the right-hand side of Figure 5. As can be seen, while the percentage of grid cells

falling into the pattern 100 is 17.71%, most of them are with significantly positive correlation skill (15.78% in 17.71%) and by contrast all of them exhibit non-significant ENSO teleconnection (17.71% in 17.71%). That is, the correlation skill is largely significantly positive but the ENSO teleconnection is generally neutral. These grid cells tend to locate in Europe and North Asia, where the influence of ENSO is limited and skillful GCM forecasts can relate to other teleconnections such as Arctic Oscillation and North Atlantic Oscillation (Hamouda et al., 2021).



250 **Figure 5: As for Figure 4 but for the pattern 100.**

The pattern 110 indicates that the overlapping information is significant and that the differing information in GCM forecasts from Niño3.4 index is also significant. The implication is that regarding global seasonal precipitation in DJF, GCM forecasts not only contain information that is contained in Niño3.4 index but also provide a considerable amount of additional information. On the left-hand side of Figure 6, some grid cells under the pattern 110 are observed in southeast Australia, eastern Africa and northeastern Asia. Comparing Figure 6 to Figure 4, it is observed that some grid cells in southern North America, northern South America and southern Africa are under the pattern 110, although many of them tend to be under the pattern 100. Around the world, the percentage of grid cells falling into the pattern 110 is 11.35%. For these grid cells, correlation skill

is predominantly significantly positive (11.25% in 11.35%) and by contrast ENSO teleconnection tends to be non-significant (7.09% in 11.35%).




Figure 6: As for Figure 4 but for the pattern 110.

265 The pattern 001 pays attention to the differing information in Niño3.4 index from GCM forecasts. As shown in Figure 7, this pattern covers 4.87% of grid cells around the world. On the left-hand side of Figure 7, it is worthwhile to note that a number of grid cells in Western Australia exhibit significantly negative ENSO teleconnection but non-significant correlation skill. The implication is that therein GCM forecasts might fail to account for the information of ENSO teleconnection. At the right-hand side of Figure 7, it is observed that most grid cells under the pattern 001 are with neutral correlation skill (4.86% in 4.87%) and that their corresponding ENSO teleconnection can be significantly negative (2.21% in 4.87%) or significantly positive (1.73% in 4.87%).

270





Figure 7: As for Figure 4 but for the pattern 001.

275

The pattern 011 indicates that both the overlapping information and the differing information in Niño3.4 index from GCM forecasts are significant. Grid cells exhibiting this pattern tend to be scattered in parts of southern North America, northern South America, Southeast Asia and southern Africa. They account for 4.38% of grid cells around the world. Among them, 1.72% exhibit significantly positive ENSO teleconnection and 2.66% significantly negative ENSO teleconnection. For these areas, the significant overlap suggests that a substantial amount of information in seasonal precipitation can be explained by both GCM forecasts and Niño3.4, while the significant differing information indicates the part that can only be explained by the Niño3.4 index.

280



285 **Figure 8:** As for Figure 4 but for the pattern 011.

The pattern 101 is shown in Figure 9. It suggests that at some grid cells, the overlapping information is not significant but the two types of differing information are significant for both GCM forecasts and Niño3.4 index. About 1.86% of grid cells fall into this pattern.

290 The pattern 111 is shown in Figure 10. It implies that at some other grid cells, the overlapping information and the two types of differing information can all be significant. It is noted that only 0.26% of grid cells around the world exhibit pattern 111.




Figure 9: As for Figure 4 but for the pattern 101.




Figure 10: As for Figure 4 but for the pattern 111.

Among the eight patterns, the pattern 000 covers the most grid cells. The left-hand side of Figure 11 shows that grid cells under the pattern 000 generally exhibit non-significant correlation skill and non-significant ENSO teleconnection. This result is in sharp contrast to the pattern 010 which indicates reasonable correspondence between correlation skill and ENSO teleconnection (Figure 4) and to the patterns 100 and 110 which suggest significantly positive correlation skill (Figures 5 and 6). Overall, the percentage of grid cells under the pattern 000 is 40.62%. These grid cells predominantly exhibit neutral correlation skill (40.30% in 40.62%) and also neutral ENSO teleconnection (40.47% in 40.62%).

305





Figure 11: As for Figure 4 but for the pattern 000.

4.3 Association of correlation skill with ENSO teleconnection

310 The results under the eight patterns are furthermore pooled in the analysis. From Figure 12, it can be observed that the eight
 patterns serve to be an effective link between correlation skill and ENSO teleconnection at the global scale. For the patterns
 that indicate significant information, the Sankey diagram at the right-hand side suggests that the percentage from the highest
 to the lowest is respectively 18.95% for the pattern 010, 17.71% for the pattern 100, 11.35% for the pattern 110, 4.87% for the
 pattern 001, 4.38% for the pattern 011, 1.86% for the pattern 101 and 0.26% for the pattern 111. More than half of the grid
 315 cells that exhibit significant correlation skill have significant overlapping information with Niño3.4, with 11.25% (8.98%) of
 grid cells under the pattern 110 (010), indicating considerable impacts of ENSO teleconnection on CFSv2 correlation skill.




320 **Figure 12: Spatial plots of correlation skill (upper left part) and ENSO teleconnection (lower left part) under the eight patterns (middle left part) at the global scale and the Sankey diagram showing the percentages of grid cells exhibiting significantly positive (P), non-significant (ns) and significantly negative (N) correlation skill/ENSO teleconnection (right part)**

GCM forecasts and Niño3.4 index generally represent two different sources of information of global precipitation. In Figure 13, GCM forecast correlation skill is plotted against ENSO teleconnection by using scatter plots. Figure 13a pools global land grid cells and employs the Viridis heatmap to indicate point density. It can be observed that the correlation skill is largely positive and fall above the horizontal line. In addition, the heatmap suggests that the correlation skill tends to increase with the increase of positive ENSO teleconnection and also with the decrease of negative ENSO teleconnection. These results suggest that the skill of GCM forecasts benefits from the prominence of ENSO teleconnection since GCMs tend to capture the influences of ENSO on the variability of global precipitation (Saha et al., 2014; Khan et al., 2017; Delworth et al., 2020; Johnson et al., 2019b; Becker et al., 2022).

325
330

The subplots in Figures 13b-i are arranged in descending order of the percentage of grid cells. Overall, a close but divergent association of correlation skill with ENSO teleconnection can be observed:

- 1) There exists significant overlapping information in GCM forecasts and Niño3.4 index under the patterns 010 (Figure 13c), 110 (Figure 13e), 011 (Figure 13g) and 111 (Figure 13i). It covers 34.94% of land grid cells, i.e., 18.95% (010) + 11.35% (110) + 4.38% (011) + 0.26% (111), around the world. From the corresponding scatter plots, it can be observed that both correlation skill and ENSO teleconnection ought to be reasonably high to facilitate significant overlapping information;
- 2) There is significant differing information in GCM forecasts from Niño3.4 index under the patterns 100 (Figure 13d), 110 (Figure 13e), 101 (Figure 13h) and 111 (Figure 13i). It covers 31.18% of global land grid cells, i.e., 17.71% (100) + 11.35% (110) + 1.86% (101) + 0.26% (111). Under these patterns, it is highlighted that the correlation skill tends to be higher than ENSO teleconnection. In particular, significantly positive correlation skill coincides with overall non-significant ENSO teleconnection under the pattern 100 in Figure 13f. Overall, these results imply that apart from ENSO, GCMs account for other hydro-climatic teleconnections to produce skilful precipitation forecasts (Saha et al., 2014; Delworth et al., 2020; Lin et al., 2020);
- 3) There is significant differing information in Niño3.4 index from GCM forecasts under the patterns 001 (Figure 13f), 011 (Figure 13g), 101 (Figure 13h) and 111 (Figure 13i). It covers 11.37% of global land grid cells, i.e., 4.87% (001) + 4.38% (011) + 1.86% (101) + 0.26% (111). Under these patterns, ENSO teleconnection is generally higher than correlation skill. Remarkable ENSO teleconnection coincides with overall non-significant correlation skill under the pattern 001 in Figure 13b. These results suggest that some ENSO teleconnection is still yet to be exploited by GCMs to improve precipitation forecast skill.
- 4) Neither the overlapping information nor the two types of differing information are significant under the pattern 000. It covers 40.62% of grid cells. From Figure 13b, it can be observed that either correlation skill or ENSO teleconnection is limited and that the corresponding scatter plot tends to cluster around the origin point. This result suggests that despite limited ENSO teleconnection, GCM forecasts still have plenty of room for improvement.



355

Figure 13: Scatter plots of the association of GCM forecast correlation skill with ENSO teleconnection at the global scale with the heatmap indicating the density of scatter points (a) and of the association of correlation skill with ENSO teleconnection under the eight patterns (b-i)

360 5 Discussion

The SOCD method is furthermore applied to investigate the eight patterns considering the effects of seasonality, lead time, lag time and significance level. The additional results are presented in the supplementary material. 1) The effect of seasonality is

shown in Figures S1 to S6. It can be observed that regions exhibiting significant ENSO teleconnections vary by season (Figures S1 to S3) and that the eight patterns remain effective in characterizing the overlapping and differing information (Figures S4 to S6). 2) The effect of lead time is illustrated in Figures S7 to S10. At the lead times of 1 and 2 months, the percentage of the pattern 010 remains the highest among the seven patterns other than 000. This result highlights the existence of significant overlapping information in DJF, particularly over southern North America, northern South America and Southern Africa. 3) The effect of the lag time of Niño3.4 index is illustrated in Figures S11 to S14. Compared to the concurrent teleconnection, the spatial distribution of the eight patterns tends to be similar for monthly Niño3.4 index at the lag times of 1 and 2 months, with a slight increase in the percentage of the pattern 000. This result confirms the temporal persistency in the Niño3.4 index (Yang et al., 2018). 4) The effect of the significance level is shown in Figures S15 to S18. As the significance level is reduced from 0.10 to 0.05 and furthermore to 0.01, the percentage of the pattern 000 evidently increases but the seven patterns that highlight significant overlapping and differing information remain.

The SOCD method is also extended to evaluate the overlapping and differing information under other GCM forecasts and hydroclimatic teleconnections. In the supplementary material, Figures S19 and S20 show the results for the CanCM4 forecasts generated at the Canadian Meteorological Center (CMC) (Merryfield et al., 2013). The CanCM4 forecasts seem to be less skilful in Europe but more skilful in the western part of Australia. Overall, the percentage of the pattern 000 is slightly higher than that for CFSv2 forecasts. These results suggest that different GCM forecasts can be complementary to each other in different regions and that they can be combined to generate more skilful forecasts (Kirtman et al., 2014; Slater et al., 2019; Schepen et al., 2020). Figures S21 and S22 present the eight patterns for the Indian Ocean Dipole (IOD) (Cai et al., 2021). It can be observed that the percentage of the pattern 010 is reduced from 18.95% to 9.41% while the percentage of the pattern 100 is increased from 17.71% to 22.83%. The indications are that CFSv2 forecasts exhibit less overlapping information with IOD and that there exists considerable differing information in CFSv2 forecasts from IOD teleconnection.

The correlation skill is one of the most popular measures of forecast skill owing to its simplicity in calculation and robustness to zero and missing values (Barnston et al., 2012; Yuan et al., 2014; Ma et al., 2016; Slater et al., 2019; Huang and Zhao, 2022). From spatial plots of correlation skill at regional or global scales, it can be observed where GCM forecasts are skilful and where GCM forecasts are not satisfactory (Ma et al., 2016; Slater et al., 2019; Delworth et al., 2020). Previously, it was observed that GCM forecasts tend to be skilful in regions subject to prominent influences of ENSO; accordingly, forecast skill is attributed to the effectiveness of GCMs in capturing ENSO-related climate dynamics (Kirtman et al., 2014; Slater et al., 2019; Lin et al., 2020). In this paper, the developed SOCD method not only confirms the significant overlapping information but also highlights that there exists significant differing information in GCM forecasts from ENSO teleconnection for 31.18% of global land grid cells and that there is significant differing information in ENSO teleconnection from GCM forecasts for 11.37% of grid cells. It is noted that the simple linear regression accounts for linear relationships. Nonlinear relationships between forecasts and observations are possible and in the future nonlinear models can be adopted in the analysis of the overlapping and differing information (Strazzo et al., 2019; Schepen et al., 2020; Li et al., 2021).

6 Conclusions

While ENSO teleconnection has been conventionally used in the forecasting of regional precipitation and streamflow, GCM forecasts are increasingly available for hydrological applications. It is important to investigate to what extent emerging GCM forecasts provide “new” information compared to conventional ENSO teleconnection. The SOCD method developed in this paper addresses this issue through the mathematical formulation of set operations. Specifically, the union operation quantifies the information of global seasonal precipitation contained in both GCM forecasts and Niño3.4 index; the intersection operation derives the overlapping information of global precipitation in GCM forecasts and Niño3.4 index; and furthermore, the difference operations illustrate two types of differing information, i.e., the differing information in GCM forecasts from Niño3.4 index and the differing information in Niño3.4 index from GCM forecasts. The significance tests of the three types of information facilitate in total eight patterns to disentangle the close but divergent association of GCM forecast correlation skill with ENSO teleconnection. GCM forecasts and Niño3.4 index generally provide two different sources of data for precipitation forecasting. While the existence of significant overlapping information suggests that they can provide some similar information, the existence of significant differing information highlights that the two data sources can also be complementary to each other. In the future, more efforts can be devoted to investigating more datasets of GCM forecasts and more hydroclimatic teleconnections to yield insights into the forecast skill of GCM forecasts and to facilitate applications of GCM forecasts to hydrological modelling and water resources management.

Acknowledgments

This research is supported by the National Key Research and Development Program of China (2021YFC3001000), the National Natural Science Foundation of China (51979295, 51861125203, 52109046 and U1911204) and the Guangdong Provincial Department of Science and Technology (2019ZT08G090).

Data Availability Statement

The forecast and observation datasets are downloaded from <https://iridl.ldeo.columbia.edu/SOURCES/Models/NMME/>. The Niño3.4 index is downloaded from <https://www.cpc.ncep.noaa.gov/data/indices/>.

Author contribution

TZ, YT, WX, HC, JW and XC designed the experiments. HC carried them out. HC and TZ developed the model code and
425 performed the simulations. TZ and HC prepared the manuscript with contributions from the co-authors.

Competing interests

The authors declare that they have no conflict of interest.

430 References

- Anghileri, D., Voisin, N., Castelletti, A., Pianosi, F., Nijssen, B., and Lettenmaier, D. P.: Value of long-term streamflow forecasts to reservoir operations for water supply in snow-dominated river catchments: VALUE OF LONG-TERM FORECASTS TO RESERVOIR OPERATIONS, *Water Resour. Res.*, 52, 4209–4225, <https://doi.org/10.1002/2015WR017864>, 2016.
- 435 Barnston, A. G., Tippett, M. K., L’Heureux, M. L., Li, S., and DeWitt, D. G.: Skill of Real-Time Seasonal ENSO Model Predictions during 2002–11: Is Our Capability Increasing?, *Bulletin of the American Meteorological Society*, 93, 631–651, <https://doi.org/10.1175/BAMS-D-11-00111.1>, 2012.
- Bauer, P., Thorpe, A., and Brunet, G.: The quiet revolution of numerical weather prediction, *Nature*, 525, 47–55, <https://doi.org/10.1038/nature14956>, 2015.
- 440 Becker, E. J., Kirtman, B. P., L’Heureux, M., Muñoz, Á. G., and Pegion, K.: A Decade of the North American Multimodel Ensemble (NMME): Research, Application, and Future Directions, *Bulletin of the American Meteorological Society*, 103, E973–E995, <https://doi.org/10.1175/BAMS-D-20-0327.1>, 2022.
- Bennett, J. C., Wang, Q. J., Li, M., Robertson, D. E., and Schepen, A.: Reliable long-range ensemble streamflow forecasts: Combining calibrated climate forecasts with a conceptual runoff model and a staged error model: LONG-RANGE
445 ENSEMBLE STREAMFLOW FORECASTS, *Water Resour. Res.*, 52, 8238–8259, <https://doi.org/10.1002/2016WR019193>, 2016.
- Cai, W., Yang, K., Wu, L., Huang, G., Santoso, A., Ng, B., Wang, G., and Yamagata, T.: Opposite response of strong and moderate positive Indian Ocean Dipole to global warming, *Nat. Clim. Chang.*, 11, 27–32, <https://doi.org/10.1038/s41558-020-00943-1>, 2021.
- 450 Cash, B. A., Manganello, J. V., and Kinter, J. L.: Evaluation of NMME temperature and precipitation bias and forecast skill for South Asia, *Clim Dyn*, 53, 7363–7380, <https://doi.org/10.1007/s00382-017-3841-4>, 2019.
- Chen, M., Shi, W., Xie, P., Silva, V. B. S., Kousky, V. E., Wayne Higgins, R., and Janowiak, J. E.: Assessing objective techniques for gauge-based analyses of global daily precipitation, *J. Geophys. Res.*, 113, D04110, <https://doi.org/10.1029/2007JD009132>, 2008.

- 455 Corti, S., Palmer, T., Balmaseda, M., Weisheimer, A., Drijfhout, S., Dunstone, N., Hazeleger, W., Kröger, J., Pohlmann, H., Smith, D., Storch, J.-S. von, and Wouters, B.: Impact of Initial Conditions versus External Forcing in Decadal Climate Predictions: A Sensitivity Experiment*, *Journal of Climate*, 28, 4454–4470, <https://doi.org/10.1175/JCLI-D-14-00671.1>, 2015.
- Delworth, T. L., Cooke, W. F., Adcroft, A., Bushuk, M., Chen, J., Dunne, K. A., Ginoux, P., Gudgel, R., Hallberg, R. W., Harris, L., Harrison, M. J., Johnson, N., Kapnick, S. B., Lin, S., Lu, F., Malyshev, S., Milly, P. C., Murakami, H., Naik, V.,
460 Pascale, S., Paynter, D., Rosati, A., Schwarzkopf, M. D., Shevliakova, E., Underwood, S., Wittenberg, A. T., Xiang, B., Yang, X., Zeng, F., Zhang, H., Zhang, L., and Zhao, M.: SPEAR: The Next Generation GFDL Modeling System for Seasonal to Multidecadal Prediction and Projection, *J Adv Model Earth Syst*, 12, <https://doi.org/10.1029/2019MS001895>, 2020.
- Efron, B. and Tibshirani, R.: Bootstrap Methods for Standard Errors, Confidence Intervals, and Other Measures of Statistical Accuracy, *Statist. Sci.*, 1, <https://doi.org/10.1214/ss/1177013815>, 1986.
- 465 Emerton, R., Cloke, H. L., Stephens, E. M., Zsoter, E., Woolnough, S. J., and Pappenberger, F.: Complex picture for likelihood of ENSO-driven flood hazard, *Nat Commun*, 8, 14796, <https://doi.org/10.1038/ncomms14796>, 2017.
- Hamlet, A. F. and Lettenmaier, D. P.: Columbia River Streamflow Forecasting Based on ENSO and PDO Climate Signals, *Journal of Water Resources Planning and Management*, 125, 333–341, [https://doi.org/10.1061/\(ASCE\)0733-9496\(1999\)125:6\(333\)](https://doi.org/10.1061/(ASCE)0733-9496(1999)125:6(333)), 1999.
- 470 Hamouda, M. E., Pasquero, C., and Tziperman, E.: Decoupling of the Arctic Oscillation and North Atlantic Oscillation in a warmer climate, *Nat. Clim. Chang.*, 11, 137–142, <https://doi.org/10.1038/s41558-020-00966-8>, 2021.
- He, X., Estes, L., Konar, M., Tian, D., Anghileri, D., Baylis, K., Evans, T. P., and Sheffield, J.: Integrated approaches to understanding and reducing drought impact on food security across scales, *Current Opinion in Environmental Sustainability*, 40, 43–54, <https://doi.org/10.1016/j.cosust.2019.09.006>, 2019.
- 475 He, X., Bryant, B. P., Moran, T., Mach, K. J., Wei, Z., and Freyberg, D. L.: Climate-informed hydrologic modeling and policy typology to guide managed aquifer recharge, *Sci. Adv.*, 7, eabe6025, <https://doi.org/10.1126/sciadv.abe6025>, 2021.
- Hidalgo, H. G. and Dracup, J. A.: ENSO and PDO Effects on Hydroclimatic Variations of the Upper Colorado River Basin, *J. Hydrometeorol*, 4, 5–23, [https://doi.org/10.1175/1525-7541\(2003\)004<0005:EAPEOH>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0005:EAPEOH>2.0.CO;2), 2003.
- Huang, Z. and Zhao, T.: Predictive performance of ensemble hydroclimatic forecasts: Verification metrics, diagnostic plots
480 and forecast attributes, *WIREs Water*, 9, <https://doi.org/10.1002/wat2.1580>, 2022.
- Huang, Z., Zhao, T., Liu, Y., Zhang, Y., Jiang, T., Lin, K., and Chen, X.: Differing roles of base and fast flow in ensemble seasonal streamflow forecasting: An experimental investigation, *Journal of Hydrology*, 591, 125272, <https://doi.org/10.1016/j.jhydrol.2020.125272>, 2020.
- Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche, S., Decremmer, D.,
485 Weisheimer, A., Balsamo, G., Keeley, S. P. E., Mogensen, K., Zuo, H., and Monge-Sanz, B. M.: SEAS5: the new ECMWF seasonal forecast system, *Geosci. Model Dev.*, 12, 1087–1117, <https://doi.org/10.5194/gmd-12-1087-2019>, 2019a.
- Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche, S., Decremmer, D., Weisheimer, A., Balsamo, G., Keeley, S. P. E., Mogensen, K., Zuo, H., and Monge-Sanz, B. M.: SEAS5: the new ECMWF seasonal forecast system, *Geosci. Model Dev.*, 12, 1087–1117, <https://doi.org/10.5194/gmd-12-1087-2019>, 2019b.

- 490 Khan, M. Z. K., Sharma, A., and Mehrotra, R.: Global seasonal precipitation forecasts using improved sea surface temperature predictions: SEASONAL PRECIPITATION FORECASTS, *J. Geophys. Res. Atmos.*, 122, 4773–4785, <https://doi.org/10.1002/2016JD025953>, 2017.
- Kirtman, B. P., Min, D., Infanti, J. M., Kinter, J. L., Paolino, D. A., Zhang, Q., van den Dool, H., Saha, S., Mendez, M. P., Becker, E., Peng, P., Tripp, P., Huang, J., DeWitt, D. G., Tippett, M. K., Barnston, A. G., Li, S., Rosati, A., Schubert, S. D.,
495 Rienecker, M., Suarez, M., Li, Z. E., Marshak, J., Lim, Y.-K., Tribbia, J., Pegion, K., Merryfield, W. J., Denis, B., and Wood, E. F.: The North American Multimodel Ensemble: Phase-1 Seasonal-to-Interannual Prediction; Phase-2 toward Developing Intraseasonal Prediction, *Bull. Amer. Meteor. Soc.*, 95, 585–601, <https://doi.org/10.1175/BAMS-D-12-00050.1>, 2014.
- Li, H., Luo, L., Wood, E. F., and Schaake, J.: The role of initial conditions and forcing uncertainties in seasonal hydrologic forecasting, *J. Geophys. Res.*, 114, D04114, <https://doi.org/10.1029/2008JD010969>, 2009.
- 500 Li, J., Wang, Z., Wu, X., Xu, C., Guo, S., Chen, X., and Zhang, Z.: Robust Meteorological Drought Prediction Using Antecedent SST Fluctuations and Machine Learning, *Water Resources Research*, 57, <https://doi.org/10.1029/2020WR029413>, 2021.
- Li, W., Duan, Q., Miao, C., Ye, A., Gong, W., and Di, Z.: A review on statistical postprocessing methods for hydrometeorological ensemble forecasting, *WIREs Water*, 4, <https://doi.org/10.1002/wat2.1246>, 2017.
- 505 Lim, E.-P., Hudson, D., Wheeler, M. C., Marshall, A. G., King, A., Zhu, H., Hendon, H. H., de Burgh-Day, C., Trewin, B., Griffiths, M., Ramchurn, A., and Young, G.: Why Australia was not wet during spring 2020 despite La Niña, *Sci Rep*, 11, 18423, <https://doi.org/10.1038/s41598-021-97690-w>, 2021.
- Lin, H., Merryfield, W. J., Muncaster, R., Smith, G. C., Markovic, M., Dupont, F., Roy, F., Lemieux, J.-F., Dirkson, A., Kharin, V. V., Lee, W.-S., Charron, M., and Erfani, A.: The Canadian Seasonal to Interannual Prediction System Version 2 (CanSIPSv2), *Weather and Forecasting*, 35, 1317–1343, <https://doi.org/10.1175/WAF-D-19-0259.1>, 2020.
- 510 Liu, X., Zhang, L., She, D., Chen, J., Xia, J., Chen, X., and Zhao, T.: Postprocessing of hydrometeorological ensemble forecasts based on multisource precipitation in Ganjiang River basin, China, *Journal of Hydrology*, 605, 127323, 2022.
- Ma, F., Ye, A., Deng, X., Zhou, Z., Liu, X., Duan, Q., Xu, J., Miao, C., Di, Z., and Gong, W.: Evaluating the skill of NMME seasonal precipitation ensemble predictions for 17 hydroclimatic regions in continental China: EVALUATING THE SKILL
515 OF NMME SEASONAL PRECIPITATION PREDICTIONS, *Int. J. Climatol.*, 36, 132–144, <https://doi.org/10.1002/joc.4333>, 2016.
- Madadgar, S., AghaKouchak, A., Shukla, S., Wood, A. W., Cheng, L., Hsu, K.-L., and Svoboda, M.: A hybrid statistical-dynamical framework for meteorological drought prediction: Application to the southwestern United States: A HYBRID
520 STATISTICAL-DYNAMICAL DROUGHT PREDICTION FRAMEWORK, *Water Resour. Res.*, 52, 5095–5110, <https://doi.org/10.1002/2015WR018547>, 2016.
- Mason, S. J. and Goddard, L.: Probabilistic Precipitation Anomalies Associated with ENSO, *Bull. Amer. Meteor. Soc.*, 82, 619–638, [https://doi.org/10.1175/1520-0477\(2001\)082<0619:PPAAWE>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<0619:PPAAWE>2.3.CO;2), 2001.
- Mei, L., Rozanov, V., Ritter, C., Heinold, B., Jiao, Z., Vountas, M., and Burrows, J. P.: Retrieval of Aerosol Optical Thickness in the Arctic Snow-Covered Regions Using Passive Remote Sensing: Impact of Aerosol Typing and Surface Reflection Model,
525 *IEEE Trans. Geosci. Remote Sensing*, 58, 5117–5131, <https://doi.org/10.1109/TGRS.2020.2972339>, 2020.

- Merryfield, W. J., Lee, W.-S., Boer, G. J., Kharin, V. V., Scinocca, J. F., Flato, G. M., Ajayamohan, R. S., Fyfe, J. C., Tang, Y., and Polavarapu, S.: The Canadian Seasonal to Interannual Prediction System. Part I: Models and Initialization, *Monthly Weather Review*, 141, 2910–2945, <https://doi.org/10.1175/MWR-D-12-00216.1>, 2013.
- 530 Peel, M. C., McMahon, T. A., and Finlayson, B. L.: Continental differences in the variability of annual runoff-update and reassessment, *Journal of Hydrology*, 295, 185–197, <https://doi.org/10.1016/j.jhydrol.2004.03.004>, 2004.
- Peng, B., Guan, K., Pan, M., and Li, Y.: Benefits of Seasonal Climate Prediction and Satellite Data for Forecasting U.S. Maize Yield, *Geophys. Res. Lett.*, 45, 9662–9671, <https://doi.org/10.1029/2018GL079291>, 2018.
- Pham, H. (Ed.): *Springer Handbook of Engineering Statistics*, Springer London, London, <https://doi.org/10.1007/978-1-84628-288-1>, 2006.
- 535 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., Chuang, H., Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M. P., van den Dool, H., Zhang, Q., Wang, W., Chen, M., and Becker, E.: The NCEP Climate Forecast System Version 2, *Journal of Climate*, 27, 2185–2208, <https://doi.org/10.1175/JCLI-D-12-00823.1>, 2014.
- Schepen, A., Everingham, Y., and Wang, Q. J.: On the Joint Calibration of Multivariate Seasonal Climate Forecasts from GCMs, *Monthly Weather Review*, 148, 437–456, <https://doi.org/10.1175/MWR-D-19-0046.1>, 2020.
- 540 Sheffield, J., Wood, E. F., Chaney, N., Guan, K., Sadri, S., Yuan, X., Olang, L., Amani, A., Ali, A., Demuth, S., and Ogallo, L.: A Drought Monitoring and Forecasting System for Sub-Sahara African Water Resources and Food Security, *Bull. Amer. Meteor. Soc.*, 95, 861–882, <https://doi.org/10.1175/BAMS-D-12-00124.1>, 2014.
- Slater, L. J., Villarini, G., and Bradley, A. A.: Evaluation of the skill of North-American Multi-Model Ensemble (NMME) Global Climate Models in predicting average and extreme precipitation and temperature over the continental USA, *Clim Dyn*, 53, 7381–7396, <https://doi.org/10.1007/s00382-016-3286-1>, 2019.
- 545 Strazzo, S., Collins, D. C., Schepen, A., Wang, Q. J., Becker, E., and Jia, L.: Application of a Hybrid Statistical–Dynamical System to Seasonal Prediction of North American Temperature and Precipitation, *Monthly Weather Review*, 147, 607–625, <https://doi.org/10.1175/MWR-D-18-0156.1>, 2019.
- Wanders, N., Bachas, A., He, X. G., Huang, H., Koppa, A., Mekonnen, Z. T., Pagán, B. R., Peng, L. Q., Vergopolan, N., Wang, 550 K. J., Xiao, M., Zhan, S., Lettenmaier, D. P., and Wood, E. F.: Forecasting the Hydroclimatic Signature of the 2015/16 El Niño Event on the Western United States, *Journal of Hydrometeorology*, 18, 177–186, <https://doi.org/10.1175/JHM-D-16-0230.1>, 2017.
- Webster, P. J. and Yang, S.: Monsoon and Enso: Selectively Interactive Systems, *Q.J Royal Met. Soc.*, 118, 877–926, <https://doi.org/10.1002/qj.49711850705>, 1992.
- 555 Wood, A. W. and Lettenmaier, D. P.: A Test Bed for New Seasonal Hydrologic Forecasting Approaches in the Western United States, *Bull. Amer. Meteor. Soc.*, 87, 1699–1712, <https://doi.org/10.1175/BAMS-87-12-1699>, 2006.
- Xu, L., Chen, N., Zhang, X., and Chen, Z.: A data-driven multi-model ensemble for deterministic and probabilistic precipitation forecasting at seasonal scale, *Clim Dyn*, 54, 3355–3374, <https://doi.org/10.1007/s00382-020-05173-x>, 2020a.
- 560 Xu, W., Fletcher, T. D., Burns, M. J., and Cherqui, F.: Real Time Control of Rainwater Harvesting Systems: The Benefits of Increasing Rainfall Forecast Window, *Water Resources Research*, 56, <https://doi.org/10.1029/2020WR027856>, 2020b.

Yang, S., Li, Z., Yu, J.-Y., Hu, X., Dong, W., and He, S.: El Niño–Southern Oscillation and its impact in the changing climate, *National Science Review*, 5, 840–857, <https://doi.org/10.1093/nsr/nwy046>, 2018.

565 Yossef, N. C., Winsemius, H., Weerts, A., van Beek, R., and Bierkens, M. F. P.: Skill of a global seasonal streamflow forecasting system, relative roles of initial conditions and meteorological forcing: Skill of a Global Seasonal Streamflow Forecasting System, *Water Resour. Res.*, 49, 4687–4699, <https://doi.org/10.1002/wrcr.20350>, 2013.

Yuan, X., Wood, E. F., and Liang, M.: Integrating weather and climate prediction: Toward seamless hydrologic forecasting: Seamless hydrologic forecast, *Geophys. Res. Lett.*, 41, 5891–5896, <https://doi.org/10.1002/2014GL061076>, 2014.

570 Yuan, X., Ma, F., Wang, L., Zheng, Z., Ma, Z., Ye, A., and Peng, S.: An experimental seasonal hydrological forecasting system over the Yellow River basin – Part 1: Understanding the role of initial hydrological conditions, *Hydrol. Earth Syst. Sci.*, 20, 2437–2451, <https://doi.org/10.5194/hess-20-2437-2016>, 2016.

Zhao, T., Bennett, J. C., Wang, Q. J., Schepen, A., Wood, A. W., Robertson, D. E., and Ramos, M.-H.: How Suitable is Quantile Mapping For Postprocessing GCM Precipitation Forecasts?, *J. Climate*, 30, 3185–3196, <https://doi.org/10.1175/JCLI-D-16-0652.1>, 2017.

575 Zhao, T., Wang, Q. J., and Schepen, A.: A Bayesian modelling approach to forecasting short-term reference crop evapotranspiration from GCM outputs, *Agricultural and Forest Meteorology*, 269–270, 88–101, <https://doi.org/10.1016/j.agrformet.2019.02.003>, 2019.

Zhao, T., Chen, H., Shao, Q., Tu, T., Tian, Y., and Chen, X.: Attributing correlation skill of dynamical GCM precipitation forecasts to statistical ENSO teleconnection using a set-theory-based approach, *Hydrology and Earth System Sciences*, 25, 5717–5732, 2021.

580