

Poor correlation between large-scale environmental flow violations and freshwater biodiversity: implications for water resource management and the freshwater planetary boundary

Chinchu Mohan^{1,2*}, Tom Gleeson^{2,3*}, James S Famiglietti^{1,4}, Vili Virkki⁵, Matti Kummu⁵, Miina Porkka^{5,6}, Lan Wang-Erlandsson^{7,8}, Xander Huggins^{1,2}, Dieter Gerten^{9,10}, Sonja C. Jähnig^{10,11}

¹ Global Institute for Water Security, University of Saskatchewan, Saskatoon, Saskatchewan, Canada.

² Department of Civil Engineering, University of Victoria, Victoria, British Columbia, Canada

³ School of Earth and Ocean Sciences, University of Victoria, Victoria, British Columbia, Canada

⁴ School of Environment and Sustainability, University of Saskatchewan, Saskatoon, Saskatchewan, Canada.

⁵ Water and Development Research Group, Aalto University, Espoo, Finland

⁶ Global Economic Dynamics and the Biosphere, Royal Swedish Academy of Sciences, Stockholm, Sweden

⁷ Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden

⁸ Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

⁹ Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, Potsdam, Germany

¹⁰ Humboldt-Universität zu Berlin, Geography Department and Integrative Research Institute on Transformations of Human–Environment Systems, Berlin, Germany

¹¹ Leibniz Institute of Freshwater Ecology and Inland Fisheries, Müggelseedamm 310, Berlin, Germany

*Correspondence to: Chinchu Mohan (chinchu.mohan@usask.ca), Tom Gleeson (tgleeson@uvic.ca)

ORCID corresponding authors: Chinchu Mohan: 0000-0001-7611-3392; Tom Gleeson: 0000-0001-9493-7707

Key Research Points

- No significant relationship between environmental flow (EF) violation and freshwater biodiversity indicators was found at global or ecoregion scales using globally consistent methods and currently available data, when not accounting for other factors affecting freshwater biodiversity.
- Several basins show a slight positive correlation between EF violation and biodiversity indicators, which could be attributed to the artificial introduction of non-native species.
- A generalized approach that incorporates EF considerations but ignores the lack of a significant EF-biodiversity relationship at large scales can underestimate the stress on the

ecosystem at smaller scales which correspond with eco-hydrological processes that determine ecological impacts from EF violation.

- Use of a globally aggregated blue water planetary boundary using biodiversity-based response variables is deceptive

Abstract

The freshwater ecosystems around the world are degrading, such that maintaining environmental flow¹ (EF) in river networks is critical to their preservation. The relationship between streamflow alterations and, respectively, EF violations², and freshwater biodiversity is well established at the scale of stream reaches or small basins ($\sim < 100 \text{ km}^2$). However, it is unclear if this relationship is robust at larger scales even though there are large-scale initiatives to legalize the EF requirement. Moreover, EFs have been used in assessing a planetary boundary³ for freshwater. Therefore, this study intends to conduct an exploratory evaluation of the relationship between EF violation and freshwater biodiversity at globally aggregated scales and for freshwater ecoregions. Four EF violation indices (severity, frequency, probability to shift to violated state, and probability to stay violated) and seven independent freshwater biodiversity indicators (calculated from observed biota data) were used for correlation analysis. No statistically significant negative relationship between EF violation and freshwater biodiversity was found at global or ecoregion scales. These findings imply the need for having a holistic bio-geo-hydro-physical approach in determining the environmental flows. While our results thus suggest that streamflow and EF may not be an only determinant of freshwater biodiversity at large scales, they do not preclude the existence of relationships at smaller scales or with more holistic EF

¹ Environmental flow (EF): “The quantity, timing, and quality of water flows required to sustain freshwater and estuarine ecosystems and the human livelihoods and well-being that depend on these ecosystems.” - Arthington et al., 2018

² EF violations: EF violations are deviations in streamflow beyond the upper and lower boundary of Environmental Flow envelopes (EFE). The EFE establish an envelope for acceptable EF deviations based on pre-industrial (1801-1860) stream discharge (See section 2.2 for more details)

³ Planetary boundary: Planetary boundary defines biogeophysical planetary scale boundaries for Earth system processes that, if violated, can irretrievably impair the Holocene-like stability of Earth system (see box 1 for more details)

methods (e.g., including water temperature, water quality, intermittency, connectivity etc.) or with other biodiversity data or metrics.

Keywords: Environmental flow violation, freshwater biodiversity, Global scale, freshwater ecoregions.

1.Introduction

Water resources are inarguably one of the most important natural resources in the Earth system for sustaining life. Nevertheless, these resources and their associated ecosystems are threatened by human actions (Bélanger and Pilling, 2019; Clausen and York, 2008; Vörösmarty et al., 2010; Wilting et al., 2017). Global freshwater covers up to 0.8% of the total Earth's surface (Gleick, 1996) and inhabits 6% of all the known species in the world including 40% of total fish diversity and nearly one third of all vertebrates (Lundberg et al., 2000). Since freshwater ecosystems have high species richness in a relatively small area and are exposed to a high level of pressure, they are more vulnerable to environmental change and human actions than any other ecosystems (Dudgeon et al., 2006). The rapid increase in the demand for natural resources is the fundamental cause for freshwater ecosystem degradation (Darwall et al., 2018). Anthropogenic climate change (Allan and Flecker, 1993; Darwall and Freyhof, 2016; Knouft and Ficklin, 2017; Meyer et al., 1999), overexploitation (Allan et al., 2005), water pollution (Albert et al., 2021; Dudgeon et al., 2006; Reid et al., 2019; Smith, 2003), flow alteration (Nilsson et al., 2005; Vorosmarty et al., 2000), habitat destruction (Dudgeon, 2001) and introduction of alien species (Gozlan et al., 2010; Vitule et al., 2009) are some of the manifestations of this increased demand which directly threatens the freshwater ecosystems. In addition, increased water impoundment in large dams and reservoirs has also led to an array of adversities to freshwater ecosystems ranging from habitat destruction to irregular flow alterations (Bergkamp et al., 2000). This situation is aggravated by increasing pressure on related Earth system functions, such as climate change and nutrient cycles, which are articulated by their respective transgressions in the planetary boundaries framework (Box 1) (Dudgeon, 2010). Freshwater ecosystem processes that were previously governed by natural Earth system facets such as temperature, rainfall, and relief are

now increasingly driven by demographic, social, and economic drivers (Clausen and York, 2008; Kabat et al., 2004; Tyson et al., 2002; Vitousek et al., 1997; Vörösmarty et al., 1997). Freshwater ecosystem health comprises both biotic factors like biodiversity and abiotic factors like habitat integrity. As any disruption in the abiotic factors is most likely to be reflected in the biotic status of the freshwater ecosystem, the scope of this paper is confined to the biotic dimension of the freshwater ecosystem (i.e., biodiversity) and not the health of the entire ecosystem.

There has been an increased recognition in recent decades for the need of maintaining a natural flow regime in streams to sustain healthy ecosystems. (Horne et al., 2017; Poff et al., 1997, 2017; Tickner et al., 2020; Tonkin et al., 2021). Despite the indispensable role of aquatic biodiversity in maintaining the quality of the system (Darwall et al., 2018), inclusion of such environmental flow (EF) in water management is often controversial, particularly in regions where freshwater availability is limited and is already a matter of severe competition. These competitions have led to an increasing trend in EF violation (insufficient streamflow than the recommended EF requirement; see section 2.1 for more details) in the past decade both in terms of severity and frequency (Virkki et al., 2022). This wakeup call has led to several international and national efforts to legalize EF requirements through large-scale EF management schemes (Arthington and Pusey, 2003; Richter et al., 1997, 2003). The Water and Nature Initiative (Smith and Cartin, 2011), the Brisbane declaration (Declaration, 2007), and the Global Action Agenda (Arthington et al., 2018) are some of these efforts. Nevertheless, there is a large gap in our understanding of the relationship between EF requirements and biodiversity responses at various spatial and temporal scales. Except for a few (Domisch et al., 2017; Xenopoulos et al., 2005; Yoshikawa et al., 2014), the majority of the studies exploring this relation were conducted at smaller scales (Anderson et al., 2006; Arthington and Pusey, 2003; Powell et al., 2008). Thus, there is a significant discrepancy in the scale at which these processes are understood versus the scale at which the policies are set (Thompson and Lake, 2010). Current knowledge of how the small-scale processes scale up (e.g., validation of large-scale EF hydrologic methods using local data) to a regional or global scale is thus limited, potentially undermining the scientific integrity of existing large-scale EF management schemes.

109

110 In order to scientifically underpin large scale EF policies, the existing assumption of the inverse
111 relationship between freshwater biodiversity response and EF violation must be tested at
112 regional and global scales (see Supplementary information S1 for more details). Therefore, in this
113 study, we evaluate the relationship between EF violation and freshwater biodiversity at two
114 different spatial scales (freshwater ecoregion, global) using four EF violation indices (frequency,
115 severity, probability to move to a violated state, and probability to stay violated) and seven
116 freshwater biodiversity indicators describing taxonomic, functional, and phylogenetic
117 dimensions of the biodiversity. The paper is not intended to be a definitive test on the
118 relationship between EF violation and aquatic biodiversity. It is rather intended to be an
119 exploratory analysis of the idea of conducting more detailed evaluations of the EF-biodiversity
120 relationship before formulating large scale EF management policies. The implications of the
121 findings for large-scale water management and the use of the relationship between
122 environmental flows and freshwater biodiversity (hereafter referred to as EF-biodiversity
123 relationship) in the planetary boundary framework (box 1) are also discussed.

124

Box 1: Introduction to blue water planetary boundary framework

The planetary boundaries framework proposed by Rockström et al. (2009) and further developed by Steffen et al. (2015) defines bio geophysical planetary scale boundaries for Earth system processes that, if violated, can irretrievably impair the Holocene-like stability of Earth system. The framework establishes scientifically determined safe operating limits for human perturbations through control and response variable relationships, under which humans and other life forms will coexist in equilibrium without jeopardizing the Earth's resilience. Nine planetary boundaries were defined to cover all independent significant Earth system processes. Out of the nine, the freshwater planetary boundary quantifies the safe limits of the terrestrial hydrosphere (Gleeson et al., 2020a, b).

The freshwater planetary boundary was originally defined using human water consumption as the control variable, set at 4000 km³/yr (with an uncertainty of 4000 to 6000 km³/yr) (Rockström et al., 2009). Gerten et al. (2013) proposed a bottom-up, spatially explicit quantification of EF violations as part of the water boundary, while Gleeson et al. (2020b) subdivided the water planetary boundary into six sub-boundaries and proposed possible control and response variables for each, with aquatic biosphere integrity (i.e., EF) as the potential control variable for a surface water sub-boundary. Quantitative evaluation of the strength and scalability of the identified control and response variables is still required.

2.Methodology and Data

The study is conducted at two spatially aggregated scales; 1) global and 2) ecoregion, for a historic time period of 30 years (1976 - 2005). All the underlying calculations were done at level 5 HydroBASIN (median basin area = 19,600 km²) (Lehner and Grill, 2013) and were aggregated to the corresponding spatial scale for further analysis. Level 5 HydroBASIN (also referred to as basin in this paper) was selected as the smallest spatial unit as it is the highest level of specificity that can be rasterized into a 0.5-degree resolution grid without significantly reducing the number of sub-basins smaller than a grid cell (Virkki et al., 2022). The EF violation indices were calculated using Virkki et al. (2022)'s novel Environmental Flow Envelope (EFE) framework, and biodiversity was represented by a combination of relative and absolute value indices.

2.1 Data

2.1.1 Streamflow data

Streamflow data used in the EFE (see section 2.2 for more details) definition were obtained from the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP) simulation phase 2b outputs of global daily discharge (available at <https://esg.pik-potsdam.de>) (Warszawski et al., 2014). Monthly streamflow data (averaged from the daily simulations) for two time periods were used in this study; 1) for the pre-industrial era (1800 - 1860), which is considered as the unaltered reference period (Poff et al., 1997) and 2) for the recent time period (1976 - 2005). These monthly

streamflow datasets were used to calculate EF violations. For calculating the EF violation indices, the estimated EFEs for each basin were obtained from Virkki et al. (2022). A total of 4 Global Hydrological Models (GHM) (H08 (Hanasaki et al., 2018), LPJmL (Schaphoff et al., 2018), PCR-GLOBWB (Sutanudjaja et al., 2018), WaterGAP2 (Müller Schmied et al., 2016)) were used to obtain the monthly streamflow data. Each GHM was forced with four different Global Circulation Models (GCM) outputs (GFDL-ESM2M (Dunne et al., 2012), HadGEM2-ES (Collins et al., 2011; Bellouin et al., 2011), IPSL-CM5A-LR (Dufresne et al., 2013), MICROC5 (Watanabe et al., 2010)). All the GHM outputs used in this study are extensively validated and evaluated in several previous studies (e.g., Zaherpour et al., 2018; Gädeke et al., 2020). Moreover, as part of the ISIMIP impact model intercomparison activity, all the GCM climate input data were bias corrected using compiled reference datasets covering the entire globe at 0.5 deg resolution (Frieler et al., 2017). Additionally, the GHM outputs are also validated using historical data to better fit reality (Frieler et al., 2017). Therefore, no additional validation of the data is done in this study.

The streamflow data were aggregated to the sub-basin scale according to level 5 HydroBASIN Version 1.0 (<https://www.hydrosheds.org/page/hydrobasins>) (Lehner and Grill, 2013). The data from ISIMIP 2b is representative of historical land use and other human influences including dams and reservoirs (Frieler et al., 2017). The maximum discharge cell value within the boundaries of each level 5 HydroBASIN is chosen to represent the outlet discharge value. Any violations within the outlet cell are regarded as indicative of the entire basin, even if conditions can differ in various areas within the level 5 HydroBASIN. As the spatial resolution of the study is level 5 HydroBASIN to allow a global analysis, we accept a certain homogenization of the local scale characteristics. See supplementary materials (see Supplementary information S.2) for more details on the datasets used in this study.

2.1.2 Freshwater biodiversity data

In addition to the streamflow data, data on fish diversity were also used in this study (Table 1). Freshwater biodiversity was evaluated using seven indices estimated from the observed biota data. The biodiversity indicators were obtained from international agencies or the literature. The

biodiversity indicators consisted of six indices of relative change in biodiversity and one index of absolute values of biodiversity.

a) Absolute biodiversity indicator

The absolute biodiversity indicator consisted of freshwater fish richness (FiR). The fish richness data was compiled and processed from 1436 published papers, books, grey literature and web-based sources published between 1960 and 2014 (Tedesco et al., 2017). They cover 3119 basins all over the world and account for 14953 fish species permanently or occasionally inhabiting freshwater systems. In addition to FiR, we used the RivFishTIME dataset by Comte et al (2021) – compiled from long-term riverine fish surveys from 46 regional and national monitoring programmes and from individual academic research efforts. Though the RivFishTIME dataset is highly spatially skewed towards the already data rich regions of Europe, North America (particularly United States of America) and Australia and temporally discontinuous, it is the only species-specific fish abundance time series data available and is useful to have an independent verification of the findings using FiR and relative biodiversity indicators.

b) Relative biodiversity indicators

The Relative biodiversity indicators consisted of six freshwater fish facets. Six key facets of freshwater fish - taxonomic, functional, and phylogenetic diversity (TR, FR, PR respectively), as well as dissimilarity of each of the three groups (TD, FD, PD respectively)- were used in this analysis to construct a holistic picture of the state of aquatic biodiversity (see Fig. 1 in Su et al., 2021 for more details on fish facets calculations). Each facet indicates the change in the corresponding biodiversity component compared to the 18th century (roughly pre-industrial era). The taxonomic facets measure the occurrence of fish in a riverine system. Functional facets are calculated using the morphological characteristics of each species that are linked to feeding and locomotive functions which in turn relates to larger ecosystem functions like food web control and nutrition transport. Phylogenetic facets measure the total length of branches linking all species from the assemblage on the phylogenetic tree. The richness component of the three categories calculates the diversity among the assemblage whereas the dissimilarity accounts for the difference between each pair of fish assemblage in one realm. All six fish facets were

calculated at basin scale (2465 river basins) covering 10682 fish species all over the world. The scale at which the fish facets are estimated, not necessarily align with the scale at which the EF violations are estimated in all cases. The basin scale facet estimates were then matched with corresponding EF violation indices using different aggregation/data matching methods (see section 2.4 for more details). All six facets are available as a single delta change in time and do not cover multiple timesteps.

Table 1. Details of different data used in this study

Data	Spatial resolution (extent)	Temporal resolution (extent)	Source/Reference
Aquatic fish richness data	30 arc second (3119 drainage basins; ~80% of Earth's land)	Temporal aggregate from data compiled from reports between 1960 and 2014	Observed/Measured data Tedesco et al. (2017)
Freshwater fish facets	Basin scale (2465 drainage basins)	Representative of 2015 (change compared to preindustrial era)	Derived from observed data Su et al. (2021)
RivFishTIME dataset ⁴	Stream reach (11386 sampling location)	1951 -2019 ⁵	Comte et al., 2021
EFE	Aggregated to Level 5 HydroBASIN (global)	Monthly (Pre-industrial: 1801-1860)	Model calculated Virkki et al. (2022)
Streamflow	Aggregated to Level 5 HydroBASIN (global)	Monthly (Pre-industrial: 1801-1860, Current: 1976-2005)	Model calculated Warszawski et al. (2014)
Basin boundaries	Level 5 HydroBASIN (global)	Not applicable	Lehner and Grill (2013)

⁴ Results only shown in Supplementary Information (see section S8 in Supplementary Information)

⁵ Variable for each species and sampling site. Each time-series has a minimum of two-year survey (mean = 8 years).

2.2 Environmental flow violation estimation

The EFE framework proposed by Virkki et al. (2022) is used to evaluate EF violations in this study. The EFE framework establishes an envelope of variability constrained by discharge limits beyond which flow in the streams may not meet the freshwater biodiversity needs (Virkki et al., 2022). EFE uses pre-industrial (1801-1860) stream discharge to establish an upper and lower boundary for EF deviations at monthly time steps. This EFE is used to define the EF violation at Level 5 HydroBASIN scale. The EF violations were calculated as median ensemble of four Global Hydrological Models (GHM) (H08, LPJmL, PCR-GLOBWB, WaterGAP2) and mean ensemble of four Global Circulation Models (GCM) (GFDL-ESM2M, HadGEM2-ES, IPSL-CM5A-LR, MICROC5). Moreover, five different EF calculation methods (Smakhtin method (Smakhtin et al., 2004), Tennant method (Tennant, 1976), Q90-Q50 (Pastor et al., 2014), Tessmann method (Tessmann, 1979) and Variable Monthly Flow method (Pastor et al., 2014)) were also used in the EFE derivation (see Supplementary Information, Table S3 for more information on EF methods) (Virkki et al., 2022). This approach addresses the uncertainty related to the outputs of models and may eliminate the largest model-related extremes that might cause results to be distorted (Virkki et al., 2022). In spite of the uncertainty in hydrological estimates generated by different models, a simple ensemble matrix often produces acceptable discharge and therefore also EF estimates at larger scales because the bias of the individual models is removed (Zaherpour et al., 2018). Moreover, all the basins with Mean Annual Flow (MAF) $< 10 \text{ m}^3/\text{s}$ were excluded due to high uncertainty in EFE and streamflow estimates (Gleeson et al., 2020a; Steffen et al., 2015; Virkki et al., 2022). After this exclusion, a total of 3906 basins were considered for further analysis. However, many low flows are seasonally observed, such that MAF may be quite large due to elevated wet season flows, with extremely low flows during a dry season (e.g., Eel River basin, California) making it difficult to model. In such cases with higher intra annual flow variability, it is appropriate to consider more detailed discharge data (seasonal/sub annual) to gain more insight into the flow modelling uncertainties.

Here we evaluate the EF violation by defining four different EF violation indices: 1) violation severity (S), violation frequency (F), probability to shift to a violated state (P.shift) and probability to stay violated (P.stay). Out of the four EF violation indicators, two (S and F) were a modification from Virkki et al. (2022) and the two (P.shift and P.stay) were calculated based on the current EFE deviations from Virkki et al. (2022). P.shift and P.stay measures the likelihood of a given year to shift or stay in a violated state. The state of a basin (violated or non-violated) was identified at an annual time step and the mean probability to shift or remain in that state is calculated.

The detailed definitions of the EF violation indicators are as follows.

1) Violation severity (S): The annual violation severity was calculated as the absolute mean of the magnitude of EF deviation from the EFE lower or upper bound in all the violated months. The magnitude of violation is based on the violation ratio proposed by Virkki et al. 2022 (See Table S4 in supplementary information). The normalized value of S is used in this study.

2) Violation frequency (F): Frequency of violation is a measure of the proportion of months a basin has violated the EFE lower or upper bound in a year. Frequency is calculated as the percentage of violated months per year. The normalized value of F is used in this study.


3) Probability to shift to a violated state (P.shift): The P.shift is defined in this paper as the probability of a basin to shift to a violated state from a non-violated state (Eq. 1). This indicator along with P.stay gives a measure of the stability of violation in each level 5 HydroBASIN. The violated/non-violated state of a basin is calculated annually based on the violations in the low flow months. If a basin violates EFE lower or upper bound for at least three consecutive months during the low flow period ($Q < 0.4MAF$) in a year, then the basin is considered to be in a violated state.

$$P. shift = \frac{\text{number of years shifted to violated state (i.e. year } i \text{ is violated and year } i-1 \text{ is not)}}{\text{total number of years}} \quad (1)$$


- 4) Probability to stay violated (P.stay): Once shifted to a violated state, the tendency of a basin to remain in that state or switch to a non-violated state is determined by this indicator. If a basin has a higher P.stay (closer to 1) then the basin continues to remain in the violated state for a longer time before switching to a non-violated state (Eq 2). Whereas the basins with lower P.stay (closer to 0) tend to remain in the violated state only for a brief period of time. In other words, the number of consecutive violated years is much lower for basins with lower P.stay value.

$$P.stay = \frac{\text{number of violated years with at least one consecutive year violated}}{\text{total number of violated years}} \quad (2)$$

(a) Environmental Flow Envelope (EFE) calculation



(b) EF violation indicators calculation



(c) EF-biodiversity relation evaluation




Fig. 1 Methodology outline for (a, b) EF violation indicators calculation and (c) EF-biodiversity relationship evaluation

2.3. Relationship between environmental flow violations and freshwater biodiversity

The relationship between freshwater biodiversity and EF violation was evaluated using regression analysis. None of the relationships explored in this study exhibited any nonlinearity and hence first order single variate and multivariate linear regression analysis was opted for this study for

reasons of parsimony and to achieve reasonable correlation accuracy. Further analysis was carried out by aggregating the level 5 HydroBASIN scale values to global level, WWF's Freshwater ecoregions major habitat type scale (results given in SI) (Abell et al., 2008) and G200 freshwater ecoregion level (Olson and Dinerstein, 2002). The G200 freshwater ecoregion is a subset of WWF's freshwater ecoregion that includes only the biodiversity hotspots. Seven freshwater ecoregions in ecologically important regions were studied, and the EF-biodiversity relationship was evaluated separately for each ecoregion type. Aggregating to major ecoregion types of accounts for some data's natural/spatial variability, in addition to using an analysis of global data.

One of the major challenges in conducting an aggregated evaluation was the discrepancy in the spatial resolution at which the EF violation indices and various biodiversity indicators and the loss of heterogeneity. Aggregation of any scale will lead to some level of homogenization of the data. A reach-by-reach evaluation will be an ideal solution to capture all the heterogeneity. However, this is not very practical for a global study due to data and computational limitations. Therefore, to partially address this challenge, two different aggregation/data matching methods were employed; case-1) matching level 5 HydroBASIN data (EF violation indices) to biodiversity data and case-2) matching biodiversity data to level 5 HydroBASIN (See supplementary information (SI); Section S5). In the first case every level 5 HydroBASIN (EF violation indices) is matched with the biodiversity data point nearest centroid. Whereas in the second case there can be three different scenarios (See SI; Fig. S4): 1) biodiversity basin is smaller than level 5 HydroBASIN; in that case all the biodiversity basins within one level 5 HydroBASIN were matched with the same EF violation value, 2) when biodiversity basin is equal in size to level 5 HydroBASIN; in this case biodiversity basins and level 5 HydroBASIN had a one-to-one match, 3) biodiversity basin is larger than level 5 HydroBASIN. In the last case, two methods were used for data mapping 1) Outlet matching: where each biodiversity basin is mapped with EF violation value from the level 5 HydroBASIN closest to the outlet and 2) Mean matching: each biodiversity basin is mapped with the mean EF violation values of all level 5 HydroBASIN within it. Data matching methods were employed to partially understand the uncertainty due to scale discrepancy between datasets. As

the results are insensitive to the aggregation method, only the results using case 1 (matching level 5 HydroBASIN data to biodiversity data) are discussed in this paper.

3.Results and Interpretations

3.1 Evaluating EF violation drivers and characteristics

The majority of basins face some kind of EF violation (either in terms of severity or frequency or with higher probabilities to shift and/or stay violated) (Fig. 2). Between 1976 and 2005, 17% and 45% of basins, respectively, experienced violation frequency (F) greater than 3 months/year and severity (S) greater than 20% from the EFE lower or upper bound (normalized violation index ≥ 0.25) (Fig.2 a, b). Additionally, 33% of basins have a higher chance of shifting ($P.\text{shift} \geq 0.5$; i.e., 33% basins have over 50% probability to shift to a violated state) to a violated state (Fig.2 c, d). EF violations are very frequent and severe in mostly arid/semi-arid regions such as the Middle East, Pakistan, India, Australia, Sahara, Sub-Saharan Africa, Southern Africa, and the southernmost part of North America. On the other hand, regions with higher probability to shift to a violated state ($P.\text{shift}$) were not limited to the low precipitation and low streamflow regions.

Although the majority of regions with high $P.\text{shift}$ values were arid or semi-arid, some exceptions included Southeastern Asia and Central South America. The non-arid regions with higher $P.\text{shift}$ also have extremely high water withdrawal in all sectors (agriculture, domestic and industry). This spatial concurrence suggests that human activities, as well as hydroclimatic influences, play a significant role in deciding a region's $P.\text{shift}$. However, once in the violated state, the flow variability regimes in the catchment determine the probability of remaining ($P.\text{stay}$) in the violated state. Catchments with highly variable flow regimes (i.e., receive most of the annual flow as floods; see SI for classification map; Fig. S2) have higher probability to stay violated once shifted whereas catchments with stable flow regimes (year-round steady high baseflow) have a higher tendency to revert to a non-violated state. An example of this behavior can be seen in the Australian basins. Though, almost all the Australian basins have a very high $P.\text{shift}$, only the highly variable flow regime northern catchments had a higher probability to stay violated. Despite

having an exceedingly high P.shift, the southern stable catchments swiftly shift back to a non-violated state.




Fig. 2 Four measures of Environmental Flow Envelope (EFE) lower or upper bound violation estimated using ensemble median of four Global hydrological models; a) Normalized frequency of violation, b) Normalized severity of violation, c) Probability to shift to a violated state from a non-violated state and d) Probability to stay violated once shifted to a violated state.

3.2 Relationship between EF violation and freshwater biodiversity

The aggregated analysis was carried out at global and ecoregion scales. Multiple aggregation methods (section 2.3) yielded comparable results, therefore only the case 1 (level 5 HydroBASIN matched with biodiversity data) results are discussed further (see supplementary material Fig. S5 and S6 for results using other aggregation methods). At the global scale, none of the biodiversity indicators correlated (significance of p value < 0.05) with any EF violation indices (Fig. 2). The biodiversity indicators were not exhibiting any strong trend in either positive or negative direction. The correlation coefficient value (R value) for the remaining biodiversity indicators

ranges only from -0.2 to 0.17 (Fig. 3 b). The three fish dissimilarity facets (TD, FD, and PD) show slight negative correlation whereas the richness facets (TR, FR, and PR) display a slight positive correlation with EF violation. The positive correlation of the richness indicators is attributed to an overall increase in the assemblage in most of the basins despite the increase in EF violation. Moreover, (relative) TR and (absolute) FiR were showing opposite trends. The positive trend in TR could be attributed to changes involving nonnative species, whereas the FiR describes the current deteriorated state. The increase in the fish assemblage over time was verified using an independent dataset RivFishTIME (see SI; Fig. S8, Fig. S9) (Comte et al., 2021). The increase in the fish richness facets primarily stems from the introduction of alien species introduced into streams for commercial purposes (Su et al., 2021). The invasion of alien species can tamper with the existing natural ecosystem equilibrium resulting in further degradation of the overall ecosystem health. The results using RivFishTIME data sets were also consistent with the findings using FiR and six relative biodiversity indicators and there was no significant correlation between EF violation indicators and fish abundance data over time (see results for five selected fish species based on data completeness and geographical distribution in Supplementary Information section S8; Fig. S8).

Correlations between EF and biodiversity are generally weak at the scale of G200 freshwater ecoregions as well (see Section 2.2, (Olson and Dinerstein, 2002)). In G200 freshwater ecoregions (see SI; Table S6 for full freshwater ecoregion results) the nature of the EF-biodiversity relationships was highly varying between different ecoregions (Fig 4). In large lakes, large rivers and small lakes, Su et al. (2021) fish richness facets were showing a strong and significant positive correlation with most of the EF violation indices. The increase in biodiversity despite increase in EF violation could be a signal of introduction of nonnative species for commercial purposes. Whereas, in large rivers, large river deltas and xeric basins, the dissimilarity indices, FiR show negative correlation. However, in most ecoregions, the EF-biodiversity relationship is insignificant (p value >0.05). Similar analysis using different aggregation/scale matching methods also yielded comparable results at G200 ecoregion scale (see Fig. S5 and Fig. S6 in Supplementary Information). In addition to this, the multivariate regression analysis results (Fig. 5) also show

very low correlation between EF violation indicators and biodiversity indices in most G200 ecoregion, except in small lakes where the coefficient of determination is between 0.25 - 0.4 for the richness indicators (TR, FR, PR). The mean coefficient of determination (r^2) is approximately 0.1. These results corroborate the above findings that EF violations are not significantly inversely correlated with biodiversity, regardless of ecoregions with the current dataset.




Fig. 3 Scatter between EF violation indices and biodiversity indices with linear fit and corresponding R value at globally aggregated scale.

Note: This figure represents results from case 1 (level 5 HydroBASIN matched with biodiversity data). The results of other aggregation methods are given in SI (Fig. S5 and S6).

Abbreviations: F - Frequency of violation; S-Severity of violation; P.shift-Probability to shift to a violated state; P.stay-Probability to stay in a violated state; FiR-Fish richness; TR-Taxonomic richness; FR-Functional richness; PR-Phylogenetic richness; TD-Taxonomic dissimilarity; FD-Functional dissimilarity; PD-Phylogenetic dissimilarity




Fig.4 (a) Spatial distribution of different G200 freshwater ecoregions and (b-h) the correlation between EF violation indices and freshwater biodiversity indicators for different G200 freshwater ecoregions.

Note: The results for all the WWF freshwater ecoregions are given in SI (SI section S.7).

Abbreviations: F - Frequency of violation; S-Severity of violation; P.shift-Probability to shift to a violated state; P.stay-Probability to stay in a violated state; FiR-Fish richness; TR-Taxonomic richness; FR-Functional richness; PR-Phylogenetic richness; TD-Taxonomic dissimilarity; FD-Functional dissimilarity; PD-Phylogenetic dissimilarity




Fig. 5 Coefficient of correlation (R^2) for multivariate regression between EF violation indicators and biodiversity indices. Each row represents on biodiversity indicator and each column represents one G200 ecoregion

4. Discussion

The findings from this study indicate that the EF-biodiversity relationship is poorly correlated at global or ecoregion scales with currently available data and methods. The most likely explanation for the lack of correlation is the overwhelming heterogeneity of the freshwater ecosystems - e.g., with some freshwater species being more susceptible to variations in flow than others (Poff and Zimmerman, 2010) - which is not adequately represented in the used spatial resolution (level 5 HydroBASIN). Moreover, when it comes to a larger-scale relationship, several other factors like climate change (Davies, 2010; Poff et al., 2002), river fragmentation (Grill et al., 2015; Herrera-R et al., 2020), large-scale habitat degradation (Moyle and Leidy, 1992), landscaping/river scaping (Allan et al., 2005), alien species (Leprieur et al., 2008, 2009; Villéger et al., 2011) and water pollution (Brooks et al., 2016; Shesterin, 2010) can also impact the freshwater ecosystem in multiple ways. Thus, at Earth system level, other interlinked factors potentially confound the impact of EF violation on biodiversity degradation.

4.1 Implications for water management

The lack of correlation between EF violation and freshwater biodiversity has implications for large-scale water management. A generalized large scale EF approach can underestimate the stress on the ecosystem at a smaller scale where the actual action is taking place. It is undeniable that adequate flow is essential for maintaining freshwater ecosystems. Nonetheless, the current generalized EF estimation methods need further refinement to adequately capture this importance. The global hydrological EF methods are often validated using locally calculated EF requirement values (Pastor et al., 2014) with the assumption of adequate scalability in the EF-biodiversity relationship. However, more holistic EF estimation methods combining hydrological, hydraulic, habitat simulation methods, and expert knowledge (Poff and Zimmerman, 2010; Shafroth et al., 2010) are essential to ensure a healthy freshwater biodiversity. The policies and decisions taken at various scales need a more dynamic framework, where different dominant drivers of ecosystem degradation can be prioritized based on particular cases. For instance, an integrated EF indicator which encompasses quantity, quality, and timeliness of water in the streams will be a better hydrologic indicator to evaluate freshwater ecosystem health than an indicator which accounts only for quantity. Moreover, when making water management decisions, care must be given to account for the temporal and spatial heterogeneity in the ecosystem dynamics.

Although there are some coordinated scientific efforts such as ELOHA (Ecological Limits of Hydrologic Alterations) (Poff et al., 2010) to provide a holistic framework for EF estimation, its scientific complexity and high implementation cost constrains its use around the world (Richter et al., 2012). For example, several European countries like Romania, Czech Republic, Serbia, and Luxembourg use a national level static method to define minimum environmental flows (Linnansaari et al., 2012). Similarly, other jurisdictions use the presumptive standards proposed by Richter et al. (2012) to establish a legal basis for EF protection. These presumptive standards limit hydrologic modifications to a percentage range of natural or historic flow variability. One example of such a case, the North Carolina's Environmental Flow Science Advisory Board uses a presumptive standard of 80-90% of the instantaneous modeled baseline flow as the EF

requirement (NCEFSAB, 2013). The limitation of such a practice is the incorrect presumption of uniformity in the EF needs over a larger region. Therefore, we recommend the application of holistic indicators at these large scales (covering all river stretches and tributaries) rather than using simplified hydrologic-only metrics of EF (violation). However, the authors also acknowledge the limits in implementation of a more dynamic EF framework in data limited regions. Programs for more monitoring and data collection and improved, more holistic modeling methods using more/better data need to be implemented in those regions. Thus, applying a holistic framework like ELOHA could be made possible and can capture the heterogeneity in the EF-biodiversity relationship.

4.2 Implications for a water planetary boundary

The current rationale in using EF in the water planetary boundary relationship is based on the assumption of its universal relationship with freshwater biodiversity. However, with the currently available data and methods the findings for EF-biodiversity relationship are inconclusive. Moreover, due to the heterogeneity of biodiversity response over time and space, the trend in any aggregate scale is likely to remain relatively constant instead of showing any discernible tipping point (Brook et al., 2013). We suggest that to reconsider the use of environmental flows in defining water planetary boundaries, given the higher degree of heterogeneity and lack of strength in the ecosystem function-biodiversity relationship. Some of the potential reasons for the reconsideration are, firstly, freshwater biodiversity may not have pan-regional or "continental-planetary" scale threshold dynamics, and its link with EF violation might be inadequate to represent the finer scale variations. Secondly, resource distribution and human impact heterogeneity suggest the need for regional boundaries as proposed by Steffen et al. (2015). Thirdly, EF calculation methods used in the current regional/planetary boundary definition are highly restricted to hydrological methods which may not be adequate to capture the biodiversity status. A regional boundary transgression can occur even well within planetary-level safe limits (Brook et al., 2013; Nykvist et al., 2017). Therefore, for an overly complex biophysical relationship like the EF-biodiversity where multiple shift states are possible, it is difficult to prioritize and manage critical regions without a regional/local boundary.

4.3 Limitations and ways forward

1) **Data scarcity:** Even though this study uses state of the art global hydrological models and best available global estimates of EF requirements, freshwater ecological data were limited to freshwater fish. Other than these, several other taxa like crayfish and other benthic invertebrates, phytoplankton, or zooplankton are also significant in determining the proper functioning of a freshwater ecosystem (AL-Budeiri, 2021; Domisch et al., 2017; Nyström et al., 1996). However, due to lack of global data, these taxa are not included in this study. To better examine the relationship, global datasets for other freshwater biodiversity metrics are urgently needed.

2) **Discrepancy in data resolution:** The spatial and temporal resolutions at which the EF violation is estimated here, and the biodiversity indicators measured/calculated are inconsistent. The basic spatial measuring unit of the biodiversity is sometimes greater or lesser than the basin size at which EF is measured. This discrepancy could have some impact on the results. However, in this study several resolution matching methods were used to account for this uncertainty. Therefore, more detailed data with better-matching scales are needed to overcome this limitation.

3) **Lack of multi-driver interaction:** In this study, we consider the impact of EF violations on biodiversity as an independent relationship. In reality, this might not be the case. Other drivers of ecosystem degradation like land use change, habitat loss, stream modifications and geographical disconnection can influence the EF-biodiversity relationship. These interactions were outside the scope of this study but should be taken into account in follow up studies.

4) **Simplified representation of human interference with freshwater systems:** The role of humans in impairing the ecosystem balance is represented here based on how human water withdrawals violate hydrologically defined EF. Other human disturbances are thus not accounted for, such as aquatic habitat degradation through change in land use, artificial introduction of

nonnative species, and non-point pollution from agriculture. Moreover, this study does not distinguish the climate driven impact on EF violation from the anthropogenic impacts.

5) Exclusion of impact of dams: The dams are indeed a large contributing factor to the results uncertainty. The dam regulated rivers may have a significantly different effect on biodiversity compared to free-flowing rivers. The ISIMIP data used to calculate EF violations considers the effects of large dams on streamflow. However, to explicitly isolate the effects of dams in this analysis from other drivers, the information on dam operation schemes for each sub-basin would be necessary and this would require a paper on its own. Therefore, the effects of the dams are incorporated in this study but are not explicitly analyzed separately from other drivers.

5. Summary and Conclusion

The relationship between EF violations and freshwater biodiversity is evaluated at globally aggregated levels in this study. No significant relationship between EF violation and freshwater biodiversity indicators was found at global or ecoregion scale using globally consistent methods and currently available data. Relationships may exist at smaller scales and could potentially be identified with more holistic EF methods including multiple factors (e.g., temperature, water quality, intermittency, connectivity) and more extensive freshwater biodiversity data. The single negative result is not a final say but it is a call for conducting more study on existing generalized and well applied methods.

The paper is not intended to be a definitive test on the relationship between EF and aquatic biodiversity but more to be an exploratory analysis to tests a widely used but rarely verified assumption on the relationship at global and ecoregion scale. The lack of correlation in the EF-biodiversity relationship found in this study suggests taking particular care when developing macro-scale EF policies (regional and above), and further implies that the conceptualization of a blue water planetary boundary ought to rest upon a broader set of relationships between hydrological processes and Earth system functioning. At larger scales, the enormous spatial and

temporal heterogeneity in EF-biodiversity relationship motivates a holistic estimation of EF grounded on ecosystem dynamics.

Data Availability

The data used in this study are temporarily made available at https://drive.google.com/drive/folders/1dXYByen5fcUqCQl3R4E0baCorpMwqN_q?usp=sharing

All data to reproduce the analysis in this manuscript will be made available via., University of Victoria data repository, DataVerse after the manuscript is accepted. Additionally, all the codes used in this study will also be made available vis. GitHub.

Author Contribution

CM, TG, JSF devised the conceptual and analysis framework of this study with inputs from MK, MP, and VV. VV performed the EFE calculation with help from MK and MP. CM performed the biodiversity data compilation and EF-biodiversity analytical evaluation with help from TG, JSF and XH. CM performed the final analysis and produced the results and visualization shown in the study, discussing together with TG, JSF, XH, MK, MP, VV and LWE. TG, JSF, MK, MP, VV, LWE, XH, DG and SCJ contributed to paper writing and the interpretation of the results. CM took the lead in writing the manuscript. All authors provided critical feedback and helped shape the research, analysis, and manuscript.

Compelling Interests

The authors declare no competing interests.

Acknowledgement

Authors acknowledge various funds that made this research possible. CM received funding from Canada First Research Excellence Fund (CFRE); MK received funding from Academy of Finland funded project WATVUL (grant no. 317320), Academy of Finland funded project TREFORM (grant no. 339834), and European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 819202). VV received funding from

Aalto University School of Engineering Doctoral Programme and European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 819202). SCJ acknowledges funding through the Leibniz Association for the project Freshwater Megafauna Futures.

Supplementary Information

The supplementary information is submitted separately.

Reference

Abell, R., Thieme, M.L., Revenga, C., Bryer, M., Kottelat, M., Bogutskaya, N., Coad, B., Mandrak, N., Balderas, S.C., Bussing, W. and Stiassny, M.L., 2008. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience*, 58(5), pp.403-414. Albert, J. S., Destouni, G., Duke-Sylvester, S. M., Magurran, A. E., Oberdorff, T., Reis, R. E., Winemiller, K. O., and Ripple, W. J.: Scientists' warning to humanity on the freshwater biodiversity crisis, *Ambio*, 50, 85–94, <https://doi.org/10.1007/s13280-020-01318-8>, 2021.

AL-Budeiri, A. S.: The role of zooplankton in the pelagic food webs of tropical lakes, PhD Thesis, University of Leicester, 2021.

Allan, J. David, and Alexander S. Flecker. "Biodiversity conservation in running waters." *BioScience* 43, no. 1: 32-43, 1993.

Allan, J. D., Abell, R., Hogan, Z., Revenga, C., Taylor, B. W., Welcomme, R. L., and Winemiller, K.: Overfishing of inland waters, *BioScience*, 55, 1041–1051, [https://doi.org/10.1641/0006-3568\(2005\)055\[1041:OOIW\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2005)055[1041:OOIW]2.0.CO;2), 2005.

Anderson, K. E., Paul, A. J., McCauley, E., Jackson, L. J., Post, J. R., and Nisbet, R. M.: Instream flow needs in streams and rivers: the importance of understanding ecological dynamics, *Frontiers in Ecology and the Environment*, 4, 309–318, [https://doi.org/10.1890/1540-9295\(2006\)4\[309:IFNISA\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)4[309:IFNISA]2.0.CO;2), 2006.

Arthington, A. H. and Pusey, B. J.: Flow restoration and protection in Australian rivers, *River research and applications*, 19, 377–395, <https://doi.org/10.1002/rra.745>, 2003.

Arthington, A. H., Bhaduri, A., Bunn, S. E., Jackson, S. E., Tharme, R. E., Tickner, D., Young, B., Acreman, M., Baker, N., Capon, S., Horne, A. C., Kendy, E., McClain, M. E., Poff, N. L., Richter, B. D., and Ward, S.: The Brisbane declaration and global action agenda on environmental flows (2018), 6, 2018.

594 B  langer, J. and Pilling, D.: The state of the world’s biodiversity for food and agriculture, FAO
595 Commission on Genetic Resources for Food and Agriculture Assessments, 2019.

596 Bellouin, N., Collins, W. J., Culverwell, I. D., Halloran, P. R., Hardiman, S. C., Hinton, T. J., Jones, C.
597 D., McDonald, R. E., McLaren, A. J., and O’Connor, F. M.: The HadGEM2 family of met office
598 unified model climate configurations, *Geoscientific Model Development*, 4, 723–757, 2011.

599 Bergkamp, G., McCartney, M., Dugan, P., McNeely, J., and Acreman, M.: Dams, ecosystem
600 functions and environmental restoration, *Thematic review II*, 1, 1–187, 2000.

601 Brook, B. W., Ellis, E. C., Perring, M. P., Mackay, A. W., and Blomqvist, L.: Does the terrestrial
602 biosphere have planetary tipping points?, *Trends in Ecology & Evolution*, 28, 396–401,
603 <https://doi.org/10.1016/j.tree.2013.01.016>, 2013.

604 Brooks, B. W., Lazorchak, J. M., Howard, M. D. A., Johnson, M.-V. V., Morton, S. L., Perkins, D. A.
605 K., Reavie, E. D., Scott, G. I., Smith, S. A., and Steevens, J. A.: Are harmful algal blooms becoming
606 the greatest inland water quality threat to public health and aquatic ecosystems?, *Environmental*
607 *toxicology and chemistry*, 35, 6–13, <https://doi.org/10.1002/etc.3220>, 2016.

608 Clausen, R. and York, R.: Global biodiversity decline of marine and freshwater fish: A cross-
609 national analysis of economic, demographic, and ecological influences, *Social Science Research*,
610 37, 1310–1320, <https://doi.org/10.1016/j.ssresearch.2007.10.002>, 2008.

611 Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes,
612 J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O’Connor, F., Rae, J., Senior, C., Sitch, S.,
613 Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System
614 model – HadGEM2, *Geoscientific Model Development*, 4, 1051–1075,
615 <https://doi.org/10.5194/gmd-4-1051-2011>, 2011.

616 Comte, L., Carvajal-Quintero, J., Tedesco, P. A., Giam, X., Brose, U., Er  s, T., Filipe, A. F., Fortin,
617 M.-J., Irving, K., Jacquet, C., Larsen, S., Sharma, S., Ruhi, A., Becker, F. G., Casatti, L., Castaldelli,
618 G., Dala-Corte, R. B., Davenport, S. R., Franssen, N. R., Garc  a-Berthou, E., Gavioli, A., Gido, K. B.,
619 Jimenez-Segura, L., Leit  o, R. P., McLarney, B., Meador, J., Milardi, M., Moffatt, D. B., Occhi, T. V.
620 T., Pompeu, P. S., Propst, D. L., Pyron, M., Salvador, G. N., Stefferud, J. A., Sutela, T., Taylor, C.,
621 Terui, A., Urabe, H., Vehanen, T., Vitule, J. R. S., Zeni, J. O., and Olden, J. D.: RivFishTIME: A global
622 database of fish time-series to study global change ecology in riverine systems, *Global Ecology*
623 *and Biogeography*, 30, 38–50, <https://doi.org/10.1111/geb.13210>, 2021.

624 Darwall, W., Bremerich, V., De Wever, A., Dell, A. I., Freyhof, J., Gessner, M. O., Grossart, H.-P.,
625 Harrison, I., Irvine, K., and J  hnig, S. C.: The Alliance for Freshwater Life: A global call to unite
626 efforts for freshwater biodiversity science and conservation, *Aquatic Conservation: Marine and*
627 *Freshwater Ecosystems*, 28, 1015–1022, 2018.

628 Darwall, W. R. and Freyhof, J.: Lost fishes, who is counting? The extent of the threat to freshwater
629 fish biodiversity, *Conservation of freshwater fishes*, 1–36, 2016.

630 Davies, P. M.: Climate change implications for river restoration in global biodiversity hotspots,
631 *Restoration Ecology*, 18, 261–268, 2010.

632 Declaration, B.: The Brisbane Declaration: environmental flows are essential for freshwater
633 ecosystem health and human well-being, in 10th International River Symposium, Brisbane,
634 Australia, 3–6, 2007.

635 Domisch, S., Portmann, F. T., Kuemmerlen, M., O’Hara, R. B., Johnson, R. K., Davy-Bowker, J.,
636 Baekken, T., Zamora-Muñoz, C., Sáinz-Bariáin, M., and Bonada, N.: Using streamflow
637 observations to estimate the impact of hydrological regimes and anthropogenic water use on
638 European stream macroinvertebrate occurrences, *Ecohydrology*, 10, e1895, 2017.

639 Dudgeon, D.: Fisheries: pollution and habitat degradation in tropical Asian rivers, *Encyclopaedia*
640 *of Global Environmental Change*, 3, 2001.

641 Dudgeon, D.: Prospects for sustaining freshwater biodiversity in the 21st century: linking
642 ecosystem structure and function, *Current Opinion in Environmental Sustainability*, 2, 422–430,
643 2010.

644 Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z.-I., Knowler, D. J., Lévêque, C.,
645 Naiman, R. J., Prieur-Richard, A.-H., Soto, D., and Stiassny, M. L.: Freshwater biodiversity:
646 importance, threats, status and conservation challenges, *Biological reviews*, 81, 163–182, 2006.

647 Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki,
648 S., Bellenger, H., and Benshila, R.: Climate change projections using the IPSL-CM5 Earth System
649 Model: from CMIP3 to CMIP5, *Climate dynamics*, 40, 2123–2165, 2013.

650 Dunne, J. P., John, J. G., Adcroft, A. J., Griffies, S. M., Hallberg, R. W., Shevliakova, E., Stouffer, R.
651 J., Cooke, W., Dunne, K. A., and Harrison, M. J.: GFDL’s ESM2 global coupled climate–carbon earth
652 system models. Part I: Physical formulation and baseline simulation characteristics, *Journal of*
653 *climate*, 25, 6646–6665, 2012.

654 Frieler, K., Lange, S., Piontek, F., Reyer, C. P., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil,
655 S., and Emanuel, K.: Assessing the impacts of 1.5 C global warming–simulation protocol of the
656 Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), *Geoscientific Model*
657 *Development*, 10, 4321–4345, 2017.

658 Gädeke, A., Krysanova, V., Aryal, A., Chang, J., Grillakis, M., Hanasaki, N., Koutroulis, A., Pokhrel,
659 Y., Satoh, Y., and Schaphoff, S.: Performance evaluation of global hydrological models in six large
660 Pan-Arctic watersheds, *Climatic Change*, 163, 1329–1351, 2020.

661 Gerten, D., Hoff, H., Rockström, J., Jägermeyr, J., Kummu, M., and Pastor, A. V.: Towards a revised
662 planetary boundary for consumptive freshwater use: role of environmental flow requirements,
663 *Current Opinion in Environmental Sustainability*, 5, 551–558, 2013.

664 Gleeson, T., Wang-Erlandsson, L., Porkka, M., Zipper, S. C., Jaramillo, F., Gerten, D., Fetzer, I.,
665 Cornell, S. E., Piemontese, L., and Gordon, L. J.: Illuminating water cycle modifications and Earth
666 system resilience in the Anthropocene, *Water Resources Research*, 56, 2020a.

667 Gleeson, T., Wang-Erlandsson, L., Zipper, S. C., Porkka, M., Jaramillo, F., Gerten, D., Fetzer, I.,
668 Cornell, S. E., Piemontese, L., and Gordon, L. J.: The water planetary boundary: interrogation and
669 revision, *One Earth*, 2, 223–234, 2020b.

670 Gleick, P. H.: Water resources, *Encyclopedia of climate, weather*, 817–823, 1996.

671 Gozlan, R. E., Britton, J. R., Cowx, I., and Copp, G. H.: Current knowledge on non-native freshwater
672 fish introductions, *Journal of fish biology*, 76, 751–786, 2010.

673 Grill, G., Lehner, B., Lumsdon, A. E., MacDonald, G. K., Zarfl, C., and Liermann, C. R.: An index-
674 based framework for assessing patterns and trends in river fragmentation and flow regulation by
675 global dams at multiple scales, *Environmental Research Letters*, 10, 015001, 2015.

676 Hanasaki, N., Yoshikawa, S., Pokhrel, Y., and Kanae, S.: A global hydrological simulation to specify
677 the sources of water used by humans, *Hydrology and Earth System Sciences*, 22, 789–817, 2018.

678 Herrera-R, G. A., Oberdorff, T., Anderson, E. P., Brosse, S., Carvajal-Vallejos, F. M., Frederico, R.
679 G., Hidalgo, M., Jézéquel, C., Maldonado, M., Maldonado-Ocampo, J. A., Ortega, H., Radinger, J.,
680 Torrente-Vilara, G., Zuanon, J., and Tedesco, P. A.: The combined effects of climate change and
681 river fragmentation on the distribution of Andean Amazon fishes, *Global Change Biology*, 26,
682 5509–5523, <https://doi.org/10.1111/gcb.15285>, 2020.

683 Horne, A. C., Webb, J. A., O'Donnell, E., Arthington, A. H., McClain, M., Bond, N., Acreman, M.,
684 Hart, B., Stewardson, M. J., and Richter, B.: Research priorities to improve future environmental
685 water outcomes, *Frontiers in Environmental Science*, 5, 89, 2017.

686 Kabat, P., Claussen, M., Dirmeyer, P. A., Gash, J. H., de Guenni, L. B., Meybeck, M., Hutjes, R. W.,
687 Pielke Sr, R. A., Vorosmarty, C. J., and Lütke-meier, S.: *Vegetation, water, humans and the climate:*
688 *A new perspective on an interactive system*, Springer Science & Business Media, 2004.

689 Knouft, J. H. and Ficklin, D. L.: The potential impacts of climate change on biodiversity in flowing
690 freshwater systems, *Annual Review of Ecology, Evolution, and Systematics*, 48, 111–133, 2017.

691 Lehner, B. and Grill, G.: Global river hydrography and network routing: baseline data and new
692 approaches to study the world's large river systems, *Hydrological Processes*, 27, 2171–2186,
693 2013.

694 Leprieur, F., Beauchard, O., Blanchet, S., Oberdorff, T., and Brosse, S.: Fish invasions in the world's
695 river systems: when natural processes are blurred by human activities, *PLoS biology*, 6, e28, 2008.

696 Leprieur, F., Brosse, S., Garcia-Berthou, E., Oberdorff, T., Olden, J. D., and Townsend, C. R.:
 697 Scientific uncertainty and the assessment of risks posed by non-native freshwater fishes, *Fish and*
 698 *Fisheries*, 10, 88–97, 2009.

699 Linnansaari, T., Monk, W. A., Baird, D. J., and Curry, R. A.: Review of approaches and methods to
 700 assess Environmental Flows across Canada and internationally, *DFO Can. Sci. Advis. Sec. Res. Doc.*,
 701 39, 74, 2012.

702 Lundberg, J. G., Kottelat, M., Smith, G. R., Stiassny, M. L., and Gill, A. C.: So many fishes, so little
 703 time: an overview of recent ichthyological discovery in continental waters, *Annals of the Missouri*
 704 *Botanical Garden*, 26–62, 2000.

705 Meyer, J. L., Sale, M. J., Mulholland, P. J., and Poff, N. L.: Impacts of climate change on aquatic
 706 ecosystem functioning and health, *JAWRA Journal of the American Water Resources Association*,
 707 1, 35, 1373–1386, 1999.

708 Moyle, P. B. and Leidy, R. A.: Loss of biodiversity in aquatic ecosystems: evidence from fish faunas,
 709 in: *Conservation biology*, Springer, 127–169, 1992.

710 Müller Schmied, H., Adam, L., Eisner, S., Fink, G., Flörke, M., Kim, H., Oki, T., Portmann, F. T.,
 711 Reinecke, R., and Riedel, C.: Variations of global and continental water balance components as
 712 impacted by climate forcing uncertainty and human water use, *Hydrology and Earth System*
 713 *Sciences*, 20, 2877–2898, 2016.

714 NCEFSAB: Recommendations for estimating flows to maintain ecological integrity in streams and
 715 rivers in North Carolina, 2013.

716 Nilsson, C., Reidy, C. A., Dynesius, M., and Revenga, C.: Fragmentation and flow regulation of the
 717 world’s large river systems, *Science*, 308, 405–408, 2005.

718 Nykvist, B., Persson, Å., Moberg, F., Persson, L., Cornell, S., and Rockström, J.: National
 719 environmental performance on planetary boundaries, A study for the Swedish Environmental
 720 Protection Agency (Stockholm Environment Institute, Stockholm), 2017.

721 Nyström, P. E. R., BRÖNMARK, C., and Graneli, W.: Patterns in benthic food webs: a role for
 722 omnivorous crayfish? *Freshwater biology*, 36, 631–646, 1996.

723 Olson, D. M. and Dinerstein, E.: The Global 200: Priority ecoregions for global conservation,
 724 *Annals of the Missouri Botanical garden*, 199–224, 2002.

725 Pastor, A. V., Ludwig, F., Biemans, H., Hoff, H., and Kabat, P.: Accounting for environmental flow
 726 requirements in global water assessments, *Hydrology and earth system sciences*, 18, 5041–5059,
 727 2014.

728 Poff, N. L. and Zimmerman, J. K.: Ecological responses to altered flow regimes: a literature review
729 to inform the science and management of environmental flows, *Freshwater biology*, 55, 194–
730 205, 2010.

731 Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., Sparks, R. E., and
732 Stromberg, J. C.: The natural flow regime, *BioScience*, 47, 769–784, 1997.

733 Poff, N. L., Brinson, M. M., and Day, J. W.: Aquatic ecosystems and global climate change,
734 Arlington, VA, 44, 1–36, 2002.

735 Poff, N. L., Richter, B. D., Arthington, A. H., Bunn, S. E., Naiman, R. J., Kendy, E., Acreman, M.,
736 Apse, C., Bledsoe, B. P., and Freeman, M. C.: The ecological limits of hydrologic alteration
737 (ELOHA): a new framework for developing regional environmental flow standards, *Freshwater*
738 *biology*, 55, 147–170, 2010.

739 Poff, N. L., Tharme, R. E., and Arthington, A. H.: Evolution of environmental flows assessment
740 science, principles, and methodologies, *Water for the Environment*, Elsevier, 203–236, 2017.

741 Powell, S. J., Letcher, R. A., and Croke, B. F. W.: Modelling floodplain inundation for
742 environmental flows: Gwydir wetlands, Australia, *Ecological Modelling*, 211, 350–362, 2008.

743 Reid, A. J., Carlson, A. K., Creed, I. F., Eliason, E. J., Gell, P. A., Johnson, P. T., Kidd, K. A.,
744 MacCormack, T. J., Olden, J. D., and Ormerod, S. J.: Emerging threats and persistent conservation
745 challenges for freshwater biodiversity, *Biological Reviews*, 94, 849–873, 2019.

746 Richter, B., Baumgartner, J., Wigington, R., and Braun, D.: How much water does a river need?,
747 *Freshwater biology*, 37, 231–249, 1997.

748 Richter, B. D., Mathews, R., Harrison, D. L., and Wigington, R.: Ecologically sustainable water
749 management: managing river flows for ecological integrity, *Ecological applications*, 13, 206–224,
750 2003.

751 Richter, B. D., Davis, M. M., Apse, C., and Konrad, C.: A presumptive standard for environmental
752 flow protection, *River Research and Applications*, 28, 1312–1321, 2012.

753 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin III, F. S., Lambin, E., Lenton, T. M.,
754 Scheffer, M., Folke, C., and Schellnhuber, H. J.: Planetary boundaries: exploring the safe operating
755 space for humanity, *Ecology and society*, 14, 2009.

756 Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke,
757 J., Jägermeyr, J., and Knauer, J.: LPJmL4—a dynamic global vegetation model with managed land—
758 Part 1: Model description, *Geoscientific Model Development*, 11 (4), 1343–1375, 2018.

759 Shafroth, P. B., Wilcox, A. C., Lytle, D. A., Hickey, J. T., Andersen, D. C., Beauchamp, V. B.,
760 Hautzinger, A., McMULLEN, L. E., and Warner, A.: Ecosystem effects of environmental flows:
761 modelling and experimental floods in a dryland river, *Freshwater Biology*, 55(1), 68–85, 2010.

762 Shesterin, I. S.: Water pollution and its impact on fish and aquatic invertebrates, Interactions:
 763 Food, Agriculture And Environment UNESCO Publishing-Eolss Publishers, Oxford, UK, 59–69,
 764 2010.

765 Smakhtin, V., Revenga, C., and Döll, P.: A pilot global assessment of environmental water
 766 requirements and scarcity, *Water international*, 29 (3), 307–317, 2004.

767 Smith, M. and Cartin, M.: Water vision to action: catalysing change through the IUCN water and
 768 nature initiative, IUCN, Gland, Switzerland. 2011.

769 Smith, V. H.: Eutrophication of freshwater and coastal marine ecosystems a global problem,
 770 *Environmental Science and Pollution Research*, 10 (2), 126–139, 2003.

771 Steffen, W., Richardson, K., Rockström, J., Cornell, S. E., Fetzer, I., Bennett, E. M., Biggs, R.,
 772 Carpenter, S. R., De Vries, W., and De Wit, C. A.: Planetary boundaries: Guiding human
 773 development on a changing planet, *Science*, 347(6223), 1259855, 2015.

774 Su, G., Logez, M., Xu, J., Tao, S., Villéger, S., and Brosse, S.: Human impacts on global freshwater
 775 fish biodiversity, *Science*, 371, 835–838, 2021.

776 Sutanudjaja, E. H., Van Beek, R., Wanders, N., Wada, Y., Bosmans, J. H., Drost, N., Van Der Ent, R.
 777 J., De Graaf, I. E., Hoch, J. M., and De Jong, K.: PCR-GLOBWB 2: a 5 arcmin global hydrological and
 778 water resources model, *Geoscientific Model Development*, 11, 2429–2453, 2018.

779 Tedesco, P. A., Beauchard, O., Bigorne, R., Blanchet, S., Buisson, L., Conti, L., Cornu, J.-F., Dias, M.
 780 S., Grenouillet, G., and Hugueny, B.: A global database on freshwater fish species occurrence in
 781 drainage basins, *Scientific data*, 4, 1–6, 2017.

782 Tennant, D. L.: Instream flow regimens for fish, wildlife, recreation and related environmental
 783 resources, *Fisheries*, 1, 6–10, 1976.

784 Tessmann, S. A.: Environmental use sector: reconnaissance elements of the western Dakotas
 785 region of South Dakota study, Water Resources Institute, South Dakota State University, 1979.

786 Thompson, R. M. and Lake, P. S.: Reconciling theory and practise: the role of stream ecology,
 787 *River Research and Applications*, 26, 5–14, 2010.

788 Tickner, D., Opperman, J. J., Abell, R., Acreman, M., Arthington, A. H., Bunn, S. E., Cooke, S. J.,
 789 Dalton, J., Darwall, W., and Edwards, G.: Bending the curve of global freshwater biodiversity loss:
 790 an emergency recovery plan, *BioScience*, 70(4), 330–342, 2020.

791 Tonkin, J. D., Olden, J. D., Merritt, D. M., Reynolds, L. V., Rogosch, J. S., and Lytle, D. A.: Designing
 792 flow regimes to support entire river ecosystems, *Frontiers in Ecology and the Environment*, 19(6),
 793 326–333, 2021.

794 Tyson, P., Odada, E., Schulze, R., and Vogel, C.: Regional-global change linkages: Southern Africa,
795 in: *Global-regional linkages in the earth system*, Springer, 3–73, 2002.

796 Villéger, S., Blanchet, S., Beauchard, O., Oberdorff, T., & Brosse, S. (2011). Homogenization
797 patterns of the world's freshwater fish faunas. *Proceedings of the National Academy of Sciences*,
798 108(44), 18003-18008.

799 Virkki, V., Alanärä, E., Porkka, M., Ahopelto, L., Gleeson, T., Mohan, C., Wang-Erlandsson, L., Flörke, M.,
800 Gerten, D., Gosling, S.N. and Hanasaki, N.: Environmental flow envelopes: quantifying global,
801 ecosystem-threatening streamflow alterations, *Hydrology and Earth System Sciences*, 1–31,

802 Vitousek, P. M., Mooney, H. A., Lubchenco, J., and Melillo, J. M.: Human domination of Earth's
803 ecosystems, *Science*, 277(5325), 494–499, 1997.

804 Vitule, J. R. S., Freire, C. A., and Simberloff, D.: Introduction of non-native freshwater fish can
805 certainly be bad, *Fish and fisheries*, 10(1), 98–108, 2009.

806 Vörösmarty, C. J., Wasson, R., and Richey, J. E.: Modelling the transport and transformation of
807 terrestrial materials to freshwater and coastal ecosystems workshop report, *International*
808 *Geosphere Biosphere Programme* [Stockholm], 1997.

809 Vorosmarty, C. J., Green, P., Salisbury, J., and Lammers, R. B.: Global water resources:
810 vulnerability from climate change and population growth, *Science*, 289(5477), 284–288, 2000.

811 Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., Glidden,
812 S., Bunn, S. E., Sullivan, C. A., and Liermann, C. R.: Global threats to human water security and
813 river biodiversity, *Nature*, 467(7315), 555–561, 2010.

814 Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., and Schewe, J.: The inter-sectoral
815 impact model intercomparison project (ISI-MIP): project framework, *Proceedings of the National*
816 *Academy of Sciences*, 111(9), 3228–3232, 2014.

817 Watanabe, M., Suzuki, T., O'ishi, R., Komuro, Y., Watanabe, S., Emori, S., Takemura, T., Chikira,
818 M., Ogura, T., and Sekiguchi, M.: Improved climate simulation by MIROC5: mean states,
819 variability, and climate sensitivity, *Journal of Climate*, 23(23), 6312–6335, 2010.

820 Wilting, H. C., Schipper, A. M., Bakkenes, M., Meijer, J. R., and Huijbregts, M. A.: Quantifying
821 biodiversity losses due to human consumption: a global-scale footprint analysis, *Environmental*
822 *Science & Technology*, 51(16), 3298–3306, 2017.

823 Xenopoulos, M. A., Lodge, D. M., Alcamo, J., Märker, M., Schulze, K., and Van Vuuren, D. P.:
824 Scenarios of freshwater fish extinctions from climate change and water withdrawal, *Global*
825 *change biology*, 11(10), 1557–1564, 2005.

826 Yoshikawa, S., Yanagawa, A., Iwasaki, Y., Sui, P., Koirala, S., Hirano, K., Khajuria, A., Mahendran,
827 R., Hirabayashi, Y., and Yoshimura, C.: Illustrating a new global-scale approach to estimating

828 potential reduction in fish species richness due to flow alteration, *Hydrology and Earth System*
829 *Sciences*, 18(2), 621–630, 2014.

830 Zaherpour, J., Gosling, S. N., Mount, N., Schmied, H. M., Veldkamp, T. I., Dankers, R., Eisner, S.,
831 Gerten, D., Gudmundsson, L., and Haddeland, I.: Worldwide evaluation of mean and extreme
832 runoff from six global-scale hydrological models that account for human impacts, *Environmental*
833 *Research Letters*, 13(6), 065015, 2018.

834