

## Responses to the reviewers

**Title:** Identification, Mapping and Eco-hydrological Signal Analysis for Groundwater-dependent Ecosystems (GDEs) in Langxi River Basin, North China (hess-2023-151)

**Authors:** Mingyang Li, Fulin Li\*, Shidong Fu, Huawei Chen, Kairan Wang, Xuequn Chen, Jiwen Huang

**Thanks to the experts giving so much valuable advice, now the revision notes are shown as following.**

### **RC1:**

This is a manuscript that deserves praise. It is difficult to concentrate on rich field experiments in the current setting. This study presents a framework for locating and mapping GDEs based on a mix of remote sensing, GIS, and hydrogeological field experiments in terms of research content and methodologies. The study's newly suggested normalized difference built-up and soil index, together with the difference between the wet index, may be used to assess changes in the water loss rate of plants at various stages of growth. After then, the spectrum of possible GDEs is defined by factors including slope, lithology, and elevation. Groundwater levels, river bed bottom elevation, plant root depth, karst springs, etc. were used to further infer different forms of GDEs. In the end, three factors were used to verify GDEs: groundwater biology, water chemical isotopes, and hydrological rhythm. Gaining a deeper understanding of a basin may be achieved by employing a variety of techniques to examine its biological and hydrological features. I would want to share a few small ideas and inquiries with the author.

**Answer:** Thank you very much for your review and the suggestions of the three experts. We have revised the paper according to each suggestion.

1. The author employed aquatic biology, hydrology, water chemistry, isotopes, and other elements to confirm. These verification findings demonstrate that spring flow and base flow in the basin have a substantial association, that the water chemistry and isotope composition of distinct water bodies vary, and that the water bodies in GDEs have comparatively separate ecosystems. It is still necessary to enhance the geographical representativeness of the aforementioned results. Is it attainable to categorize and validate GDEs in space, despite the fact that they are challenging to locate and define precisely?

**Answer:** Thank you very much for your comment. We will reply to your questions in the following points:

Identifying different types of GDEs is of practical significance for stressed GDE habitats, limiting groundwater overexploitation, and implementing ecological replenishment. It is feasible to identify potential GDEs solely using remote sensing technology, but relatively accurate GDEs mapping requires hydrogeological surveys,

hydrochemical isotope tracing analysis, and groundwater animal validation.

The Langxi River Basin (LRB) is a typical continuous carbonate-type karst basin in northern China. Generally speaking, this kind of basin is scattered and relatively independent. The small area contains complex topography and groundwater exchange relationships. Therefore, similar research often selects basins with a small area, which is convenient for detailed investigation. The research methods have operational commonality in similar basins around the world. For example, this study is based on previous methods and aims at the characteristics of the underlying surface of the LRB, trying to propose a more complete research framework and improved methods. In this study, we targeted the classification and identification of GDEs according to the types of GDEs in LRB. Because different types of GDEs are distributed in LRB, including karst springs and hyporheic zones, they are more typical in northern China. In other places, affected by groundwater overexploitation and high-intensity human activities, GDEs have been seriously damaged. Even though large karst springs exist in some places, they do not form an obvious ecosystem. Therefore, first of all, we believe that choosing LRB as a typical research area is representative of the karst areas in northern China.

In this study, we propose a more systematic and comprehensive method compared to current identification and mapping research. We are not simply doing GDE classification research, because the classification is relatively simple and can be identified in the wild. And our research mainly focuses on recognition and mapping.

Based on previous research, we proposed a framework for identifying, mapping and verifying the distribution range of GDEs. This system can be improved and transplanted to other regions. For example, Barron et al. (2014) used NDVI and NDWI to classify potential GDEs in Australia. This is not significantly feasible in LRB, but this does not mean that its idea is not feasible. We selected appropriate remote sensing indices NDBSI and NDVI and jointly identified potential GDEs in LRB, and the results showed that the discrimination effect was very good. Therefore, we believe that this study is more focused on proposing such a system for identifying, mapping and verifying GDEs, which can be used in specific research areas with appropriate improvements. One thing to note is that we are not yet able to accurately determine the boundaries of GDEs. We can only make a comprehensive judgment based on surface vegetation and water bodies. A more precise division requires further exploration.

2. The author used data from 2020 to 2021 to divide the scope of GDEs in the Langxi River Basin. This seems to be because the author conducted the experiment during this period. Does this method also work at other times?

Answer: Thank you for your question. First of all, the distribution range of GDEs will change with the amount and range of groundwater recharge. The amount and extent of groundwater recharge will slowly change with the impacts of climate change and human activities. When natural disasters such as earthquakes, coal mining, or human activities cause serious changes in underground aquifers, the distribution range of GDEs may change significantly in a short period of time. During the period when the

stratum does not change significantly, long-term groundwater level monitoring data shows that the changes in groundwater levels in the entire LRB and its vicinity are very weak. Therefore, we believe that the distribution range identification method of GDEs proposed in this article is representative. It should be noted that when the groundwater level fluctuates significantly or the stratigraphy changes significantly, we recommend re-evaluating the relevant indices and parameters in the study area and this system.

We have added relevant discussion of this situation in the Discussion section of the paper.

**In the revised paper:**

Line 563 to 571:


The quantity and range of groundwater recharge will affect the GDEs' distribution range. The effects of climate change and human activity will gradually alter the volume and scope of groundwater recharge. The distribution range of GDEs can alter dramatically in a short amount of time when major changes are made to subterranean aquifers by natural disasters like earthquakes, coal mining, or human activity. It should be noted that when the groundwater level fluctuates significantly or the stratigraphy changes significantly, we recommend re-evaluating the relevant indices and parameters in the study area and this system. Long-term groundwater level monitoring data indicates that the changes in groundwater levels in the entire LRB and its vicinity are very weak during the period when the stratum does not change significantly. Therefore, the Four diagnostic criteria framework proposed in the study can effectively identify GDEs in areas such as grasslands, deserts, plains, and karsts in arid, semi-arid, and sub-humid areas.

3. The Langxi River Basin is a typical study area selected by the author. Is the GDEs identification and mapping framework proposed by the author also applicable to other river basins? I think this is also an interesting question for other readers.


Answer: We understand that readers will consider whether this approach can actually be applied to other basins. We subsequently conducted relevant experiments in another Chinese karst area, the Tanglang River Basin (TRB), and used this framework to classify GDEs. We will show you some of the results here.

Since the following content may involve some of our unpublished content, we have watermarked the images. We hope you understand.

First, we collected and organized relevant data on TRB and divided plains and mountains. Land use types, groundwater levels, water gaining and losing reaches, river bed elevation, and vegetation root systems of the TRB were investigated. The vegetation index at the end of the dry season and the wet season is used to classify crops, plants that lose water quickly, plants that lose water slowly, evergreen vegetation and buildings, etc.



The GDEs distribution characteristics of TRB can be further calculated by using the spatial kernel density method used in this study.



It should be noted that the identification, drawing and verification framework we proposed has been transplanted from LRB to TRB and has also been improved accordingly. First of all, because there are a large number of population centers and cities in TRB, we added the identification of construction land to the identification framework of GDEs. Secondly, compared with LRB, TRB has more karst caves. During the verification process of water bodies, we added work such as sampling of cave fissure water and aquatic life inside and outside caves. This can make the verification more convincing. In subsequent research, by incorporating hydrological (ecological hydrological) models that consider groundwater, and conducting source analysis of water chemistry, isotopes, and sediment, a series of research accuracy such as the identification and mapping of GDEs can be improved.

Some other formatting questions or suggestions:

1. Line 275 should add relevant introduction to the base flow segmentation method.

Answer: We have added a brief introduction to baseflow segmentation at the corresponding location in the text.

**In the revised paper:**

Page 9 Line 266 to 269:

Base flow segmentation is a method used in hydrology to separate streamflow data into its base flow and surface runoff components. Base flow generally represents the groundwater contribution to streamflow, while surface runoff comes from precipitation events and other surface sources. There are several methods for base flow segmentation, including hydrograph separation, chemical separation,

**hydrometric separation, et.** The study utilizes the straight-line secant method, which involves horizontally dividing the peak of the flow process line using a horizontal line. It is stipulated that the contribution of surface runoff lies above the horizontal cutting line, while the contribution of base flow lies below the horizontal cutting line. The value of the horizontal line, which represents the runoff, can be determined as the minimum flow during the dry season, the minimum daily average flow during the dry season, or the minimum monthly average flow for the year. During the non-rainy season when the karst aquifer is recharged, the flow of the spring will gradually decrease in size until it matches the recharge rate of the aquifer, as there is no additional recharge from precipitation. The equation for the flow attenuation process can be written as (Rodríguez et al., 2017).

2. It is recommended to use tables to express the data part of Line 310

**Answer: Thanks to your suggestion. We have added Table 1 to sort out the types, names, resolutions, sources of data sets used in this study, and the bands used for remote sensing data.**

**In the revised paper:**

Page 11 Line 316:

**In this paper, the data mainly includes remote sensing data, and hydrogeological survey data (Table 1).**

Page 33 Line 803:

**Table 1: Remote sensing and hydrogeological survey data used in the research.**

Data Type	Data Name	Resolution	Resources	Bands used in research
<b>Remote sensing Data</b>	USGS Landsat 8 Level 2, Collection 2, Tier 1	30 m	Google Earth Engine (GEE)	B2-B7 (Blue, Green, Red, Near infrared, Shortwave infrared 1 and Shortwave infrared 2)
	NASA SRTM Digital Elevation	30 m	GEE	Elevation and Slope (Calculated by elevation)
<b>Hydrogeological Survey Data</b>	Chinese stratigraphic lithology dataset	1:2,500,000	China Geological Survey	Geological lithology, geological body boundary, amphibole schist, crater point, et.
	The maximum root depth	1:5,000,000	Harmonized World Soil Database	-
	River bed level	Point scale	Field surveys	-
	Groundwater level	Point scale	Field surveys	-
	Water hydrochemical and groundwater fauna sampling	Point scale	Field surveys	-

Note: Please see Supplementary Table 2 for remote sensing data sources.

3. Line 495 Please adjust the font size

**Answer:** Thank you for your suggestion. We have adjusted the font size.


4. Line 565 In the conclusion, the author uses the full names of K-GDE, S-GDE and V-GDE. In fact, the abbreviation has been used in the previous article, and it is recommended to use the abbreviation here.

**Answer:** Thank you for your opinion, but we have our own considerations here. As the finishing touch of a paper, the conclusion plays the role of naming the main content and summarizing the article. In this article, we reorganize the classification of GDE based on the characteristics of the study area. Using the full names of the three GDEs at the end allows readers to accept this definition (concept) more clearly. Therefore, we believe that the full names of K-GDE, S-GDE and V-GDE can be used here.

5. The study area map of Line 710 in Figure 1 should be redesigned. It's not pretty now.

**Answer:** We modified Figure 1, adjusted the layout, and replaced the more difficult-to-understand geological symbols with abbreviations.

**In the revised paper:**



**Figure 1: The location, lithology, topography, spring water, groundwater level survey points, hydrochemical groundwater biological sampling points of the Langxi River Basin. HRAD: Holocene fluvial alluvial deposits; UPGL: Upper Pleistocene gravel layer; UCL: Upper Cambrian limestone; UCSLA: Upper Cambrian shale-limestone amalgamation.**

## RC2:

The manuscript by Li et al. focuses on the identification and analysis of groundwater-dependent ecosystems (GDEs) in a specific region. The authors propose a four-diagnostic criteria framework for identifying GDEs based on remote sensing, GIS data dredging, and hydrogeological surveys. Overall, I think this work will be of interest to the HESS readership but needs some reworking to be considered further for publication. I make a few suggestions for improvement below:

**Answer:** Thank you very much for your review and the suggestions. We have revised the paper according to each suggestion. In response to your opinions, we have made the following point-to-point modifications, which can be summarized as:

(1) We have adjusted the arrangement of the conclusion section to make the research findings more like an article rather than a report listing the research results. The three chapters are now: 3.1 Hydrogeological investigation of three types of GDEs; 3.2 Distribution and mapping of GDEs; 3.3 Ecohydrological signals of GDEs.

(2) We have strengthened the connection with the method in the results section, and related the formulas in the method to the results. On the basis of adjusting the order of the results part, the description in the results was significantly modified to make the entire article more fluent.

(3) We have redrawn Figure 2, Figure 8 and Figure 10, adding a lot of details. This makes our images and texts more relevant.

(4) In the discussion section, we added a comparative analysis with two existing similar studies, analyzed the advantages and disadvantages of this study and existing studies, and further illustrated the advancement of the Four diagnostic criteria framework.

1. The result section lacks clarity and structure. It resembles a report rather than a concise summary of findings. Some materials in the results, particularly those pertaining to the authors' methodologies or rationales, such as those indicated on Page 11 Lines 337-339, Page 14 Lines 407-408, and Page 14 Lines 421-423, should be relocated to the method section to enhance organization and coherence.

**Answer:** Thank you for your opinion. To improve the clarity of the results section of the article, we modified revised the arrangement of chapters. The results section of this article now has five chapters. Please allow us to introduce the structural ideas of the results section of this article. The results part of this article is arranged according to the framework of the research method in Figure 2. First, Section 3.1 introduces the hydrogeological investigation results for three types of GDEs. Section 3.2 introduces the distribution and mapping of GDEs. Section 3.3 introduces the ecohydrological signals of GDEs to verify whether the GDEs distribution is correct.

Then, we have rewritten the results section in relation to the methods and rationale of Chapter 2 to enhance organization and coherence.

(1) Page 11 Lines 337-339 of the **original text**

As the first sentence of Section 3.2, in order to better connect the methods and results,

it should also serve as the beginning to guide the rest of the text. We've rewritten this.

**In the revised paper:**

Page 13 Line 387-383

Based on the research framework in Figure 3, the study first uses remote sensing indicators such as terrain, NDVI, NDWI, and DWN to identify characteristics such as waters, bare land, wetlands, and vegetation to determine the potential distribution of GDE. Supplementary Figure 2 displays the distribution characteristics of NDVI and NDWI in the study area at the end of the dry and wet seasons. In the central and southern plains of the study area, NDVI remained high at the end of the wet season (Supplementary Figure 2a) and slightly decreased at the end of the dry season (Supplementary Figure 2b), indicating that the vegetation in this area primarily experiences rapid drying.

(2) Page 14 Lines 407-408 of the **original text**

We did not clearly express what we wanted to describe in the original text. We've rewritten this.

**In the revised paper:**

Page 12 Line 344-347

The research acquired information about the maximum vegetation root depth, river bottom elevation, and groundwater level of the LRB (Figure 5a) by the gathering of GIS data and basin hydrogeological survey. We qualitatively evaluated the gaining and losing river portions using data on the elevation of the river bed and the depth of the groundwater table (Figure 5b).

The analysis of underground water table depths reveals that the shallow water table area (0 to 5m) is primarily located in the middle of the basin where the tributaries converge, and numerous karst springs are situated nearby. The vegetation in LRB is predominantly composed of deciduous broad-leaved forest and deciduous open shrubs, with relatively developed underground root systems owing to the year-round flow of rivers. The maximum root depth in the basin ranges from 1.5 to 2.5 m, with some exceeding 4.5 m. Areas with deeper root depths are present on both sides of the river. In the lower reaches of the Langxi River near the Yellow River, the roots of vegetation are relatively shallow, consistent with the spatial distribution of surface lithology. However, the reason for the shallow roots is not caused by the surface lithology. In fact, it is due to the abundance of water in the area, and a large number of farmlands are distributed here. The riverbed bottom level changes more gently from upstream to downstream compared to the change in elevation. Based on the identification of potential GDEs, the study was able to accurately divide S-GDE and V-GDE.

(3) Page 14 Lines 421-423 of the **original text**

This paragraph in the original article is not closely related to the previous article, so we have rewritten this paragraph.

**In the revised paper:**

Page 14 Line 430-436

Based on the previous ecological and hydrogeological survey results, the study subdivided potential GDEs into S-GDEs, V-GDEs and K-GDEs according to the four diagnostic criteria framework (Figure 3), and used the spatial kernel density function for mapping. The distribution results of GDEs in the watershed are shown in Figure 8. Green, orange and magenta represent V-GDEs, K-GDEs and S-GDEs respectively. The depth of the color represents the spatial core density of the GDE. When the climate is dry, areas with lower spatial core density will gradually no longer belong to the scope of the GDE. The GDEs in the basin are mainly located in the central and western parts of the LRB, covering an area of approximately 49 km<sup>2</sup>, which accounts for 29% of the basin's total area.

(4) We have also made corresponding modifications to some sentences that were not mentioned by experts in this article.

**In the revised paper:**

Page 1 Line 23-33

In order to verify the reliability of GDE distribution, the study verified the determination of GDEs through hydrological rhythm analysis, hydrochemical characteristics analysis of various water bodies in the basin, and ecohydrological signals such as groundwater invertebrates. The hydrological rhythm analysis in Shuyuan section showed that the proportion of base flow to river flow is about 54.15% and S-GDEs still receive spring water recharge even in the extremely dry season. And the analysis of hydrochemical sampling from the karst aquifer, Quaternary aquifer, spring water and surface reservoir water reveals that GDEs are also recharged by groundwater. More important, we also found a distinctive ecohydrological signal of GDEs is the presence of millimeter-sized groundwater fauna living in the different types of GDEs. In addition, the study believes that the use of isotope and environmental DNA technology to analyze the hydrological-sediment-biological connectivity between groundwater and GDE is the future development direction of this field.

2. The current version lacks substantial supporting information on ecohydrological signals aiding in mapping GDEs. Section 3.4 primarily focuses on the spatial distribution of GDEs within the study area. Sections 3.5 and 3.6 delve into detailed descriptions of hydro-biogeochemical features observed in GDEs. However, there appears to be a weak connection between these descriptions and the verification of various GDEs, as depicted in Figure 2. Strengthening this connection is essential for a more cohesive presentation of the study's findings regarding the role of ecohydrological signals in GDE mapping.

Answer: Thank you for your opinion, your opinion is very important to our modifications. We recognized the shortcomings of the original version of the weak connection of ecohydrological signals in GDEs drawing and made the following modifications:

(1) The section shown in Figure 2 has important connections with Section 3.3.1. Section 3.3.1 describes the lateral recharge relationship of groundwater to the Langxi River. In response to this, this study conducted a geological survey on the Shuyuan Spring and Langxi River sections, and explored the relationship between the groundwater level and Shuyuan Spring, as well as relevant information about the stratigraphy of the sections. In the text, we reformulate the connection between Figure 2 and Section 3.3.1. And in order to increase relevance, we modified Figure 2 to more clearly represent the three types of GDEs present in the modified typical section.

**In the revised paper:**

Page 15 Line 446-451

The Shuyuan Spring and Langxi River profile (hereinafter referred to as the Shuyuan Spring profile, Figure 2) shows the location and elevation relationship between groundwater and the Langxi River profile. The eastern Cambrian Zhushadong-Zhangxia Formation limestone is rich in groundwater buried deeper than the river. Shuyuan Spring, one of the descending springs in the basin, gushes out at the intersection of the Quaternary sedimentary layer and the Cambrian Zhushadong-Zhangxia Formation limestone. At the same time, groundwater also recharges to the Langxi River along the stratigraphic fissures.

**In the revised paper:**

Figure 2.




Figure 2: Hydrogeological profile of Shuyuan spring in LRB. The dotted line shows the characteristics of the water table in the geological section. The geological types in the figure are Q: Quaternary sedimentary layer; EjZ: Cambrian Zhushadong-Zhangxia Formation limestone; CcM: Cambrian Gushan-Chaomidian Formation limestone.

(2) Sections 3.3.2 and 3.3.3 respectively use the hydrochemical characteristics of different water bodies and the groundwater fauna to verify the identification of GDEs. In the original text, we did not reflect the corresponding connection well.

In the revised manuscript, we first modified Figure 10 in Section 3.3.2. We classified the water chemical characteristics corresponding to the distribution of the three GDEs in Figure 8 and discussed them accordingly in the text.

**In the revised paper:**

Page 16 Line 494-503:

Combined with the distribution range of the three GDEs and the water sample collection locations in Figure 8, we can clearly see that the hydrochemical characteristics of water bodies of the same GDE type have obvious clustering relationships. The overlapping relationship of hydrochemical characteristics in GDE groups is consistent with the distribution characteristics of GDEs types at the spatial scale of the basin (Figure 10a), which is also reflected in the cluster analysis diagram (Figure 10b). In addition, descending springs like Shuyuan Spring are not only closely related to karst aquifers, but also have good hydraulic connections with Quaternary aquifers. On the other hand, Huiquan Reservoir is a surface reservoir stored in a river barrage and receives a large amount of groundwater recharge. Therefore, it exhibits hydrochemical characteristics similar to spring water and Quaternary pore water. It can be seen that the interaction between surface water and groundwater in this area is strong, and there are obvious differences in the hydrochemical characteristics of GDE, but there are also certain similarities.

**In the revised paper:**

Page 30 Line 790:





Figure 8: Langxi River Basin GDEs distribution area and ecohydrological signals.

**In the revised paper:**

Page 32 Line 800:



(b) Cluster analysis dendrogram




Figure 10: Hydrochemical characteristics of ten water samples in LRB.

**In the revised paper:**

Page 17 Line 509-515:

Finding groundwater fauna near karst caves is easy because there is an abundance of food, making it easier for them to survive. Hu Spring, a natural karst cave, is home to three species of Chironomid larvae; *Anisogammarus* sp.; *Radix lagotis* were found in it. Typically, these faunas mainly comprise of Arthropods, Coelenterates and Mollusks that live in groundwater throughout their entire life cycle, and are known as stygobites, a true groundwater fauna. Combining the spatial distribution range of GDEs in Figure 8 of this study and the groundwater fauna in Tables 3 and 4, it can be seen that these groundwater fauna not only appear in the range of K-GDEs with caves and karst springs, but also in the range of some S-GDEs and a small amount of V-GDEs.

3. The proposed framework should be compared to other existing approaches to highlight its potential advantages and limitations. This discussion is crucial for providing insights into the novelty and effectiveness of the proposed methodology.

Answer: Thank you for your opinion. In the original article, we discussed the differences between this study and Barron et al. (2014)'s study, and explored the applicability of NDVI and NDWI in the identification of GDEs. In response to the experts' opinions, we conducted a comparative analysis with existing similar research from the perspective of input variables of the identification model and verification of GDEs. In the discussion section, it was proposed that this study can continue to supplement input to improve the accuracy of GDEs identification, and the superiority of this study in using hydrological rhythms, hydrochemical characteristics, and groundwater fauna for GDEs verification.

**In the revised paper:**

Page 18 Line 543-554:

Compared with the study of El-Hokayem et al. (2023) on identifying GDEs in Italy, this study did not include climate factors into the identification indicators. Instead, it focused more on elements of the local watershed itself, such as water body characteristics and groundwater biological characteristics. This is because the study area of El-Hokayem et al. (2023) belongs to the coastal area, and the vegetation responds more sensitively to climate elements. Our research prefers to identify and establish direct connections between groundwater-surface water-surface/near-surface ecosystems through hydrochemical characteristics, and then verify the identification scope of GDEs.

The research results of Duran-Llacer et al. (2022) using more than ten terrain and remote sensing indicators to identify and map GDEs in Chile showed that increasing the input variables of the identification model can help improve the accuracy of identification. This can be used as a reference for the application of the identification framework proposed in this study in other basins. The multi-index verification system of hydrological rhythm, water chemistry, and groundwater fauna used in this study can verify the accuracy of GDEs identification from multiple dimensions, which has basically not been used in existing research. And this will help provide stronger support for the results for verification in future relevant studies.

Specific comments Page 4, Line 100: Delete ", etc".

Answer: Here, there are two etc. in the original text, which may cause ambiguity to readers. We have corrected the inappropriate expression.

**In the revised paper:**

Page 3-4 Line 97-100:

These characteristics can often be regarded as a specific signal for monitoring ecosystem status and linking the functions of organisms to ecohydrological processes, such as the rhythm of hydrometeorological elements, hydrogeochemical

characteristics, and biological indicators (just like biodiversity, connectivity), etc.

Page 6, Line 183: Please clarify the scale behind select a 10-degree angle.

Answer: In the original article, we briefly stated "The determination of this parameter can be manually adjusted based on one-third of the average slope of the basin until the plains and mountains are clearly distinguished." In response to experts' suggestions, we first put the method of parameter determination and the results of selecting different parameters into a supplementary file. This will allow readers to better understand my approach.

**In the revised paper:**

Page 6 Line 184-186:

where,  $grid_{plain}$  represents the grid divided into plains;  $\Delta_{slope}$  is the threshold of the maximum plain slope, in this paper we take  $\Delta_{slope} = 10^\circ$ . The determination of this parameter can be manually adjusted based on one-third of the average slope of the basin until the plains and mountains are clearly distinguished (See Supplementary material I).

**In the Supplementary material I:**

Below, Figure S1 shows our results of plain and hill classification using 4 different slopes. It can be seen that except for minor differences in details, overall slope does not have a great impact on plains and hills in LRB. This is due to the fact that the mean elevation limits the role of slope in this basin.




Figure S1. Different slopes distinguish the plain and hilly results of LRB.

Page 10, Line 285: Delete "2"

Answer: Thanks for your opinion, but the 2 here is not a typo. We have modified the way this sentence is expressed. The original text means that we use HNO<sub>3</sub> to adjust the pH value of the solution to 2.

**In the revised paper:**

Page 10 Line 290-291:

Subsequently, all the water samples were passed through a 0.45 µm filter, and the liquid samples were acidified the pH to 2 using pure HNO<sub>3</sub> to prevent the precipitation of metals before metal analysis.

Page 16, Line 495: Font size is not consistent.

Answer: Thank you for your suggestion. We have adjusted the font size.

**Notification to the authors:**

1.Coloured or marked text in \*.pdf manuscript file is not allowed. Please provide a clean version of \*.pdf manuscript file (with black text) with the next revision.




Answer: Thank you for your review. In the first draft, we colored citations in blue for easier reading. In accordance with journal rules, we have adjusted the text to black in the revised manuscript.

2. It seems that table is included as figure #11. If it is so, it must be re-labelled as table and the references in the manuscript text must be adjusted accordingly. A table may be inserted as an image, but still be called as a table.

Answer: As requested we have modified Figure 11 into a table, it is now Table 4.

**In the revised paper: Page 35:**

**Table 4: Groundwater fauna samples. (a) *Galba.jervia.sp*; (b) *Radix lagotis Schrank*; (c) *Chironmidae*; (d) *Gyraulus.sp*; (e) *Dytiscidae.sp*; (f) *Anisogammarus.sp*.**

Neocaridina denticulata sinensis			Radix lagotis Schrank		
	Phylum:	Arthropoda		Phylum:	Mollusca
	Class:	Malacostraca		Class:	Gastropoda
	Order:	Decapoda		Order:	-
	Family:	Atyidae		Family:	Lymnaeidae
	Genus:	<i>Caridina</i>		Genus:	<i>Radix</i>
Chironmidae			Gyraulus.sp		
	Phylum:	Arthropoda		Phylum:	Mollusca
	Class:	Insecta		Class:	Gastropoda
	Order:	Diptera Nematocera		Order:	-
	Family:	Culicomorpha		Family:	Planorbidae
	Genus:	<i>Chironomoidea</i>		Genus:	<i>Gyraulus</i>
Dytiscidae.sp			Anisogammarus.sp		
	Phylum:	Arthropoda		Phylum:	Arthropoda
	Class:	Insecta		Class:	Malacostraca
	Order:	Coleoptera		Order:	Amphipoda
	Family:	Dytiscidae		Family:	Anisogammaridae
	Genus:	<i>Cybister</i>		Genus:	<i>Anisogammarus</i>

3. Please ensure that the colour schemes used in your maps and charts allow readers with colour vision deficiencies to correctly interpret your findings. Please check your figures using the Coblis – Color Blindness Simulator (<https://www.color-blindness.com/coblis-color-blindness-simulator/>) and revise the colour schemes accordingly.

**Answer:** All images in this article have been verified using Coblis – Color Blindness Simulator. We believe that the information in the picture can be clearly identified in the three modes of Anomalous Trichromacy, Dichromatic view, and Monochromatic view. If the editorial department believes that our color matching still does not meet the regulations, please contact us and we will redraw it in the next version. Many Thanks!

4. For the next revision, please check if your figures containing photos require a copyright statement/image credit and add it to the figures (or captions) ([https://publications.copernicus.org/for\\_authors/manuscript\\_preparation.html#figurestables](https://publications.copernicus.org/for_authors/manuscript_preparation.html#figurestables) -> Reproduction and reuse of figures and tables). If these figures were entirely created by the authors, there is no need to add a copyright statement or credit. In that case it is important that you confirm this explicitly by email.

**Answer:** We have read the publication rules. We have modified one image for which reproduction rights have not been granted and certify that all images are created by us.

5. For the next revision, please make sure that information about the contribution of each of the authors of the manuscript is presented in the "Author contribution" section of the \*.pdf manuscript.

**Answer:** As requested we have included an author contribution section in the paper.

**In the revised paper:**

Author contribution

Mingyang Li and Fulin Li developed the initial and final versions of this manuscript and analyzed the data. Shidong Fu, Huawei Chen, Kairan Wang, Xuequn Chen, and Jiwen Huang contributed their expertise and insights to oversee the analysis.