

Evaluation of reanalysis soil moisture products using Cosmic Ray Neutron Sensor observations across the globe

Yanchen Zheng^{1,2}, Gemma Coxon¹, Ross Woods², Daniel Power², Miguel Angel Rico-Ramirez², David McJannet³, Rafael Rosolem^{2,4}, Jianzhu Li⁵ and Ping Feng⁵

5 ¹School of Geographical Sciences, University of Bristol, Bristol, UK.

²School of Civil, Aerospace and Design Engineering, University of Bristol, Bristol, UK.

³CSIRO Environment, EcoSciences Precinct, Dutton Park, Queensland, Australia.

⁴Cabot Institute for the Environment, University of Bristol, Bristol, UK.

⁵State Key Laboratory of Hydraulic Engineering Simulation and Safety, Tianjin University, Tianjin, China.

10 *Corresponding to:* Yanchen Zheng (yanchen.zheng@bristol.ac.uk)

Abstract. Reanalysis soil moisture products are valuable for diverse applications but their quality assessment is limited due to scale discrepancies when compared to traditional in-situ point-scale measurements. The emergence of Cosmic Ray Neutron Sensors (CRNS) with field-scale soil moisture estimates (~250m radius, up to 0.7m deep) is more suitable for the product evaluation owing to its larger footprint. In this study, we perform a comprehensive evaluation of eight widely-used reanalysis soil moisture products (ERA5-Land, CFSv2, MERRA2, JRA55, GLDAS-Noah, CRA40, GLEAM and SMAP L4 datasets) against 135 CRNS sites from the COSMOS-UK, COSMOS-Europe, COSMOS USA and CosmOz Australia networks. We evaluate the products using six metrics capturing different aspects of soil moisture dynamics. Results show that all reanalysis products generally exhibit good temporal correlation with the measurements, with the median of temporal correlation coefficient (R) values spanning from 0.69 to 0.79, though large deviations are found at sites with seasonally varying vegetation cover. Poor performance is observed across products for soil moisture anomalies timeseries, with R values varying from 0.46 to 0.66. The performance of reanalysis products differs greatly across regions, climate, land covers and topographic conditions. In general, all products tend to overestimate in arid climates and underestimate in humid regions as well as grassland. Most reanalysis products perform poorly in steep terrain. Relatively low temporal correlation and high *Bias* are detected in some sites from west of the UK, which might be associated with relatively low bulk density and high soil organic carbon. Overall, ERA5-Land, CRA40, CFSv2, SMAP L4 and GLEAM exhibit superior performance compared to MERRA2, GLDAS-Noah and JRA55. We recommend ERA5-Land and CFSv2 could be used in humid climates, whereas SMAP L4 and CRA40 perform better in arid regions. SMAP L4 has good performance for cropland, while GLEAM is more effective in shrubland regions. Our findings also provide insights on directions for improvement of soil moisture products for product developers.

1 Introduction

30 Soil moisture plays a key role in water and energy interactions between the atmosphere and land surface (Zeng et al., 2015; Kim et al., 2018; Ling et al., 2021), which controls many physical processes in hydrology, meteorology and agriculture, such as, evapotranspiration, infiltration, runoff generation, drought development, crop growth, among others. Accurate and timely soil moisture information is critical for a wide range of environmental analyses such as hydrological and climate modelling (Yee et al., 2017; Al-Yaari et al., 2014; Brocca et al., 2012; Zheng et al., 2021), flood and drought predictions (Martínez-Fernández et al., 2016; Massari et al., 2018; Ford and Quiring, 2019; Massari et al., 2014), water resources and agriculture management (Chawla et al., 2020; Karthikeyan et al., 2020).

To date, soil moisture data are available from a variety of sources. Reanalysis products provide soil moisture data over long time periods (Li et al., 2005; Baatz et al., 2021) and typically merge soil moisture observations and land surface model output by adopting data assimilation techniques, which often results in better soil moisture estimation than satellite products (Naz et al., 2020; Beck et al., 2021; Mahto and Mishra, 2019). At present, reanalysis products have been employed in a wide range of fields such as hydrological model initialization (Zheng et al., 2020), flood modelling (Mcclean et al., 2023; El Khalki et al., 2020; Zheng et al., 2023), drought monitoring (Chen et al., 2019; El Khalki et al., 2020) and climatology research (Miralles et al., 2014). Currently, many reanalysis products exist including ERA5-Land (Muñoz-Sabater 2019, Muñoz-Sabater et al. 2021), CFSv2 (Saha et al. 2011, Saha et al. 2014), MERRA2 (GMAO 2015, Gelaro et al. 2017), JRA55 (JMA 2013, Kobayashi et al. 2015), GLDAS-Noah (Rodell et al. 2004, Beaudoin 2020), CRA40 (Liu et al. 2017, Li et al. 2021), GLEAM (Miralles et al. 2011, Martens et al. 2017) datasets and SMAP Level 4 datasets (Reichle et al., 2019; Reichle et al., 2017a; Reichle et al., 2017b) etc (one should note that technically speaking GLDAS-Noah and GLEAM datasets are global land model-based products, we termed them as ‘reanalysis products’ in this paper for consistency). The quality of these reanalysis products is of significant interest to researchers and their performance against point soil moisture observations has provided valuable guidance on potential applications and further improvement (Li et al., 2020; Xu et al., 2021; Zheng et al., 2022; Beck et al., 2021; Chen and Yuan, 2020; Ling et al., 2021). However, due to the heterogeneity in soil properties, topography and climate condition, low-density point measurements are not entirely representative of larger scale soil moisture information (Gruber et al., 2013). Previous works have extensively reported the limited evaluation reliability between point measurements and soil moisture products, because the observed discrepancies can be attributed to the spatial sampling error rather than the intrinsic error of soil moisture products (Dorigo et al., 2015; Crow et al., 2012; Gruber et al., 2013; Stillman et al., 2016; Miralles et al., 2010).

Cosmic-ray neutron sensors (CRNS) are a more recent soil moisture measurement technique, compared to other traditional methods, that can measure area-average soil moisture at the field scale by capturing the variations in water content in the soil profile by fast neutron detection (Zreda et al., 2008). The neutron counting rates data from the CRNS can be converted into soil moisture via conversion equations from Desilets et al. (2010), Dong et al. (2014) and Hawdon et al. (2014). Additional influences on the neutron counting rate, besides soil moisture, need to be accounted for including; atmospheric pressure (Zreda et al., 2012; Hawdon et al., 2014), incoming high energy neutron intensity (Desilets et al., 2006), atmospheric water vapour (Rosolem et al., 2013), and above ground biomass (Rivera Villarreyes et al., 2011; Baatz et al., 2015). CRNS sensor calibration is also a crucial step, which requires multiple soil samples taken from within the sensor footprint, oven-dried then weighted and averaged to give field-scale accurate soil moisture estimates (Köhli et al. 2015; Schrön et al. 2017; Power et al., 2021). The horizontal footprint of the CRNS varies approximately between 400m and 600m in diameter (Zreda et al., 2008; Zreda et al., 2012; Evans et al., 2016; Desilets and Zreda, 2013; Schrön et al., 2017), while the vertical measurement depths depend strongly on soil moisture content ranging from 0.1 m (under wet conditions) to 0.7 m (under dry conditions) (Franz et al., 2012; Rosolem et al., 2014). Given that the spatial variations in soil moisture and other factors such as micro-topography and land cover can be considered within the footprint area, the CRNS measurements are better suited for the evaluation of satellite and reanalysis products compared to point measurements whose signal tends to be more strongly associated with soil properties (Montzka et al., 2017; Dong et al., 2014; Peng et al., 2021; Kim et al., 2015; Desilets et al., 2010). Soil moisture estimated by CRNS presents a more compatible spatial scale with recent efforts to promote hyper-resolution large-scale hydrological and land surface models (Iwema et al., 2017; Wood et al., 2011; Bierkens et al., 2015).

The establishment of CRNS networks across the globe is ongoing. Following the development of a first national scale CRNS sensors network in the United States, called Cosmic-Ray Soil Moisture Observing System (COSMOS) (Zreda et al., 2012), other countries, such as Australia (Hawdon et al., 2014), Germany (Zacharias et al., 2011; Bogena, 2016), UK (Evans et al.,

2016; Cooper et al., 2021) and India (Upadhyaya et al., 2021), have also started to establish their national networks. Several studies have evaluated a variety of soil moisture data products against CRNS measurements over different regions such as United States (Kim et al., 2015), UK (Peng et al., 2021), Australia (Renzullo et al., 2014), Germany (Schmidt et al., 2022) and India (Upadhyaya et al., 2021). Yet, most studies only use a few individual CRNS sites (Renzullo et al., 2014; Kędzior and Zawadzki, 2016; Montzka et al., 2017; Mwangi et al., 2020), which hampers a comprehensive assessment at large scale. While some studies have evaluated soil moisture products against numerous CRNS sites (Kim et al., 2015; Montzka et al., 2017; Duygu and Akyürek, 2019), these studies did not consider the deviations across multiple CRNS networks caused by different calibration and neutron correction methods. That is because despite having now more than 200 CRNS sites across global operation providing soil moisture data (Andreasen et al., 2017), there has not been a community-wide consensus on best practices for sensor calibration and signal correction methods shared across the different networks. This has resulted in a non-harmonized dataset among networks to support large or global scale soil hydrology analysis (Rosolem et al., 2013; Hawdon et al., 2014; Power et al., 2021).

This work provides, for the first time, a systematic evaluation of frequently used reanalysis soil moisture products against a global dataset of harmonized CRNS measurements. We analyse the reanalysis products with contrasting climate, soil properties, land cover and topography to provide insights on explaining the differences in performance. Finally, we provide recommendations to researchers for selecting suitable reanalysis soil moisture products.

2 Data description

2.1 CRNS measurements

In this paper, we collected CRNS data from numerous networks globally and ensured the data were processed in a harmonized way to serve as the reference for evaluating reanalysis products. The geographical locations of CRNS sites collected in this study are shown in Figure 1. A total of 180 CRNS data were collected from COSMOS-UK (51 sites) (Stanley et al., 2021), COSMOS-Europe (66 sites) (Bogena et al., 2022), COSMOS USA (45 sites) (Zreda et al., 2012) and Australian CosmOz network (18 sites) (Hawdon et al., 2014). Details of each CRNS network are summarized in Table 1.

2.1.1 CRNS sensor calibration


The inverse relationship between fast neutrons and hydrogen atoms means that as neutron counts rise (fall), we know that the moisture content of the soil is decreasing (increasing). However, in order to convert this signal into volumetric soil moisture values, calibration of each sensor is required. This involves obtaining multiple samples of soil moisture profiles within the sensor footprint that are together combined to provide an average moisture content (Zreda et al., 2012). Each sample is subjected to oven-drying, providing us with gravimetric soil moisture values, which can be converted to volumetric soil moisture when multiplied by the dry soil bulk density of the soil sample. As our understanding of the sensor signal has grown, improvements to this calibration step have been developed which have been shown to provide more accurate results. In particular, revised weighting schemes have been derived that consider the increased sensitivity of the signal to soil moisture nearer the sensor (Köhli et al., 2015; Schrön et al., 2017), as well as research showing the benefit in conducting multiple calibration campaigns across different seasons (Iwema et al., 2015). Ultimately this calibration step will provide us with the so called N_0 number (i.e., the theoretical neutron count found in absolutely dry conditions), which is calculated by comparing the averaged field scale soil moisture value derived through the sampling campaign with the count rate at the time of sampling. This N_0 number is used to derive the ratio between the actual counting rate (N) and the theorised maximum counting rate (N_0)

115 in the Desilets et al (2010) equation for converting neutrons to soil moisture values. It's important to note therefore, that changes in this number, or differences in how this number is derived, can lead to biases in soil moisture values.

The COSMOS-UK and COSMOS-EUROPE datasets each use the aforementioned revised weighting schemes to calibrate the sensors (Bogena et al., 2022; Cooper et al., 2021). To ensure comparability the COSMOS-USA and CosmOz sites were updated to utilise the revised scheme using crspy (Power et al., 2021), which was possible thanks to the openly available calibration data provided by each of the networks (Zreda et al., 2012, Hawdon et al., 2014). When multiple calibration days were available calibration would be taken on more than one day, with the N0 number being the value that reduced the error across all calibration days. It should be noted that recently the CosmOz network updated their data to utilise the revised weighting scheme, however there is still a difference in incoming neutron intensity correction, necessitating harmonization through crspy (<https://cosmoz.csiro.au/about>, last accessed 12/01/2024). More detailed of CRNS data reprocessing can be found in section 2.1.2. The calibrated N0 values, along with information on how many calibration days were used are given in the supporting information (see 'CRNSsiteDataNEWR1.xlsx' file).

120

125



130

Figure 1: Locations of CRNS sites collected in this study. The aridity index (I_m) global map derived from (Knoben et al., 2018) is used as a reference. The shape of the dots represents the CRNS sites from different networks, while the colour of the dots denotes the land cover type of each site.

Table 1: Details of four CRNS network used in this study.

| Network name | Number of sites collected | Period of data collection | Key references |
|------------------|---------------------------|---------------------------|---|
| COSMOS-UK | 51 sites | 2013-2019 | (Cooper et al., 2021; Evans et al., 2016; Stanley et al., 2021) |
| COSMOS-Europe | 66 sites | 2010-present | (Bogena and Ney, 2021; Bogena et al., 2022) |
| COSMOS USA | 45 sites | 2009-present | (Zreda et al., 2012) |
| CosmOz Australia | 18 sites | 2010-present | (Hawdon et al., 2014; Mcjannet et al., 2021) |

2.1.2 Reprocessing of CRNS data

To remove the possible influence of different CRNS processing methodologies, CRNS data were processed using the Cosmic-Ray Sensor PYthon tool (crspy) to ensure a harmonized methodology (Power et al., 2021). The correction of the
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995

2.2 Reanalysis soil moisture products

Eight widely used reanalysis products that provide soil moisture data are evaluated in this study. The reanalysis products
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995

Table 2: Overview of eight reanalysis soil moisture products used in this study.

| Product name | Spatial resolution | Temporal resolution | Temporal coverage | Vertical soil layers | Unit | Reference |
|--------------|--------------------|---------------------|-------------------|--------------------------------------|--------------------------------|---|
| ERA5-Land | 0.1° | Hourly | 1950-present | 0-7cm, 7-28cm, 28-100cm, 100-289 cm | m ³ /m ³ | (Muñoz-Sabater et al., 2021; Muñoz-Sabater, 2019) |
| CFSv2 | 0.5° | Hourly | 2011-present | 0-10cm, 10-40cm, 40-100cm, 100-200cm | % | (Saha et al., 2014; Saha et al., 2011) |
| MERRA-2 | 0.5°×0.625° | Hourly | 1980-present | 0-5cm, 0-100cm, 0-bedrock (~1.3m) | m ³ /m ³ | (Gelaro et al., 2017; Gmao, 2015) |

| | | | | | | |
|---------------------|---------------|----------|-----------------|---|--------------------------------|--|
| JRA-55 | 0.563°×0.562° | 3-hourly | 1958-present | total of 3 soil layers, varying depths | kg/m ³ | (Jma, 2013; Kobayashi et al., 2015) |
| GLDAS- Noah v2.1 | 0.25° | 3-hourly | 2000-present | 0-10cm, 10-40cm, 40- 100cm, 100-200cm | kg/m ² | (Rodell et al., 2004; Beauloing, 2020) |
| CRA40 | 0.5° | Daily | 1979-2020 | 0-10cm, 10-40cm, 40- 100cm, 100-200cm | m ³ /m ³ | (Li et al., 2021; Liu et al., 2017) |
| GLEAM- 3.5a | 0.25° | Daily | 1980-2020 | surface (0-10cm), root-zone (varying depths) | m ³ /m ³ | (Martens et al., 2017; Miralles et al., 2011) |
| SMAP L4 | 9km (~0.1°) | 3-hourly | 2015.03-present | surface (0-5cm), root-zone (0-100cm) | m ³ /m ³ | (Reichle et al., 2022; Reichle et al., 2019) |

160 2.3 Ancillary data preparation

To provide insights into the possible reasons for differences in reanalysis products performance, we collated data on 11 possible factors representing climate (i.e., aridity, seasonality, snow, mean annual temperature and mean annual precipitation), soil properties (i.e., bulk density, soil organic carbon and lattice water), vegetation (land cover) and topography conditions (i.e., altitude and slope).

165 Three climate indices (i.e., aridity, aridity seasonality and fraction of precipitation as snow) derived by Knoben et al. (2018) were adopted, which have been proved to be more effective than Köppen-Geiger classification for revealing the climatic influence especially on streamflow signatures. In their methods, aridity I_m is calculated based on Thornthwaite's moisture index MI (Willmott and Feddema, 1992) by using equations (1)-(2), in which $P(t)$ and $E_p(t)$ are mean monthly precipitation and potential evapotranspiration values from the CRU TS v3.23 dataset (Harris et al., 2014). The range of aridity index is [-1, 170 1], where -1 indicates the most arid conditions and 1 denotes the most humid conditions. More details and equations for other two climate indices can be found in Knoben et al. (2018). Additionally, mean annual temperature (MAT) and mean annual precipitation (MAP) for each CRNS site are retrieved from ERA5-Land product as it has the highest spatial resolution.

$$MI(t) = \begin{cases} 1 - \frac{E_p(t)}{P(t)} & , \quad P(t) > E_p(t) \\ 0 & , \quad P(t) = E_p(t) \\ \frac{P(t)}{E_p(t)} - 1 & , \quad P(t) < E_p(t) \end{cases} \quad (1)$$

$$I_m = \frac{1}{12} \sum_{t=1}^{t=12} MI(t) \quad (2)$$

175 Soil properties data, i.e., bulk density, soil organic carbon content and lattice water, are provided in metadata from Power et al. (2021) and Bogena et al. (2022). Soil organic carbon represents the total organic carbon in the soil at the site, while lattice water represents the hydrogen contained in the mineral structures of the soil. In studies from Power et al. (2021) and Bogena et al. (2022), local measurements of soil properties data are collected for the majority of CRNS sites (bulk density: 98% sites; soil organic carbon content: 94% sites, lattice water: 98% sites), while the global raster-based SoilGrids soil dataset (Hengl et al., 2017) was used to provide data for the sites with missing measurements.

180 For the land cover attributes at each CRNS site, we reclassify the different land cover classes from Power et al. (2021) and Bogena et al. (2022) into a harmonised land cover classification including four land cover types: forest, cropland, shrubland

and grassland (see Table A1 in Appendix A). The land cover data collected in crspy is obtained from the ESA CCI Land Cover dataset (Esa Land Cover Cci Project Team and Defourny, 2019). A small proportion of COSMOS-Europe sites with unclear land cover classes (e.g., plantation, reforestation, orchard and heathland) that are hard to reclassify were checked in the high-resolution Sentinel-2 10m land use and land cover map (Karra et al., 2021).

185 Finally, metadata from Power et al. (2021) and Bogena et al. (2022) offer altitude information for each CRNS site. We used the 90 m MERIT DEM data (Yamazaki et al., 2017) to provide the topographic slope. To reduce the spatial scale mismatch of the topographic slope between CRNS site point location and its horizontal footprints, we calculated the average slope of the area with a radius of 250m centered on the CRNS site.

3 Methods

190 3.1 Data processing

3.1.1 Temporal resolution and coverage

Due to the nature of the CRNS technology, the hourly measurements might contain higher uncertainty compared to daily measurements (Zreda et al., 2008; Desilets et al., 2010; Iwema et al., 2021), in general, high measurement uncertainty can be compensated by applying simple daily averaging (Davies et al., 2022). Additionally, some reanalysis products only provide daily data, thus we aggregate sub-daily reanalysis products data over the beginning of the day (00:00 hours UTC) and the proceeding 24 hour period, then perform the evaluation with the CRNS measurements at daily scale. Since CRA40 and GLEAM data provide data until the end of 2020, all available soil moisture data for each product from the start date of CRNS measurements to 2020 are used for the evaluation. To ensure the reliability of evaluation with enough soil moisture data, only the CRNS sites available with at least two years (730 days) of observations are selected for the analysis to avoid deviations under short-term extreme weather conditions. Therefore, a total of 135 (i.e., UK: 45 sites; mainland Europe: 41 sites; USA: 38 sites; Australia: 11 sites) out of 180 CRNS sites are used in the study. Note that since the evaluation against COSMOS-UK network has been specifically compared, the analysis for Europe only compares the COSMOS-Europe sites from mainland European regions, omitting the UK. The basic information of these selected 135 CRNS sites is listed in Table S2.

195
200

3.1.2 Spatial scale matching

205 The selected reanalysis products in this study exhibit a great variety of spatial resolutions (Table 2). The measurements from each CRNS site are compared with the reanalysis product grid cells in which the CRNS site is situated. In some cases, more than one CRNS site is located within one reanalysis product grid cell. Many studies take the average of multiple CRNS sites data first before the comparison with the grid cell data (Kim et al., 2015; Miralles et al., 2014). Yet, when many CRNS sites are located densely within the same grid cell (e.g., 13 CRNS German sites are located within one grid cell of CRA40 reanalysis product), different overlapping time periods across sites make it difficult to take an average. In this study, to maximize the use of CRNS data, we individually compare and calculate the statistical metric for the multiple CRNS sites data that are located in the same grid cell with the corresponding grid cell multiple times. Then, the Brunke ranking method (details introduced below in section 3.3) is used to comprehensively compare the product performance based on these statistic metrics.

210

3.1.3 Vertical footprint matching

215 For CRNS measurements, the vertical sensing depth has a strong dependency on actual soil moisture. The wetter the soil, the shallower the signal. The neutron signal exhibits the highest sensitivity to the uppermost layers and decays nearly exponentially from the surface downwards (Köhli et al., 2015; Zreda et al., 2008). The effective vertical sensing depth D86, defined as the

depth within which 86% of neutrons probed the soil (Köhli et al., 2015), ranging from 10 to 70 cm deep varies at each time step. By contrast, reanalysis products normally provide data for multiple soil layers.

220 To solve the inconsistency of vertical footprint between CRNS measurements and reanalysis products, the revised vertical weighting function, which is initially proposed to calculate vertical weighted averages of point measurements for sensor calibration (Schrön et al., 2017), is used to determine the weights for each soil layer of the reanalysis product. This revised vertical weighting function, assigning weights to soil layers, outperformed on better temporal correlation with CRNS measurements than that of other vertical processing methods (Figure S3). Following the same procedure, we assign weights
225 for each soil layer of the reanalysis product at different depths d and calculate the weighted average to compare with CRNS measurements. The formula for the revised vertical weighting function w_i is given in equation (3)-(4). The function to calculate the vertical average of soil layers i with values θ_i and weights w_i is shown in equation (5).

$$w_i = e^{-2d/D} \quad (3)$$

$$D \equiv D_{86}(r^*, \theta, \rho_{bulk}) \quad (4)$$

$$wt(\theta, w) = \frac{\sum_{i=1}^n w_i \theta_i}{\sum_{i=1}^n w_i} \quad (5)$$

Where D represents the effective penetration depth D_{86} . The variation of D_{86} is related to the adjusted distance r^* from the sensor centre (which is influenced by atmospheric pressure, (Schrön et al., 2017)), soil moisture wetness θ and soil bulk density
230 ρ_{bulk} . Since D_{86} is provided along with CRNS measurements at each time step, the weights w_i at different depths d can be obtained with the exponential function. n denotes the total number of reanalysis product soil layers up to D_{86} . The units of all the soil moisture values θ_i from reanalysis products and measurements are transferred into m^3/m^3 for comparison in this paper.

3.1.4 Data post-processing

Neutron signals can be substantially affected by snow cover, resulting in unreliable soil moisture measurements. Yet, at most
235 of the CRNS sites, there is a lack of measured snow data. Consequently, we follow the same procedure adopted in COSMOS-Europe (Bogena et al., 2022) to discard the soil moisture data affected by the presence of snow for other CRNS networks. Snow water equivalent data from ERA5-Land product is used to detect the snow events. CRNS data are excluded from the analysis when the 24 h moving average of the snow water equivalent data exceeds 1 mm.

3.2 Statistical metrics

240 The statistical metrics we used in this study include the Pearson correlation coefficient (R), Pearson correlation coefficient for seasonal (R_{sea}) and anomaly (R_{ano}) soil moisture timeseries, the mean square error (MSE), the unbiased root mean square error ($ubRMSE$), and the *Bias*. These statistical metrics are widely used in soil moisture data evaluation and capture different aspects of soil moisture dynamics (Peng et al., 2021; Cui et al., 2018; González-Zamora et al., 2019; Peng et al., 2015; Albergel et al., 2012; Yee et al., 2017; Zheng et al., 2022; Xu et al., 2021; Al-Yaari et al., 2014).

$$R = \frac{COV(SM_{product}, SM_{CRNS})}{\sigma_{product} \sigma_{CRNS}} \quad (6)$$

$$MSE = \overline{(SM_{product} - SM_{CRNS})^2} \quad (7)$$

$$ubRMSE = \sqrt{\overline{((SM_{product} - \overline{SM}_{product}) - (SM_{CRNS} - \overline{SM}_{CRNS}))^2}} \quad (8)$$

$$Bias = \overline{SM_{product}} - \overline{SM_{CRNS}} \quad (9)$$

245 Where cov denotes the covariance of both variables. $SM_{product}$ is the reanalysis soil moisture product, and SM_{CRNS} is the soil moisture derived from CRNS measurements. σ is the standard deviation of soil moisture values. The overbar represents the mean operator.

250 The R metric measures how well the soil moisture derived from CRNS measurements and reanalysis products correspond in terms of temporal correlation. Since the spatial scale mismatch differences between site measurements and soil moisture reanalysis data are inevitable (Beck et al., 2021; Miralles et al., 2010; Gruber et al., 2020), the comparisons in R metrics are considered to be the most reliable (Kim et al., 2015). To quantify the temporal dynamic performance of the soil moisture timeseries at different time scales, the original soil moisture timeseries data are decomposed into the seasonal signals and anomalies (Zheng et al., 2022; Beck et al., 2021; Kim et al., 2015; Peng et al., 2015; Al-Yaari et al., 2014; Li et al., 2020). The seasonal cycle data are derived by taking a moving average with a window size of 31-day over the soil moisture data time period coverage. Then, the anomaly timeseries are calculated by removing the seasonal signals from the original soil moisture data. The moving mean is extracted only if more than 16 days with available soil moisture values are present in the 31-day window. The Pearson correlation coefficients calculated for seasonal and anomaly soil moisture timeseries data are denoted as R_{sea} and R_{ano} , respectively.

3.3 Brunke ranking method

260 This study aims to provide recommendations for researchers in choosing suitable reanalysis soil moisture products. To comprehensively quantify the performances for 8 reanalysis products in terms of all 6 statistical metrics, the Brunke ranking scheme (Brunke et al., 2003) is adopted, which is a frequently used soil moisture products ranking method (Deng et al., 2021; Yang et al., 2020; Wang and Zeng, 2012; Deng et al., 2020; Decker et al., 2012).

265 For each statistical metric at each site, the 8 reanalysis products are ranked and assigned a score from 1 to 8, with 1 given to the products with the best performance (e.g., the lowest value of MSE , $ubRMSE$ and $Bias$ or highest correlation) and 8 given to the lowest performance (e.g., the largest value of MSE , $ubRMSE$ and $Bias$ or lowest correlation). It should be noted that if the metric values are missing for some reanalysis products at one site due to insufficient timeseries or missing values in a specific grid cell, the ranking score is given from 1 to the number of available products metric values (Wang and Zeng, 2012). To obtain the overall ranking score of each product, the ranking scores are further averaged across all 6 metrics for all sites.

270 4 Results

4.1 Rank of reanalysis products for different regions

275 Figure 2 displays the Brunke ranking results for 8 reanalysis products against CRNS measurements in terms of 6 statistical metrics (Table S3). More details for each metric can be found in the supporting information (Figure S4, S5). Overall, the performance of reanalysis products varies across different regions. In the UK, CFSv2 exhibits good performance in terms of R , R_{sea} and $ubRMSE$. GLEAM also ranks relatively high in multiple metrics, especially R_{sea} , MSE and $Bias$. As for mainland Europe, ERA5-Land performs well in terms of R , R_{sea} , R_{ano} and $ubRMSE$. JRA55 shows minimum $Bias$ and better MSE relative to other datasets in Europe. In the USA, CRA40 shows superior performance in terms of $Bias$ and MSE , while GLEAM provides better R , R_{sea} and $ubRMSE$. CFSv2 performs relatively poor in the USA. As for Australia, SMAP L4 exhibits good temporal correlation with both original and seasonal measured soil moisture timeseries and also performs well in MSE , $ubRMSE$. JRA55 presents the lowest rank in terms of R , R_{sea} and $ubRMSE$.




Figure 2: Brunke ranking results for a total of 8 products performance in terms of 6 statistical metrics across different regions, i.e., (a) UK, (b) mainland Europe, (c) USA and (d) Australia (AUS). Each coloured line represents a reanalysis product. For all metrics, the farther away of the line in this plot from the centre (i.e., closer towards black arrow) indicates good performance (i.e., the lowest value of MSE , $ubRMSE$ and $Bias$ or highest correlation).

285

Figure 3 summarizes the spatial distribution of the average performance for all 8 soil moisture reanalysis products in terms of R and $Bias$. The spatial map of the rest of the metrics can be found in Figure S6 and Figure S7. Around 70% of the sites exhibit good temporal correlation with CRNS measurements, with R average values of 8 soil moisture products larger than 0.7, and the median of R value across all sites reaches 0.74. Few sites in the west of the UK, southwest and north middle of the USA show worse performance with $R < 0.5$. In terms of $Bias$, all reanalysis soil moisture products tend to underestimate in the west of the UK, central Europe, northeastern USA and southeastern Australia, while overestimation is observed in northern Europe, northern Australia and most sites in western USA. The spatial distribution of R_{sea} is similar to that of R (Figure S6a, b and c). The performance of R_{ano} is generally lower than the correlation of original and seasonal soil moisture time series. No clear spatial pattern is observed in the USA and Australia in terms of MSE and $ubRMSE$, while several sites with high MSE and $ubRMSE$ values are notable in the UK (Figure S7).

290

295




Figure 3: Spatial distribution of all 8 products average performance in terms of statistical metrics R and $Bias$. The green letter G denotes the statistical metric values with good performance. The size of the dots in the map indicates the length of the measurements (i.e., number of days).

300 Soil moisture timeseries comparison between reanalysis soil moisture products and CRNS measurements for 4 representative sites are presented in Figure 4. Site WADDN in the UK is a grassland site from the COSMOS-UK network. The soil moisture values from reanalysis products for this site closely follow the temporal trend of the CRNS measurements, exhibiting high temporal correlation with the $\bar{R} = 0.85$ (Figure 4a). In contrast, the UK site RDMER is selected to demonstrate the variations in timeseries data for sites with low bulk density and high organic soils (Figure 4b). All reanalysis soil moisture products show less variability of temporal dynamics and exhibit low performance in describing soil moisture anomalies.

305 Moreover, the large deviation in $Bias$ between reanalysis soil moisture products and CRNS measurements is notable except for ERA5-Land and SMAP L4. Even though the reanalysis soil moisture products and CRNS measurements both provide accurate soil moisture information in an ideal situation, the mismatch in spatial resolution is still inevitable, which might also result in large bias (Montzka et al., 2017; Kim et al., 2015; Peng et al., 2021). Higher spatial resolution might reduce the bias, thus the smaller bias of ERA5-Land and SMAP L4 is likely attributed to the finer spatial resolution. In addition, a great variability of soil moisture data at hourly time step is observed in RDMER site (grey scatters in Figure 4b), while the hourly data in WADDN exhibits reasonable temporal dynamic. This also indicates that it is challenging to capture the temporal soil moisture variations at hourly scale for sites with low bulk density and high organic soils. This might be related to the fact that organic soils contain hydrogen and also the variations of organic soil content could lead to uncertainties in soil moisture calibration (Dimitrova-Petrova et al., 2021; Peng et al., 2021; Bogena et al., 2013).




Figure 4: Time series comparison between CRNS measurements and reanalysis products over 4 sites (i.e., (a) site WADDN from the UK; (b) site RDMER from the UK; (c) site Hauser Farm South from the USA; (d) site Rosemount from the USA). The locations of these 4 sites are listed in Figure 3.

320 Figure 4(c) and (d) show the timeseries comparison of two US sites with low average R values. The timeseries from reanalysis soil moisture products for site Hauser Farm South generally capture the temporal variations against CRNS observations. Yet,

the reanalysis soil moisture products do not dry out as much as CRNS measurements during the dry periods. Accordingly, the reanalysis soil moisture products fail to represent the accurate soil moisture condition in the sites located in the dry regions.

Site Rosemount from the USA is affected by snow cover in winter, thus the CRNS measurements during the snow events are discarded. Since the metrics are calculated only when CRNS observations and reanalysis soil moisture products are both available, the low average R value indicates poor temporal correlation during the growing season rather than the effect of snow. It is clear that the sites in USA with low temporal dynamic are normally from the cropland or shrubland land cover type (with the comparison of Figure 3b and Figure 1b). The seasonal variations of the biomass signals in cropland and shrubland might be the reasons that affect the deviations and low temporal correlation between CRNS measurements and reanalysis soil moisture products, as currently the harmonization of all sites does not explicitly account for changes in biomass.

4.2 Possible reasons for the differences in performance

All selected CRNS sites are adopted to investigate the different performances of reanalysis products under various conditions. The boxplots in Figure 5 show the distribution of site performance for each reanalysis product with the influential factors filled in colour. In particular, bulk density, soil organic carbon, aridity and mean annual precipitation are four informative variables in explaining the reanalysis performance. The distribution of some other possible factors (i.e., seasonality, snow), which shows insignificant influence, is not presented in this paper. Yet, data of all 11 possible factors for each site is provided in the supplementary information for those seeking further investigation. From Figure 5, the sites in the dry regions exhibit better performance in terms of R_{ano} than that of humid regions. These boxplots also demonstrate the performance for each reanalysis soil moisture product by adopting all CRNS sites data. For direct comparison, the median value of a given statistical metric across all available sites is often used to reflect the product performance (Beck et al., 2021; Deng et al., 2020). It is notable that GLDAS performs worse in terms of R_{ano} (Figure 5c).




Figure 5: Influential factors in explaining the reanalysis product performance. The shape of the dots denotes the CRNS sites from different networks (UK: circle; mainland Europe: square; USA: triangle; Australia: diamond), while the colour of the dots represents the values of possible factors for each site. The green letter G denotes the good performance of the statistical metric values.

4.2.1 Soil properties

With the comparison of the statistical metrics average performance map (Figure 3) and the spatial map of bulk density and soil organic carbon for all sites used in this study (Figure S8), it is clear that all reanalysis products exhibit low temporal correlation and high bias in the sites with low bulk density and high soil organic carbon. Almost all statistical metrics have lower performance on these sites (Figure 5b, 5d, 5e and Figure S9). These sites with low bulk density and high soil organic carbon exhibiting negative *Bias* are mainly from the humid region, especially the UK (Figure 5f).

4.2.2 Land cover

All reanalysis products tend to have lower performance in terms of R , R_{sea} and $ubRMSE$ metrics at shrubland and several sites at cropland, indicating that the reanalysis products exhibit poor performance in regions characterized by high mean annual temperature, low mean annual precipitation, and high altitude (Figure 6). The average Brunke ranking scores show that GLEAM performs best at shrubland (Table S3). The performance for each reanalysis product under four land cover types is presented in Figure 7. As for the forest land cover type, reanalysis products show small errors in terms of $ubRMSE$, but perform worse in MSE (Figure 7e, d). The *Bias* in grassland from a total of 6 reanalysis products is primarily negative, which means that the reanalysis products tend to underestimate the soil moisture observations in grassland (Figure 7f). CFSv2 performs best in grassland in terms of R and $ubRMSE$. For cropland land cover type, ERA5-Land relatively captures well the temporal dynamic of soil moisture timeseries (i.e., R , R_{sea} and R_{ano}) and $ubRMSE$, whereas SMAP L4 also shows good performance in terms of R_{ano} .





Figure 6: Average statistical metrics R , R_{sea} and $ubRMSE$ performance of all 8 products under different land cover types, altitude, mean annual precipitation and mean annual temperature. The shape of the dots denotes the CRNS sites from different networks (UK: circle; mainland Europe: square; USA: triangle; Australia: diamond). The colour of dots in (a) denotes different land cover types, while the colour in other subplots represents the statistical metric values. MAT stands for mean annual temperature, and MAP is mean annual precipitation.



370 **Figure 7: Statistical metric performance for all products under four land cover types (Forest: 33 sites; Cropland: 41 sites; Shrubland: 20 sites; Grassland: 41 sites). The values of the dots represent the median metric values of the sites in a given land cover type, the error bar of each dot denotes the variability of the metric values. The green letter G stands for the good performance of the statistical metric values.**


4.2.3 Climate

375 Figure 8 displays the performance of each reanalysis product under three climate conditions (humid, temperate, and arid). In general, compared to the humid and temperate climate, all reanalysis products perform noticeably better in terms of R_{ano} in arid climates but overestimate the CRNS measurements in terms of $Bias$ (Figure 5c and 5f). In contrast, the $Bias$ of reanalysis products for humid regions is primarily negative, indicating underestimation in humid regions. Moreover, large errors are observed in terms of R_{ano} and $ubRMSE$ from all reanalysis products in humid climates, highlighting that the soil moisture anomalies information is difficult to capture accurately by reanalysis products. In temperate regions, reanalysis products exhibit good performance in terms of $ubRMSE$ and $Bias$. Specifically, MERRA2 performs worse in humid and temperate climates particular in terms of R , R_{sea} and $ubRMSE$. CRA40 and SMAP L4 show better performance in arid regions, especially for the metrics R_{ano} and MSE , whereas JRA55 exhibit large errors in terms of R , R_{sea} and $ubRMSE$ in arid regions.


380

4.2.4 Slope

385 Most reanalysis soil moisture products perform worse in areas of steep terrain, especially the metrics including R , R_{ano} , MSE , $ubRMSE$ and $Bias$ (Figure 9). The values in R_{sea} under different topographic slopes from several reanalysis products (i.e., ERA5-Land, CFSv2, GLDAS, JRA55, and SMAP L4) are close to each other, which indicates the performance in describing the temporal dynamic of seasonal soil moisture pattern does not depend on the terrain slope (Figure 9b).



390 **Figure 8: Statistical metric performance for all products under three climate conditions (Humid: 53 sites; Temperate: 42 sites; Arid: 40 sites). The values of the dots represent the median metric values of the sites in a given climate zone, while the error bar of each dot denotes the variability of the metric values. The green letter G stands for the good performance of the statistical metric values. Im denotes the aridity index, which is described and shown in Figure 1.**



395 **Figure 9: Statistical metric performance for all products under different topographic slopes (steep terrain: 55 sites; flat terrain: 80 sites). The values of the dots represent the median metric values of the sites in a given terrain slope, the error bar of each dot denotes the variability of the metric values. The green letter G indicates the good performance of the statistical metric values.**

5 Discussion

5.1 Reanalysis products performance

400 This study has selected 135 CRNS sites across numerous networks globally and ensured the data were processed in a harmonized way to perform the assessment. We found that the performance of reanalysis soil moisture products varies in different regions, and this can be explained by climate, soil properties, land cover and topography conditions. Our findings provide recommendations in choosing reanalysis soil moisture products for use and insights on how to improve the accuracy of the evaluated datasets.

405 We find that while all reanalysis soil moisture products generally exhibit good agreement in the temporal correlation of soil moisture original timeseries (Figure 5a), large deviations in temporal correlation during the growing season are observed in cropland and shrubland (Figure 4d and Figure 6). Low performance in cropland and shrubland might be attributed to strong biomass signal seasonal variations, which affects the accuracy of CRNS measurements. Seasonally varying vegetation cover or high amounts of vegetation biomass are found to be key sources of uncertainty in CRNS measurements (Andreasen et al., 2017; Zreda et al., 2012; Franz et al., 2013b; Bogena et al., 2013; Iwema et al., 2021). Montzka et al. (2017) also identified the challenges in evaluating satellite soil moisture products against CRNS data at sites with seasonally changing vegetation cover. Besides, the lower correlation of reanalysis or satellite soil moisture products over densely vegetation regions are reported in studies from (Hagan et al., 2019; Beck et al., 2021; Kim et al., 2015; Kim et al., 2020), indicating the need for improving the vegetation parameters in land surface model or soil moisture retrieving algorithms (Baatz et al., 2015).

415 Furthermore, the low skill in capturing the temporal correlation of soil moisture anomalies timeseries is detected (Figure 5c), which means that reanalysis soil moisture products generally show poor response to precipitation events. Similar findings are reported in previous work (Hagan et al., 2019; Ling et al., 2021; Naz et al., 2020). Hagan et al. (2019) evaluated 7 reanalysis products (i.e., ERA5, ERA-Interim, MERRA1, MERRA2, MERRA-Land, Noah 1.0, Noah 2.5) against ground measurements and showed the R_{ano} of all soil moisture products are below 0.6. It is found that GLDAS has a weak ability in capturing the soil moisture anomalies (Deng et al., 2020; Naz et al., 2020), which is consistent with the notable R_{ano} performance of GLDAS in our results. Moreover, Beck et al. (2017) evaluated 22 reanalysis and satellite precipitation datasets. Their study revealed that these precipitation products also tend to capture the monthly variation well but have lower performance in shorter timescales (i.e., Pearson correlation coefficient calculated for 3-day means, R_{3day}). This aligns with our findings that the reanalysis products tend to reproduce the seasonal pattern of the variables well but that it is hard to capture the anomalies.

425 Similar to previous studies, our results identify that climate, topographic slope, soil properties and land cover types are all influential factors in explaining the performance of the reanalysis products (Deng et al., 2020; Beck et al., 2021; Li et al., 2020; Hagan et al., 2019; Decker et al., 2012). In particular, our results show that all reanalysis soil moisture products tend to underestimate the soil moisture values in the sites located in humid regions, while overestimation is observed in arid climate (observed in comparison of Figure 1 and Figure 3d, e, f). Direct timeseries comparison over representative sites also confirms that reanalysis products generally exhibit negative bias and less variation in humid climate (Figure 4b), whereas all products are rarely dry out below $0.1 \text{ m}^3/\text{m}^3$ during the dry periods in arid climate (Figure 4c). This might be related to the fact that the soil moisture values from reanalysis products behave as an indicator of the soil wetness with a minimum threshold for soil moisture, for example, acting as residual or limiting soil wetness fraction (Koster et al., 2009). Currently, most of the reanalysis soil moisture product evaluation studies are mainly regional scale analysis, especially for China (Wu et al., 2021; Zheng et al., 2022; Xing et al., 2021; Qin et al., 2017; Yang et al., 2020; Cheng et al., 2019). Several studies have found that reanalysis products tend to overestimate soil moisture conditions in Qinghai-Tibet Plateau, northern China and Mongolia (regions with arid/semi-arid climate dominated) (Ling et al., 2021; Wen et al., 2014; Zheng et al., 2022; Xing et al., 2021; Yang et al., 2020),

or reanalysis products show worse performance in arid areas (Ling et al., 2021; Yang et al., 2021). Moreover, we also observed that most reanalysis products perform poorly in steep terrain, which is supported by previous studies (Beck et al., 2021; Kim et al., 2015; Ma et al., 2019; Yang et al., 2021; Nicolai-Shaw et al., 2015; Li et al., 2020).

Our results reveal that all reanalysis products show lower performance in terms of all statistical metrics at the sites with low bulk density and high soil organic carbon (Figure S9), which are particularly from the humid regions in the UK. Similar findings are reported in the evaluation of soil moisture products against COSMOS-UK data by Peng et al. (2021). This suggests that the inaccurate soil properties parameters in land surface models might be the cause of the large errors, as the soil organic carbon could exert a great impact on soil thermal as well as hydraulic properties, leading to deviations in soil moisture (Zhu et al., 2019; Chen et al., 2012; Lawrence and Slater, 2008; Hagan et al., 2019; Ling et al., 2021). The influence of soil organic carbon on the performance of satellite and reanalysis soil moisture products is frequently reported (Yang et al., 2020; Jonard et al., 2018; Qin et al., 2017; Xing et al., 2021). Moreover, relatively high uncertainties of CRNS hourly measurements over organic soils or in humid region (observed in Figure 4b) might also be the reason for explaining the lower performance for these UK sites. Although the impact of soil organic carbon on CRNS observations has been taken into account according to the crspy tool (Power et al., 2021) used in this study, as mentioned in previous studies, it is more difficult to obtain the accurate soil moisture estimation and quantify the uncertainties of CRNS observations at sites with high soil organic carbon or in humid climate (Iwema et al., 2021; Bogena et al., 2013; Peng et al., 2021; Sigouin et al., 2016; Zhu et al., 2014; Franz et al., 2013a; Dimitrova-Petrova et al., 2021).

5.2 Recommendations for selecting suitable reanalysis products

To provide recommendations for the users, we classified the reanalysis products into three categories according to the Brunke ranking scores (Table S3, Figure 10), which are calculated based on all 6 statistic metrics. We provide recommendations based on these scores regarding which product demonstrates better overall performance under various regions, climate, land cover and topographic slopes. A lower Brunke ranking score indicates the superior overall product performance, suggesting users to prioritize its selection for their analysis. The soil moisture products with top two lowest Brunke ranking scores for each category are labelled with “Ok”, indicating relatively good performance, whereas the soil moisture reanalysis products ranked last are labelled with “Lower performance”. In particular, the soil moisture products with Brunke ranking scores less than 3.2 (i.e., the product with this threshold score indicates its average ranking is within or around the top 3 in terms of all 6 metrics across most of the sites in this category) are labelled with “Higher performance”. The recommendations only highlight the notable performance of products (i.e., “Higher performance” or “Lower performance”) across regions, climate, land covers and topographic conditions. Yet, it should be noted that the differences in the median value of statistical metrics across all reanalysis products are relatively small, especially for the metric MSE and $ubRMSE$ (Figure 5). Figure 10 presents the intercomparison across different reanalysis soil moisture products, which does not mean that the product labelled with “Lower performance” is not acceptable.

Overall, ERA5-Land, CRA40, CFSv2, SMAP L4 and GLEAM could be good choices for global analysis, which generally exhibit better performance in most circumstances than MERRA2, GLDAS-Noah and JRA55 (Figure 10). Performances of reanalysis products vary across different regions. ERA5-Land ranks best in terms of R , R_{sea} , R_{ano} and $ubRMSE$ in mainland Europe (Figure 2). CFSv2 and GLEAM are good alternatives for UK, while SMAP L4 and ERA5-Land performs better in Australia, especially in describing the temporal correlation against the observations. CRA40 and SMAP L4 are the top two datasets that are suitable for the USA compared to other products.




Figure 10: Recommendations for choosing 8 reanalysis soil moisture products under various regions, climate, land cover and topographic slope conditions based on the average Brunke ranking scores.

Some products show excellent performance under specific climate, land cover or topographic conditions. For instance, CFSv2 can be an alternative for use in humid climate, temperate climate or regions in grassland, while SMAP L4 is suitable for arid climate. SMAP L4 also shows the good performance in cropland, which is consistent with the finding from Tavakol et al. (2019). Similar performance is observed in CRA40 and GLEAM, which both exhibit superior performance in shrubland. Although all reanalysis products show large bias in humid region, ERA5-Land and CFSv2 are the optimal choice for humid climate. ERA5-Land also performs well in forest, cropland, grassland and steep or flat terrain. High spatial resolution might be the reason for explaining the superior performance of ERA5-Land. In contrast, products with coarser spatial resolution (>0.5°), i.e., MERRA2 and JRA55, exhibit relatively poor performance, which is also observed by Li et al. (2020), Mahto and Mishra (2019). GLDAS-Noah ranks last for representing soil moisture conditions under forest, shrubland or steep terrain. Deng et al. (2020) also observed worse performance of GLDAS in many land cover types, especially the underestimation of soil moisture in forest. Reasons for low performance between reanalysis products and observations are diverse and complex. The performance of GLDAS-Noah and MERRA2 in predicting soil moisture could be related with the quality of meteorological forcing data and the soil property database (Zheng et al., 2022; Beck et al., 2021). Many other factors that are not included in this study might also contribute to the influence, e.g., land surface model structures and parameterization schemes (Deng et al., 2020; Yang et al., 2020; Xu et al., 2021). Future studies are encouraged to investigate the impact of these factors.

5.3 Limitations

The proposed method for resolving spatial scale and vertical footprint matching currently represents the most reasonable solution after comparing several available approaches. Further studies are encouraged to find the ideal solution especially for grids of multiple CRNS sites with different overlapping time periods and how to process reanalysis products data at various depths while considering the CRNS sites effective depth. In addition, the sites used in this study are from COSMOS-UK,

500 COSMOS-Europe, COSMOS USA and CosmOz Australia networks. Accordingly, the evaluation results are more applicable
in these regions and regions with similar climates, soil properties, land cover and topographic conditions. It should be noted
that the number of reanalysis product grid cells for evaluating the performance in Europe is limited (Table 3) as the majority
of CRNS sites in the COSMOS-Europe network are concentrated in a small area of Germany. For some reanalysis products
with large grid cells (e.g., CRA40), a total of 13 CRNS German sites are located within one grid cell, whereas the density of
505 sites is relatively sparse for the rest of Europe; e.g., 19 CRA40 cells are evaluated against 41 CRNS sites, giving a ratio of
0.46. The ratio of the corresponding reanalysis product grid cells used for evaluation in Europe is relatively low ranging from
0.46 to 0.78, while the lowest ratio of reanalysis product grid cells used for evaluation for the UK, USA and Australia is 0.73,
0.89 and 0.91, respectively. Additionally, the evaluation over Australia is also not very reliable due to lack of representativeness,
because huge parts of Australia are not sampled by CRNS. Thus, the evaluation results for mainland Europe and Australia are
510 less reliable than that of the other regions. Furthermore, it is observed that the ratio also varies across different reanalysis
products. The finer the spatial resolution, the lower the possibility that multiple CRNS sites located within the same grid cell.
Consequently, a large proportion of grid cells is used in ERA5-Land over different regions, while that ratio for CRA40, CFSv2,
MERRA2 and JRA55 is normally the smallest.

Table 3: The ratio of the reanalysis product grid cells and used CRNS sites for each region.

| Regions | ERA5-Land | CFSv2 | MERRA2 | JRA55 | GLDAS | CRA40 | GLEAM | SMAP L4 |
|-------------------|-----------|-------|--------|-------|-------|-------|-------|---------|
| UK (45 sites) | 1.00 | 0.78 | 0.82 | 0.78 | 0.98 | 0.73 | 0.98 | 1.00 |
| Europe (41 sites) | 0.78 | 0.49 | 0.49 | 0.54 | 0.63 | 0.46 | 0.63 | 0.76 |
| USA (38 sites) | 0.97 | 0.92 | 0.89 | 0.92 | 0.95 | 0.89 | 0.95 | 0.97 |
| AUS (11 sites) | 1.00 | 0.91 | 1.00 | 1.00 | 1.00 | 0.91 | 1.00 | 1.00 |

515 Regarding temporal coverage, our assessment of SMAP L4 product is limited by the period of record which begins in 2015.
Since 70% of our CRNS sites were established before 2015, the evaluation period for SMAP L4 is shorter than that for other
reanalysis products, with 10 CRNS sites exhibiting temporal overlaps of less than 2 years. SMAP L4 is included in this analysis
because it is a state-of-the-art data assimilation soil moisture reanalysis product in recent years, yet it should be noted that it is
evaluated over a shorter time period compared to the other reanalysis products.

520 In addition, the statistical metrics describing *Bias* might not be as reliable as the ones quantifying the temporal correlation.
Metrics such as R , R_{sea} and R_{ano} , which measures how well the reanalysis product soil moisture timeseries data consistent
with the soil moisture timeseries temporal dynamic, are the most reliable statistical metrics and also of interest for the majority
of soil moisture products applications (Beck et al., 2021; Kim et al., 2015; Gruber et al., 2020). By contrast, the reliability of
Bias metric might be affected, as the inherent scale discrepancy in soil depth and footprint between CRNS and reanalysis
525 product grid cells still remains a limit (Crow et al., 2012; Miralles et al., 2010; Albergel et al., 2012; Kim et al., 2015; Montzka
et al., 2017; Peng et al., 2021). The timeseries comparison in RDMER UK site indicates that the finer spatial resolution might
relieve the *Bias* caused by the scale mismatch (Figure 4b).

6 Conclusion

To assist researchers in choosing suitable reanalysis soil moisture products, this study systematically evaluates 8 reanalysis
530 soil moisture products against soil moisture field measurements from 135 CRNS sites across numerous networks globally,
which are processed in a harmonized way. The performance of reanalysis products under diverse soil properties, climates, land
cover and topographic conditions are also investigated.

All reanalysis products generally exhibit good agreement in terms of temporal correlation with the median of R values over 0.7, whereas the lower performance with R_{ano} values are detected, indicating the weaker ability of capturing the soil moisture anomalies. In particular, GLDAS has the lowest R_{ano} values (0.46) across all sites. Low correlations of reanalysis products are observed in cropland or shrubland with seasonally varying vegetation cover. As for the *Bias*, reanalysis soil moisture products tend to overestimate in arid regions and underestimate in humid regions as well as grassland. It is also notable that reanalysis products exhibit worse performance in steep terrain.

Performances of reanalysis soil moisture products differ among regions, climate, soil properties, land cover and topography conditions. CFSv2 exhibits good performance across the UK and Europe. ERA5-Land performs well in Europe and Australia. SMAP L4 is suitable to be used in Australia and USA. GLEAM is a viable choice for the UK, whereas CRA40 can be an alternative in representing soil moisture conditions for the USA. Generally, ERA5-Land and CFSv2 show superior performance in humid climate, while SMAP L4 and CRA40 are recommended for arid regions. MERRA2 is less effective in humid and temperate climate, whereas JRA55 performs poorly in arid climate. Besides, GLEAM performs best over shrubland, followed by CRA40. ERA5-Land and CRA40 are suitable for steep terrain. For users seeking one product for global analysis, ERA5-Land, CRA40, CFSv2, SMAP L4 and GLEAM are viable options, as these products generally show better performance. Yet, it is important to acknowledge that due to the availability of CRNS observations, the findings of this study are more applicable in UK, Europe, USA, Australia and regions with similar conditions.

We also find that all reanalysis products fail to provide good performance in all statistical metrics at the sites with low bulk density and high soil organic carbon. These sites are mainly from the humid regions, i.e., the UK. This might be attributed to the limitation in representing the process over low bulk density and high organic soils in the land surface model. It is also possible that CRNS technology is challenging to provide accurate soil moisture information over these soil properties in humid regions.

Appendix A

In this paper, we used the following classification (Table A1) to reclassify the land cover data from two works to four land cover types.

Table A1: Land cover classes from Power et al. (2021) and Bogena et al. (2022) along with the reclassified land cover types used in this work.

| No. | Land cover types used in this study | Land cover classes in metadata from Power et al. (2021) | Land cover classes in COSMOS-Europe Bogena et al. (2022) |
|-----|-------------------------------------|---|--|
| 1 | Forest | tree_needleleaved_evergreen_closed_to_open tree_broadleaved_deciduous_open tree_broadleaved_deciduous_closed_to_open tree_mixed tree_cover_flooded_fresh_or_brakish_water mosaic_tree_and_shrub tree_broadleaved_evergreen_closed_to_open | Forest Reforestation Plantation |
| 2 | Cropland | mosaic_cropland cropland_rainfed_herbaceous_cover cropland_irrigated | Cropland Orchard |
| 3 | Shrubland | shrubland | Shrubland |

| | | | |
|---|-----------|---------------------|--------------------------------|
| | | mosaic_herbaceous | Heathland |
| | | shrubland_deciduous | |
| 4 | Grassland | grassland | Grassland Sparse Vegetation |

Code/Data availability

560 The code of Cosmic-Ray Sensor PYthon tool (crspy) is available at <https://github.com/danpower101/crspy>. COSMOS-UK is freely available at <https://catalogue.ceh.ac.uk/documents/b5c190e4-e35d-40ea-8f8e-598da03a1185>. COSMOS-Europe is available at <https://doi.org/10.34731/x9s3-kr48>. COSMOS USA is available via <http://cosmos.hwr.arizona.edu/>. CosmOz Australia is available via <https://doi.org/10.25901/5e7ab81af0394>. ERA5-Land data is accessible at <https://www.ecmwf.int/en/era5-land>. CFSv2 data can be downloaded from <https://rda.ucar.edu/datasets/ds094.1/>. MERRA-2

565 data can be found from <https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2/>. JRA55 data is given at <https://rda.ucar.edu>. GLDAS-Noah v2.1 data is available from <https://disc.gsfc.nasa.gov/>. CRA40 can be obtained from <http://data.cma.cn/>. GLEAM v3.5a is available at <https://www.gleam.eu/>. SMAP L4 can be downloaded from <https://nsidc.org/data/spl4smau/versions/7>. The detailed information ('CRNSsiteDataNEW1.xlsx') and calculated statistical metrics ('StatisticMetricNEW1.xlsx') for each site can be found in the supplementary files.

570 Author contribution

Y Zheng conceptualized the work, Y Zheng, G Coxon and R Woods designed the study. D Power, R Rosolem, M Rico-Ramirez and D McJannet contributed to process CRNS observations. Y Zheng processed the reanalysis soil moisture products and performed the analysis. J Li and P Feng provided resources and funding. All authors contributed to the investigation and discussion of the results and writing, editing of the manuscript.

575 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

The financial support from China Scholarship Council (No.202006250090) for Y Zheng is gratefully acknowledged. Y Zheng and G Coxon were funded by a UKRI Future Leaders Fellowship award [MR/V022857/1]. J Li would like to thank National natural science foundation of China (Grant number 52279022). P Feng was supported by National natural science foundation of China (Grant number 52079086). R Rosolem and D Power would also like to acknowledge a number of funding agencies including the Engineering and Physical Sciences Research Council (EPSRC) Water Informatics: Science and Engineering Centre for Doctoral Training (WISE-CDT; grant no. EP/L016214/1), additional support from the MOSAIC Digital Environment Feasibility Study (grant no. NE/T005645/1), and the COSMIC-SWAMP IoT Enabled Cosmic Ray Sensors for

580 Irrigation Monitoring (grant no. NE/W004364/1), all of which are funded by the Natural Environment Research Council (NERC), and also from the International Atomic Energy Agency of the United Nations (IAEA/UN; project no. CRP D12014). Funding for the CosmOz network was provided through the Terrestrial Ecosystem Research Network (TERN) which is part of the National Collaborative Research Infrastructure Strategy (NCRIS) program. The authors express their great thanks to Magdalena Szczykulska, Elizabeth Cooper, Jonathan Evans from UKCEH and COSMOS-UK team for providing the N0

585

590 calibration data. The authors declare that they have no known competing financial interests or personal relationships that could
have appeared to influence the work reported in this paper.

References

- Al-Yaari, A., Wigneron, J.-P., Ducharne, A., Kerr, Y., De Rosnay, P., De Jeu, R., Govind, A., Al Bitar, A., Albergel, C., and Munoz-Sabater, J.: Global-scale evaluation of two satellite-based passive microwave soil moisture datasets (SMOS and AMSR-E) with respect to Land Data Assimilation System estimates, *Remote Sens. Environ.*, 149, 181-195, <https://doi.org/10.1016/j.rse.2014.04.006>, 2014.
- Albergel, C., De Rosnay, P., Gruhier, C., Muñoz-Sabater, J., Hasenauer, S., Isaksen, L., Kerr, Y., and Wagner, W.: Evaluation of remotely sensed and modelled soil moisture products using global ground-based in situ observations, *Remote Sens. Environ.*, 118, 215-226, <https://doi.org/10.1016/j.rse.2011.11.017>, 2012.
- 600 Andreasen, M., Jensen, K. H., Desilets, D., Franz, T. E., Zreda, M., Bogena, H. R., and Looms, M. C.: Status and perspectives on the cosmic-ray neutron method for soil moisture estimation and other environmental science applications, *Vadose Zone J.*, 16, 1-11, <https://doi.org/10.2136/vzj2017.04.0086>, 2017.
- Baatz, R., Bogena, H., Hendricks Franssen, H. J., Huisman, J., Montzka, C., and Vereecken, H.: An empirical vegetation correction for soil water content quantification using cosmic ray probes, *Water Resour. Res.*, 51, 2030-2046, <https://doi.org/10.1002/2014WR016443>, 2015.
- 605 Baatz, R., Hendricks Franssen, H., Euskirchen, E., Sihi, D., Dietze, M., Ciavatta, S., Fennel, K., Beck, H., De Lannoy, G., and Pauwels, V.: Reanalysis in Earth system science: Toward terrestrial ecosystem reanalysis, *Rev. Geophys.*, 59, e2020RG000715, <https://doi.org/10.1029/2020RG000715>, 2021.
- Beaudoing, H. a. M. R., NASA/GSFC/HSL: GLDAS Noah Land Surface Model L4 3 hourly 0.25 x 0.25 degree V2.1, Greenbelt, Maryland, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC) [dataset], 10.5067/E7TYRXPJKWOQ, 2020.
- Beck, H. E., Pan, M., Miralles, D. G., Reichle, R. H., Dorigo, W. A., Hahn, S., Sheffield, J., Karthikeyan, L., Balsamo, G., and Parinussa, R. M.: Evaluation of 18 satellite-and model-based soil moisture products using in situ measurements from 826 sensors, *Hydrol. Earth Syst. Sci.*, 25, 17-40, <https://doi.org/10.5194/hess-25-17-2021>, 2021.
- 615 Beck, H. E., Vergopolan, N., Pan, M., Levizzani, V., van Dijk, A. I. J. M., Weedon, G. P., Brocca, L., Pappenberger, F., Huffman, G. J., and Wood, E. F.: Global-scale evaluation of 22 precipitation datasets using gauge observations and hydrological modeling, *Hydrol. Earth Syst. Sci.*, 21, 6201-6217, 10.5194/hess-21-6201-2017, 2017.
- Bierkens, M. F., Bell, V. A., Burek, P., Chaney, N., Condon, L. E., David, C. H., de Roo, A., Döll, P., Drost, N., and Famiglietti, J. S.: Hyper-resolution global hydrological modelling: what is next? "Everywhere and locally relevant", *Hydrol. Process.*, 29, 310-320, <https://doi.org/10.1002/hyp.10391>, 2015.
- 620 Bogena, H., Huisman, J., Baatz, R., Hendricks Franssen, H. J., and Vereecken, H.: Accuracy of the cosmic-ray soil water content probe in humid forest ecosystems: The worst case scenario, *Water Resour. Res.*, 49, 5778-5791, <https://doi.org/10.1002/wrcr.20463>, 2013.
- Bogena, H. R.: TERENO: German network of terrestrial environmental observatories, *Journal of large-scale research facilities JLSRF*, 2, A52-A52, <https://doi.org/10.17815/jlsrf-2-98>, 2016.
- 625 Bogena, H. R. and Ney, P.: COSMOS-Europe: A European network of Cosmic-Ray Neutron Soil Moisture Sensors, Forschungszentrum Jülich, <https://doi.org/10.34731/x9s3-kr48> 2021.
- Bogena, H. R., Schrön, M., Jakobi, J., Ney, P., Zacharias, S., Andreasen, M., Baatz, R., Boorman, D., Duygu, M. B., and Eguibar-Galán, M. A.: COSMOS-Europe: a European network of cosmic-ray neutron soil moisture sensors, *Earth Syst. Sci. Data*, 14, 1125-1151, <https://doi.org/10.5194/essd-14-1125-2022>, 2022.
- 630 Brocca, L., Moramarco, T., Melone, F., Wagner, W., Hasenauer, S., and Hahn, S.: Assimilation of surface- and root-zone ASCAT soil moisture products into rainfall-runoff modeling, *IEEE Trans. Geosci. Remote*, 50, 2542-2555, 10.1109/TGRS.2011.2177468, 2012.
- Brunke, M. A., Fairall, C. W., Zeng, X., Eymard, L., and Curry, J. A.: Which bulk aerodynamic algorithms are least problematic in computing ocean surface turbulent fluxes?, *J. Climate*, 16, 619-635, [https://doi.org/10.1175/1520-0442\(2003\)016<0619:WBAAAL>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0619:WBAAAL>2.0.CO;2), 2003.
- 635 Chawla, I., Karthikeyan, L., and Mishra, A. K.: A review of remote sensing applications for water security: Quantity, quality, and extremes, *J. Hydrol.*, 585, 124826, <https://doi.org/10.1016/j.jhydrol.2020.124826>, 2020.
- Chen, S., Gan, T. Y., Tan, X., Shao, D., and Zhu, J.: Assessment of CFSR, ERA-Interim, JRA-55, MERRA-2, NCEP-2 reanalysis data for drought analysis over China, *Clim. Dynam.*, 53, 737-757, <https://doi.org/10.1007/s00382-018-04611-1>, 2019.
- 640 Chen, Y. and Yuan, H.: Evaluation of nine sub-daily soil moisture model products over China using high-resolution in situ observations, *J. Hydrol.*, 588, 125054, <https://doi.org/10.1016/j.jhydrol.2020.125054>, 2020.
- Chen, Y., Yang, K., Tang, W., Qin, J., and Zhao, L.: Parameterizing soil organic carbon's impacts on soil porosity and thermal parameters for Eastern Tibet grasslands, *Sci. China Earth Sci.*, 55, 1001-1011, <https://doi.org/10.1007/s11430-012-4433-0>, 2012.
- 645

- Cheng, M., Zhong, L., Ma, Y., Zou, M., Ge, N., Wang, X., and Hu, Y.: A study on the assessment of multi-source satellite soil moisture products and reanalysis data for the Tibetan Plateau, *Remote Sens.*, 11, 1196, <https://doi.org/10.3390/rs11101196>, 2019.
- 655 Cooper, H. M., Bennett, E., Blake, J., Blyth, E., Boorman, D., Cooper, E., Evans, J., Fry, M., Jenkins, A., and Morrison, R.: COSMOS-UK: national soil moisture and hydrometeorology data for environmental science research, *Earth Syst. Sci. Data*, 13, 1737-1757, <https://doi.org/10.5194/essd-13-1737-2021>, 2021.
- Crow, W. T., Berg, A. A., Cosh, M. H., Loew, A., Mohanty, B. P., Panciera, R., de Rosnay, P., Ryu, D., and Walker, J. P.: Upscaling sparse ground-based soil moisture observations for the validation of coarse-resolution satellite soil moisture products, *Rev. Geophys.*, 50, <https://doi.org/10.1029/2011RG000372>, 2012.
- 660 Cui, C., Xu, J., Zeng, J., Chen, K.-S., Bai, X., Lu, H., Chen, Q., and Zhao, T.: Soil moisture mapping from satellites: An intercomparison of SMAP, SMOS, FY3B, AMSR2, and ESA CCI over two dense network regions at different spatial scales, *Remote Sens.*, 10, 33, <https://doi.org/10.3390/rs10010033>, 2018.
- Davies, P., Baatz, R., Bogena, H. R., Quansah, E., and Amekudzi, L. K.: Optimal Temporal Filtering of the Cosmic-Ray Neutron Signal to Reduce Soil Moisture Uncertainty, *Sensors*, 22, 9143, 2022.
- 665 Decker, M., Brunke, M. A., Wang, Z., Sakaguchi, K., Zeng, X., and Bosilovich, M. G.: Evaluation of the reanalysis products from GSFC, NCEP, and ECMWF using flux tower observations, *J. Climate*, 25, 1916-1944, <https://doi.org/10.1175/JCLI-D-11-00004.1>, 2012.
- Deng, M., Meng, X., Lu, Y., Li, Z., Zhao, L., Hu, Z., Chen, H., Shang, L., Wang, S., and Li, Q.: Impact and Sensitivity Analysis of Soil Water and Heat Transfer Parameterizations in Community Land Surface Model on the Tibetan Plateau, *J. Adv. Model. Earth Sy.*, 13, e2021MS002670, <https://doi.org/10.1029/2021MS002670>, 2021.
- Deng, Y., Wang, S., Bai, X., Wu, L., Cao, Y., Li, H., Wang, M., Li, C., Yang, Y., and Hu, Z.: Comparison of soil moisture products from microwave remote sensing, land model, and reanalysis using global ground observations, *Hydrol. Process.*, 34, 836-851, <https://doi.org/10.1002/hyp.13636>, 2020.
- 670 Desilets, D. and Zreda, M.: Footprint diameter for a cosmic-ray soil moisture probe: Theory and Monte Carlo simulations, *Water Resour. Res.*, 49, 3566-3575, <https://doi.org/10.1002/wrcr.20187>, 2013.
- Desilets, D., Zreda, M., and Ferré, T. P.: Nature's neutron probe: Land surface hydrology at an elusive scale with cosmic rays, *Water Resour. Res.*, 46, <https://doi.org/10.1029/2009WR008726>, 2010.
- 675 Desilets, D., Zreda, M., and Prabu, T.: Extended scaling factors for in situ cosmogenic nuclides: new measurements at low latitude, *Earth Planet. Sci. Lett.*, 246, 265-276, <https://doi.org/10.1016/j.epsl.2006.03.051>, 2006.
- Dimitrova-Petrova, K., Rosolem, R., Soulsby, C., Wilkinson, M. E., Lilly, A., and Geris, J.: Combining static and portable Cosmic ray neutron sensor data to assess catchment scale heterogeneity in soil water storage and their integrated role in catchment runoff response, *J. Hydrol.*, 601, 126659, <https://doi.org/10.1016/j.jhydrol.2021.126659>, 2021.
- 680 Dong, J., Ochsner, T. E., Zreda, M., Cosh, M. H., and Zou, C. B.: Calibration and validation of the COSMOS rover for surface soil moisture measurement, *Vadose Zone J.*, 13, 1-8, <https://doi.org/10.2136/vzj2013.08.0148>, 2014.
- Dorigo, W., Gruber, A., De Jeu, R., Wagner, W., Stacke, T., Loew, A., Albergel, C., Brocca, L., Chung, D., and Parinussa, R.: Evaluation of the ESA CCI soil moisture product using ground-based observations, *Remote Sens. Environ.*, 162, 380-395, <https://doi.org/10.1016/j.rse.2014.07.023>, 2015.
- 685 Duygu, M. B. and Akyürek, Z.: Using cosmic-ray neutron probes in validating satellite soil moisture products and land surface models, *Water*, 11, 1362, <https://doi.org/10.3390/w11071362>, 2019.
- El Khalki, E. M., Trambly, Y., Massari, C., Brocca, L., Simonneaux, V., Gascoin, S., and Saidi, M. E. M.: Challenges in flood modeling over data-scarce regions: how to exploit globally available soil moisture products to estimate antecedent soil wetness conditions in Morocco, *Nat. Hazard Earth Sys.*, 20, 2591-2607, <https://doi.org/10.5194/nhess-20-2591-2020>, 2020.
- 690 ESA Land Cover CCI project team and Defourny, P.: ESA Land Cover Climate Change Initiative (Land_Cover_cci): Global Land Cover Maps, Version 2.0.7, Centre for Environmental Data Analysis [data set], available at: <https://catalogue.ceda.ac.uk/uuid/b382ebe6679d6644b6678b6670e6668ea6674ef6674b6701c> (last access: 6611 November 2021), 2019 (data available at: <http://cci.esa.int/data>, last access: 6610 January 2021), 2019.
- Evans, J., Ward, H., Blake, J., Hewitt, E., Morrison, R., Fry, M., Ball, L., Doughty, L., Libre, J., and Hitt, O.: Soil water content in southern England derived from a cosmic-ray soil moisture observing system-COSMOS-UK, *Hydrol. Process.*, 30, 4987-4999, <https://doi.org/10.1002/hyp.10929>, 2016.
- 695 Ford, T. W. and Quiring, S. M.: Comparison of contemporary in situ, model, and satellite remote sensing soil moisture with a focus on drought monitoring, *Water Resour. Res.*, 55, 1565-1582, <https://doi.org/10.1029/2018WR024039>, 2019.
- Franz, T., Zreda, M., Rosolem, R., and Ferre, T.: A universal calibration function for determination of soil moisture with cosmic-ray neutrons, *Hydrol. Earth Syst. Sci.*, 17, 453-460, <https://doi.org/10.5194/hess-17-453-2013>, 2013a.
- 700 Franz, T. E., Zreda, M., Ferre, T., Rosolem, R., Zweck, C., Stillman, S., Zeng, X., and Shuttleworth, W.: Measurement depth of the cosmic ray soil moisture probe affected by hydrogen from various sources, *Water Resour. Res.*, 48, <https://doi.org/10.1029/2012WR011871>, 2012.
- Franz, T. E., Zreda, M., Rosolem, R., Hornbuckle, B. K., Irvin, S. L., Adams, H., Kolb, T. E., Zweck, C., and Shuttleworth, W. J.: Ecosystem-scale measurements of biomass water using cosmic ray neutrons, *Geophys. Res. Lett.*, 40, 3929-3933, <https://doi.org/10.1002/grl.50791>, 2013b.
- 705

- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., and Reichle, R.: The modern-era retrospective analysis for research and applications, version 2 (MERRA-2), *J. Climate*, 30, 5419-5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 710 GMAO, G. M. a. A. O.: MERRA-2 tavgM_2d_Ind_Nx: 2d, Monthly mean, Time-Averaged, Single-Level, Assimilation, Land Surface Diagnostics V5.12.4, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC), 2015.
- González-Zamora, Á., Sánchez, N., Pablos, M., and Martínez-Fernández, J.: CCI soil moisture assessment with SMOS soil moisture and in situ data under different environmental conditions and spatial scales in Spain, *Remote Sens. Environ.*, 225, 469-482, <https://doi.org/10.1016/j.rse.2018.02.010>, 2019.
- 715 Gruber, A., Dorigo, W. A., Zwieback, S., Xaver, A., and Wagner, W.: Characterizing coarse-scale representativeness of in situ soil moisture measurements from the International Soil Moisture Network, *Vadose Zone J.*, 12, 1-16, <https://doi.org/10.2136/vzj2012.0170>, 2013.
- Gruber, A., De Lannoy, G., Albergel, C., Al-Yaari, A., Brocca, L., Calvet, J.-C., Colliander, A., Cosh, M., Crow, W., and Dorigo, W.: Validation practices for satellite soil moisture retrievals: What are (the) errors?, *Remote Sens. Environ.*, 244, 111806, <https://doi.org/10.1016/j.rse.2020.111806>, 2020.
- 720 Hagan, D. F. T., Parinussa, R. M., Wang, G., and Draper, C. S.: An evaluation of soil moisture anomalies from global model-based datasets over the people's republic of China, *Water*, 12, 117, <https://doi.org/10.3390/w12010117>, 2019.
- Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic observations—the CRU TS3. 10 Dataset, *Int. J. Climatol.*, 34, 623-642, <https://doi.org/10.1002/joc.3711>, 2014.
- 725 Hawdon, A., McJannet, D., and Wallace, J.: Calibration and correction procedures for cosmic-ray neutron soil moisture probes located across Australia, *Water Resour. Res.*, 50, 5029-5043, <https://doi.org/10.1002/2013WR015138>, 2014.
- Hengl, T., Mendes de Jesus, J., Heuvelink, G. B., Ruiperez Gonzalez, M., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M. N., Geng, X., and Bauer-Marschallinger, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PLoS one*, 12, e0169748, <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- 730 Iwema, J., Rosolem, R., Rahman, M., Blyth, E., and Wagener, T.: Land surface model performance using cosmic-ray and point-scale soil moisture measurements for calibration, *Hydrol. Earth Syst. Sci.*, 21, 2843-2861, 10.5194/hess-21-2843-2017, 2017.
- Iwema, J., Schrön, M., Koltermann Da Silva, J., Schweiser De Paiva Lopes, R., and Rosolem, R.: Accuracy and precision of the cosmic-ray neutron sensor for soil moisture estimation at humid environments, *Hydrol. Process.*, 35, e14419, <https://doi.org/10.1002/hyp.14419>, 2021.
- 735 JMA: JRA-55: Japanese 55-year Reanalysis, Daily 3-Hourly and 6-Hourly Data, Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory [dataset], 10.5065/D6HH6H41, 2013.
- Jonard, F., Bircher, S., Demontoux, F., Weihermüller, L., Razafindratsima, S., Wigneron, J.-P., and Vereecken, H.: Passive L-band microwave remote sensing of organic soil surface layers: A tower-based experiment, *Remote Sens.*, 10, 304, <https://doi.org/10.3390/rs10020304>, 2018.
- 740 Karra, K., Kontgis, C., Statman-Weil, Z., Mazzariello, J. C., Mathis, M., and Brumby, S. P.: Global land use / land cover with Sentinel 2 and deep learning, 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, 11-16 July 2021, 4704-4707, 10.1109/IGARSS47720.2021.9553499, 2021.
- Karthikeyan, L., Chawla, I., and Mishra, A. K.: A review of remote sensing applications in agriculture for food security: Crop growth and yield, irrigation, and crop losses, *J. Hydrol.*, 586, 124905, <https://doi.org/10.1016/j.jhydrol.2020.124905>, 2020.
- 745 Kędzior, M. and Zawadzki, J.: Comparative study of soil moisture estimations from SMOS satellite mission, GLDAS database, and cosmic-ray neutrons measurements at COSMOS station in Eastern Poland, *Geoderma*, 283, 21-31, <https://doi.org/10.1016/j.geoderma.2016.07.023>, 2016.
- Kim, H., Parinussa, R., Konings, A. G., Wagner, W., Cosh, M. H., Lakshmi, V., Zohaib, M., and Choi, M.: Global-scale assessment and combination of SMAP with ASCAT (active) and AMSR2 (passive) soil moisture products, *Remote Sens. Environ.*, 204, 260-275, <https://doi.org/10.1016/j.rse.2017.10.026>, 2018.
- 750 Kim, H., Wigneron, J.-P., Kumar, S., Dong, J., Wagner, W., Cosh, M. H., Bosch, D. D., Collins, C. H., Starks, P. J., and Seyfried, M.: Global scale error assessments of soil moisture estimates from microwave-based active and passive satellites and land surface models over forest and mixed irrigated/dryland agriculture regions, *Remote Sens. Environ.*, 251, 112052, <https://doi.org/10.1016/j.rse.2020.112052>, 2020.
- 755 Kim, S., Liu, Y. Y., Johnson, F. M., Parinussa, R. M., and Sharma, A.: A global comparison of alternate AMSR2 soil moisture products: Why do they differ?, *Remote Sens. Environ.*, 161, 43-62, <https://doi.org/10.1016/j.rse.2015.02.002>, 2015.
- Knoben, W. J., Woods, R. A., and Freer, J. E.: A quantitative hydrological climate classification evaluated with independent streamflow data, *Water Resour. Res.*, 54, 5088-5109, <https://doi.org/10.1029/2018WR022913>, 2018.
- 760 Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H., Kobayashi, C., and Endo, H.: The JRA-55 reanalysis: General specifications and basic characteristics, *Journal of the Meteorological Society of Japan. Ser. II*, 93, 5-48, <https://doi.org/10.2151/jmsj.2015-001>, 2015.
- Köhli, M., Schrön, M., Zreda, M., Schmidt, U., Dietrich, P., and Zacharias, S.: Footprint characteristics revised for field-scale soil moisture monitoring with cosmic-ray neutrons, *Water Resour. Res.*, 51, 5772-5790, <https://doi.org/10.1002/2015WR017169>, 2015.
- 765

- Koster, R. D., Guo, Z., Yang, R., Dirmeyer, P. A., Mitchell, K., and Puma, M. J.: On the nature of soil moisture in land surface models, *J. Climate*, 22, 4322-4335, <https://doi.org/10.1175/2009JCLI2832.1>, 2009.
- Lawrence, D. M. and Slater, A. G.: Incorporating organic soil into a global climate model, *Clim. Dynam.*, 30, 145-160, 2008.
- 770 Li, C., Zhao, T., Shi, C., and Liu, Z.: Assessment of precipitation from the CRA40 dataset and new generation reanalysis datasets in the global domain, *Int. J. Climatol.*, <https://doi.org/10.1002/joc.7127>, 2021.
- Li, H., Robock, A., Liu, S., Mo, X., and Viterbo, P.: Evaluation of reanalysis soil moisture simulations using updated Chinese soil moisture observations, *J. Hydrometeorol.*, 6, 180-193, <https://doi.org/10.1175/JHM416.1>, 2005.
- Li, M., Wu, P., and Ma, Z.: A comprehensive evaluation of soil moisture and soil temperature from third-generation atmospheric and land reanalysis data sets, *Int. J. Climatol.*, 40, 5744-5766, <https://doi.org/10.1002/joc.6549>, 2020.
- 775 Ling, X., Huang, Y., Guo, W., Wang, Y., Chen, C., Qiu, B., Ge, J., Qin, K., Xue, Y., and Peng, J.: Comprehensive evaluation of satellite-based and reanalysis soil moisture products using in situ observations over China, *Hydrol. Earth Syst. Sci.*, 25, 4209-4229, <https://doi.org/10.5194/hess-25-4209-2021>, 2021.
- Liu, Z., Shi, C., Zhou, Z., Jiang, L., Liang, X., Zhang, T., Liao, J., Liu, J., Wang, M., and Yao, S.: CMA global reanalysis (CRA-40): Status and plans, *Proc. 5th International Conference on Reanalysis*, 13-17, 2017.
- 780 Ma, H., Zeng, J., Chen, N., Zhang, X., Cosh, M. H., and Wang, W.: Satellite surface soil moisture from SMAP, SMOS, AMSR2 and ESA CCI: A comprehensive assessment using global ground-based observations, *Remote Sens. Environ.*, 231, 111215, <https://doi.org/10.1016/j.rse.2019.111215>, 2019.
- Mahto, S. S. and Mishra, V.: Does ERA-5 outperform other reanalysis products for hydrologic applications in India?, *J. Geophys. Res.-Atmos.*, 124, 9423-9441, <https://doi.org/10.1029/2019JD031155>, 2019.
- 785 Martens, B., Miralles, D. G., Lievens, H., Schalie, R. v. d., De Jeu, R. A., Fernández-Prieto, D., Beck, H. E., Dorigo, W. A., and Verhoest, N. E.: GLEAM v3: Satellite-based land evaporation and root-zone soil moisture, *Geosci. Model Dev.*, 10, 1903-1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.
- Martínez-Fernández, J., González-Zamora, A., Sánchez, N., Gumuzzio, A., and Herrero-Jiménez, C.: Satellite soil moisture for agricultural drought monitoring: Assessment of the SMOS derived Soil Water Deficit Index, *Remote Sens. Environ.*, 177, 277-286, <https://doi.org/10.1016/j.rse.2016.02.064>, 2016.
- 790 Massari, C., Camici, S., Ciabatta, L., and Brocca, L.: Exploiting satellite-based surface soil moisture for flood forecasting in the Mediterranean area: State update versus rainfall correction, *Remote Sens.*, 10, 292, <https://doi.org/10.3390/rs10020292>, 2018.
- Massari, C., Brocca, L., Barbetta, S., Papathanasiou, C., Mimikou, M., and Moramarco, T.: Using globally available soil moisture indicators for flood modelling in Mediterranean catchments, *Hydrol. Earth Syst. Sci.*, 18, 839-853, <https://doi.org/10.5194/hess-18-839-2014>, 2014.
- 795 McClean, F., Dawson, R., and Kilsby, C.: Intercomparison of global reanalysis precipitation for flood risk modelling, *Hydrol. Earth Syst. Sci.*, 27, 331-347, 2023.
- McJannet, D., Stenson, M., Sommer, A., and Hawdon, A.: CosmOz— The Australian Cosmic-ray Soil Moisture Sensor Network, Version 1.0.0, Terrestrial Ecosystem Research Network (TERN) [data set], <https://doi.org/10.25901/5e7ab81af0394>, 2021.
- 800 Miralles, D. G., Crow, W. T., and Cosh, M. H.: Estimating spatial sampling errors in coarse-scale soil moisture estimates derived from point-scale observations, *J. Hydrometeorol.*, 11, 1423-1429, <https://doi.org/10.1175/2010JHM1285.1>, 2010.
- 805 Miralles, D. G., Holmes, T. R. H., De Jeu, R. A. M., Gash, J. H., Meesters, A. G. C. A., and Dolman, A. J.: Global land-surface evaporation estimated from satellite-based observations, *Hydrol. Earth Syst. Sci.*, 15, 453-469, <https://doi.org/10.5194/hess-15-453-2011>, 2011.
- Miralles, D. G., Van Den Berg, M. J., Gash, J. H., Parinussa, R. M., De Jeu, R. A., Beck, H. E., Holmes, T. R., Jiménez, C., Verhoest, N. E., and Dorigo, W. A.: El Niño–La Niña cycle and recent trends in continental evaporation, *Nat. Clim. Change*, 4, 122-126, <https://doi.org/10.1038/nclimate2068>, 2014.
- 810 Montzka, C., Bogena, H. R., Zreda, M., Monerris, A., Morrison, R., Muddu, S., and Vereecken, H.: Validation of spaceborne and modelled surface soil moisture products with cosmic-ray neutron probes, *Remote Sens.*, 9, 103, <https://doi.org/10.3390/rs9020103>, 2017.
- Muñoz-Sabater, J.: ERA5-Land hourly data from 1981 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [dataset], <https://doi.org/10.24381/cds.e2161bac>, 2019.
- 815 Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., and Hersbach, H.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13, 4349-4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- Mwangi, S., Zeng, Y., Montzka, C., Yu, L., and Su, Z.: Assimilation of cosmic-ray neutron counts for the estimation of soil ice content on the eastern Tibetan Plateau, *J. Geophys. Res.-Atmos.*, 125, e2019JD031529, <https://doi.org/10.1029/2019JD031529>, 2020.
- 820 Naz, B. S., Kollet, S., Franssen, H.-J. H., Montzka, C., and Kurtz, W.: A 3 km spatially and temporally consistent European daily soil moisture reanalysis from 2000 to 2015, *Scientific data*, 7, 1-14, <https://doi.org/10.1038/s41597-020-0450-6>, 2020.
- Nicolai-Shaw, N., Hirschi, M., Mittelbach, H., and Seneviratne, S. I.: Spatial representativeness of soil moisture using in situ, remote sensing, and land reanalysis data, *J. Geophys. Res.-Atmos.*, 120, 9955-9964, <https://doi.org/10.1002/2015JD023305>, 2015.
- 825

- Peng, J., Niesel, J., Loew, A., Zhang, S., and Wang, J.: Evaluation of satellite and reanalysis soil moisture products over southwest China using ground-based measurements, *Remote Sens.*, 7, 15729-15747, <https://doi.org/10.3390/rs71115729>, 2015.
- Peng, J., Tanguy, M., Robinson, E. L., Pinnington, E., Evans, J., Ellis, R., Cooper, E., Hannaford, J., Blyth, E., and Dadson, S.: Estimation and evaluation of high-resolution soil moisture from merged model and Earth observation data in the Great Britain, *Remote Sens. Environ.*, 264, 112610, <https://doi.org/10.1016/j.rse.2021.112610>, 2021.
- Power, D., Rico-Ramirez, M. A., Desilets, S., Desilets, D., and Rosolem, R.: Cosmic-Ray neutron Sensor PYTHON tool (crspy 1.2.1): an open-source tool for the processing of cosmic-ray neutron and soil moisture data, *Geosci. Model Dev.*, 14, 7287-7307, <https://doi.org/10.5194/gmd-14-7287-2021>, 2021.
- Qin, Y., Wu, T., Wu, X., Li, R., Xie, C., Qiao, Y., Hu, G., Zhu, X., Wang, W., and Shang, W.: Assessment of reanalysis soil moisture products in the permafrost regions of the central of the Qinghai–Tibet Plateau, *Hydrol. Process.*, 31, 4647-4659, <https://doi.org/10.1002/hyp.11383>, 2017.
- Reichle, R. H., G. De Lannoy, R.D. Koster, W.T. Crow, J.S. Kimball, Q. Liu, and Bechtold, M.: SMAP L4 Global 3-hourly 9 km EASE-Grid Surface and Root Zone Soil Moisture Analysis Update, Version 7, Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center, <https://nsidc.org/data/spl4smau/versions/7>, 2022.
- Reichle, R. H., De Lannoy, G. J., Liu, Q., Ardizzone, J. V., Colliander, A., Conaty, A., Crow, W., Jackson, T. J., Jones, L. A., and Kimball, J. S.: Assessment of the SMAP level-4 surface and root-zone soil moisture product using in situ measurements, *J. Hydrometeorol.*, 18, 2621-2645, 2017a.
- Reichle, R. H., De Lannoy, G. J., Liu, Q., Koster, R. D., Kimball, J. S., Crow, W. T., Ardizzone, J. V., Chakraborty, P., Collins, D. W., and Conaty, A. L.: Global assessment of the SMAP Level-4 surface and root-zone soil moisture product using assimilation diagnostics, *J. Hydrometeorol.*, 18, 3217-3237, 2017b.
- Reichle, R. H., Liu, Q., Koster, R. D., Crow, W. T., De Lannoy, G. J. M., Kimball, J. S., Ardizzone, J. V., Bosch, D., Colliander, A., Cosh, M., Kolassa, J., Mahanama, S. P., Prueger, J., Starks, P., and Walker, J. P.: Version 4 of the SMAP Level-4 Soil Moisture Algorithm and Data Product, *J. Adv. Model. Earth Sy.*, 11, 3106-3130, <https://doi.org/10.1029/2019MS001729>, 2019.
- Renzullo, L. J., Van Dijk, A., Perraud, J.-M., Collins, D., Henderson, B., Jin, H., Smith, A., and McJannet, D.: Continental satellite soil moisture data assimilation improves root-zone moisture analysis for water resources assessment, *J. Hydrol.*, 519, 2747-2762, <https://doi.org/10.1016/j.jhydrol.2014.08.008>, 2014.
- Rivera Villarreyes, C., Baroni, G., and Oswald, S. E.: Integral quantification of seasonal soil moisture changes in farmland by cosmic-ray neutrons, *Hydrol. Earth Syst. Sci.*, 15, 3843-3859, <https://doi.org/10.5194/hess-15-3843-2011>, 2011.
- Rodell, M., Houser, P., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C.-J., Arsenault, K., Cosgrove, B., Radakovich, J., and Bosilovich, M.: The global land data assimilation system, *Bull Am. Meteorol. Soc.*, 85, 381-394, <https://doi.org/10.1175/BAMS-85-3-381>, 2004.
- Rosolem, R., Shuttleworth, W., Zreda, M., Franz, T. E., Zeng, X., and Kurc, S.: The effect of atmospheric water vapor on neutron count in the cosmic-ray soil moisture observing system, *J. Hydrometeorol.*, 14, 1659-1671, <https://doi.org/10.1175/JHM-D-12-0120.1>, 2013.
- Rosolem, R., Hoar, T., Arellano, A., Anderson, J. L., Shuttleworth, W. J., Zeng, X., and Franz, T. E.: Translating aboveground cosmic-ray neutron intensity to high-frequency soil moisture profiles at sub-kilometer scale, *Hydrol. Earth Syst. Sci.*, 18, 4363-4379, <https://doi.org/10.5194/hess-18-4363-2014>, 2014.
- Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., Chuang, H.-y., and Iredell, M.: The NCEP climate forecast system version 2, *J. Climate*, 27, 2185-2208, <https://doi.org/10.1175/JCLI-D-12-00823.1>, 2014.
- Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y., Chuang, H., Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M. P., H., v. d. D., Q., Z., W., W., M., C., and Becker, E.: NCEP Climate Forecast System Version 2 (CFSv2) Selected Hourly Time-Series Products. Research Data Archive at the National Center for Atmospheric Research [dataset], 2011.
- Schmidt, T., Schrön, M., Li, Z., and Peng, J.: Intercomparison of current soil moisture products from remote sensing and modeling over COSMOS field sites in Germany, *Copernicus Meetings*, <https://doi.org/10.5194/egusphere-egu22-10336>, 2022.
- Schrön, M., Köhli, M., Scheffele, L., Iwema, J., Bogena, H. R., Lv, L., Martini, E., Baroni, G., Rosolem, R., Weimar, J., Mai, J., Cuntz, M., Rebmann, C., Oswald, S. E., Dietrich, P., Schmidt, U., and Zacharias, S.: Improving calibration and validation of cosmic-ray neutron sensors in the light of spatial sensitivity, *Hydrol. Earth Syst. Sci.*, 21, 5009-5030, <https://doi.org/10.5194/hess-21-5009-2017>, 2017.
- Sigouin, M. J., Dyck, M., Si, B. C., and Hu, W.: Monitoring soil water content at a heterogeneous oil sand reclamation site using a cosmic-ray soil moisture probe, *J. Hydrol.*, 543, 510-522, <https://doi.org/10.1016/j.jhydrol.2016.10.026>, 2016.
- Stanley, S., Antoniou, V., Askquith-Ellis, A., Ball, L. A., Bennett, E. S., Blake, J. R., Boorman, D. B., Brooks, M., Clarke, M., Cooper, H. M., Cowan, N., Cumming, A., Evans, J. G., Farrand, P., Fry, M., Hitt, O. E., Lord, W. D., Morrison, R., Nash, G. V., Rylett, D., Scarlett, P. M., Swain, O. D., Szczykulska, M., Thornton, J. L., Trill, E. J., Warwick, A. C., and Winterbourn, B.: Daily and sub-daily hydrometeorological and soil data (2013-2019) [COSMOS-UK], NERC Environmental Information Data Centre [dataset], 2021.
- Stillman, S., Zeng, X., and Bosilovich, M. G.: Evaluation of 22 precipitation and 23 soil moisture products over a semiarid area in southeastern Arizona, *J. Hydrometeorol.*, 17, 211-230, <https://doi.org/10.1175/JHM-D-15-0007.1>, 2016.
- Tavakol, A., Rahmani, V., Quiring, S. M., and Kumar, S. V.: Evaluation analysis of NASA SMAP L3 and L4 and SPoRT-LIS soil moisture data in the United States, *Remote Sens. Environ.*, 229, 234-246, 2019.

- Upadhyaya, D. B., Evans, J., Muddu, S., Tomer, S. K., Al Bitar, A., Yeggina, S., Morrison, R., Fry, M., Tripathi, S. N., and Mujumdar, M.: The Indian COSMOS Network (ICON): validating L-band remote sensing and modelled soil moisture data products, *Remote Sens.*, 13, 537, <https://doi.org/10.3390/rs13030537>, 2021.
- 890 Wang, A. and Zeng, X.: Evaluation of multireanalysis products with in situ observations over the Tibetan Plateau, *J. Geophys. Res.-Atmos.*, 117, <https://doi.org/10.1029/2011JD016553>, 2012.
- Wen, X., Lu, H., Li, C., Koike, T., and Kaihotsu, I.: Inter-comparison of soil moisture products from SMOS, AMSR-E, ECWMF and GLDAS over the Mongolia Plateau, *Land Surface Remote Sensing II*, 103-110, <https://doi.org/10.1117/12.2068952>, 2014.
- Willmott, C. J. and Matsuura, M.: A more rational climatic moisture index, *The Professional Geographer*, 44, 84-88, 1992.
- 895 Wood, E. F., Roundy, J. K., Troy, T. J., Van Beek, L., Bierkens, M. F., Blyth, E., de Roo, A., Döll, P., Ek, M., and Famiglietti, J.: Hyperresolution global land surface modeling: Meeting a grand challenge for monitoring Earth's terrestrial water, *Water Resour. Res.*, 47, <https://doi.org/10.1029/2010WR010090>, 2011.
- Wu, Z., Feng, H., He, H., Zhou, J., and Zhang, Y.: Evaluation of soil moisture climatology and anomaly components derived from ERA5-land and GLDAS-2.1 in China, *Water Resour. Manag.*, 35, 629-643, <https://doi.org/10.1007/s11269-020-02743-w>, 2021.
- 900 Xing, Z., Fan, L., Zhao, L., De Lannoy, G., Frappart, F., Peng, J., Li, X., Zeng, J., Al-Yaari, A., and Yang, K.: A first assessment of satellite and reanalysis estimates of surface and root-zone soil moisture over the permafrost region of Qinghai-Tibet Plateau, *Remote Sens. Environ.*, 265, 112666, <https://doi.org/10.1016/j.rse.2021.112666>, 2021.
- Xu, L., Chen, N., Zhang, X., Moradkhani, H., Zhang, C., and Hu, C.: In-situ and triple-collocation based evaluations of eight global root zone soil moisture products, *Remote Sens. Environ.*, 254, 112248, <https://doi.org/10.1016/j.rse.2020.112248>, 2021.
- 905 Yamazaki, D., Ikeshima, D., Tawatari, R., Yamaguchi, T., O'Loughlin, F., Neal, J. C., Sampson, C. C., Kanae, S., and Bates, P. D.: A high - accuracy map of global terrain elevations, *Geophys. Res. Lett.*, 44, 5844-5853, <https://doi.org/10.1002/2017GL072874>, 2017.
- Yang, S., Li, R., Wu, T., Hu, G., Xiao, Y., Du, Y., Zhu, X., Ni, J., Ma, J., and Zhang, Y.: Evaluation of reanalysis soil temperature and soil moisture products in permafrost regions on the Qinghai-Tibetan Plateau, *Geoderma*, 377, 114583, <https://doi.org/10.1016/j.geoderma.2020.114583>, 2020.
- 910 Yang, Y., Zhang, J., Bao, Z., Ao, T., Wang, G., Wu, H., and Wang, J.: Evaluation of multi-source soil moisture datasets over central and eastern agricultural area of China using in situ monitoring network, *Remote Sens.*, 13, 1175, <https://doi.org/10.3390/rs13061175>, 2021.
- 915 Yee, M. S., Walker, J. P., Rüdiger, C., Parinussa, R. M., Koike, T., and Kerr, Y. H.: A comparison of SMOS and AMSR2 soil moisture using representative sites of the OzNet monitoring network, *Remote Sens. Environ.*, 195, 297-312, <https://doi.org/10.1016/j.rse.2017.04.019>, 2017.
- Zacharias, S., Bogena, H., Samaniego, L., Mauder, M., Fuß, R., Pütz, T., Frenzel, M., Schwank, M., Baessler, C., and Butterbach-Bahl, K.: A network of terrestrial environmental observatories in Germany, *Vadose Zone J.*, 10, 955-973, <https://doi.org/10.2136/vzj2010.0139>, 2011.
- 920 Zeng, J., Li, Z., Chen, Q., Bi, H., Qiu, J., and Zou, P.: Evaluation of remotely sensed and reanalysis soil moisture products over the Tibetan Plateau using in-situ observations, *Remote Sens. Environ.*, 163, 91-110, <https://doi.org/10.1016/j.rse.2015.03.008>, 2015.
- Zheng, J., Zhao, T., Lü, H., Shi, J., Cosh, M. H., Ji, D., Jiang, L., Cui, Q., Lu, H., and Yang, K.: Assessment of 24 soil moisture datasets using a new in situ network in the Shandian River Basin of China, *Remote Sens. Environ.*, 271, 112891, <https://doi.org/10.1016/j.rse.2022.112891>, 2022.
- 925 Zheng, Y., Coxon, G., Woods, R., Li, J., and Feng, P.: A framework for estimating the probability distribution of event runoff coefficient in ungauged catchments, *Water Resour. Res.*, 59, e2022WR033227, 2023.
- Zheng, Y., Li, J., Zhang, T., Rong, Y., and Feng, P.: Exploring the Application of Flood Scaling Property in Hydrological Model Calibration, *J. Hydrometeorol.*, 22, 3255-3274, <https://doi.org/10.1175/JHM-D-21-0123.1>, 2021.
- 930 Zheng, Y., Li, J., Dong, L., Rong, Y., Kang, A., and Feng, P.: Estimation of Initial Abstraction for Hydrological Modeling Based on Global Land Data Assimilation System-Simulated Datasets, *J. Hydrometeorol.*, 21, 1051-1072, <https://doi.org/10.1175/JHM-D-19-0202.1>, 2020.
- Zhu, D., Ciais, P., Krinner, G., Maignan, F., Jornet Puig, A., and Hugelius, G.: Controls of soil organic matter on soil thermal dynamics in the northern high latitudes, *Nat. Commun.*, 10, 1-9, <https://doi.org/10.1038/s41467-019-11103-1>, 2019.
- 935 Zhu, Z., Tan, L., Gao, S., and Jiao, Q.: Observation on soil moisture of irrigation cropland by cosmic-ray probe, *IEEE Geoscience and Remote Sensing Letters*, 12, 472-476, 10.1109/LGRS.2014.2346784, 2014.
- Zreda, M., Desilets, D., Ferré, T., and Scott, R. L.: Measuring soil moisture content non-invasively at intermediate spatial scale using cosmic-ray neutrons, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL035655>, 2008.
- 940 Zreda, M., Shuttleworth, W., Zeng, X., Zweck, C., Desilets, D., Franz, T., and Rosolem, R.: COSMOS: The cosmic-ray soil moisture observing system, *Hydrol. Earth Syst. Sci.*, 16, 4079-4099, <https://doi.org/10.5194/hess-16-4079-2012>, 2012.