

# Spatial distribution and controls of snowmelt runoff in a sublimation-dominated environment in the semiarid Andes of Chile

Álvaro Ayala<sup>1</sup>, Simone Schauwecker<sup>1</sup>, Shelley MacDonell<sup>1, 2</sup>

<sup>1</sup> Centro de Estudios Avanzados en Zonas Áridas (CEAZA), La Serena, 1700000, Chile

5 <sup>2</sup> Waterways Centre for Freshwater Management, Lincoln University and the University of Canterbury, Christchurch, New Zealand

*Correspondence to:* Álvaro Ayala (alvaro.ayala@ceaza.cl)

**Abstract.** Sublimation is the main ablation component of snow ~~and ice~~ in the upper areas of the semiarid Andes (~26-32°S and ~69-71°W). This region reaches up to 6000 m, is characterized by scarce precipitation, high solar radiation receipt, and low air humidity, and has been affected since 2010 by a severe drought. In this study, we suggest that most of the snowmelt runoff originates from specific areas with topographic and meteorological features that permit large snow accumulation, limited removal by sublimation and sufficient energy for snowmelt. To test this hypothesis, We we quantify the spatial distribution of snowmelt runoff and sublimation in a catchment of the semiarid Andes using a process-based snow model that is forced with field data and validated with field data, satellite derived indices of snow presence and an independent SWE reconstruction product. Results from mModel simulations over a two-year period reproduce point-scale records of snow depth and SWE and are also in good agreement with an independent SWE reconstruction product as well as satellite distributed patterns obtained from satellite derived products, such as snow cover area and indices of snow winter absence and summer persistence. We estimate that 50% of snowmelt runoff is produced by 21-29+8-28% of the catchment area, which we define as “snowmelt hotspots”. Snowmelt hotspots are located at mid-to-lower elevations of the catchment on wind-sheltered, low-angle slopes. Snowmelt hotspots are located at elevations between 4200 and 4800 m, have easterly aspects, low slope angles, and high snow accumulation and persistence. Our findings show that sublimation is not only the main ablation component, but it also plays an important role shaping the spatial variability of total annual snowmelt. Snowmelt hotspots might be connected. We suggest that snowmelt hotspots play a key hydrological role when connecting with other hydrological features of arid and semiarid dry mountain regions, such as areas of groundwater recharge, rock glaciers and mountain peatlands. We, and recommend a more detailed snow and hydrological monitoring of these sites, especially in the current and projected scenarios of scarce precipitation.

## 1 Introduction

30 Snowmelt is typically the largest runoff contributor in high-elevation mountain regions and their adjacent areas (Mankin et al., 2015; Kraaijenbrink et al., 2021). The annual volume of snowmelt runoff generated from each catchment is the result of the interactions between the topography, the land cover and the physical processes that control snowfall, snow transport, the surface energy balance and the internal changes of the snowpack (Lehning et al., 2008; Mott et al., 2018; Pomeroy et al., 1998). A good understanding of these processes and how to model them is key to quantify water supply availability for several 35 populated regions around the world (Freudiger et al., 2017; Hock et al., 2017). A particularly critical case of snowmelt dependency is that of arid and semiarid high-elevation mountain ranges, such as the semiarid Andes, Central Asia, and Southwestern USA (Huning and AghaKouchak, 2020). The climate of these mountain ranges is characterized by low temperatures and little precipitation (~~hereafter we refer to these ranges as cold and dry mountain regions~~), and pose characteristic challenges to estimate snowmelt runoff, such as episodic precipitation events (Schauwecker et al., 2022), shallow 40 snowpacks (Zhang and Ishikawa, 2008) and high sublimation rates (Stigter et al., 2018).

Sublimation from ~~the~~ snow cover is the direct transition of water from solid to vapor state and it occurs as surface sublimation (e.g. Hood et al., 1999), or as blowing (or drifting) snow sublimation (e.g. Groot Zwaaftink et al., 2013). Both surface and blowing snow sublimation ~~reduces~~ the mass of the snow cover and can significantly affect the water balance of various regions around the world, such as the Canadian Prairies (Pomeroy and Li, 2000), Antarctica (Palm et al., 2017), and the semiarid 45 Andes (Réveillet et al., 2020; Gascoïn et al., 2013). Additionally, turbulent latent heat fluxes associated with the solid to vapor transition use energy from the snowpack, lowering its temperature and decreasing the energy available for melting~~lowers the temperature of the snow surface, thus decreasing the energy available for melting~~. This process can have impacts on glacier mass balance (Ayala et al., 2017a; Stigter et al., 2018) and the performance of temperature-index melt models (Ayala et al., 2017b; Litt et al., 2019). Blowing snow sublimation depends on the amount of transported snow and sublimation rates and is 50 usually important in the mass balance of open and windswept environments (Pomeroy and Li, 2000), and wind-exposed mountain ridges (Strasser et al., 2008).

The interplay of physical processes controlling the evolution of the seasonal snowpack results in a large spatial variability of snowmelt and surface sublimation~~Both snowmelt and surface sublimation are determined by the surface energy balance of the snow cover. While the spatial distribution of snow accumulation is controlled by preferential deposition, wind redistribution and gravitational transport~~ (Mott et al., 2010; Freudiger et al., 2017), the energy balance is controlled by~~The spatial variability of the energy balance can be very large and is controlled by snow accumulation,~~ solar radiation and snow-atmosphere interactions that evolve during the ablation season (Mott et al., 2018; Pomeroy et al., 2003) and generate large differences in snowmelt rates across a certain domain (Pohl et al., 2006; DeBeer and Pomeroy, 2017). By the end of the ablation season, the snow cover often reaches a patchy state in which the advection of sensible and latent heat from snow-free areas plays a key 60 role for snow and ice ablation (Mott et al., 2020; van der Valk et al., 2022). Solving the energy balance of ~~the~~ snow cover over complex and steep terrain is a very complex task, as it depends on the availability of distributed meteorological forcing data

and adequate parameterizations of the physical processes controlling radiation, turbulent fluxes, snow microstructure and snow metamorphism (Vionnet et al., 2012; Lehning et al., 2002). Despite this complexity, in hydrological models the temporal and spatial variability of the energy balance is usually averaged over elevation bands or other hydrological units, but with negative effects on the representation of snow ablation and runoff at the basin scale (Dornes et al., 2008; Luce et al., 1998). Moreover, In cold and dry mountain regions in the case of dry mountain environments, turbulent latent heat fluxes and sublimation can represent a large part of the snow cover energy and mass balance (Jackson and Prowse, 2009; Zhang and Ishikawa, 2008), suggesting a particularly large spatial variability of snowmelt and runoff generation. However, snow sublimation is frequently neglected in hydrological models (e.g. Ragettli et al., 2014; Seibert et al., 2017).

70 The semiarid Andes is a good example of a snowmelt-dependent region where snow sublimation can significantly reduce the mass of the winter snow cover and complicate the calculation of snowmelt runoff. Several studies have estimated snow mass and energy balances (Gascoïn et al., 2013; Réveillet et al., 2020; Voordendag et al., 2021), and have provided useful insights into processes occurring in this region, as well as into the challenges of estimating distributed snowmelt and sublimation in this environment. However, these studies have largely focused on either model or input uncertainties (Réveillet et al., 2020; Voordendag et al., 2021), or snow redistribution (Gascoïn et al., 2013), rather than melt processes and hydrological implications importance. From another a geostatistical perspective, Mendoza et al. (2020) analyzed the spatial properties of a set of Lidar snow depth measurements across several catchments of central Chile and found at the strong relation between snow depth and local topographic and land cover properties in semiarid catchments of central Chile. Given that snow represents 85% of streamflow variability in semiarid catchments (Masiokas et al., 2006) and is a useful predictor of streamflow (Sproles et al., 2016), it is of vital importance to connect distributed snow mass and energy balance processes to runoff generation hydrological impacts in their full complexity and spatial variability.

In this work, we hypothesize that the meteorological and topographical conditions of the semiarid Andes result in large areas where snow surface sublimation losses dominate over snowmelt, thus delimiting relatively small areas from where most of the catchment snowmelt runoff is generated. We argue that the typically large spatial variability of snowmelt in mountain terrain is further enlarged in dry environments by large sublimation rates that almost fully deplete the snow mass available for melt at wind-exposed sites. We refer to the areas producing most of the snowmelt runoff these sites as “snowmelt hotspots” and we define them as those forming the minimum area in a snow-covered catchment from where at least 50% of the snowmelt runoff is generated. To test this hypothesis, we calculate spatially distributed amounts of snowmelt and snow sublimation using a process-based snow evolution model in a 79 km<sup>2</sup> catchment of the semiarid Andes of Chile over a two-year period (April 2019 to March 2021). The model is forced with in-situ meteorological data, and it is verified against point observations of snow variables and a set of remote sensing independent satellite-derived products. Our main objectives are to i) quantify the snow mass balance components, ii) determine the spatial distribution of snowmelt runoff and proof the existence of snowmelt hotspots determine the existence and location of snowmelt hotspots, and iii) identify the main characteristics of the snowmelt hotspots and discuss their possible connection with other hydrological components. As our selected study catchment contains

95 glacierized areas, we also include estimates of ice melt and glacier runoff. We expect that our results will provide new insights

for studies focusing on snowmelt runoff generation in ~~cold and~~ dry mountain ~~regions environments~~ and its connection with other components and processes, such as glaciers, permafrost, groundwater recharge and the spatial distribution of vegetation.

## 2 Study area

100 The semiarid Andes of Chile extend from approximately 26°S to 32°S (~69-71°W) reaching elevations ~~of up to~~ more than 6000 m a.s.l. Its climate is defined as cold semi-arid with dry summers in the low-lying areas up to approximately 3000 m a.s.l. and as tundra climate in the upper areas (Sarricolea et al., 2017). While winters (June-August) are cold with occasional precipitation events, summers (December-February) are hot and dry with low cloudiness and intense solar radiation (Favier et al., 2009). Precipitation has a large interannual variability mostly associated with El Niño Southern Oscillation (ENSO) (Arias et al., 2021; Montecinos and Aceituno, 2003). Annual precipitation in the lowlands increases from ca. 100 to 300 mm a<sup>-1</sup> towards  
105 south and is ~~2-33-5~~ times higher ~~at 3000 m a.s.l. near the peak of the Andes~~ (Scaff et al., 2017; Favier et al., 2009), ~~but it usually does not increase significantly above 3000 m.a.s.l.~~ Cloud cover and air moisture are usually very low, which determines very high values of solar radiation and potential evapotranspiration (MacDonell et al., 2013). ~~Snowmelt is the main runoff contributor in this region, even though~~ Although mass losses produced by snow sublimation can reach up to 50 or 80% of the annual snowfall (Réveillet et al., 2020), the hydrological regime of the rivers in this region is mostly nival (Favier et al., 2009).  
110 Glacier runoff is ~~also~~ an important contributor during droughts and at the end of summer (Gascoin et al., 2011; Ragetti et al., 2014).




Figure 1: (a): Location of the Corrales catchment in Chile, (b) Map of the Corrales catchment including Tapado Glacier and the Tapado (TAP), Tapado Glacier (TGL) and Paso Agua Negra (PAN) meteorological stations, (c) Hypsometry of the basin, (d) Slope aspect frequency distribution in the catchment, (e) Winter snow accumulation close to La Laguna reservoir (~3200 m a.s.l.) ([Photo: La Laguna Reservoir personnel, 17/06/2020](#)), (f) Tapado AWS with Tapado Glacier in the background ([Photo: Cristian Orrego, 22/11/2010](#)).

115

This study focuses on the Corrales catchment, a 79 km<sup>2</sup> catchment (30.16°S, 69.88°W) located in the upper areas of the semiarid Andes of Chile, approximately 180 km east of La Serena city (Figure 1a-b). The elevation of the catchment ranges from 3800 to 5600 m a.s.l., with ca. 50% of the catchment located above 4700 m a.s.l. (Figure 1c). The orientation of the slopes is relatively uniform across the catchment (Figure 1d). Meteorological records from the nearby La Laguna meteorological station (30.20°S, 70.04°W, 3160 m a.s.l., maintained by the Chilean Water Directorate (DGA)) show an annual mean precipitation of 160 mm a<sup>-1</sup> (with a coefficient of variation of 0.68) and an average temperature of 8.1°C in the period 1965-

120

125 2020. Snow varies spatially across the catchment (Figure 1e), and resultant runoff leaving the Corrales catchment is stored in  
La Laguna Reservoir (storage capacity of 40 Mm<sup>3</sup>) from where it is regulated for irrigation and other uses. Above 3000 m a.s.l.  
vegetation is almost completely absent except for peatlands next to streams (Valois et al., 2020) and shrubs on the mountain  
slopes (Kalthoff et al., 2006). The Corrales catchment contains Tapado Glacier, which is the largest glacier on the Chilean side  
of this region (1.67 km<sup>2</sup> in 2019) (DGA, 2022). Tapado Glacier has lost  $25.2 \pm 4.6\%$  of its area since 1956, with an accelerated  
130 negative mass balance since 2000 (Robson et al., 2022), which is in line with other glaciers in the area (Pitte et al., 2022). In  
addition, the Corrales catchment contains a few small debris-free glaciers located above 5000 m a.s.l. (Figure 1b), and several  
rock glaciers (not shown) that might play a relevant hydrological role by active layer thawing and by storing and releasing  
meltwater from snow and ice(not shown) (Schaffer et al., 2019; Navarro et al., 2023a). Since 2009, the Corrales catchment has  
been instrumented with hydro meteorological equipment and several glaciological field campaigns have been carried out,  
135 focusing on the collection of groundwater, glaciological and hydrological data (e.g., Figure 1f).

### 3 Data

#### 3.1 Field data

We use meteorological data from a network of stations in the Corrales basin to force a snow evolution model that is presented  
in Section 4. The meteorological network consists of three Automatic Weather Stations (AWSs): Tapado (TAP), Tapado  
140 Glacier (TGL) and Paso Agua Negra (PAN) (Figure 1 and Table 1). The AWSs are maintained by the Centro de Estudios  
Avanzados en Zonas Áridas (CEAZA), based in La Serena, Chile, and are visited several times each year between October  
and May when the sites is-are accessible by road. TAP and PAN consist of meteorological towers (Figure 1f, S1a) and transmit  
near-real time data via satellite communication. TGL is a HOBO U30 station installed on a tripod next to the lower debris-free  
section of Tapado Glacier that is visited with a lower frequency (2 to 3 times per year) due to logistical limitations (Figure  
145 S1b). Data from TGL is downloaded manually.

Hourly precipitation is derived from the cumulative precipitation data recorded at TAP (Table 1). As the cumulative  
precipitation record contains noise, we apply a simple method that discriminates actual precipitation from sensor noise (see  
Section S1). The method is based on the identification of positive changes in the daily precipitation cumulative record that  
lead to increases in the 5-day moving average of the same series. Additionally, based on a comparison with the SWE sensor  
150 and results from previous studies at TAP (Réveillet et al., 2020; Voordendag et al., 2021) and other regions (MacDonald and  
Pomeroy, 2007), we increase precipitation amounts by 30% due to snow undercatch by the sensor and precipitation  
underestimation due to differences between the exact location of the precipitation sensor and the meteorological tower where  
snow depth and SWE are recorded. Following these corrections, we obtain total precipitation amounts of 309 mm and 330 mm  
in the hydrological years 2019-2021 and 2020-2021 (from April to March), respectively (Figure 2a). Although annual  
155 precipitation was higher in 2020-2021, precipitation in the main snow season (April-September) was higher in 2019 (238 mm  
compared with 196 mm). Annual precipitation at TAP was 4 to 5 times higher than at La Laguna DGA, where a total of 63

~~and 82 mm were registered in 2019-2020 and 2020-2021, respectively.~~ Although we do not have a long-term precipitation at this particular site, we verified that, according to La Laguna DGA precipitation records (Información Oficial Hidrometeorológica y de Calidad de Aguas en Línea, 2023), our study years (2019-2020 and 2020-2021) corresponded to a dry period, with an annual exceedance probability of ca. 80% for both years.

Figure 2 shows a summary of the data collected by the three AWSs during the study period and Table 1 includes ~~more~~ additional details, such as the sensors and their installation heights. Apart from occasional gaps at each station, the largest period of missing data is found at TAP between December 2020 and January 2021. Daily mean air temperature at the elevation of TAP remained above 0°C for several months (November to April), with a maximum near 10°C in January 2020 (Figure 2a), although there was a large data gap in summer 2021. Since the correlation between the three air temperature records is very high (~~correlation coefficients are higher than 0.97~~ $R^2 \rightarrow 0.95$ ), missing data was filled by establishing linear relationships between the three stations. Figure 2b shows the variability of snow depth, SWE and surface albedo at TAP. Snow depth reached up to 1 m in ~~the~~-winter 2019, but only 0.8 m in ~~the~~-winter 2020. In general, snow depth decreases rapidly during the days following each snowfall event. SWE and surface albedo closely followed snow depth variations. We found that the decay of surface albedo was faster in spring 2019 than in 2020. Mean daily solar radiation reached values ~~over up to more than~~ 400 W m<sup>-2</sup> with frequent drops associated with cloudiness, especially ~~during~~in the winter period (Figure 2c). The seasonal variability of incoming longwave radiation was much lower than that of solar radiation and varies between 180 and 300 W m<sup>-2</sup> (Figure 2c). Daily mean relative humidity at TAP remained ~~mostly~~ below 50% for most of the time, increasing above this value only a few ~~times~~days during the study period (Figure 2d). Relative humidity at the other stations was similar (not shown). Air pressure varied between 550 and 580 hPa with large and relatively constant values in summer and low and oscillating values in winter (Figure 2d). Wind speed at PAN was 2.6 times larger than at TAP and TGL, with a large predominance of westerly~~n~~ winds, indicating that the latter are located at wind-sheltered locations (Figure 2e and 2f).

In addition to the meteorological data, we use the inflow to the La Laguna reservoir as a reference for streamflow variations in response to snow and ice melt. These data were derived from the water balance of the reservoir and were provided by the organization in charge of the operation of the reservoir (Junta de Vigilancia del río Elqui y sus afluentes). Finally, we use a dataset of surface elevation changes of Tapado Glacier to validate ice melt estimates produced by the snow evolution model (Section 4). These observations correspond to ablation stake readings collected in the summer periods 2019-2020 and 2020-2021, and snow density measurements at the end of the corresponding spring periods (Figure S2 and Table S1). ~~The stakes were manually drilled on the ablation zone of the glacier and were visited approximately every month during the ablation season (Figure S2 and Table S1). The visit and maintenance of the ablation stakes was difficult due to the development of snow and ice penitentes of 3-4 m height (Figure S3), which also led to the collapse of some monitored sites. Penitentes are cone-shaped snow and ice structures formed in dry environments by differential ablation rates along their height (Lliboutry, 1954). The ablation stakes network consists of traditional single point measurements and one 5-m ablation transect where we registered surface lowering every 20 cm to account for the differential ablation of the surface associated with the penitentes following Nicholson et al. (2016).~~


**Table 1: AWSs in the Corrales catchment**

Station	Name	East <u>(WGS84</u> <u>UTM</u> <u>19S)</u>	South <u>(WGS84</u> <u>UTM</u> <u>19S)</u>	Elevation <u>(m a.s.l.)</u>	Monitored variables (*)	Instrument	Sensor height (m)
TAP	Tapado	412546	6663325	4306	T <sub>a</sub> -RH	Vaisala HMP45C	4.0
					WS <sub>a</sub> -WD	RM Young 5103 Wind Monitor	5.0
					P	Geonor T-200B 1000 mm	1.5
					Sin <sub>a</sub> -Sout <sub>a</sub> -Lin	Kipp and Zonen CNR4	3.5
					SD	Luft SHM31	3.5
					SWE	Campbell CS725	3.0
TGL	Tapado Glacier	411121	6664158	4727	T <sub>a</sub> -RH	HOBO S-THB-M002	2.0
					WS	HOBO S-WSA-M003	2.0
					WD	HOBO S-WDA-M003	2.0
					Sin <sub>a</sub> -Sout	HOBO S-LIB M003	2.0
PAN	Paso Agua Negra	420534	6659795	4774	T <sub>a</sub> -RH	Vaisala HMP155	3.0
					WS <sub>a</sub> -WD	Campbell CSAT3	5.0
					Pa	Vaisala PTB110	2.0

(\*): T: Air temperature, RH: Relative humidity, WS: Wind speed, WD: Wind direction, P: Precipitation, Sin: Incoming solar radiation, Sout: Reflected solar radiation, Lin: Incoming longwave radiation, SD: Snow depth, SWE: Snow water equivalent,

195 Pa: Air pressure.





**Figure 2: Meteorological time series recorded at Tapado (TAP), Tapado Glacier (TGL) and Paso Agua Negra (PAN) during the study period. If not indicated, the meteorological variables were recorded at TAP. (a) Precipitation (left axis) and air temperature (right axis), (b) Snow depth, SWE, and albedo, (c) Incoming shortwave and longwave radiation, (d) Relative humidity and air pressure, (e) Wind speed, (f) Wind roses.**

## 3.2 Snow products

We use ~~two~~-three spatially distributed datasets that describe the interannual and seasonal variations of the snow cover to evaluate the results of the snow evolution model (Section 4). The first dataset corresponds to satellite-based maps of snow absence and persistence indices (Wayand et al., 2018) in the period 2019-2021, the second to SWE daily maps that were reconstructed using a data assimilation scheme for the subtropical Andes in the period 1985-2015 (Cortés and Margulis, 2017). Finally, we use Sentinel-2 images to derive snow cover area (SCA).

### 3.2.1 Maps of snow absence (SA) and snow persistence (SP) indices

We use the snow indices defined by Wayand et al. (2018) to characterize the spatial patterns of snow cover in the study area. The snow absence (SA) index is defined as the fraction of time in which snow is absent during the accumulation period, whereas the snow persistence (SP) index is defined as the fraction of time in which snow is present during the melt period. In their study, Wayand et al. (2018) developed a method to estimate SA and SP from optical satellite images obtained by Sentinel-2 and Landsat-8 missions. The method was implemented in the Google Earth Engine platform, which is freely available. In our study, we adapt the definition of the snow accumulation and melt periods to the Southern Hemisphere (accumulation: April-September and melt: October-March) and apply the method to the Corrales catchment. For simplicity we refer to the indices as snow winter absence and snow summer persistence. A summary of the utilized images is given in Table 2 (see Table S2 for the full list of images).

**Table 2: Summary of satellite images used for the calculation of the snow indices (SA and SP)**

Snow index	Months	Product	Period	Number of images
Snow <u>winter</u> absence	April to September	Landsat-8	13-04-2019 to 22-09-2020	18
		Sentinel-2	03-06-2019 to 10-09-2020	15
Snow <u>summer</u> persistence	October to March	Landsat-8	06-10-2019 to 17-03-2021	21
		Sentinel-2	15-11-2019 to 14-03-2021	23

### 3.2.2 Maps of reconstructed SWE

We analyze the spatial variability of snow accumulation in the study area using the SWE reconstruction developed by Cortés and Margulis (2017) (hereafter C&M2017). This dataset consists of daily maps of SWE with a spatial resolution of 180 m over the extratropical Andes (27-37°S) in the period 1985-2015. The maps were calculated using a data assimilation scheme that combines results from a Land Surface Model and Landsat optical images. The data assimilation scheme explicitly addresses the uncertainty of input variables by weighting simulation ensembles to fit observed fractional snow cover area (fSCA) from the satellite images. Reconstructed SWE maps compare well with manual records along the Andes (Cortés and Margulis, 2017; Cortés et al., 2016) and melt season runoff records (Álvarez-Garretón et al., 2018). Later results have shown that these SWE

maps are suitable for other analyses, such as the impact of atmospheric rivers on snow accumulation (Saavedra et al., 2020) and the validation of hydrological models (Ayala et al., 2020). In our study, we summarize the information provided by the SWE reconstruction product by calculating the interannual median and coefficient of variation of SWE at the time of annual maximum accumulation in the catchment (hereafter SWE<sub>max</sub> and SWE<sub>max</sub>CV). The equation for the coefficient of variation at each grid cell is:

$$CV = \frac{\mu}{\sigma}, \tag{1}$$

where  $\mu$  is the mean and  $\sigma$  is the standard deviation of peak annual SWE during the period 1985-2015.

### **3.3.3 Sentinel-2 NDSI**

~~Additionally, we~~ We use the Normalized Difference Snow Index (NDSI) derived by the Sentinel-2 Level 2A processing (Level-2A Algorithm Overview, 2023) to calculate SCA over the study catchment and compare it with our model simulations (see Section 4). We manually selected a total of 103 cloud-free NDSI maps in the period April 2019-March 2021 using the EO Browser (<https://apps.sentinel-hub.com/eo-browser/>).

### **3.3 Other data**

~~Catchment topography is extracted from a 30 m resolution Digital Elevation Model (DEM) produced by NASADEM (NASA JPL, 2020). Additionally, we use the Normalized Difference Snow Index (NDSI) derived by the Sentinel 2 Level 2A processing (Level 2A Algorithm Overview, 2023) to calculate SCA over the study catchment and compare it with our model simulations (see section 4). We manually selected a total of 103 cloud free NDSI maps in the period April 2019 March 2021 using the EO Browser (<https://apps.sentinel-hub.com/eo-browser/>).~~

## **4 SnowModel**

### **4.1 Model description**

We use SnowModel to calculate snowmelt, sublimation, and other snow-related variables in the Corrales catchment. SnowModel is a numerical model for the spatially-distributed simulation of snow evolution with an explicit consideration of the main physical processes that shape the seasonal snowpack (Liston and Elder, 2006a). The model consists of four modules: a module that interpolates meteorological variables recorded at specific locations to a two-dimensional grid using specially developed algorithms (MicroMet) (Liston and Elder, 2006b); a module that solves the energy balance and yields surface sublimation and snowmelt (EnBal) (Liston, 1995); a module that solves the internal changes of the snowpack, such as refreezing, densification and metamorphism (SnowPack) (Liston and Hall, 1995); and a module that solves the snow transport and blowing snow sublimation due to saltation and suspension of the snow (SnowTran-3D) (Liston et al., 1998). The model does not include a representation of snow gravitational transport (avalanches). SnowModel has been successfully tested in

several snow environments around the world (Liston et al., 2007; Mernild et al., 2016), including the semiarid Andes (Gascoin et al., 2013; Réveillet et al., 2020; Voordendag et al., 2021). Readers are referred to other studies for a full description of the model (Liston and Elder, 2006a) and its updates (Mernild et al., 2018; Merkouriadi et al., 2021).

260 In SnowModel, snowmelt either refreezes ~~in~~ lower layers or drains as runoff when the snowpack has already reached a ripe state. In our study, we analyze results from snowmelt runoff (snowmelt leaving the snowpack) and snow surface sublimation (extracted from the surface turbulent latent heat flux). Rain on snow contributes to a snow density increase until reaching a maximum density of 550 kg m<sup>-3</sup>. Beyond that limit, rain on snow is added to snowmelt runoff, but we do not consider it in the snowmelt runoff variable. We define the sublimation ratio-fraction (SublRatioSublFrac) as shown in Eq. (24):

$$265 \text{ SublFrac} = \frac{\text{Snow surface sublimation}}{\text{Snow surface sublimation} + \text{Snowmelt runoff}} \quad (24)$$

Glaciers can be included in SnowModel as a type of surface and ice is melted once the snow has completely disappeared from the surface. Debris-covered and rock glaciers are not included in the model.

## 4.2 Setup

270 We run SnowModel in the Corrales basin using a 3 h timestep in the period April 2019-March 2021. The domain of the model runs is a rectangle that contains the Corrales basin with an additional buffer of 500 m in all four main directions (N-S-E-W). We use a DEM of a spatial resolution of 50 m that is bilinearly resampled from a ~30 m resolution Digital Elevation Model (DEM) produced by NASADEM (NASA JPL, 2020)–NASADEM (see Section 3.3). The land cover of the model domain consists entirely of bare soil except for the debris-free glaciers shown in Figure 1b. The DEMs and related variables are analyzed using Topotoolbox functions in MATLAB (Schwanghart and Scherler, 2014).

275 **Table 3: Parameters values used in the SnowModel simulations**

Module	Parameter	Value or source	Units
General	Spatial resolution	50	m
	Time step	3	h
	Number of grid cells (East, North)	270, 272	
Meteorological inputs	Precipitation	TAP	mm
	Air temperature	TAP, TGL, PAN	°C
	Relative humidity	TAP, TGL, PAN	%
	Wind speed and direction	TAP, TGL, PAN	m s <sup>-1</sup> and °
	Solar radiation	TAP	W m <sup>-2</sup>
	Incoming longwave radiation	TAP	W m <sup>-2</sup>
	Air pressure	PAN	Pa
MicroMet	<u>Precipitation factor</u>	<u>0.7, 1.0, 1.3</u>	
	Curvature length scale	500 <u>(default)</u>	m

	Slope weight for wind distribution	0.25, 0.58 (default), 0.75	
	Curvature weight for wind distribution	0.25, 0.42 (default), 0.75	
	Monthly mean air temperature lapse rates	From January to December: 8.1,8.1,7.7,6.8,5.5,4.7, 4.4,5.9,7.1,7.8,8.1,8.2 (default)	°C km <sup>-1</sup>
	Precipitation and wind speed lapse rate	0	
	<u>Rain-snow air temperature threshold</u>	2 (default)	°C
Enbal	Albedo decay melt conditions	0.024	s <sup>-1</sup>
	Albedo decay cold conditions	0.008	s <sup>-1</sup>
	<u>IceGlacier</u> albedo	0.3	
	Albedo fresh snow	0.9	
	Soil albedo	0.14	
	Precipitation threshold for albedo reset	0.006 (default)	m
	Aerodynamic surface roughness length for snow ( $z_0$ )	0.001 (default), 0.005, 0.010	m
SnowPack	<del>Number</del> <u>Maximum number of snow of</u> layers	6	
SnowTran-3D	Threshold surface shear velocity	0.25 (default)	ms <sup>-1</sup>

We run the model using the parameters shown in Table 3 ~~mostly default model parameters (Table 3)~~ and without performing any calibration except for the albedo decay rates, which were manually set to fit the albedo changes observed at TAP (Figure S4). The model is forced using the meteorological field data described in Section 3.1. The wind speed records were adjusted from the corresponding sensor height to a height of 2 m using a logarithmic wind profile and an aerodynamic surface roughness length for snow of 5 mm. Although there is an annual precipitation lapse rate from the lowlands of the Coquimbo Region up to La Laguna DGA station (3160 m a.s.l.), we used a value of zero because we do not have enough data to support a precipitation lapse rate above that elevation, particularly within the relatively small area of the Corrales catchment. In general, snow distribution at high-elevation catchments is governed mostly by wind transport (e.g. Lehning et al., 2011).

### 285 4.3 Ensemble runs

We produce ensemble runs ~~with the characteristics described in Table 3 and~~ considering three different values for three selected variables: a precipitation factor, the aerodynamic surface roughness length for snow (hereafter  $z_0$ ) and two wind distribution weights based on the slope and curvature of the terrain in the MicroMet module (Table 3). Previous snow simulations in this area have resulted to be most sensitive to these variables (Réveillet et al., 2020; Voordendag et al., 2021). We selected

290 precipitation factors that can be interpreted as an uncertainty range for snow undercatch. The aerodynamic surface roughness  
length (~~hereafter  $z_0$~~ ) is a key parameter controlling the mass and energy exchanges between the surface and the atmosphere  
and has been defined as the height above a surface at which the extrapolated horizontal wind-speed profile reaches zero (Brock  
et al., 2006). The selected values for  $z_0$  vary within typical ranges for snow and ice surfaces (Brock et al., 2006; Fitzpatrick et  
al., 2019). The slope and curvature distribution weights increase wind speed in the presence of windward and convex slopes  
295 and decrease it in the case of leeward and convex ones (Liston et al., 1998). According to the MicroMet module, the slope and  
curvature wind distribution weights should sum to one. The selected values for these variables were chosen to explore the  
sensitivity of snow ablation to these parameters in the semiarid Andes. ~~Previous snow simulations in this area have resulted to~~  
~~be most sensitive to these variables (Réveillet et al., 2020; Voordendag et al., 2021).~~ We test all the different combinations of  
these values, obtaining a total of 27 simulations that are used to assess model sensitivity and uncertainty. Table 4 provides a  
300 summary of the values and variables in the ensemble runs.

**Table 4: Summary of SnowModel ensemble runs**


Parameters and configuration	Values
Precipitation factor	0.7
	1.0
	1.3
Aerodynamic surface roughness length for snow ( $z_0$ ) (m)	0.001
	0.005
	0.010
Slope (curvature) weight	0.25 (0.75)
	0.58 (0.42)
	0.75 (0.25)
Other parameters	As in Table 3

## 5 Results

### 5.1 Analysis of snow products

305 The analysis of snow winter absence (SA), snow summer persistence (SP) and the interannual median of peak annual SWE  
(SWEmax) shows that the north-west section of the catchment present larger values of snow accumulation and persistence  
than the areas to the east. While SA and SWEmax values on the north-west of the catchment are lower than 0.6 (Figure 3a)  
and larger than 400 mm a<sup>-1</sup> (Figure 3g), respectively, the areas to the east present the opposite behavior (SA>0.8 and  
SWEmax<200 mm a<sup>-1</sup>). We further connect these patterns to the underlying topography by means of polar plots (Figure 3b-e-  
h). These plots show that while the lowest values of SA (SA<0.6) are found at sites higher than 4000 m a.s.l. with a SW, S or

310 SE aspect, the highest values ( $SA > 0.8$ ) are found at sites with a NW, N or NE aspect. Approximately 40% of the catchment  
present SA values lower than 0.6 (Figure 3c). The elevation band with the largest values of  $SWEmax$  is that between 4500 and  
5000 m a.s.l. with a S-SE aspect (Figure 3i), which correspond to the areas close to Tapado Glacier (around 10% of the  
catchment area, Figure 3g-i). These areas also have a very consistent interannual behavior (Figure S5). During the melt period,  
the catchment was mostly snow-free (Figure 3d), except for sites mostly above 5000 m a.s.l. with a SW-W aspect (Figure 3e),  
315 and only about 10% of the catchment area present values above 0.2 (Figure 3f). The maps of the SA and SP indices show the  
areas that are prone to remaining snow free during the accumulation period and the areas where snow persists during the melt  
period, respectively (Figure 3). While high SA values are found at ridges and sites located at the east of the catchment, low  
values are found at valley bottoms that are mostly located at the west of the catchment (Figure 3a). Polar plots in Figure 3b  
show the relationship between elevation, aspect and the SA index. In general, sites higher than 4000 m a.s.l. with a SW-S-SE  
320 orientation are snow covered during the accumulation period ( $SA < 0.5$ ). On the contrary, sites located with a N, NW and NE  
aspects have the highest values of SA, which suggests a strong role of solar radiation even in the period of the year with the  
lowest radiation values. Approximately 15% of the catchment is snow free during the accumulation period ( $SA$  close to one,  
Figure 3c). The map of SP shows that the Corrales catchment is mostly snow free during summer (Figure 3d), except for sites  
above 4500 m a.s.l. with S and SW aspects ( $SP > 0.2$ , Figure 3e). Approximately 45% of the catchment is almost always snow-  
325 free during the melt period ( $SP$  close to zero, Figure 3f). Glaciers appear as areas of minimum SA (Figure 3a) and maximum  
SP (Figure 3d).




330 **Figure 3: Snow winter absence (SA), snow summer persistence (SP) and the interannual median of peak annual SWE (SWEmax) and SP indices in the Corrales basin. While SA and SP are derived following Wavand et al. (2018), SWEmax is calculated from the Cortés and Margulis (2017) dataset, -calculated over the study period- using the methodology developed by Wayand et al. (2018).**  
 335 **(a) Map of SA, (b) Polar plot of SA, (c) Histogram of SA values, (d) Map of SP, (e) Polar plot of SP, (f) Histogram of SP values, (g) Map of SWEmax, (h) Polar plot of SWEmax, (i) Histogram of SWEmax. In the polar plots (b-e-h) the angles and the inverse radial distance represent the aspect and the elevation, respectively. While SA and SP refer to the percentage of time when snow is absence or present (a-b and d-e) in the accumulation (April-September) and melt (October-March) periods, respectively, the histogram represent the spatial distribution of the variables across the catchment. Glaciers outlines and contours are shown in (a), (d) and (g). Blank areas in (a) represent sites where the SA index cannot be calculated due to an insufficient number of cloud-free images in April-September.**

340 In Figure 4 we analyze the SWE reconstruction of Cortés and Margulis (2017) for our study catchment in the period 1985-2015. While Figures 4a-c present the interannual median of the annual maximum SWE, Figures 4d-f show the coefficient of variation of the same variable. The highest values of maximum SWE are found over westerly areas, close to Tapado Glacier, whereas maximum SWE values are very low in eastern sections (Figure 4a). Figure 4b shows that maximum SWE values are



highest at sites between 4500 and 5000 m a.s.l. with SW-S-SE aspects and between 5000 and 5500 m a.s.l. with a S aspect. This is an indication of sites with preferential snow deposition. The snowpack of the catchment is relatively shallow with only ca. 5% of the area having maximum SWE > 200 mm w.e. (Figure 4c). The coefficient of variation of maximum SWE reaches values up to 4.5, particularly at the east of the catchment, at sites > 4500 m a.s.l. with N aspects (Figures 4d-e). Almost 75% of the catchment area has a coefficient of variation > 1, which is in line with the high interannual variability of precipitation in this region. Glaciers are located in areas of high snow accumulation (Figure 4a) and low coefficients of variation (Figure 4d).




**Figure 4: Annual maximum snow accumulation in the Corrales basin during 1985-2015 according to the daily SWE reconstruction by Cortés and Margulis (2017). (a) Map of the interannual median, (b) Polar plot of the interannual median, (c) Histogram of the interannual median, (d) Map of the coefficient of variation, (e) Polar plot of the coefficient of variation, (f) Histogram of the coefficient of variation. Glaciers outlines are shown in (a) and (d).**

## 5.2 Verification of SnowModel results

We verify SnowModel results by comparing them against the available field data and the snow products analyzed in Section 5.1. In Figure 5, we compare results of SnowModel ensemble runs against measurements of sAt the point scale, snow depth and SWE records at TAP compare well against the ensemble runs for the corresponding grid cell (Figure 4). Model results are in good agreement with both independent records. The simulated changes of snow depth following precipitation events are similar to observations, particularly during periods of rapid depletion following each precipitation event. Rapid depletion periods are likely due to compaction, as well as mechanical removal and sublimation caused by frequent strong wind gusts and rapid drops in humidity occurring post precipitation. We discarded the possibility that these drops are caused by melt

because air temperature was mostly below zero~~very low~~ following events. We ~~also~~ observe differences in the modeled and observed peak snow depth and SWE that are likely caused by discrepancies between the precipitation amounts registered by the ~~Geonor-precipitation~~ sensor and the snow records. The snow disappearance dates ~~is-are~~ well approximated in 2019 (October) and 2020 (September and October)~~October 2019, September 2020 and October 2020~~ in the snow depth record (Figure 45a), but the SWE observations show a slower disappearance rate (Figure 4b).




**Figure 45: Time series of the observed ~~and simulated~~ snow variables at TAP and the ensemble runs for the corresponding grid cell.** (a) Snow depth, (b) SWE. The red lines and areas represent the median values and interquartile ranges 25 and 75 of the SnowModel ensemble runs, respectively. The blue lines correspond to the observations.


When we compare the SnowModel ensemble runs against the selected snow products (Sections 3.2 and 5.1) we find that our simulations correctly reproduce the main spatial patterns of the snow cover throughout the year, but tend to overestimate the winter snow cover area and smooth the spatial variability of snow during summer. A comparison of observed and simulated snow variables at the distributed scale is presented in Figure 6. Figure 6a shows SCA from the ensemble runs (red area and line) against SCA derived from Sentinel 2A (blue points). The SCAs from the ensemble runs (red areas and line) coincides well with the Sentinel-2A SCA values (blue points) at several times throughout the study period (obtaining scores of  $R^2=0.66$  and  $RMSE=14.8\%$ ), but the snow cover area is sometimes overestimated in winter and tends to disappear/disappear too early in the simulations (Figure 5a). Figures 5b-c show the comparison between SA and SP derived from satellite images and from SnowModel. The SnowModel values of SA and SP follow the same definitions of the satellite-derived indices but using the results of the simulations. With respect, In correspondence to the polar plots presented in Figure 3b-e-h~~Figures 3 and 4~~, each

point in Figures 56b-c-d represents the SA<sub>ave</sub> and SP<sub>ave</sub> and SWE<sub>max</sub> average value of all the grid cells contained in a 500-m elevation band (from 3500 to 6000 m, 5 values in total) and 45° aspect (from N to NW, 8 values in total), whereas the error bars represent SA<sub>ave</sub> and SP<sub>ave</sub> the standard deviation over the same grid cells. There is a relative good correspondence ~~good correlation~~ between observed and simulated values ( $R^2=0.47$  for SA and  $R^2=0.674$  for SP), ~~although the absolute numbers do not coincide~~ but we observe a bias in SA (BIAS = -0.26) and an offset of approximately 0.5 in SP. Both effects could be explained by snow-free sites across the catchment that are not well reproduced by the model. This is a consequence of comparing satellite derived indices from images that are available only every 2 to 3 weeks, and model derived indices that are calculated using every 3h time step. In Figure 6b, An extreme case of this situation are the two data points that do not align ~~to~~ in the general linear relationship of SA<sub>ave</sub> (lower right corner in Figure 5b), which correspond to sites located ~~to~~ above 5500 m a.s.l. with an E and SE aspect. Figure 56d shows the same plots but comparing ~~the interannual medians of simulated and reference SWE<sub>max</sub>~~ (Cortés and Margulis, 2017) ~~SWE at the time of annual maximum accumulation~~. While the reference SWE<sub>max</sub> ~~is interannual median of the reference values are~~ calculated from the 1985-2015 period, simulated SWE<sub>max</sub> ~~the interannual median of simulated values~~ is calculated using the two hydrological years of the modelling period (2019-2020 and 2020-2021). As the reference and simulated datasets correspond to different study periods, this comparison is evaluated mostly in relative terms. The  $R^2$  obtained by comparing ~~correlation between~~ both records is lower than that obtained for ~~of~~ the snow indices ( $R^2=0.2523$ ), likely due to difficulties of SnowModel to reproduce the shallow and deep snowpacks produced by the reference dataset.

Finally, we also verify that the results of SnowModel are in good agreement with the set of ablation stakes summer readings ~~collected on Tapado Glacier (Figure S6). In Figure 7 we compare simulated values of surface mass balance on the lowest section of the debris free Tapado Glacier (below 4800 m a.s.l.) against the set of ablation stakes readings (Section 3.1). The surface glacier mass balance of this area is calculated using snow accumulation and erosion (snowfall and wind transport), snow ablation (surface sublimation, blowing snow sublimation and snowmelt runoff) and ice melt and sublimation. As the ablation measured at the stakes and that of their corresponding model grid cells might differ due to topographic differences~~ ~~between the sites and that of the 50 m resolution DEM, we prefer to compare all the stakes readings against spatially averaged values over the glacier tongue.~~ Although there are ~~some~~ differences between observed and simulated values, the main trends throughout the season are similar, with a higher ablation rate between November and February and a reduced rate after March.



410 **Figure 56: Comparison of the observed reference (Wayand and C&M2017) and simulated (SnowModel) snow**  
**variables at the scale of the Corrales catchment. (a) Snow cover area (SCA), (b) snow winter absence (SA), (c) snow**  
**summer persistence (SP), and (d) interannual median of peak annual SWE (SWE<sub>max</sub>) annual maximum snow**  
**accumulation. In (a) the red lines and areas represent the median values and interquartile ranges 25 and 75 of the**  
**SnowModel ensemble runs, respectively. The blue lines-dots correspond to the observations. In b-d the blue points**  
**and error bars represent the mean and standard deviations values, respectively, within a 500-m elevation band (from**  
415 **3500 to 6000 m a.s.l.) and a 45° aspect range (from 22.5° to 337.5°) as the polar plots in Figures 3b-c-h and 4. As**  
**evaluation metrics we use the coefficient of determination ( $R^2$ ), the root mean square error (RMSE) and the mean**  
**bias (BIAS).**




420 **Figure 7: Comparison of the simulated cumulative mass balance of the ablation area of Tapado Glacier against**  
**ablation stakes readings in the period October–April of years 2019–2020 and 2020–2021. The red and blue lines and**  
**areas represent the median values and interquartile ranges 25 and 75 of the SnowModel ensemble runs, respectively.**  
**The markers correspond to the ablation stakes readings and the dashed lines are connecting lines that help**  
**visualization. To help comparison, we set the first marker of each line at the simulated mass balance value in that**  
425 **date. The cumulative mass balance is set at zero on April 1 of each year.**

### 5.3 Snow mass balance and runoff generation


430 In this section we address the main objectives of the study by using the SnowModel simulations to quantify the snow mass balance and to describe the spatial distribution of snowmelt and sublimation fluxes. The catchment-average snow mass balance derived from the SnowModel ensemble runs is shown in Figure 6  
Figure 8 shows the catchment average snow mass fluxes in the study period calculated as the median of the SnowModel ensemble runs. Precipitation occurred mostly in autumn and winter (April to June) with some extraneous events in September 2019, January 2020, October 2020 and March 2021.  
435 Estimated Rainfall–rainfall was much lower than snowfall and was mostly restricted to summer and autumn. Snow surface sublimation was the process that removed most of the snow mass and dominated ablation during winter and spring. Snowmelt runoff and ice melt played a secondary role at the annual level, but their relative importance increased in spring and summer, respectively. From November to March, ice melt was almost the only runoff contributor. Fluxes derived from the transport of snow (wind-transported snow and blowing snow sublimation) were smaller er than surface sublimation and took place only during winter or other periods with fresh snow on the surface. While blowing snow sublimation is always a mass loss (negative values), wind-transported snow can add or remove mass to the catchment (positive or negative values), but in the study period it was mostly a mass loss. The inflow to La Laguna reservoir is shown in Figure 6 to understand the response of streamflow  
440 to snow and ice melt, but as the reservoir is located approximately 24 km downstream of the Corrales outlet it also includes

the contribution from other sub-catchments. We observe a certain correspondence between snow and ice melt from Corrales and La Laguna inflow in summer 2020, but this is not that evident in summer 2021. An interesting feature is that the dry winter of 2020 resulted in a notoriously lower streamflow in the next spring and summer.



445 **Figure 68: Results from SnowModel showing Mmonthly mass fluxes in the Corrales catchment as stacked bars. The**  
**bars correspond to the median value of the ensemble runs. Liquid and solid precipitation are considered as inputs to**  
**the catchment (plotted as positive bars), whereas ice melt, snowmelt runoff, surface sublimation and blowing snow**  
**sublimation are considered as losses from the catchment (plotted as negative bars). Wind-transported snow is the**  
**sum of suspension and saltation and can be both an input and an output (plotted as a black line). The inflow to La**  
450 **Laguna reservoir (normalized by the Corrales catchment area) is plotted as a reference for streamflow variations in**  
**the study region, but it includes the contribution from other sub-catchments downstream of the Corrales outlet.**


Next, we analyze the spatial distribution of snowmelt runoff and other fluxes that remove mass from the snowpack. During winter, snowmelt is minimum, and the snowpack loses mass mostly through sublimation wind transport. We find that the spatial heterogeneity of wind-related snow fluxes is large, with low values of wind transport and blowing snow sublimation on the north-west section of the catchment and larger values to the east. This is explained by the contrasting wind exposure of western and eastern areas (see wind speed records in Figure 2). On the east of the catchment, snow erosion dominates over snow deposition (positive values), ranging from negative to positive values, with a large small-scale variability (Figure 7a). On average, snow erosion is largest at sites between 4500 and 5500 with a W-NW-N aspect (Figure 7c). Blowing snow sublimation, on the other hand, varies from low to large values from west to east, reaching up to 100 mm a<sup>-1</sup> above 4500 m a.s.l. at the border with Argentina (Figure 7b). The elevation bands that present the largest values of blowing snow sublimation are mostly located above 4500 m a.s.l. and have a N-NE aspect (Figure 7d).



**Figure 7: Annual snow mass fluxes associated with wind transport calculated as the medians of the SnowModel ensemble runs. (a) Map of wind transport, (b) Blowing snow sublimation, (c) Polar plot of wind transport and (d) Polar plot of blowing snow sublimation.**

Figure 9 presents maps and polar plots of the ensemble runs medians of snowmelt runoff, snow surface sublimation and sublimation ratio. Once the snow metamorphism starts in response to internal exchanges of energy and vapor, wind transport is reduced, and the snowpack is more favorable to surface sublimation and snowmelt runoff. We find that surface sublimation was the biggest loss of snow mass, showing large values at high-elevation sites on the western areas of the catchment (Figure 8b). Snowmelt runoff shows a heterogeneous distribution with large values at wind-protected valleys in the north-west section of the catchment and very low values to the east. Snowmelt runoff is concentrated at wind sheltered sites located on the west side of the catchment (Figure 89a). On average, the elevation band with the largest values of snowmelt runoff was that particularly at sites located between 4000 and 4500 m a.s.l. with E and SE aspects (Figure 89d). Snow surface sublimation occurs mostly at wind exposed locations (Figure 9b), largely dominating total ablation on the east side of the catchment (large values of sublimation ratio in Figure 9c). Figure 9e shows that snow surface sublimation increases with elevation reaching its

maximum values at sites located above 5500 m a.s.l. with a W, NW and SW aspect. The sublimation fraction is above 60% across the entire domain, and above 80% at high-elevation sites on the west, where surface sublimation is very large, and to the eastern areas of the catchment, where snowmelt runoff is almost zero (Figure 8c). The lowest values of sublimation ratio, i.e. where snowmelt runoff is more important than sublimation, are slightly above 50% and are located in the valleys on the west side of the catchment (Figure 9c), mostly at sites between 4000 and 4500 m a.s.l. with E and SW aspects (Figure 9f). Glaciers appear as sites dominated by snow surface sublimation with a large sublimation fraction larger than 80% (Figure 9e8c). However, in terms of runoff volume, we find that ice melt corresponds to 60% of the runoff contribution from the cryosphere total ablation (snowmelt and ice melt), which is equivalent to 55 mm a<sup>-1</sup> (or 4.3 Mm<sup>3</sup> a<sup>-1</sup>, about 10% of the maximum capacity of La Laguna reservoir).




**Figure 89:** Maps and polar plots of snow mass fluxes calculated as the medians of the SnowModel ensemble runs for 2019-2021. (a) and (d) Snowmelt runoff, (b) and (e) Snow surface sublimation, (c) and (f) Sublimation ratio.

The contrast between runoff generation from (debris-free) glaciers and the rest of the catchment is further analyzed in Figure 9. In this figure, Figure 10, we compare snowmelt runoff and snow surface sublimation over the catchment and its glaciers using boxplots that explicitly consider the uncertainty of the simulations calculated with the (SnowModel ensemble runs). Snow surface sublimation is the largest mass loss from the catchment reaching values between 100 and 250 mm a<sup>-1</sup> (Figure 910a) and represents between 71 to 90% of total ablation (Figure 910b). Over glaciers, surface sublimation is larger than over the rest of the catchment, but snowmelt runoff is lower (Figure 9a), resulting in a very large sublimation fraction (Figure



9b). We find that snowmelt runoff at the catchment level and ice melt have similar mean values (between 34 and 55 mm a<sup>-1</sup>), but the uncertainties derived from our ensemble runs are much larger for snowmelt runoff than for ice melt (Figure 9a). In Figure 10a, the boxplots for debris free glaciers are made using values averaged over glacier grid cells. These results show that snow surface sublimation over glaciers is larger than the catchment average, but the sublimation ratio is similar (Figure 10a b). We find that 50% of the snowmelt runoff is generated from 18–28% of the catchment area (Figure 10c), which we identify as snowmelt hotspots. Another interesting result is that 50% of the catchment produces ca. 80% of the total snowmelt runoff (Figure 10c). In Figure 10d, we show a map of the snowmelt hotspots calculated as the minimum area that produces 50% of the snowmelt runoff in the catchment (calculated from the median of the SnowModel ensemble runs). Using this definition, none of the debris free glaciers correspond to snowmelt hotspots.

The relationship between snowmelt runoff and snow surface sublimation to the selected inputs in Table 3 is assessed using an R-squared analysis of the ensemble runs (Table 4). Ensemble variability in snowmelt runoff is mostly explained by  $z_0$  ( $R^2=0.72$ ) and less so by precipitation ( $R^2=0.11$ ). Snow surface sublimation ensemble variability presents an almost opposite behavior being mostly explained by precipitation ( $R^2=0.65$ ) and less so by  $z_0$  ( $R^2=0.29$ ). Interestingly, we find that snowmelt runoff is better explained by  $z_0$  than snow surface sublimation is. In fact, as snow surface sublimation is the largest flux in the snow mass balance of the catchment, this variable is mostly explained by total precipitation. Sublimation fraction is largely dependent on  $z_0$  ( $R^2=0.85$ ), which controls the magnitude of turbulent latent heat fluxes. Consequently, sublimation ratio is not related to total precipitation, and it is almost completely dependent on  $z_0$  ( $R^2=0.85$ ).



**Figure 9: (a-b) Boxplots of snowmelt runoff, snow surface sublimation, ice melt and sublimation fraction in the Corrales catchment (red boxplots) and debris-free glaciers (blue boxplots). In (a) ice melt values are normalized by**

the catchment area. The edges of the boxes represent the 25<sup>th</sup> and 75<sup>th</sup> percentiles, the horizontal lines represent the median and the dots corresponds to the mean. Outliers are shown with a black cross.

520

**Table 4: R-squared values of linear regressions between SnowModel parameters and outputs**

		<u>Model inputs</u>		
		<u>Precipitation factor</u>	<u>Aerodynamic surface roughness length for snow (<math>z_0</math>)</u>	<u>Wind factor</u>
<u>Model outputs</u>	<u>Snowmelt runoff</u>	<u>0.11</u>	<u>0.72</u>	<u>0.00069</u>
	<u>Snow surface sublimation</u>	<u>0.65</u>	<u>0.29</u>	<u>0.00019</u>
	<u>Sublimation fraction</u>	<u>0.03</u>	<u>0.85</u>	<u>0.00039</u>

#### **5.4 Snowmelt hotspots**

525

We further assess the spatial distribution of runoff generation based on the existence of sites that generate most of the snowmelt runoff in the study catchment. To depict the large heterogeneity of snowmelt runoff we plot the cumulative percentage of snowmelt runoff as a function of the cumulative area (Figure 10a). We find that 50% of the snowmelt runoff is generated from 21-29% of the catchment area (Figure 10ae), which we identify as snowmelt hotspots (Figure 10b). For this, we first rank the grid cells based on their snowmelt runoff and then we define as snowmelt hotspots the first grid cells of the ranking that produce 50% of the total snowmelt runoff. Another interesting result is that we also find that 50% of the catchment produces ca. 80% of the total snowmelt runoff (Figure 10be). This heterogeneity is larger than that of the annual peak SWE (blue areas and line in Figure 10b) and is an indication that snow removal by wind transport and sublimation further increases the spatial variability of the resulting seasonal snowmelt runoff. In Figure 10bd, we show a map of the snowmelt hotspots calculated as the minimum area that produces 50% of the snowmelt runoff in the catchment (calculated from the median of the SnowModel ensemble runs). Interestingly, using this definition, despite the large runoff contribution of ice melt, none of the debris-free glaciers are located on areas labeled as correspond to snowmelt hotspots.


530

535

To determine which factors control the location of snowmelt hotspots, we Figure 11 shows the importance of different factors that control the location of snowmelt hotspots by comparing the cumulative distribution functions of different several descriptive variables of snowmelt hotspots and the rest of the catchment (Figure 11). We find that Snowmelt hotspots are located below 5000 m a.s.l., (mostly between 4200 and 4800 m a.s.l., Figure 11a), have a E, NE and SE aspect (ca. 80% of the snowmelt hotspots have an aspect lower than 180°, Figure 11b) and lower slope angles than the rest of the catchment (Figure 11c). In terms of snow variables, snowmelt hotspots show lower values of SA (Figure 11d), but relatively similar values of SP (Figure 11e) in comparison with the rest of the catchment. For this analysis, simulated SA and SP were calculated using every timestep in the simulation period (in contrast to Figure 5 where we use only the timesteps with available satellite


540

images). snow Snow depth values at the time of maximum accumulation was of at least 0.7 m (Figure 11fd), with SA values lower than 0.3 (Figure 11e) and SP values of at least 0.03 (Figure 11f).



545


**Figure 10: (a) Cumulative simulated snowmelt runoff and maximum SWE against cumulative area, (b) Map of snowmelt hotspots. In (b) the red lines and areas represent the median values and interquartile ranges 25 and 75 of the SnowModel ensemble runs, respectively.**



550

555

**Figure 10: (a-b) Boxplots of snowmelt runoff, snow surface sublimation, ice melt and sublimation ratio in the Corrales catchment (red boxplots) and debris-free glaciers (blue boxplots), (c) Cumulative snowmelt and area, (d) Map of snowmelt hotspots. Boxplots are built using the SnowModel ensemble runs. In (a) ice melt values are normalized by the catchment area. The edges of the boxes represent the 25<sup>th</sup> and 75<sup>th</sup> percentiles, the horizontal lines represent the median and the black dots corresponds to the mean. Outliers are shown with a red cross. In (c) the red lines and areas represent the median values and interquartile ranges 25 and 75 of the SnowModel ensemble runs, respectively.**




**Figure 11: Cumulative distribution functions (CDFs) of selected topographic and snow variables for snowmelt hotspots and the rest of the catchment (except hotspots). In upper part of (b) we add cardinal directions in letters for reference. (d) and (e) were made using the simulated SA and SP (f) was made using the ensemble median of the snow depth at the time of maximum accumulation.**

**Table 5: R-squared values of linear regressions between SnowModel parameters and outputs**

		Model inputs		
		Precipitation factor	Aerodynamic surface roughness length for snow ( $z_0$ )	Wind factor
Model outputs	Snowmelt runoff	0.11	0.72	0.00069
	Snow surface sublimation	0.65	0.29	0.00019
	Sublimation ratio	0.03	0.85	0.00039

In Figure 12 we assess the snow fluxes derived from the snow transport by wind. Wind-transported snow is close to zero on the west side of Corrales catchment and varies between positive (snow deposition) and negative (snow erosion) values to the east (Figure 12a). Snow erosion (negative values) dominates over deposition at sites located between 4500 and 5500 m a.s.l.

570 with W, NW and N aspects (Figure 12c). Blowing snow sublimation is also close to zero on the west side of the catchment and reaches values of up to  $-100 \text{ mm a}^{-1}$  to the east (Figure 12b), particularly at sites above 4500 m a.s.l. with W, NW, N and NE aspects (Figure 12d).



575 **Figure 12: Annual maps and polar plots of snow mass fluxes calculated as the medians of the SnowModel ensemble runs. (a) Snowmelt runoff, (b) Snow surface sublimation, (c) Sublimation ratio.**

## 6 Discussion

### 6.1 Verification of model results

580 The comparison between simulated and reference datasets, both at the point and distributed scales suggest that our SnowModel ensemble runs correctly simulate the most important features of the snow cover in the Corrales catchment, but we identify a number of issues that we discuss in this section. At the point scale, The-the seasonal amounts of snow observed at TAP are well reproduced by SnowModel, but the simulations fail to match the magnitude of some of the accumulation peaks, particularly in terms of snow depth during the-winter 2019 (Figure 5a). Additionally, Most-most of the depletion curves are well captured by the model simulations, except for the depletion curve in the spring 2020, which does not follow the same

585 rates shown by the snow depth and SWE (Figure 5a-b). As suggested by Voordendag et al. (2021) when analyzing the TAP station site in winter 2017, these differences in precipitation peaks and depletions rates are likely given by the uncertainty in the precipitation forcing and the parametrization of fresh snow density and albedo. As Snowmodel does not consider the advection of sensible and latent heat from snow-free areas, we suggest that this might be an additional factor that explain differences between the observed and simulated snow depletion rates (Mott et al., 2011). Additionally Interestingly, model results reproduce the main patterns of glacier ablation on the tongue of Tapado Glacier (Figure S5), although more detailed parameterizations are certainly needed to simulate the energy and mass balance on penitentes fields in their full complexity (Lhermitte et al., 2014; Sinclair and MacDonell, 2015; Nicholson et al., 2016).

Model results are also in good agreement with distributed datasets in terms of general spatial trends, but simulations show some differences with the snow products that are likely related to an insufficient small-scale spatial variability of simulated snow variables. There is a general underestimation of SA that can be explained by the simulated accumulation of snow at wind-exposed sites where snow is quickly removed (Figure 5b), and there is also a very large spatial variability of SP that is not reproduced by the model (Figure 5c). We find that the snow indices proposed by Wayand et al. (2018) are particularly useful to evaluate our results as they provide additional information on the spatial distribution and duration of the snow, in contrast to analyses based solely on the percentage of SCA in the study basin. The ~~correlation-R<sup>2</sup> obtained between for the~~ observed and simulated SA is lower than that obtained for SP ( $R^2=0.479$  versus  $R^2=0.676$ , Figure 5b-c). This result is somehow expected because SA is controlled by accumulation processes that are difficult to reproduce in snow models, such as wind fields over complex terrain and preferential deposition (Mott et al., 2010; Freudiger et al., 2017; Hock et al., 2017). When comparing outputs from the Canadian Hydrological Model (CHM) against satellite-derived SP, Vionnet et al. (2021) obtained evaluation metrics that are similar to those in this study (Pearson correlation coefficients between 0.69 and 0.75). The poorer performance of the model in the accumulation season can also be observed in the ~~low correlation~~ weak relationship ( $R^2=0.23$ ) between the annual maximum accumulation from the model and the SWE reconstruction dataset that we use as reference (Cortés and Margulis, 2017). This ~~low correlation~~ weak relationship is in part explained by the large spread of simulated maximum SWE values when compared against low values of the reference dataset (Figure 5d). This is likely caused by the simulation of low wind speeds and large snow accumulation at sites dominated, in reality, by snow erosion and high wind speeds. On the other hand, a correct representation of SP is more easily achieved because this index is more related to ablation processes that depend on variables that are easier to reproduce, such as solar radiation and air temperature gradients (Wayand et al., 2018). However, despite the ~~good correlation~~ high R<sup>2</sup> values between the ~~obtained from the comparison of~~ observed and simulated SP, the snow cover disappears earlier in the simulations than in the satellite images (Figure 5a). This is likely caused by insufficient snow accumulation-, or an accelerated ablation simulated by the model at high-elevation sites, whereas in reality snow can remain on the surface until the start of the summer (December).

A key advantage of our study is the use of well-distributed AWS data to force the snow simulations, particularly the contrasting wind records of PAN and the rest of the AWSs, which are located at wind-exposed and wind-sheltered locations, respectively. These contrasting wind records allow the simulation of very different snow conditions at western and eastern areas of the

620 catchment. In fact, some differences between our simulations and those of Réveillet et al. (2020), who also used AWS forcing  
in the same catchment, might be caused by the gaps in unavailability of those PAN records during long time periods of their  
study period. For example, we obtain much larger sublimation ratios compared with their simulations resulting from a similar  
AWS forcing (~85% versus ~35%). Similarly, Gascoïn et al. (2013) obtained a sublimation fraction ratio of ca. 70% applying  
SnowModel to an instrumented site in Pascua-Lama (2600–5630 m a.s.l., 29°S) over the 2008 winter. Another good indication  
625 for our simulations are the results of point-scale studies on glaciers of the semiarid Andes, such as those from Ginot et al.  
(2006) and MacDonell et al. (2013), which obtained year-round sublimation ratios around 80% on the upper areas of Tapado  
and Guanaco (5324 m a.s.l., 29.34°S, 70.01°W) glaciers, respectively. On the other hand, Ayala et al. (2017b) obtained a  
sublimation fraction ratio of 12% on the tongue of Tapado Glacier, but those results were obtained from a two-month summer  
period.

## 6.2 Runoff generation and Snowmelt-snowmelt hotspots and hydrological significance

630 In relation to the main hypothesis of this study, i.e. the existence of snowmelt hotspots in the semiarid Andes, we have identified  
the areas that produce half of the snowmelt runoff in the Corrales catchment, and described their main topographic and  
meteorological characteristics. It is highly likely that our model simulations based on ensemble runs correctly identify the  
general location of the snowmelt hotspots, but as the model has difficulties to simulate the full small-scale variability of the  
snow cover (see Section 6.1), the exact location of the sites producing the largest snowmelt amounts is difficult to find. We  
635 estimate that these areas correspond to 21-2918-28% of the catchment area. Although we use a threshold of 50% to define the  
snowmelt hotspots, alternative definitions might be applied in the future depending on the objectives of each study. These  
results are in line with other studies showing how the heterogeneity of snow processes affect runoff generation (DeBeer and  
Pomeroy, 2017). The hydrological relevance of snowmelt hotspots can also be connected with the results from Badger et al.  
(2021), who that used idealized simulations to show that the heterogeneity of the snow cover delays snowmelt runoff and  
640 creates areas of snow persistence. We suggest that in other arid or semiarid catchments, snowmelt hotspots could be expected  
for leeward slopes with relatively low slope angles and moderate elevations. Snow persistence might be an indicator for such  
hydrologically relevant sites. However, to precisely identify snowmelt hotspots and quantify their contribution in other regions,  
a modeling approach combined with satellite information should be applied such as the one presented in this work.  
Snowmelt hotspots in the semiarid Andes are a clear illustration of the large spatial variability of physical processes (from  
645 accumulation to heat exchange) that ultimately control snowmelt runoff (Mott et al., 2018). First, the sheltering and exposure  
to strong winds during and after snowfall events determine large differences in snow accumulation, from areas of preferential  
deposition, such as glacier accumulation zones, to large areas of strong snow erosion by the wind and shallow snowpacks  
(Figures 3 and 4). In this direction, other phenomena, such as mesoscale lowering of the 0°C isotherm have also been suggested  
as important for snow accumulation in this type of environment (Schauwecker et al., 2022). When snow surface conditions are  
650 favorable to wind transport, snowdrift and blowing snow sublimation can mobilize and remove large amounts of snow mass  
from the catchment (Figures 6 and 7). These processes are not spatially uniform and can define areas of snow removal, such



as the eastern section of Corrales. Second, the high surface sublimation rates (of up to 300 mm a<sup>-1</sup>, Figure 89) remove snow and use large amounts of energy from the snowpack that would be otherwise available for melt. The combination of these two factors define an heterogeneous snow cover in which snow-free areas appear very early in the melting season (ca. 50% of the catchment area in August, Figure 6a), likely contributing to snowmelt through the advection of sensible and latent heat (Liston, 1995; Mott et al., 2020; van der Valk et al., 2022). A logical step towards improving our estimates of snowmelt runoff in this type of environment would be the calculation of the contribution of local heat advection from snow-free areas to snowmelt rates, but with a focus on turbulent latent heat fluxes. Other studies have estimated a contribution from heat advection to melt of up to 40% in alpine, prairie and Arctic environments (Mott et al., 2018). The removal of snow by sublimation processes is thus a key feature of dry mountain environments as it adds heterogeneity to the resulting spatial distribution of the seasonal snowmelt runoff. If sublimation rates were lower, the spatial distribution of the seasonal snowmelt runoff are expected to be very similar to that of the end-of-winter SWE, although the snowmelt timing can be very variable from site to site due to differences in melt rates. According to our simulations, snowmelt hotspots are located at sites dominated by preferential deposition, reduced snow transport and low sublimation rates in relation with other wind-exposed areas of the catchment.

~~In regions where snow sublimation is less important in the water balance or restricted to specific locations (e.g. Strasser et al., 2008), snowmelt runoff is expected to originate in areas with large snow accumulation. We here show that this is not necessarily the case in environments with high sublimation rates. The existence of snowmelt hotspots is relevant. This is an important finding~~ for hydrological modelling and water management, since not all sub catchments with snow cover during winter contribute the same way to available water downstream during e.g. the irrigation season. Future snow monitoring driven by the development of seasonal streamflow forecasts in the semiarid Andes could benefit from considering what are sites with the largest snowmelt contribution, because the relationships between streamflow and snow accumulation will be more sensitive at those sites. A better understanding of the factors driving differences between snowmelt hotspots and other areas could be achieved by installing meteorological stations that allow for the calculation of all the energy balance components. Future snow monitoring in the semiarid Andes should thus focus especially on sites where large part of snowmelt is generated and their connection with other hydrological components. In fact, s

Snowmelt hotspots might play a key hydrological role in connection with other components of this landscape, such as peatlands, groundwater recharge and rock glaciers. Valois et al. (2020) investigated the hydrological dynamics of a mountain peatland in a catchment close to Corrales and found that these features connect snowmelt at high-elevation areas with downstream agriculture and human needs, but there are no estimates of the runoff recharge from the melting of the cryosphere.

In this direction, we recommend the assessment of the spatial connectivity between snowmelt hotspots and peatlands. The infiltration of meltwater into rock glaciers is another possible link between snowmelt hotspots and the hydrology of the catchments (Schaffer et al., 2019; Pourrier et al., 2014). Recently, Navarro et al. (2023b) presented a set of geophysical measurements of the internal structure of Tapado debris-cover and rock glacier that reveal an intricate hydrological network that connects surface meltwater with massive ice lenses and ice-rich permafrost, but the dynamics of this connection is yet to be understood in detail. For example, it is not clear how the regulating role of rock glaciers changes in the presence of snow

rich or poor years. Interestingly, At a larger scale, Álvarez-Garretón et al. (2021) showed that the hydrological memory of snow-dominated catchments between 30 and 35-°S are strongly influenced by the infiltration of snowmelt at high-elevation. Exploratory drillings have shown the presence of groundwater near the outlet of the Corrales catchment, which might be indicative of a recharge area. Finally, a brief analysis of satellite images acquired after the disappearance of the seasonal snow cover (Figure S7) suggests that snowmelt hotspots areas tend to have higher values of soil moisture and might contribute to the development of riparian vegetation. In any case, this connection is not evident and needs to be verified and extended by field measurements and more in-depth analyses of remote sensing products.

Although glacier runoff is an important component in the water balance of the catchment and the semiarid Andes (Gascoin et al., 2011; Rodriguez et al., 2016), we found that the main glacier in our study catchment, Tapado Glacier, is not located in an area defined as snowmelt hotspot in this study. However, in line with the results of Gascoin et al. (2013), the snow products analyzed in this study and the results of the SnowModel simulations show that Tapado Glacier is located in an area of preferential deposition and low snow erosion rates (Figures 3 and 7, 4 and 12). Furthermore, our results show that snow sublimation in glacierized areas is very high (Figure ~~10b~~9b), suggesting that sublimation and its associated turbulent latent heat fluxes significantly reduce summer ablation by reducing the energy available for melt. On the other hand, once the snow disappears, the ice surface is quickly melted (ice sublimation calculated by SnowModel was much lower than ice melt), creating large amounts of runoff that reach up to ~~60~~4% of the runoff contributions in the catchment. These results confirm that glaciers are key contributors to runoff in this region, and the most important ones during dry periods. As the ice melt volume estimate of  $4.3 \text{ Mm}^3 \text{ a}^{-1}$  is relatively high in comparison with other studies (Robson et al. (2022) estimated that the average annual volume change of Tapado Glacier in the period 2012-2015 was of  $0.6 \text{ Mm}^3$ ), we recommend the application of more detailed schemes to solve the energy and mass balance of glaciers in the Corrales catchment.

Although the advantageous location of the AWSs allowed us to capture the main patterns of the spatial variability of the meteorological forcing variables, the total number of stations is still insufficient to capture the full complexity of snow processes in this dry environment. Additionally, future modelling studies will benefit from the inclusion of heat advection from snow-free areas (van der Valk et al., 2022) and time-dependent parameterizations of  $z_0$  (Réveillet et al., 2020; Voordendag et al., 2021). As the period analyzed in this study was very dry, the presence and role of snowmelt hotspots should be analyzed during wet years, where a more uniform distribution of snow accumulation could be expected. However, the hydrological processes that we analyzed in this study might become more frequent in the future, in line with the projections for this region in the context of climate change.

## 7 Conclusions

In this study we hypothesize and test the existence of areas providing most of the snowmelt runoff in a sublimation-dominated environment-catchment in the semiarid Andes of Chile. For this we used a process-based snow model (SnowModel) to simulate the evolution of the seasonal snow cover over a two-year period in a  $79 \text{ km}^2$  catchment (Corrales catchment) located in the

upper areas of the semiarid Andes of Chile. Our simulations are analyzed using field data and a set of independent satellite-derived snow products that describe the spatial properties of the snow cover. Our main conclusions are as follows:

1. Snow surface sublimation is the dominant snow ablation component across the study catchment, representing between 71 and ~~90~~4% of total ablation from the surface (snowmelt plus snow surface sublimation). In winter and after snowfalls, blowing snow sublimation is also important and dominates over wind-exposed areas in the east section of the catchment.
2. We estimate that 50% of the snowmelt runoff is produced by ~~21-29~~18-28% of the catchment area, which we define as “snowmelt hotspots”. As in any mountain terrain, the large spatial heterogeneity of seasonal snowmelt runoff is a consequence of complex snow accumulation patterns, but in dry mountain environments, this heterogeneity is further enlarged by large sublimation rates, which reduce snow available for melt at wind-exposed sites.
3. Snowmelt hotspots in our study sites are mostly located at elevations between 4200 and 4800 m, have easterly aspects, low slope angles, and high snow accumulation ~~and persistence.~~

As suggested by previous research in the semiarid Andes, glaciers of the study catchment play a key role during dry years, providing most of the runoff during the end of the summer. Snowmelt hotspots can be a useful concept to understand runoff generation in arid and semiarid mountain regions ~~this type of environment~~ as it might help to identify the areas of hydrological connections between the components of the catchment, such as the snow cover, groundwater, peatlands and rock glaciers. Additionally, snowmelt hotspots are good candidates for the installation of hydro-meteorological stations, particularly in situations where only limited financial resources are available for the monitoring of large, remote regions. This study was carried out during a severe drought that affects the semiarid Andes since 2010. As current climate projections for this region suggest that droughts will be more frequent in the future, we expect that our results will be relevant in the context of climate change. In particular, they can be used to better understand the hydrological response of the semiarid Andes and to improve the planning and management of its water resources.

## 740 **Acknowledgments**

ÁA acknowledges ANID-Fondecyt postdoctoral project 3190732. We thank the project ANID-FONDEF IDeA I+D 2021 ID21I10129, ANID-CENTROS REGIONALES R20F0008 and all the people involved in the data collection on Tapado Glacier, particularly Ignacio Diaz. CEAZAmet is thanked for support with meteorological data acquisition. Gonzalo Cortés is kindly acknowledged for providing the SWE reconstruction dataset (Cortés and Margulis, 2017).

## 745 **Data availability**

Tapado and Paso Agua Negra meteorological [raw](#) data are freely available at [www.cezamet.cl](http://www.cezamet.cl). [The dataset and scripts used in this study are available at 10.5281/zenodo.8029996.](#) ~~Tapado AWS appears in the webpage as El Tapado. Tapado Glacier meteorological data will be included in the same webpage.~~

## **Author contributions**

750 **ÁA** designed the study with the collaboration of **SM**. **ÁA** led the collection at the Tapado Glacier field site. **ÁA** performed the SnowModel simulations and main analysis with the collaboration of **SS**. **ÁA** interpreted results and wrote the manuscript with the help of **SM** and **SS**.

## **Competing interests**

The authors declare that they have no conflict of interest.

## 755 **References**

- Álvarez-Garretón, C., Mendoza, P. A., Pablo Boisier, J., Addor, N., Galleguillos, M., Zambrano-Bigiarini, M., Lara, A., Puelma, C., Cortes, G., Garreaud, R., McPhee, J., and Ayala, A.: The CAMELS-CL dataset: Catchment attributes and meteorology for large sample studies-Chile dataset, *Hydrol. Earth Syst. Sci.*, 22, 5817–5846, <https://doi.org/10.5194/hess-22-5817-2018>, 2018.
- 760 Álvarez-Garretón, C., Pablo Boisier, J., Garreaud, R., Seibert, J., and Vis, M.: Progressive water deficits during multiyear droughts in basins with long hydrological memory in Chile, *Hydrol. Earth Syst. Sci.*, 25, 429–446, <https://doi.org/10.5194/hess-25-429-2021>, 2021.
- Arias, P. A., Garreaud, R., Poveda, G., Espinoza, J. C., Molina-Carpio, J., Masiokas, M., Viale, M., Scaff, L., and van Oevelen, P. J.: Hydroclimate of the Andes Part II: Hydroclimate Variability and Sub-Continental Patterns, *Front. Earth Sci.*, 8, 1–25, <https://doi.org/10.3389/feart.2020.505467>, 2021.
- 765 Ayala, Á., Pellicciotti, F., Peleg, N., and Burlando, P.: Melt and surface sublimation across a glacier in a dry environment: Distributed energy-balance modelling of Juncal Norte Glacier, Chile, *J. Glaciol.*, 63, 803–822, <https://doi.org/10.1017/jog.2017.46>, 2017a.
- Ayala, Á., Pellicciotti, F., MacDonell, S., McPhee, J., and Burlando, P.: Patterns of glacier ablation across North-Central Chile:   
770 Identifying the limits of empirical melt models under sublimation-favorable conditions, *Water Resour. Res.*, 53, 5601–5625, <https://doi.org/10.1002/2016WR020126>, 2017b.
- Ayala, Á., Farías-Barahona, D., Huss, M., Pellicciotti, F., McPhee, J., and Farinotti, D.: Glacier runoff variations since 1955

- in the Maipo River basin, in the semiarid Andes of central Chile, *Cryosph.*, 14, 2005–2027, <https://doi.org/10.5194/tc-14-2005-2020>, 2020.
- 775 Badger, A. M., Bjarke, N., Molotch, N. P., and Livneh, B.: The sensitivity of runoff generation to spatial snowpack uniformity in an alpine watershed: Green Lakes Valley, Niwot Ridge Long-Term Ecological Research station, *Hydrol. Process.*, 35, 1–14, <https://doi.org/10.1002/hyp.14331>, 2021.
- Brock, B. W., Willis, I. C., and Sharp, M. J.: Measurement and parameterization of aerodynamic roughness length variations at Haut Glacier d’Arolla, Switzerland, *J. Glaciol.*, 52, 281–297, <https://doi.org/10.3189/172756506781828746>, 2006.
- 780 Cortés, G. and Margulis, S.: Impacts of El Niño and La Niña on interannual snow accumulation in the Andes: Results from a high-resolution 31 year reanalysis, *Geophys. Res. Lett.*, 44, 6859–6867, <https://doi.org/10.1002/2017GL073826>, 2017.
- Cortés, G., Giroto, M., and Margulis, S. a.: Snow process estimation over the extratropical Andes using a data assimilation framework integrating MERRA data and Landsat imagery, *Water Resour. Res.*, 52, <https://doi.org/10.1002/2014WR015716>, 2016.
- 785 DeBeer, C. M. and Pomeroy, J. W.: Influence of snowpack and melt energy heterogeneity on snow cover depletion and snowmelt runoff simulation in a cold mountain environment, *J. Hydrol.*, 553, 199–213, <https://doi.org/10.1016/j.jhydrol.2017.07.051>, 2017.
- Información Oficial Hidrometeorológica y de Calidad de Aguas en Línea: <https://snia.mop.gob.cl/BNAConsultas/reportes>, last access: 20 January 2023.
- 790 DGA: Inventario Público de Glaciares, actualización 2022, <https://dga.mop.gob.cl/Paginas/InventarioGlaciares.aspx>, 2022.
- Dornes, P. F. P., Pomeroy, J. J. W., Pietroniro, A., Carey, S. K., and Quinton, W. L.: Influence of landscape aggregation in modelling snow-cover ablation and snowmelt runoff in a sub-arctic mountainous environment, *Hydrol. Sci. J.*, 53, 725–740, <https://doi.org/10.1623/hysj.53.4.725>, 2008.
- Level-2A Algorithm Overview: <https://sentinels.copernicus.eu/web/sentinel/technical-guides/sentinel-2-psi/level-2a/algorithm>, last access: 20 January 2023.
- 795 Favier, V., Falvey, M., Rabatel, A., Praderio, E., and Lopez, D.: Interpreting discrepancies between discharge and precipitation in high-altitude area of Chile’s Norte Chico region (26–32°S), *Water Resour. Res.*, 45, <https://doi.org/10.1029/2008WR006802>, 2009.
- Fitzpatrick, N., Radić, V., and Menounos, B.: A multi-season investigation of glacier surface roughness lengths through in situ and remote observation, *Cryosphere*, 13, 1051–1071, <https://doi.org/10.5194/tc-13-1051-2019>, 2019.
- 800 Freudiger, D., Kohn, I., Seibert, J., Stahl, K., and Weiler, M.: Snow redistribution for the hydrological modeling of alpine catchments, *Wiley Interdiscip. Rev. Water*, 4, e1232, <https://doi.org/10.1002/wat2.1232>, 2017.
- Gascoïn, S., Kinnard, C., Ponce, R., Lhermitte, S., MacDonell, S., and Rabatel, A.: Glacier contribution to streamflow in two headwaters of the Huasco River, Dry Andes of Chile, *Cryosph.*, 5, 1099–1113, <https://doi.org/10.5194/tc-5-1099-2011>, 2011.
- 805 Gascoïn, S., Lhermitte, S., Kinnard, C., Bortels, K., and Liston, G. E.: Wind effects on snow cover in Pascua-Lama, Dry Andes of Chile, *Adv. Water Resour.*, 55, 25–39, <https://doi.org/10.1016/j.advwatres.2012.11.013>, 2013.

- Ginot, P., Kull, C., Schotterer, U., Schwikowski, M., and Gäggeler, H. W.: Glacier mass balance reconstruction by sublimation induced enrichment of chemical species on Cerro Tapado (Chilean Andes), *Clim. Past*, 2, 21–30, <https://doi.org/10.5194/cp-2-21-2006>, 2006.
- 810 Groot Zwaafink, C. D., Mott, R., and Lehning, M.: Seasonal simulation of drifting snow sublimation in Alpine terrain, *Water Resour. Res.*, 49, 1581–1590, <https://doi.org/10.1002/wrcr.20137>, 2013.
- Hock, R., Hutchings, J. K., and Lehning, M.: Grand Challenges in Cryospheric Sciences: Toward Better Predictability of Glaciers, Snow and Sea Ice, *Front. Earth Sci.*, 5, 1–14, <https://doi.org/10.3389/feart.2017.00064>, 2017.
- Hood, E., Williams, M., and Cline, D.: Sublimation from a seasonal snowpack at a continental, mid-latitude alpine site, *Hydrol. Process.*, 1797, 1781–1797, 1999.
- 815 Huning, L. S. and AghaKouchak, A.: Global snow drought hot spots and characteristics, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 19753–19759, <https://doi.org/10.1073/PNAS.1915921117>, 2020.
- Jackson, S. and Prowse, T.: Spatial variation of snowmelt and sublimation in a high-elevation semi-desert basin of western Canada, *Hydrol. Process.*, 2627, 2611–2627, <https://doi.org/10.1002/hyp>, 2009.
- 820 Kalthoff, N., Fiebig-Wittmaack, M., Meißner, C., Kohler, M., Uriarte, M., Bischoff-Gauß, I., and Gonzales, E.: The energy balance, evapo-transpiration and nocturnal dew deposition of an arid valley in the Andes, *J. Arid Environ.*, 65, 420–443, <https://doi.org/10.1016/j.jaridenv.2005.08.013>, 2006.
- Kraaijenbrink, P. D. A., Stigter, E. E., Yao, T., and Immerzeel, W. W.: Climate change decisive for Asia’s snow meltwater supply, *Nat. Clim. Chang.*, 11, 591–597, <https://doi.org/10.1038/s41558-021-01074-x>, 2021.
- 825 Lehning, M., Bartelt, P., Brown, B., and Lehning, M.: A physical SNOWPACK model for the Swiss avalanche warning: Part II. Snow microstructure, *Cold Reg. Sci. Technol.*, 35, 147–167, 2002.
- Lehning, M., Löwe, H., Ryser, M., and Raderschall, N.: Inhomogeneous precipitation distribution and snow transport in steep terrain, *Water Resour. Res.*, 44, W07404, <https://doi.org/10.1029/2007WR006545>, 2008.
- Lehning, M., Grünewald, T., and Schirmer, M.: Mountain snow distribution governed by an altitudinal gradient and terrain roughness, *Geophys. Res. Lett.*, 38, 1–5, <https://doi.org/doi:10.1029/2011GL048927>, 2011.
- 830 Lhermitte, S., Abermann, J., and Kinnard, C.: Albedo over rough snow and ice surfaces, *Cryosph.*, 8, 1069–1086, <https://doi.org/10.5194/tc-8-1069-2014>, 2014.
- Liston, G. E.: Local advection of momentum, heat, and moisture during the melt of patchy snow covers, *J. Appl. Meteorol.*, 1995.
- 835 Liston, G. E. and Elder, K.: A distributed snow-evolution modeling system (snowmodel), *J. Hydrometeorol.*, 7, 1259–1276, <https://doi.org/10.1175/JHM548.1>, 2006a.
- Liston, G. E. and Elder, K.: A meteorological distribution system for high-resolution terrestrial modeling (MicroMet), *J. Hydrometeorol.*, 7, 217–234, <https://doi.org/10.1175/JHM486.1>, 2006b.
- Liston, G. E. and Hall, D. K.: An energy-balance model of lake-ice evolution, *J. Glaciol.*, 41, 373–382, 1995.
- 840 Liston, G. E., Sturm, M. H., En, G., Lr Ston, E., and Sturm, M. H.: A snow-transport model for complex terrain, *J. Glaciol.*,

- 44, 498–516, <https://doi.org/https://doi.org/10.3198/1998JoG44-148-498-516>, 1998.
- Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. A., Berezovskaya, S., and Tabler, R. D.: Simulating complex snow distributions in windy environments using SnowTran-3D, *J. Glaciol.*, 53, 241–256, <https://doi.org/10.3189/172756507782202865>, 2007.
- 845 Litt, M., Shea, J., Wagnon, P., Steiner, J., Koch, I., Stigter, E., and Immerzeel, W.: Glacier ablation and temperature indexed melt models in the Nepalese Himalaya, *Sci. Rep.*, 9, 1–13, <https://doi.org/10.1038/s41598-019-41657-5>, 2019.
- Luce, C. H., Tarboton, D. G., and Cooley, K. R.: The influence of the spatial distribution of snow on basin-averaged snowmelt, *Hydrol. Process.*, 12, 1671–1683, [https://doi.org/10.1002/\(SICI\)1099-1085\(199808/09\)12:10/11<1671::AID-HYP688>3.0.CO;2-N](https://doi.org/10.1002/(SICI)1099-1085(199808/09)12:10/11<1671::AID-HYP688>3.0.CO;2-N), 1998.
- 850 MacDonald, J. P. and Pomeroy, J. W.: Gauge Undercatch of Two Common Snowfall Gauges in a Prairie Environment, *Proc. 64th East. Snow Conf. St. John's, Canada.*, 119–126, 2007.
- MacDonell, S., Kinnard, C., Mölg, T., Nicholson, L., and Abermann, J.: Meteorological drivers of ablation processes on a cold glacier in the semi-arid Andes of Chile, *Cryosph.*, 7, 1513–1526, <https://doi.org/10.5194/tc-7-1513-2013>, 2013.
- Mankin, J. S., Viviroli, D., Singh, D., Hoekstra, A. Y., and Diffenbaugh, N. S.: The potential for snow to supply human water  
855 demand in the present and future, *Environ. Res. Lett.*, 10, <https://doi.org/10.1088/1748-9326/10/11/114016>, 2015.
- Masiokas, M. H., Villalba, R., Luckman, B. H., Le Quesne, C., and Aravena, J. C.: Snowpack variations in the central Andes of Argentina and Chile, 1951-2005: Large-scale atmospheric influences and implications for water resources in the region, *J. Clim.*, 19, 6334–6352, <https://doi.org/10.1175/JCLI3969.1>, 2006.
- Mendoza, P. A., Shaw, T. E., McPhee, J., Musselman, K. N., Revuelto, J., and MacDonell, S.: Spatial Distribution and Scaling  
860 Properties of Lidar-Derived Snow Depth in the Extratropical Andes, *Water Resour. Res.*, 56, e2020WR028480, <https://doi.org/https://doi.org/10.1029/2020WR028480>, 2020.
- Merkouriadi, I., Lemmetyinen, J., Liston, G. E., and Pulliainen, J.: Solving Challenges of Assimilating Microwave Remote Sensing Signatures With a Physical Model to Estimate Snow Water Equivalent, *Water Resour. Res.*, 57, 1–24, <https://doi.org/10.1029/2021WR030119>, 2021.
- 865 Mernild, S. H., Liston, G. E., Hiemstra, C. A., Malmros, J. K., Yde, J. C., and McPhee, J.: The Andes Cordillera. Part I: snow distribution, properties, and trends (1979-2014), *Int. J. Climatol.*, <https://doi.org/10.1002/joc.4804>, 2016.
- Mernild, S. H., Liston, G. E., Hiemstra, C. A., Yde, J. C., and Casassa, G.: Annual river runoff variations and trends for the Andes Cordillera, *J. Hydrometeorol.*, 19, 1167–1189, <https://doi.org/10.1175/JHM-D-17-0094.1>, 2018.
- Montecinos, A. and Aceituno, P.: Seasonality of the ENSO-related rainfall variability in central Chile and associated  
870 circulation anomalies, *J. Clim.*, 16, 281–296, [https://doi.org/10.1175/1520-0442\(2003\)016<0281:SOTERR>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0281:SOTERR>2.0.CO;2), 2003.
- Mott, R., Schirmer, M., Bavay, M., Grünewald, T., and Lehning, M.: Understanding snow-transport processes shaping the mountain snow-cover, *Cryosph.*, 4, 545–559, <https://doi.org/10.5194/tc-4-545-2010>, 2010.
- Mott, R., Egli, L., Grünewald, T., Dawes, N., Manes, C., Bavay, M., and Lehning, M.: Micrometeorological processes driving snow ablation in an Alpine catchment, *Cryosph.*, 5, 1083–1098, <https://doi.org/10.5194/tc-5-1083-2011>, 2011.

- 875 Mott, R., Vionnet, V., and Grünewald, T.: The Seasonal Snow Cover Dynamics: Review on Wind-Driven Coupling Processes, *Front. Earth Sci.*, 6, <https://doi.org/10.3389/feart.2018.00197>, 2018.
- Mott, R., Stiperski, I., and Nicholson, L.: Spatio-temporal flow variations driving heat exchange processes at a mountain glacier, *Cryosph.*, 14, 4699–4718, <https://doi.org/10.5194/tc-14-4699-2020>, 2020.
- NASA JPL: NASADEM Merged DEM Global 1 arc second V001 [Data set],  
880 [https://doi.org/10.5067/MEASURES/NASADEM/NASADEM\\_HGT.001](https://doi.org/10.5067/MEASURES/NASADEM/NASADEM_HGT.001), 2020.
- Navarro, G., MacDonell, S., and Valois, R.: A conceptual hydrological model of semiarid Andean headwater systems in Chile, *Prog. Phys. Geogr.*, 0, 1–19, <https://doi.org/10.1177/03091333221147649>, 2023a.
- Navarro, G., Valois, R., MacDonell, S., de Pasquale, G., and Díaz, J. P.: Internal structure and water routing of an ice-debris  
landform assemblage using multiple geophysical methods in the semiarid Andes, 1–14,  
885 <https://doi.org/10.3389/feart.2023.1102620>, 2023b.
- Nicholson, L. I., Petlicki, M., Partan, B., and MacDonell, S.: 3D surface properties of glacier penitentes over an ablation season, measured using a Microsoft Xbox Kinect, *Cryosph.*, 10, 1897–1913, <https://doi.org/10.5194/tc-2015-207>, 2016.
- Palm, S. P., Kayetha, V., Yang, Y., and Pauly, R.: Blowing snow sublimation and transport over Antarctica from 11 years of CALIPSO observations, *Cryosph.*, 11, 2555–2569, <https://doi.org/10.5194/tc-11-2555-2017>, 2017.
- 890 Pitte, P., Masiokas, M., Gargantini, H., Ruiz, L., Berthier, E., Ferri Hidalgo, L., Zalazar, L., Dussailant, I., Viale, M., Zorzut, V., Corvalán, E., Scarpa, J. P., Costa, G., and Villalba, R.: Recent mass-balance changes of Agua Negra glacier (30°S) in the Desert Andes of Argentina, *J. Glaciol.*, 1–13, <https://doi.org/10.1017/jog.2022.22>, 2022.
- Pohl, S., Marsh, P., and Liston, G. E.: Spatial-temporal variability in turbulent fluxes during spring snowmelt, Arctic, Antarct. Alp. Res., 38, 136–146, [https://doi.org/10.1657/1523-0430\(2006\)038\[0136:SVITFD\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2006)038[0136:SVITFD]2.0.CO;2), 2006.
- 895 Pomeroy, J. W. and Li, L.: Prairie and arctic areal snow cover mass balance using a blowing snow model, *J. Geophys. Res.*, 105, 26619, <https://doi.org/10.1029/2000JD900149>, 2000.
- Pomeroy, J. W., Gray, D. M., Shook, K. R., Toth, B., Essery, R. L. H., Pietroniro, A., and Hedstrom, N.: An evaluation of snow accumulation and ablation processes for land surface modelling, *Hydrol. Process.*, 12, 2339–2367, [https://doi.org/https://doi.org/10.1002/\(SICI\)1099-1085\(199812\)12:15<2339::AID-HYP800>3.0.CO;2-L](https://doi.org/https://doi.org/10.1002/(SICI)1099-1085(199812)12:15<2339::AID-HYP800>3.0.CO;2-L), 1998.
- 900 Pomeroy, J. W., Toth, B., Granger, R. J. J., Hedstrom, N. R. R., and Essery, R. L. H. L. H.: Variation in surface energetics during snowmelt in a subarctic mountain catchment, *J. Hydrometeorol.*, 4, 702–719, [https://doi.org/10.1175/1525-7541\(2003\)004<0702:VISED>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0702:VISED>2.0.CO;2), 2003.
- Pourrier, J., Jourde, H., Kinnard, C., Gascoïn, S., and Monnier, S.: Glacier meltwater flow paths and storage in a geomorphologically complex glacial foreland: The case of the Tapado glacier, dry Andes of Chile (30°S), *J. Hydrol.*, 519,  
905 1068–1083, <https://doi.org/10.1016/j.jhydrol.2014.08.023>, 2014.
- Ragetti, S., Cortés, G., McPhee, J., and Pellicciotti, F.: An evaluation of approaches for modelling hydrological processes in high-elevation, glacierized Andean watersheds, *Hydrol. Process.*, 28, 5674–5695, <https://doi.org/10.1002/hyp.10055>, 2014.
- Réveillet, M., MacDonell, S., Gascoïn, S., Kinnard, C., Lhermitte, S., and Schaffer, N.: Impact of forcing on sublimation



- simulations for a high mountain catchment in the semiarid Andes, *Cryosph.*, 14, 147–163, [https://doi.org/10.5194/tc-14-147-](https://doi.org/10.5194/tc-14-147-2020)  
910 2020, 2020.
- Robson, B. A., MacDonell, S., Ayala, Á., Bolch, T., Nielsen, P. R., and Vivero, S.: Glacier and Rock Glacier changes since the 1950s in the La Laguna catchment, Chile, *Cryosph.*, 16, 647–665, <https://doi.org/10.5194/tc-2021-200>, 2022.
- Rodriguez, M., Ohlanders, N., Pellicciotti, F., Williams, M. W., and McPhee, J.: Estimating runoff from a glacierized catchment using natural tracers in the semi-arid Andes cordillera, *Hydrol. Process.*, 30, 3609–3626,  
915 <https://doi.org/10.1002/hyp.10973>, 2016.
- Saavedra, F., Cortés, G., Viale, M., Margulis, S., and McPhee, J.: Atmospheric Rivers Contribution to the Snow Accumulation Over the Southern Andes (26.5° S–37.5° S), *Front. Earth Sci.*, 8, 1–11, <https://doi.org/10.3389/feart.2020.00261>, 2020.
- Sarricolea, P., Herrera-Ossandon, M., and Meseguer-Ruiz, Ó.: Climatic regionalisation of continental Chile, *J. Maps*, 13, 66–73, <https://doi.org/10.1080/17445647.2016.1259592>, 2017.
- 920 Scaff, L., Rutllant, J. A., Rahn, D., Gascoin, S., and Rondanelli, R.: Meteorological Interpretation of Orographic Precipitation Gradients along an Andes West Slope Basin at 30°S (Elqui Valley, Chile), *J. Hydrometeorol.*, 18, 713–727, <https://doi.org/10.1175/JHM-D-16-0073.1>, 2017.
- Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E., and Valois, R.: Rock glaciers as a water resource in a changing climate in the semiarid Chilean Andes, *Reg. Environ. Chang.*, 2, <https://doi.org/10.1007/s10113-018-01459-3>, 2019.
- 925 Schauwecker, S., Palma, G., MacDonell, S., Ayala, Á., and Viale, M.: The Snowline and 0°C Isotherm Altitudes During Precipitation Events in the Dry Subtropical Chilean Andes as Seen by Citizen Science, Surface Stations, and ERA5 Reanalysis Data, *Front. Earth Sci.*, 10, 1–19, <https://doi.org/10.3389/feart.2022.875795>, 2022.
- Schwanghart, W. and Scherler, D.: Short Communication: TopoToolbox 2 - MATLAB-based software for topographic analysis and modeling in Earth surface sciences, *Earth Surf. Dyn.*, 2, 1–7, <https://doi.org/10.5194/esurf-2-1-2014>, 2014.
- 930 Seibert, J., Vis, M. J. P., Kohn, I., Weiler, M., and Stahl, K.: Technical Note: Representing glacier dynamics in a semi-distributed hydrological model, *Hydrol. Earth Syst. Sci. Discuss.*, 1–20, <https://doi.org/10.5194/hess-2017-158>, 2017.
- Sinclair, K. E. and MacDonell, S.: Seasonal evolution of penitente glaciochemistry at Tapado Glacier, Northern Chile, *Hydrol. Process.*, 30, n/a-n/a, <https://doi.org/10.1002/hyp.10531>, 2015.
- Sproles, E. A., Kerr, T., Orrego Nelson, C., and Lopez Aspe, D.: Developing a Snowmelt Forecast Model in the Absence of  
935 Field Data, *Water Resour. Manag.*, 30, 2581–2590, <https://doi.org/10.1007/s11269-016-1271-4>, 2016.
- Stigter, E. E., Litt, M., Steiner, J. F., Bonekamp, P. N. J., Shea, J. M., Bierkens, M. F. P., and Immerzeel, W. W.: The Importance of Snow Sublimation on a Himalayan Glacier, *Front. Earth Sci.*, 6, 1–16, <https://doi.org/10.3389/feart.2018.00108>, 2018.
- Strasser, U., Bernhardt, M., Weber, M., Liston, G. E., and Mauser, W.: Is snow sublimation important in the alpine water balance?, *Cryosph.*, 2, 53–66, 2008.
- 940 van der Valk, L. D., Teuling, A. J., Girod, L., Pirk, N., Stoffer, R., and van Heerwaarden, C. C.: Understanding wind-driven melt of patchy snow cover, *Cryosph.*, 16, 4319–4341, <https://doi.org/10.5194/tc-16-4319-2022>, 2022.

- Valois, R., Schaffer, N., Figueroa, R., Maldonado, A., Yáñez, E., Hevia, A., Carrizo, G. Y., and MacDonell, S.: Characterizing the water storage capacity and hydrological role of mountain peatlands in the arid andes of North-Central Chile, *Water*, 12, 945 <https://doi.org/10.3390/W12041071>, 2020.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M. M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geosci. Model Dev.*, 5, 773–791, <https://doi.org/10.5194/gmd-5-773-2012>, 2012.
- Vionnet, V., Marsh, C. B., Menounos, B., Gascoin, S., Wayand, N. E., Shea, J., Mukherjee, K., and Pomeroy, J. W.: Multi-950 scale snowdrift-permitting modelling of mountain snowpack, *Cryosphere*, 15, 743–769, <https://doi.org/10.5194/tc-15-743-2021>, 2021.
- Voordendag, A., Réveillet, M., MacDonell, S., and Lhermitte, S.: Snow model comparison to simulate snow depth evolution and sublimation at point scale in the semi-arid Andes of Chile, *Cryosph.*, 15, 4241–4259, <https://doi.org/10.5194/tc-15-4241-2021>, 2021.
- 955 Wayand, N. E., Marsh, C. B., Shea, J. M., and Pomeroy, J. W.: Globally scalable alpine snow metrics, *Remote Sens. Environ.*, 213, 61–72, <https://doi.org/10.1016/j.rse.2018.05.012>, 2018.
- Zhang, Y. and Ishikawa, M.: Sublimation from thin snow cover at the edge of the Eurasian cryosphere in Mongolia, *Hydrol. Process.*, 3575, 3564–3575, <https://doi.org/10.1002/hyp>, 2008.