

1 Mixed-cultivated grasslands enhance runoff generation and reduce soil loss in the restoration of  
2 degraded alpine hillsides

3 Yulei Ma<sup>1</sup>, Yifan Liu<sup>1,2</sup>, Jesús Rodrigo-Comino<sup>3</sup>, Manuel López-Vicente<sup>4</sup>, Gao-Lin Wu<sup>1,2,5,\*</sup>

4 <sup>1</sup> *State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, College of Soil*  
5 *and Water Conservation Science and Engineering (Institute of Soil and Water Conservation),*  
6 *Northwest A & F University, Yangling, Shaanxi 712100, China*

7 <sup>2</sup> *Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water*  
8 *Resource, Yangling, Shaanxi 712100, China*

9 <sup>3</sup> *Departamento de Análisis Geográfico Regional y Geografía Física, Facultad de Filosofía y Letras,*  
10 *Campus Universitario de Cartuja, University of Granada, Granada, Spain*

11 <sup>4</sup> *Group Aquaterra, Interdisciplinary Centre of Chemistry and Biology, CICA-UDC, Universidade da*  
12 *Coruña, 15071 A Coruña, Spain*

13 <sup>5</sup> *CAS Center for Excellence in Quaternary Science and Global Change, Xi'an, 710061, China*

14 **\*Correspondence author:** Gao-Lin Wu, e-mail: wugaolin@nwsuaf.edu.cn (G.L. Wu).

15 **Correspondence address:** State Key Laboratory of Soil Erosion and Dryland Farming on the Loess  
16 Plateau, Northwest A & F University, No 26, Xinong Road, Yangling, Shaanxi 712100, P.R. China

17 Phone: +86- (29) 87012884 Fax: +86- (29) 87016082

18 **ABSTRACT**

19 Vegetation restoration is among the most effective measures for controlling runoff and soil erosion  
20 resulting from human activity. Nevertheless, few studies have been undertaken to analyze the effects  
21 of grassland restoration on maintaining ~~the stability of the~~ local runoff ~~amount~~, especially, in alpine  
22 degraded hillsides where mixed-cultivated grasslands predominate. In this research, ~~we conducted in-~~  
23 ~~situ monitoring using~~ runoff plots were established to investigate the impact of three mixed-cultivated  
24 grasslands, each sowing two grass species per plot: *Deschampsia cespitosa* and *Elymus nutans* (*DE*),  
25 *Poa pratensis* L.cv. Qinghai and *Elymus nutans* (*PE*), and *Poa pratensis* L.cv. Qinghai and  
26 *Deschampsia cespitosa* (*PD*); on a 20-degree slope, assessing the activation and volume of surface  
27 runoff and the magnitude of soil loss in alpine degraded hillsides over three years: 2019, 2020 and  
28 2022. A severely degraded meadow (*SDM*) plot was used as control. The findings indicated that  
29 mixed-cultivated grasslands can effectively maintain runoff and reduce soil loss as planting age  
30 increases. Between 2019 and 2022, the values of the average runoff depth for *DE*, *PE*, *PD* and *SDM*  
31 were 0.47, 0.55, 0.45 and 0.27 mm, respectively. Despite the increase in runoff, grassland restoration  
32 favored soil conservation: the net soil loss per unit area of *SDM* was 1.4, 1.3 and 1.9 times greater  
33 than that in *DE*, *PE* and *PD*, respectively. The key factors affecting soil loss and runoff were rainfall  
34 amount, duration and intensity (60-min intensity). We conclude that the results of this study can serve  
35 as scientific guides to design efficient policy decisions for planning the most effective vegetation  
36 restoration in the severely degraded hillside alpine meadows. To improve the effectiveness ~~boost the~~  
37 ~~benefits~~ of grassland restoration, we suggest that protective measures should be prioritized during the  
38 initial planting stage of cultivated grasslands.

39 **Keywords:** Alpine meadow; Degraded hillside; Mixed-cultivated grassland; land management; runoff;  
40 soil erosion.

## 41 **1 Introduction**

42 Grasslands are an essential component of terrestrial ecosystems and habitats for the development of  
43 animal livestock (O'Mara, 2012). They make significant contributions to biodiversity conservation,  
44 climate mitigation, carbon sequestration, and water supply and regulation (Bardgett et al., 2021).  
45 Despite the importance of grasslands, about half of them are degraded globally –5% of them  
46 undergoing severe degradation (based on net primary productivity)–, and this issue has become a  
47 major concern for landscape conservation (Gang et al., 2014; Török et al., 2021). Global grassland  
48 net primary productivity (NPP) has declined by 58.84 Tg C per year. Grassland degradation causes  
49 the loss of up to 90% of the soil structure that facilitates water movement (infiltration) and retention  
50 (water-holding capacity) in soils (Wick et al., 2016), reduces carbon storage potential (Liebig et al.,  
51 ~~2009~~2013), and impedes soil functioning. Moreover, degraded grasslands are prone to severe soil  
52 erosion, especially in mountainous areas. For example, in the Swiss alpine uplands, water erosion  
53 ranges from 0.14 to 1.25 t ha<sup>-1</sup> month<sup>-1</sup> according to the phenological stage of the grasses (Schmidt et  
54 al., 2019); and in the gully slope of the Loess Plateau, the average amount of soil erosion was 306.7  
55 t ha<sup>-1</sup> month<sup>-1</sup> during the 2018–2020 grass growing season (Zhu et al., 2021).

56 Precipitation is the main water source of soil moisture supply in semi-arid areas and the  
57 conversion of precipitation to runoff is one of the major contributors to river streamflow (Leung et al.  
58 2015; Li et al., 2024). In some previous experiences, it was observed that vegetation restoration  
59 reduced surface runoff and decreased sediment production, which led to lower river levels,  
60 threatening the health of river ecosystems (Dijk et al., 2007). A recent study conducted by Wu et al.  
61 (2020) proposed sustainable management strategies for semi-arid areas with a positive trade-off  
62 between surface runoff maintenance and erosion control. However, very few studies have addressed  
63 to date the effects of restored grasslands in maintaining surface runoff and preventing soil erosion

64 (Minea et al., 2022). This topic is particularly important for alpine grasslands, which play a vital role  
65 in the supply of fresh water and the development of livestock husbandry (Cui et al., 2022). Therefore,  
66 it is necessary to assess the impacts of grassland restoration on runoff generation and soil protection.

67 Vegetation restoration is widely considered as one of the most effective methods for controlling  
68 runoff and soil erosion worldwide (Anache et al., 2018; [Vanacker et al., 2022](#)). The effects of  
69 vegetation cover properties on runoff and soil loss reduction are strongly connected to plant species,  
70 leaf and branch coverage, above-ground biomass, litter biomass, and root systems (Liu et al., 2022;  
71 Freschet and Roumet, 2017; Gyssels et al., 2005; Zhu et al., 2021). Furthermore, the processes of  
72 runoff and soil loss are significantly influenced by the improvement of soil characteristics with  
73 vegetation restoration (Schwarz et al., 2015; Gyssels et al., 2005). The interaction between vegetation  
74 and soil could stabilize the topsoil and alter soil properties (Saxton and Rawls, 2006; Ma et al., 2023a).  
75 Vegetation restoration promotes the formation of soil aggregates, decreases soil bulk density,  
76 enhances organic matter and nutrients and improves soil porosity, resulting in high soil hydraulic  
77 conductivity and field capacity (Qiu et al., 2022; Saxton and Rawls, 2006). The above-interlinked soil  
78 properties alter soil hydrological properties and ultimately influence hillslope and watershed  
79 hydrology, such as runoff and soil erosion (Lu et al., 2020; Qiu et al., 2022). While vegetation  
80 restoration holds the potential to be a key method of environmental restoration under human  
81 management, the inappropriate selection of species can negatively impact the sustainability of local  
82 economic and environmental development (Huang et al., 2017; 2019). For example, cultivated  
83 grasslands were already advocated as a sensible solution for the conservation of soil and water, as  
84 well as the regrowth of vegetation in semi-arid mountain areas (Liu et al., 2022; Wu et al., 2010).  
85 Grass communities with multiple stratified structures are better at maintaining surface runoff and  
86 decreasing soil loss than those with a single composition and structure (Mohammad and Adam, 2010).

87 Surface runoff –also known as stormwater runoff or overland flow– reaches the stream in the form  
88 of sheet, rill and gully flow (Rumynin, 2015). The conversion of rainfall to overland flow depends on  
89 the rainfall intensity, the soil hydrological properties, such as (non-)saturated hydraulic conductivity,  
90 matrix flux potential and field capacity, and initial soil water content (López-Vicente and Navas, 2012;  
91 Gyssels et al., 2005; De Baets et al., 2007). Because runoff is the primary driver of water erosion on  
92 hillslopes and serves as the main agent for sediment transport, reducing the conversion of rainfall to  
93 runoff is regarded as an effective way to control water erosion, ~~such as~~ through vegetation restoration  
94 (Zhou et al., 2016; Zhu et al., 2021). On the other hand, in arid and semi-arid regions, surface runoff  
95 is the major water supply source to the river streamflow, thereby it is vital for ensuring the  
96 sustainability of ecosystems and human activities (Liu et al., 2020; Robinson et al., 2003). Therefore,  
97 restoration efforts in areas with low rainfall should be oriented to maintain runoff while reducing its  
98 level of sediment concentration.

99 Soil erosion can be reduced by various factors, including the above- and below-ground biomass  
100 of grasses, litter cover, and root systems (De Baets et al., 2007; [Durán Zuazo and Rodríguez](#)  
101 [Pleguezuelo, 2008; Gyssels and Poesen, 2003; Wen et al., 2024](#)). Grasslands can control water erosion  
102 relying on the role of the aboveground biomass in dissipating flow energy (Bochet and García-Fayos,  
103 2004), living roots in decreasing soil detachment capacity (Zhang et al., 2013), grass plant cover in  
104 intercepting rainfall (Liu et al., 2019), and litter cover in enhancing rainwater infiltration (Liu et al.,  
105 2022). Moreover, the interweaving of plant roots can remarkably alter the physical properties of the  
106 topsoil, enhancing its resistance to erosion (Schwarz et al., 2015; Wang et al., 2018). The impact of  
107 grassroots on the soil characteristics can be summarized as follows: i) increasing the stability of soil  
108 aggregates through aggregating fine soil particles into macroaggregates; ii) enhancing soil cohesion  
109 through interweaving with the soil; and iii) decreasing soil bulk density through increasing soil

110 porosity (Wu et al., 2019; Gyssels et al., 2005). For example, numerous recent studies have confirmed  
111 that a grass with shallow yet dense fibrous root system appears to be more effective at controlling  
112 water erosion than grass with good ground cover but low root density (De Baets et al., 2007; Bochet  
113 et al., 2006).

114 Alpine meadows, especially in the Qinghai-Tibetan Plateau, constitute the predominant  
115 ecosystem in China and the world, accounting for 44% and 6% of total grassland areas, respectively  
116 (Wang et al., 2016). Over 50% of the alpine meadows have been subject to an increasing degree of  
117 degradation (Bardgett et al., 2021), with the extent of degradation depending on the meadow patch  
118 coverage resulting from the fragmentation of alpine meadows (Fig. 1b). Severely degraded meadows  
119 (also known as “black beach” and “black soil-type degraded meadow”) formed after the mastic  
120 epipedon, typically 10 to 15 cm deep, was fully removed by overgrazing and rodent activity exposing  
121 the sub-soil (Fig. 1c; Ma et al., 2023a; Shang et al., 2008). Severely degraded meadows amounted to  
122 about 30% of the total area of alpine meadows on the Qinghai-Tibetan Plateau (Shang et al., 2008).  
123 Recent studies by Niu et al. (2021) and Ma et al. (2023b) have observed that fragmentation of alpine  
124 meadows and severely degraded meadows could reduce surface runoff and enhance soil erosion ~~in~~  
125 ~~alpine meadow hillslopes.~~

126 The Qinghai-Tibetan Plateau serves as the headwaters for many of Asia's major rivers (Xu, 2018).  
127 The eastern and southern parts of the Qinghai-Tibet Plateau are influenced by the monsoon, and  
128 rainfall is the primary source of streamflow (Cuo et al., 2014). The long-term and widespread  
129 degradation of hillside alpine meadows has disrupted the soil water balance, reducing surface runoff  
130 (Niu et al., 2019; Ma et al., 2023b). This, in turn, diminishes river streamflow, ultimately constraining  
131 the sustainable development of both local and downstream regions. The importance of artificial  
132 grassland in restoring alpine degraded meadow is widely accepted (Wen-Li et al., 2018; Wu et al.,

133 2010). Artificial grassland –also known as tamed grassland, sowed grassland and cultivated  
134 grassland– refers to fields that have been broken up and replanted with exotic grasses and forbs and  
135 utilized for hay crop production or cattle grazing (Fisher et al., 2018). The establishment of artificial  
136 grassland on severely degraded areas provides a dual benefit by boosting productivity and improving  
137 the ecological environment of alpine grasslands (Shang et al., 2008; Liu et al., 2022).

138 While previous studies have often focused on carbon sequestration capacity, vegetation  
139 characteristics, soil quality and productivity of cultivated grassland (Wang et al., 2013; Bai and  
140 Cotrufo et al., 2022), there has been a limited examination of the impacts of mixed-cultivated  
141 grasslands on the provision of runoff and prevention of soil erosion on the alpine hillsides. Recently,  
142 Liu et al. (2022) evaluated the effects of plant morphological characteristics on runoff and soil erosion  
143 in different mixed-cultivated grassland under natural rainfall events. Here, we present novel research  
144 to examine the ability of ~~alpine hillsides~~ cultivated grasslands to regulate runoff and soil loss and  
145 evaluate the effect of~~through~~ three different mixed-cultivated grasslands: *Deschampsia cespitosa* and  
146 *Elymus nutans* (DE), *Poa pratensis* L.cv. Qinghai and *Elymus nutans* (PE), and *Poa pratensis* L.cv.  
147 Qinghai and *Deschampsia cespitosa* (PD), compared to a severely degraded meadow (SDM) by a  
148 three-year field experiment. In particular, this study aimed to (1) assess the temporal variations in soil  
149 and water loss of DE, PE and PD grasslands during the growing season and under natural rainfall;  
150 and (2) determine the key factors influencing the mixed-cultivated grasslands in controlling runoff  
151 and soil erosion. This study has realistic implications for understanding the contribution of mixed-  
152 cultivated grasslands restoration on soil erosion control in degraded alpine hillsides.

153

## 154 2 Materials and methods

## 155 2.1 Study area

156 This study was carried out in the representative area of Zhique Village (33°40'01" N and 99°43'06" E,  
157 elevation over 4200 m a.s.l), Dari County, Qinghai province, which served as a field experimental  
158 site and model area for the restoration of severely degraded alpine meadow on the Qinghai-Tibetan  
159 Plateau (Fig. 1a). The climate conditions correspond to a typical highland one with low temperatures  
160 throughout the year, i.e., not showing the typical four-season pattern (spring, summer, autumn, winter),  
161 but rather just two main seasons: cold and warm. In the study region, the average annual temperature  
162 is -3.1 °C, with monthly variations from -14.7 °C in January to 7.5 °C in July (values corresponded  
163 to the period 1981-2018; data source: European Centre for Medium-Range Weather Forecasts). The  
164 average annual precipitation is 416 mm, with the majority of it falling from July to September, based  
165 on Climate Hazards Group InfraRed Precipitation with Station data (CHIRPS). The majority of the  
166 precipitation and the warm season falls during the vegetation growth period (from May to September),  
167 favoring optimal conditions for the development of plants. The soil type in the study area is classified  
168 as Mat Cryi-gelic Cambisols (IUSS-WRB, 2015). Currently, the remnant vegetation in this site is  
169 composed of an alpine shrub (*Salix cupularis* and *Potentilla fruticose*), alpine meadow (*Kobresia*  
170 *pygmaea*, *Kobresia humilis* and *Kobresia capillifoli*) and swamp meadow (*Carex atrofusca*, *Poa*  
171 *annua* and *Carex parva*).

172 Soil erosion in the degraded alpine meadows is severe, becoming the primary source of sediment  
173 delivered to streams in the study area (Liu et al., 2022). The mattic epipedon of alpine meadow has  
174 experienced fragmentation and even disappearance (Fig. 1b), eventually forming a severely degraded  
175 meadow (Fig. 1c). Before implementing the grassland restoration project, i.e., Subsidy and Incentive  
176 System for Grassland Conservation, the average soil erosion rate and the total erosion in the study  
177 area were 13.63 t ha<sup>-1</sup> y<sup>-1</sup> and 323.58 × 10<sup>6</sup> t y<sup>-1</sup>, respectively (Zhao et al., 2021). Severely degraded



178 meadows were restored via mixed-cultivated grasslands –fields were ploughed and replanted with  
179 two grass species– and moderately degraded meadows were restored by broadcast sowing on the  
180 hillslopes during the implementation of the grassland restoration project. The grass species used for  
181 the projects have excellent characteristics like strong trampling tolerance, good palatability, abundant  
182 leaf quantity and developed rhizome, such as *Poa pratensis L. cv. Qinghai*, *Deschampsia cespitosa*  
183 and *Elymus nutans* (Shang et al., 2008).

184

## 185 **2.2 Experimental design and measurement**

186 The degraded hillslopes are the main component of runoff generation and confluence areas on the  
187 Qinghai-Tibetan Plateau. Hence, the grass species chosen for mixed-cultivated grasslands not only  
188 must it be grazing-tolerant and good forage, but also prevent soil loss and maintain surface runoff.  
189 Potential grass species should also be fully acclimated to harsh alpine climate and have  
190 complementary morphological characteristics and living habits (Liu et al., 2022). The community  
191 established by matching of grasses morphological characteristics and habits has a hierarchical vertical  
192 cover structure and little inter- or intraspecific competition. Following the above-mentioned  
193 guidelines for choosing grass species, we ultimately decided on three species (*Deschampsia cespitosa*,  
194 *Poa pratensis L. cv. Qinghai* and *Elymus nutans*) from the most widely utilized grass species.  
195 *Deschampsia cespitosa* is a cool-season bunching grass native to alpine environments. It typically  
196 forms a low, dense tussock (to 30–50 cm tall) of very thin (0.5 cm wide), arching, flat to inrolled,  
197 dark green grass blades (to 5 cm long). *Deschampsia cespitosa*, a common bottom grass, has 70% of  
198 its grass stems growing between 0 and 30 cm tall. *Elymus nutans* is a common and important plant  
199 species in the alpine meadows of the Qinghai-Tibetan Plateau (Chen et al., 2009). It is a valuable  
200 fodder grass in alpine locations that has been extensively employed for animal production, disturbed

201 grassland restoration, and artificial grassland construction due to its resilience to cold, drought and  
202 pests (Ren et al., 2010). *Elymus nutans* is a herbaceous perennial species with sparsely tufted culms  
203 that can grow to heights of 70 to 100 cm (Liu et al., 2022). *Poa pratensis* L. cv. *Qinghai* is the common  
204 and dominant species native to the Qinghai-Tibetan Plateau. It is an excellent species that has been  
205 selected and cultivated to restore degraded alpine meadows. Also, *Poa pratensis* L. cv. *Qinghai* is an  
206 herbaceous perennial species with erect or geniculate base culms that grow 20–60 cm tall.

207 To reveal the effects of mixed-cultivated grasslands in controlling runoff and soil loss on hillsides,  
208 field observation of mixed grass plots designed by us was conducted from the 2019 to 2022 growing  
209 seasons. Therefore, one plot with severely degraded meadow (*SDM*) as a control and three plots with  
210 two mixed grass seeds per plot of *Deschampsia cespitosa* and *Elymus nutans* (*DE*), *Poa pratensis*  
211 L. cv. *Qinghai* and *Elymus nutans* (*PE*), and *Poa pratensis* L. cv. *Qinghai* and *Deschampsia cespitosa*  
212 (*PD*) were selected as the testing site (Fig. 1d). All four runoff plots were spaced 1m apart and were  
213 located on the same hillside with the same elevation and soil texture. All plots were bounded by steel  
214 plates (30 cm high and 2 mm thick sheet) and built during May 2019, with an area of 10 m<sup>2</sup> (2 m wide  
215 and 5 m long parallel to the maximum slope gradient). To collect only runoff and soil loss from the  
216 runoff plot, the steel plate was put vertically into the soil to a depth of about 10 cm, with the remainder  
217 sticking out from the soil surface. At the outlet of each plot, a steel runoff collection and calibrated  
218 tank (75 L) were set up to gather sediment and runoff. To prevent the collected runoff from being lost  
219 to evaporation, the calibrated tank was set inside a sealed vat (Fig. 1d).

220 In addition, the grass seeding for each runoff plot was completed in May 2019. For the runoff  
221 plots, grass seeds were distributed to a depth of less than 1 cm in strips at 20 cm intervals following  
222 plowing. The seeding rate was set at 6.0 g m<sup>-2</sup> for *Poa pratensis* L. cv. *Qinghai* and *Deschampsia*  
223 *cespitosa* and 4.5 g m<sup>-2</sup> for *Elymus nutans* to ensure a constant number of plants based on germination

224 and seedling emergence rates. None of the runoff plots experienced any human disturbance during  
225 the observation period (2019–2022), including grazing, harvesting, and excavation.

226

### 227 **2.3 Rainfall, runoff and soil loss measurement**

228 A Vantage pro 2<sup>TM</sup> weather station (Davis Instruments Corp., USA) with a measurement accuracy of  
229 4% was positioned next to the experimental plots to monitor precipitation intensity and duration (Fig.  
230 1). A precipitation event was defined by the occurrence of a no-rain interval lasting more than 3 h  
231 between them. A total of 42 precipitation events were recorded from 2019 to 2022 throughout the  
232 growing season. Snow was not collected, and only rainfall was recorded during the growing season  
233 (from June 15 to August 25). Precipitation characteristics of each event, including amount ( $P$ ),  
234 duration ( $RD$ ), and maximum intensities of 60 minutes ( $RI_{60}$ ) were recorded. Average rainfall intensity  
235 ( $ARI$ ) was calculated by dividing the total rainfall amount by the duration of the rainfall event. After  
236 each rainfall-runoff event, both runoff and sediment were collected right away. The water level in the  
237 calibrated tank was first measured to calculate the runoff volume. Then, runoff was fully mixed inside  
238 the calibrated tank using a stirring bar to thoroughly whirl, and two 500 ml bottles were used to obtain  
239 mixture samples of sediment and runoff. When the calibrated tank had less than 1000 ml of runoff  
240 sample, all runoff was collected. Lastly, the calibrated tank was cleaned in order to collect sediment  
241 and runoff for the subsequent rainfall-runoff event. The mixture samples in the bottle were transported  
242 back to the lab to be filtered on filter paper with a pore size of 30–50  $\mu\text{m}$ . The filter paper with  
243 sediment was oven-dried to a consistent weight at 105 °C. The ratio of soil loss amount to runoff  
244 volume in the mixed samples was applied to calculate the sediment concentration. Finally, runoff  
245 volume and sediment concentration were multiplied to calculate soil loss in each plot.

246 We collected runoff and soil erosion data during the growing season for the years 2019 to 2022.

247 Data for 2021 could not be collected due to the prevention and control strategies for coronavirus  
248 (COVID-19). Soil erosion and runoff were portrayed in this work by soil erosion per unit area ( $\text{g m}^{-2}$ )  
249 and runoff depth (mm). The runoff depth ( $R$ ) and soil erosion per unit area ( $S$ ) could be calculated  
250 using the following formulas:

$$R = \frac{V_R}{A} \times 10^3 \quad (1)$$

$$S = \frac{S_T}{A} \quad (2)$$

251 where  $V_R$  is the volume of runoff from runoff plots ( $\text{m}^3$ ),  $S_T$  is the total amount of soil erosion from  
252 runoff plots (g), and  $A$  is the area of runoff plot ( $\text{m}^2$ ).

253

## 254 **2.4 Vegetation and soil properties measurement**

255 Vegetation cover ( $VC$ ), including dead (standing litter) and living vegetation, was measured monthly  
256 from 2019 to 2022 growing seasons using a steel wire frame ( $50 \text{ cm} \times 50 \text{ cm}$ ) subdivided into 25  
257 plots of  $10 \text{ cm} \times 10 \text{ cm}$ . Fig. 2 exhibited the change in vegetation coverage for all runoff plots from  
258 2019 to 2022. After collecting runoff samples each year, the quadrats ( $50 \text{ cm} \times 50 \text{ cm}$ ) were positioned  
259 in the up-, mid-, and down-slope areas. Litter in each quadrant was collected and oven-dried to  
260 determine litter biomass ( $LB$ ) (Zhu et al., 2021). The litter collection for 2019 was not completed due  
261 to the seeding of mixed-cultivated grasslands in May 2019, and the litter collection for 2020 and 2021  
262 was collected at the end of the runoff collection for the current year. Undisturbed soil samples were  
263 taken in the 0–10 cm soil layers using steel rings in 2022. All soil samples were saturated and then  
264 weighed ( $W_{sat}$ ). Then the saturated soil samples were placed on the dry sand layer to drain water for  
265 about 2 h and 8 h, and weighed ( $W_{2h}$  and  $W_{8h}$ ). Finally, soil samples were dried in an oven at  $105 \text{ }^\circ\text{C}$   
266 for 24 h and then weighed ( $W_{dr}$ ). Based on the above measurement, soil bulk density ( $BD$ ,  $\text{g cm}^{-3}$ ),  
267 total porosity ( $TP$ , %), capillary porosity ( $CP$ , %), non-capillary porosity ( $NCP$ , %), and soil water

268 content at field capacity ( $FC$ , %) were determined as follows:

$$FC = \frac{(W_{8h} - W_{dr})}{(W_{dr} - W_{sr})} \quad (3)$$

$$BD = \frac{(W_{dr} - W_{sr})}{V} \quad (4)$$

$$TP = \left(1 - \frac{BD}{ds}\right) \times 100 \quad (5)$$

$$CP = \frac{(W_{2h} - W_{dr})}{V} \quad (6)$$

$$NCP = TP - CP \quad (7)$$

269 where  $W_{sr}$  is the weight of the ~~eu~~ steel ring (g),  $ds$  is the soil particle density (generally being 2.65  
270 g cm<sup>-3</sup>), and  $V$  is the volume of the ~~cutting~~ ring (100 cm<sup>3</sup>).

271 In addition, root mass density ( $RMD$ ) was obtained using a root drill, followed by washing with  
272 water and drying in the oven. Four undisturbed samples were collected in each quadrat using a steel  
273 ring (6.18 cm diameter and 2.0 cm height), and they were applied to a direct shear (ZJ type). The soil  
274 cohesion was obtained by the Mohr–Coulomb theory (Labuz and Zang, 2012).

275

## 276 **2.5 Calculating the reduction effect of runoff and soil loss**

277 Four metrics were employed to assess the efficiencies of the mixed-cultivated grasslands in regulating  
278 runoff and soil loss, which were: The runoff reduction benefit ( $RRB$ , %), sediment concentration  
279 reduction benefit ( $CRB$ , %), soil erosion reduction benefit ( $SRB$ , %), and the percentage of runoff  
280 reduction ratio to soil loss reduction ratio ( $RRSR$ ) (Zhao et al., 2014). High values of  $RRB$ ,  $SRB$  or  
281  $CRB$  indicated that vegetation was able to reduce runoff, soil erosion or sediment concentration  
282 compared to the rates observed in the control plot (severely degraded meadow). In addition, a low  
283  $RRSR$  implied that vegetation was more beneficial in minimizing soil erosion than in minimizing  
284 runoff (Liu et al., 2020). These indices were calculated as follows:

$$RRB = \frac{R_c - R_v}{R_c} \times 100 \quad (8)$$

$$SRB = \frac{S_c - S_v}{S_c} \times 100 \quad (9)$$

$$CRB = \frac{C_c - C_v}{C_c} \times 100 \quad (10)$$

$$RRSR = \frac{RRB}{SRB} \times 100 \quad (11)$$

285 where  $R_c$  and  $R_v$  are the runoff depths of the degraded meadow plot and plots covered by mixed-  
 286 cultivated grasslands;  $S_c$  and  $S_v$  are the soil loss per unit area of the degraded meadow plot and  
 287 plots covered by mixed-cultivated grasslands;  $C_c$  and  $C_v$  are the sediment concentrations of the  
 288 degraded meadow plot and plots covered by mixed-cultivated grasslands, respectively.

289

## 290 **2.6 Statistical analyses**

291 All data were analyzed using SPSS statistics software (IBM, USA, version 26.0). The Kolmogorov–  
 292 Smirnov test was used to test the normality of data. Duncan’s multiple range tests of one-way analysis  
 293 of variance (ANOVA) were applied to test for significant differences between soil and vegetation  
 294 characteristics, runoff depth, soil erosion amount, and runoff and soil loss reduction ratio under  
 295 various mixed-cultivated grasslands at 0.05 significance levels. Also, path analysis is a form of  
 296 multiple regression statistical analysis that is used to evaluate causal models by examining the  
 297 relationships between runoff, soil loss and soil and vegetation properties. By using this method, one  
 298 can identify the major factors influencing runoff and soil loss and determine the direct and indirect  
 299 effects of soil and vegetation properties on runoff and soil loss.

300

## 301 **3 Results**

### 302 **3.1 Mixed-cultivated grasslands modified runoff yield amount and soil loss**

303 Mixed-cultivated grasslands increased runoff and reduced soil erosion. One-way analysis of variance  
304 (ANOVA) revealed that runoff significantly ( $p < 0.05$ ) increased after the severely alpine degraded  
305 hillside was restored by the mixed-cultivated grassland (Fig. 3). During the three evaluated growing  
306 seasons (2019, 2020 and 2022), the average runoff depth for *DE*, *PE*, *PD* and *SDM* was 0.47, 0.55,  
307 0.45 and 0.27 mm, respectively. The average runoff depths of *SDM* in 2019, 2020, and 2022 were  
308 0.23, 0.34 and 0.25 mm, respectively, all significantly ( $p < 0.05$ ) lower than (except for 2020) the  
309 average runoff of mixed-cultivated grassland *DE*, *PE* and *PD*, which measured 0.44, 0.59 and 0.50  
310 mm in 2019, 0.55, 0.51 and 0.38 mm in 2020, 0.43, 0.54 and 0.40 mm in 2022 (Fig. 3a). Regarding  
311 soil conservation, the amount of soil loss in grasslands was significantly influenced by planting age.  
312 As depicted in Fig. 3b, soil loss in *DE*, *PE* and *PD* (except for *DE* in 2019) were significantly ( $p <$   
313  $0.05$ ) higher in 2019 and 2020 (the first and second years of planting) than those in the fourth year of  
314 planting (2022). In 2020, soil loss produced by *DE*, *PE*, and *PD* was significantly higher ( $p < 0.05$ )  
315 than that of *SDM*. Satisfactorily, the three mixed-cultivated grasslands did exhibit a clear reduction  
316 in soil loss compared to *SDM* in 2022 (albeit not significantly), with soil loss per unit area for *SDM*  
317 being 1.4, 1.3 and 1.9 times higher than those for *DE*, *PE* and *PD*, respectively. No significant  
318 difference ( $p > 0.05$ ) was observed in runoff depth and soil loss between *DE*, *PE* and *PD* in 2019,  
319 2020 and 2022. The results showed that any of the three mixed-cultivated grasslands (*DE*, *PE* and  
320 *PD*) could be effective in controlling soil loss and maintaining runoff.

321

### 322 **3.2 Specific runoff and soil loss reduction ratios of the cultivated grasslands**

323 Fig. 4 illustrates the runoff, soil loss and sediment concentration reduction ratio after planting various  
324 mixed-cultivated grasslands. Lower *RRB* values indicated a better ability to maintain runoff for  
325 mixed-cultivated grasslands, while higher *SRB* and *CRB* values indicated better effectiveness of

326 grasslands in soil loss reduction. The mean *RRB* values of the grass community *DE*, *PE* and *PD* were  
327 -79.3%, -130.4% and -48.5% in 2019, -36.9%, -53.5% and -21.5% in 2020, and -115.4%, -156.1%  
328 and -87.6% in 2022, respectively (Fig. 4a). Regardless of the combination of the above-mentioned  
329 grass species, the average increase ratio of runoff in 2022 (the fourth years of planting) was  
330 significantly ( $p < 0.05$ ) higher than that in 2019 and 2020 (the first and second years of planting). The  
331 *SRB* of the three mixed-cultivated grasslands (*DE*, *PE* and *PD*) increased with increasing planting  
332 age. It is worth noting that the average *SRB* values in the grassland communities of *DE*, *PE* and *PD*  
333 were 18.0%, 24.3% and 31.9% in 2022, respectively (Fig. 4b). The *SRE* values of *DE*, *PE* and *PD* in  
334 2022 were significantly ( $p < 0.05$ ) higher than those of 2019, whereas *SRE* values between 2020 and  
335 2022 was significant ( $p < 0.05$ ) for *DE* but not ( $p > 0.05$ ) for *PE* and *PD*. Additionally, *CRB* for all  
336 mixed-cultivated grasslands in 2022 was significantly ( $p < 0.05$ ) higher than that in 2019 and 2020.  
337 The mean *CRB* values of the cultivated-grassland communities *DE*, *PE* and *PD* increased from -120.9%  
338 to 55.8%, from -112.4% to 59.7%, and from -94.3% to 62.1% from 2019 to 2022, respectively (Fig.  
339 4c). Regardless of the age of the grasslands, the value of *RRSR* was less than 1, suggesting that the  
340 soil erosion reduction effect of the grasslands was higher than its runoff reduction effect (Fig. 4d). No  
341 significant differences ( $p > 0.05$ ) appeared in *RRB*, *SRB*, *CRB* and *RRSR* between *DE* *PE* and *PD* in  
342 2019, 2020 and 2022.

343

### 344 **3.3 Key factors affecting runoff and soil loss**

345 Precipitation characteristics and vegetation features played a significant role in influencing the  
346 hydrological response of the soil. In this study, path analysis was applied to identify the key factors  
347 affecting soil loss. The results of this analysis indicated that the sum of path coefficients of *RI<sub>60</sub>*, *RD*,  
348 *P* and *VC* were 0.31, 0.36, 0.40 and 0.32, respectively (Table 1). This suggests that *P*, *RD*, *VC* and



349  $RI_{60}$  had positive effects on runoff amount, with  $P$  being the most influential factor. Direct influences  
350 on runoff were primarily attributed to  $ARI$  and  $RD$ , with direct path coefficients of 0.37 and 0.67,  
351 respectively. Meanwhile, the influences of  $P$  and  $LB$  on runoff were mainly indirect, with indirect  
352 path coefficients of 0.57 and 0.25, respectively. For instance,  $P$ , in combination with other factors,  
353 particularly  $RI_{60}$  and  $RD$ , contributed significantly to runoff.

354 Soil loss was significantly influenced by  $R$ ,  $RI_{60}$ ,  $ARI$  and  $LB$ . The sum of path coefficients of  $R$ ,  
355  $RI_{60}$ ,  $ARI$  and  $LB$  were 0.52, 0.20, 0.28 and -0.25, respectively (Table 2). These results show that  $R$ ,  
356  $RI_{60}$  and  $ARI$  had a promotional effect, whereas  $LB$  had an inhibitory effect on soil loss. Meanwhile,  
357  $R$  and  $P$  had a direct positive influence on soil erosion, with direct path coefficients of 0.60 and 0.28,  
358 whereas  $RI_{60}$  and  $RD$  had a direct negative influence on soil erosion, with direct path coefficients of  
359 -0.29 and -0.41 (Table 2). In addition, the direct and indirect path coefficients both indicated that  $LB$   
360 had an inhibitory influence on the soil loss per unit area, with values of -0.10 and -0.25, respectively.

361

## 362 **4 Discussion**

### 363 **4.1 Benefits of mixed-cultivated grasslands on soil conservation and runoff maintenance**

364 The mixed-cultivated grasslands ( $DE$ ,  $PE$  and  $PD$ ) effectively maintained runoff and minimized soil  
365 loss (Fig. 4). This finding is similar to those of [some previous studies](#) ~~conducted checking different~~  
366 ~~grassland communities~~ (Liu et al., 2019; Liu et al., 2022). In this study, the mixed-cultivated  
367 grasslands significantly increased surface runoff compared to the  $SDM$ . The difference in runoff  
368 between mixed-cultivated grasslands and  $SDM$  may be attributed to the soil infiltration rate. Mixed-  
369 cultivated grasslands had more abundance of fibrous roots in the topsoil compared with  $SDM$  (Fig.  
370 5), and those fine roots reduced infiltration by occupying the soil pore (Leung et al., 2015). In

371 comparison to *SDM*, soil non-capillary porosity (*NCP*) and field capacity (*FC*) of *DE*, *PE* and *PD*  
372 significantly decreased by 46%, 32% and 48%, and increased by 55%, 59% and 48%, respectively  
373 (Fig. 5). This implied that *SDM* was restored to mixed-cultivated grasslands with lower permeability  
374 and better water retention. ~~This was further evidence that infiltration was responsible for the~~  
375 ~~difference in runoff between the mixed-cultivated grasslands and *SDM*.~~

376 Soil loss in all three mixed-cultivated grassland communities (*DE*, *PE* and *PD*) was higher than  
377 that in the *SDM* during the first- and second years following planting. However, by the fourth year,  
378 the *SDM* exhibited higher soil loss than the three mixed-cultivated grasslands (Fig. 3). These changes  
379 in soil erosion were dominantly attributed to the developing of the root system and improvement of  
380 soil structure (Zhu et al., 2021). The seeding method of plowing led to a disruption of soil structure,  
381 resulting in increased soil loss in the three mixed-cultivated grasslands during the initial stages of  
382 planting. ~~The loosening of the soil structure caused by the seeding method of plowing resulted in a~~  
383 ~~greater soil loss of the three mixed-cultivated grasslands than the *SDM* at the beginning of the planting.~~

384 We confirmed that the age of plantation was a key factor in understanding the inter-annual changes  
385 of soil erosion. This idea was also demonstrated in other types of primary land uses such as woody  
386 crops or young forests (Rodrigo-Comino et al., 2018). Nevertheless, we hypothesize that grassland  
387 topsoil demonstrated a stronger resilience to erosion as its root system grew, which had a  
388 reinforcement impact on the soil and led to lower soil loss in the fourth year of planting than that of  
389 the *SDM*. The topsoil (0–10 cm) of the grasslands had significantly different soil properties from the  
390 *SDM* in the fourth year after planting, as detailed in Table 3. In comparison to *SDM*, the root mass  
391 density and soil cohesion of grasslands *DE*, *PE* and *PD* increased by 672%, 890% and 589%, and by  
392 379%, 282% and 315%, respectively.

393

#### 394 **4.2 Effect of rainfall and vegetation characteristics on runoff and soil loss**

395 Surface runoff and erosion process is influenced and constrained by rainfall depth, intensity and  
396 duration, and by vegetation cover (*VC*) as well (Mohamadi and Kavian, 2015; Bochet et al., 2006).  
397 In this study, the *VC* had a directly promoted effect on surface runoff. Moreover, this result was in  
398 line with the finding of Niu et al. (2021), who reported that the surface runoff increased with the  
399 grassland coverage. Our results also indicated that rainfall amount (*P*) could have an indirect effect  
400 on surface runoff through rainfall duration (*RD*) and maximum intensities of 60 minutes (*RI<sub>60</sub>*). This  
401 implies that heavier and longer-lasting rainfall events were more likely to lead to surface runoff  
402 generation (Dos Santos et al., 2017). The findings demonstrated that runoff depth (*R*) and the average  
403 rainfall intensity (*ARI*) were the most and second most influential factors in promoting soil erosion  
404 (Table 2). The primary cause for this is that runoff velocity increases with higher precipitation  
405 intensity (Wang et al., 2013), which likely enhances the capacity of soil detachment and transport by  
406 surface runoff (Zhu et al., 2021). Furthermore, litter biomass (*LB*) had a direct and negative impact  
407 on soil loss (Table 2), indicating that the effectiveness of grasslands in reducing soil loss increased as  
408 litter biomass increased. Liu et al. (2022) found that the soil loss rate decreased with increasing litter  
409 biomass in grassland. Plant litter can intercept precipitation, reducing rainfall kinetic energy and  
410 splash erosion, while also increasing surface roughness (Liu et al., 2017; Xia et al., 2019) All these  
411 processes favor a reduction in runoff yield and soil loss rates.

412 Grass combinations with different ~~The matching of~~ morphological characteristics ~~of plants~~ can  
413 effectively reduce soil loss (Liu et al., 2022). In this study, the reduction in soil loss in the early stages  
414 of mixed-cultivated grassland planting (2019 and 2020) was attributed to grassland cover and plant  
415 morphological characteristics. *Deschampsia cespitosa*, *Poa pratensis* L.cv. Qinghai and *Elymus*  
416 *nutans* are dense clump type, rhizomatic-sparse clump type, and sparse clump perennial grasses,

417 respectively. In addition, *Deschampsia cespitosa* and *Poa pratensis* L.cv. Qinghai are bottom grasses,  
418 while *Elymus nutans* belongs to the top grass. The mix of dense and sparse grasses (*DE* and *PD*), and  
419 mix of top and bottom grasses (*DE* and *PE*) can complement each other morphologically and  
420 structurally, thereby more effectively reducing the kinetic energy of raindrops (Liu et al., 2022). *Poa*  
421 *pratensis* L.cv. Qinghai, a rhizomatic grass, also has abundant root systems intertwined with the soil,  
422 increasing soil cohesion and consequently reducing soil detachment capacity (Wang et al., 2018).  
423 Overall, in this study, the morphological and root characteristics of mixed-cultivated grasslands  
424 reduced runoff velocity, influenced water infiltration process and decreased soil erodibility.

425

### 426 **4.3 Implications for grasslands restoration on degraded alpine hillsides**

427 Our findings demonstrated that mixed-cultivated grasslands with complementing morphological  
428 features and habits can be effective at maintaining runoff and reducing soil erosion. Three mixed-  
429 cultivated grasslands (*DE*, *PE* and *PD*) exhibited an effective role in controlling soil loss on the  
430 degraded alpine hillside. However, at the start of planting, the mixed planted grassland had a greater  
431 soil erosion than the severely degraded meadow, whereas ~~the function of reducing~~ soil loss was  
432 ~~reached-reduced~~ in the 4<sup>th</sup> year of planting (Figs. 2 and 3). This suggested that protection measures,  
433 such as mesh covering and anti-trampling, may be taken into account to reduce soil loss in the initial  
434 planting stage of cultivated grassland in alpine hillsides (Liu et al., 2022). Moreover, grass may also  
435 be planted with a no-till system to avoid the initial increase of soil erosion at the initial phases of  
436 cultivated grassland by destroying soil structure (Karayel and Sarauskis, 2019). In addition, spring  
437 meltwater is the main driver of soil erosion in degraded alpine meadows in alpine regions, which  
438 greatly increases turbidity of rivers (Shi et al., 2020). The restoration of severely degraded hillslope  
439 meadows increased vegetation cover and soil ability, both of which could have an inhibitory impact

440 against meltwater erosion (Liu et al., 2022). To better understand the effects of cultivated grassland  
441 on meltwater erosion, future experiments under natural freezing and thawing conditions need to be  
442 conducted and monitored.

443 Cultivated grasslands, considered a crucial component of vegetation restoration, have been  
444 extensively utilized in the rehabilitation of degraded alpine hillsides (Shang et al., 2008). Nevertheless,  
445 plant restoration is not necessarily beneficial to the long-term viability of on- and off-site ecosystems'  
446 functions, including natural succession and river ecosystems. Therefore, the selected vegetation types  
447 ought to be advantageous for the ecosystem's sustainability, both on- and off-site, such as maintaining  
448 river streamflow and unrestricted natural succession. The seed prices of cultivated grass communities  
449 of *Deschampsia cespitosa* and *Elymus nutans*, *Poa pratensis* L.cv. Qinghai and *Elymus nutans*, and  
450 *Poa pratensis* L.cv. Qinghai and *Deschampsia cespitosa* were about \$690, \$750 and \$480 per ha,  
451 respectively, in Xining, Qinghai Province, in 2019. Planting properly mixed-cultivated grassland on  
452 the alpine degraded hillsides can achieve both environmental and economic benefits. This study  
453 proved that mixed-cultivated grasslands could maintain runoff and decrease soil loss.

## 454 **5 Conclusions**

455 Based on the measured data during the 2019, 2020 and 2022 growing seasons, the planting of mixed-  
456 cultivated grassland on severely degraded hillside alpine meadow effectively maintained surface  
457 runoff and decreased soil loss, especially after the mixed-cultivated grassland played a positive role  
458 in consolidating the soil surface. The benefits were statistically significant compared with the control  
459 plot, but differences between the three types of cultivated grasslands were not significant. Planting  
460 the mixed-cultivated grasslands after ploughing loosened the soil structure and thus increased  
461 sediment concentration in runoff during the first stage after planting. Subsequently, sediment

462 concentration decreased with the growth of the root system of the mixed-cultivated grasslands,  
463 improving root-soil cohesion due to the root architecture. To guarantee that mixed-cultivated  
464 grasslands can perform the aforementioned functions, protective measures should be implemented  
465 during the initial planting stage to support their healthy growth. Our results also suggested that mixed-  
466 cultivated grasslands with different but complementary morphology and structure and abundant fine  
467 root systems were effective in maintaining surface runoff and reducing soil erosion. Precipitation  
468 amount, duration, maximum 60-minute intensity, vegetation coverage and composition were the  
469 predominant factors affecting surface runoff and soil loss. The erosion resistance contribution of the  
470 above-ground community characteristics and below-ground roots along the cultivated time could  
471 maintain a relatively high surface runoff and decrease sediment concentration. These findings have  
472 potential implications for understanding the contribution of mixed-cultivated grasslands restoration  
473 on soil erosion control in the degraded hillsides of alpine areas.

474

475 *Data availability.* The data that support the findings of this study are available on request from the  
476 corresponding author.

477

478 *Author contributions.* Yulei Ma: Investigation, Formal analysis, Methodology, Software, Writing -  
479 original draft. Yifan Liu: Investigation, Formal analysis, Writing - review & editing. Jesús Rodrigo-  
480 Comino: Interpretation of data, Writing - review & editing. Manuel López-Vicente: Interpretation of  
481 data, Writing - review & editing. Gao-Lin Wu: Conceptualization, Funding acquisition, Supervision,  
482 Writing - original draft, review & editing.

483

484 *Competing interests.* The authors declare that they have no known competing financial interests or

485 personal relationships that could have appeared to influence the work reported in this paper.

486

487 *Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional  
488 claims in published maps and institutional affiliations

489 *Acknowledgments.* We thank [Nunzio Romano](#), Gall Corinna, Vanacker Veerle and Qianjin Liu for  
490 their constructive comments and suggestions on this manuscript.

491 *Financial support.* This research was funded by the National Natural Science Foundation of China  
492 (NSFC41930755, NSFC32230068), the Strategic Priority Research Program of the Chinese Academy  
493 of Sciences (XDB40000000), and the Second Stage's Research and Technique Extending Project of  
494 Sanjiangyuan Ecological Protection and Building in Qinghai (2019-S-1).

495

## 496 **References**

497 Anache, J.A.A., Flanagan, D.C., Srivastava, A., and Wendland, E.C.: Land use and climate change  
498 impacts on runoff and soil erosion at the hillslope scale in the Brazilian Cerrado, *Sci. Total*  
499 *Environ.*, 622–623, 140–151, <https://doi.org/10.1016/j.scitotenv.2017.11.257>, 2018.

500 Bai, Y., Cotrufo, M.F.: Grassland soil carbon sequestration: Current understanding, challenges, and  
501 solutions, *Science*, 377(6606), 603–608. <https://doi.org/10.1126/science.abo2380>, 2022.

502 Bardgett, R.D., Bullock, J.M., Lavorel, S., Manning, P., Schaffner, U., Ostle, N., Chomel, M.,  
503 Durigan, G., Fry, E.L., Johnson, D., Lavallee, J.M., Le Provost, G., Luo, S., Png, K., Sankaran,  
504 M., Hou, X.Y., Zhou, H.K., Ma, L., Ren, W.B., Li, X.L., Ding, Y., Li, Y.H., and Shi, H.X.:

505        Combating global grassland degradation, *Nat. Rev. Earth Env.*, 2(10), 720–735,  
506        <https://doi.org/10.1038/s43017-021-00207-2>, 2021.

507    Bochet, E., and García-Fayos, P.: Factors controlling vegetation establishment and water erosion on  
508        motorway slopes in Valencia, Spain, *Restor. Ecol.*, 12(2), 166–174, [https://doi.org/](https://doi.org/10.1111/j.1061-2971.2004.0325.x)  
509        10.1111/j.1061-2971.2004.0325.x, 2004.

510    Bochet, E., Poesen, J., and Rubio, J.L.: Runoff and soil loss under individual plants of a semi-arid  
511        Mediterranean shrubland: influence of plant morphology and rainfall intensity, *Earth Surf. Proc.*  
512        *Land.*, 31(5), 536–549, <https://doi.org/10.1002/esp.1351>, 2006.

513    Chen, S., Ma, X., Zhang, X., and Chen, Z.: Genetic variation and geographical divergence in *Elymus*  
514        *nutans* Griseb. (Poaceae: Triticeae) from West China, *Biochem. Syst. Ecol.*, 37(6), 716–722,  
515        <https://doi.org/10.1016/j.bse.2009.12.005>, 2009.

516    Cui, Z., Liu, Y.F., Liu, Y., Leite, P.A.M., Shi, J.J., Shi, Z.H., and Wu, G.L.: Fragmentation alters the  
517        soil water conservation capacity of hillside alpine meadows on the Qinghai-Tibetan Plateau,  
518        *Geoderma*, 428, 116133, <https://doi.org/10.1016/j.geoderma.2022.116133>, 2022.

519    Cuo, L., Zhang, Y.X., Zhu, F.X., and Liang, L.Q.: Characteristics and changes of streamflow on the  
520        Tibetan Plateau: A review, *J. Hydrol-Reg. Stud.*, 2, 2014, 49–68,  
521        <https://doi.org/10.1016/j.ejrh.2014.08.004>, 2014

522    De Baets, S., Poesen, J., Knapen, A., Barberá, G.G., and Navarro, J.A.: Root characteristics of  
523        representative Mediterranean plant species and their erosion-reducing potential during  
524        concentrated runoff, *Plant Soil*, 294(1–2), 169–183, [https://doi.org/10.1007/s11104-007-9244-](https://doi.org/10.1007/s11104-007-9244-2)  
525        2, 2007.

526    Dijk, A. I. J. M., and Keenan, R. J: Planted forests and water in perspective, *Forest Ecol. Manag.*,  
527        251, 1–9, <https://doi.org/10.1016/j.foreco.2007.06.010>, 2007.



528 Dos Santos, J.C.N., de Andrade, E.M., Medeiros, P.H.A., Guerreiro, M.J.S., and de Queiroz Palácio,  
529 H.A.: Effect of Rainfall Characteristics on Runoff and Water Erosion for Different Land Uses  
530 in a Tropical Semiarid Region, *Water Resour. Manag.*, 31(1), 173–185, [https://doi.org/](https://doi.org/10.1007/s11269-016-1517-1)  
531 10.1007/s11269-016-1517-1, 2017.

532 [Durán Zuazo, V.H., and Rodríguez Pleguezuelo, C.R.: Soil-erosion and runoff prevention by plant](#)  
533 [covers. A review. \*Agron. Sustain. Dev.\*, 28\(1\), 65–86, <https://doi.org/10.1051/agro:2007062>,](#)  
534 [2008.](#)

535 Fisher, R.J., Sawa, B., and Prieto, B.: A novel technique using LiDAR to identify native-dominated  
536 and tame-dominated grasslands in Canada, *Remote Sens. Environ.*, 218, 201–206,  
537 <https://doi.org/10.1016/j.rse.2018.10.003>, 2018.

538 Freschet, G.T., and Roumet, C.: Sampling roots to capture plant and soil functions, *Funct. Ecol.*, 31(8),  
539 1506–1518, <https://doi.org/10.1111/1365-2435.12883>, 2017.

540 Gang, C.C., Zhou, W., Chen, Y.Z., Wang, Z.Q., Sun, Z.G., Li, J.L., Qi, J.G., and Odeh, I.:  
541 Quantitative assessment of the contributions of climate change and human activities on global  
542 grassland degradation, *Environ. Earth Sci.*, 72(11), 4273–4282, [https://doi.org/10.1007/s12665-](https://doi.org/10.1007/s12665-014-3322-6)  
543 014-3322-6, 2014.

544 [Gyssels, G. and Poesen, J.: The importance of plant root characteristics in controlling concentrated](#)  
545 [flow erosion rates, \*Earth Surf. Proc. Land.\*, 28, 371–384. <https://doi.org/10.1002/esp.447>, 2003.](#)

546 Gyssels, G., Poesen, J., Bochet, E., and Li, Y., Impact of plant roots on the resistance of soils to  
547 erosion by water: a review, *Progr. Phys. Geogr.*, 29(2), 189–217, [https://doi.org/10.1191/](https://doi.org/10.1191/0309133305pp443ra)  
548 0309133305pp443ra, 2005.

549 Huang, Z., Tian, F.P., Wu, G.L., Liu, Y., and Dang, Z.Q.: Legume grasslands promote precipitation  
550 infiltration better than gramineous grasslands in arid regions, *Land Degrad. Dev.*, 28(1), 309–

551 316, <https://doi.org/10.1002/ldr.2635>, 2017.

552 Huang, Z., Liu, Y.F., Cui, Z., Liu, Y., Wang, D., Tian, F.P., and Wu, G.L.: Natural grasslands  
553 maintain soil water sustainability better than planted grasslands in arid areas, *Agr. Ecosyst.*  
554 *Environ.*, 286(1), 106683, <https://doi.org/10.1016/j.agee.2019.106683>, 2019.

555 Karayel, D., and Šarauskis, E.: Environmental impact of no-tillage farming, *Environ. Res. Eng.*  
556 *Manag.*, 75(1), 7–12, <http://dx.doi.org/10.5755/j01.arem.75.1.20861>, 2019.

557 Labuz, J.F., and Zang, A. Mohr–Coulomb Failure Criterion, *Rock Mech. Rock Eng.*, 45, 975–979,  
558 <https://doi.org/10.1007/s00603-012-0281-7>, 2012.

559 Leung, A.K., Garg, A., Coe, J.L., Ng, C.W.W., and Hau, B.C.H.: Effects of the roots of *Cynodon*  
560 *dactylon* and *Schefflera heptaphylla* on water infiltration rate and soil hydraulic conductivity,  
561 *Hydrol. Process.*, 29(15), 3342–3354. <https://doi.org/10.1002/hyp.1045>, 2015.

562 [Liebig, M.A., Kronberg, S.L., Hendrickson, J.R., Dong, X., and Gross, J.R.: Carbon dioxide efflux](#)  
563 [from long-term grazing management systems in a semiarid region, \*Agr. Ecosyst. Environ.\*, 164,](#)  
564 [137–144, <https://doi.org/10.1016/j.agee.2012.09.015>, 2013.](#)

565 Li, J., Wu, J., Yu, J., Wang, K., Li, J., Cui, Y., Shangguan Z.P., and Deng, L.: Soil enzyme activity  
566 and stoichiometry in response to precipitation changes in terrestrial ecosystems. *Soil Biol.*  
567 *Biochem.*, 191, 109321, <https://doi.org/10.1016/j.soilbio.2024.109321>, 2024.

568 Liu, W.J., Luo, Q.P., Lu, H.J., Wu, J.E., and Duan, W.P.: The effect of litter layer on controlling  
569 surface runoff and erosion in rubber plantations on tropical mountain slopes, SW China, *Catena*,  
570 149, 167–175, <https://doi.org/10.1016/j.catena.2016.09.013>, 2017.

571 Liu, Y., Li, S.Y., Niu, Y.L., Cui, Z., Zhang, Z.C., Wang, Y.L., Ma, Y.S., L'opez-Vicente, M., and  
572 Wu, G.L.: Effectiveness of mixed cultivated grasslands to reduce sediment concentration in  
573 runoff on hillslopes in the Qinghai-Tibetan Plateau, *Geoderma*, 422, 115933, <https://doi.org/>

574 10.1016/j.geoderma.2022.115933, 2022.

575 Liu, Y.F., Dunkerley, D., López-Vicente, M., Shi, Z.H., and Wu, G.L.: Trade-off between surface  
576 runoff and soil erosion during the implementation of ecological restoration programs in semiarid  
577 regions: A meta-analysis, *Sci. Total Environ.*, 712, 136477, [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.scitotenv.2019.136477)  
578 [scitotenv.2019.136477](https://doi.org/10.1016/j.scitotenv.2019.136477), 2020.

579 Liu, Y.F., Liu, Y., Wu, G.L., and Shi, Z.H.: Runoff maintenance and sediment reduction of different  
580 grasslands based on simulated rainfall experiments., *J. Hydrol.*, 572, 329–335, [https://doi.org/10.](https://doi.org/10.1016/j.jhydrol.2019.03.008)  
581 [1016/j.jhydrol.2019.03.008](https://doi.org/10.1016/j.jhydrol.2019.03.008), 2019.

582 Liu, Y., Zhao, L.R., Liu, Y.F., Huang, Z., Shi, J.J., Wang, Y.L., Ma, Y.S., Lucas-Borja, M.E., López-  
583 Vicente, M., and Wu, G.L.: Restoration of a hillslope grassland with an ecological grass species  
584 (*Elymus tangutorum*) favors rainfall interception and water infiltration and reduces soil loss on  
585 the Qinghai-Tibetan Plateau, *Catena*, 219, 106632, [https://doi.org/10.1016/j.catena.2022.](https://doi.org/10.1016/j.catena.2022.106632)  
586 [106632](https://doi.org/10.1016/j.catena.2022.106632), 2022.

587 [Li, W., Wang, J.L., Zhang, XJ., Shi, SL., Cao, W. X.: Effect of degradation and rebuilding of artificial](#)  
588 [grasslands on soil respiration and carbon and nitrogen pools on an alpine meadow of the](#)  
589 [Qinghai-Tibetan Plateau, \*Ecol. Eng.\*, 111, 134–142,](#)  
590 <https://doi.org/10.1016/j.ecoleng.2017.10.013>, 2018.

591 López-Vicente, M., and Navas, A.: A new distributed rainfall-runoff (DR2) model based on soil  
592 saturation and runoff cumulative processes, *Agricultural Water Management*, 104, 128–141.  
593 <https://doi.org/10.1016/j.agwat.2011.12.007>, 2012.

594 Lu, J.R., Zhang, Q., Werner, A.D., Li, Y.L., Jiang, S.Y., and Ta, Z.Q.: Root-induced changes of soil  
595 hydraulic properties – a review, *J. Hydrol.*, 589, 125203 [https://doi.](https://doi.org/10.1016/j.jhydrol.2020.125203)  
596 [org/10.1016/j.jhydrol.2020.125203](https://doi.org/10.1016/j.jhydrol.2020.125203), 2020.

- 597 Ma, Y.L., Liu, Y.F., López-Vicente, M., and Wu G.L.: Divergent shift of normal alpine meadow  
598 towards shrub and degraded meadows reduces soil water retention and storage capacity, *J.*  
599 *Hydrol.*, 625, 130109, <https://doi.org/10.1016/j.jhydrol.2023.130109>, 2023a.
- 600 Ma, Y.L., Liu, Y.F., López-Vicente, M., and Wu, G.L.: Divergent shift of normal alpine meadow  
601 exacerbated soil loss of hillslope alpine meadows based on field experiments, *Int. Soil Water*  
602 *Conserv. Res.*, In Press, <https://doi.org/10.1016/j.iswcr.2023.11.007>, 2023b.
- 603 Minea, G., Mititelu-Ionuș, O., Gyasi-Agyei, Y., Ciobotaru, N., and Rodrigo-Comino, J.: Impacts of  
604 grazing by small ruminants on hillslope hydrological processes: A review of European current  
605 understanding, *Water Resour. Res.*, 58, e2021WR030716, <https://doi.org/10.1029/2021WR>,  
606 2022.
- 607 Mohamadi, M.A., and Kavian, A.: Effects of rainfall patterns on runoff and soil erosion in field plots,  
608 *Int. Soil Water Conse.*, 3(4), 273–281, <http://dx.doi.org/10.1016/j.iswcr.2015.10.001>, 2015.
- 609 Mohammad, A.G., and Adam, M.A.: The impact of vegetative cover type on runoff and soil erosion  
610 under different land uses, *Catena*, 81(2), 97–103, <http://dx.doi.org/10.1016/j.catena.2010.01.008>,  
611 2010.
- 612 Niu, Y.L., Li, S.Y., Liu, Y., Shi, J.J., Wang, Y.L., Ma, Y.S., and Wu, G.L.: Regulation of alpine  
613 meadow patch coverage on runoff and sediment under natural rainfall on the eastern Qinghai-  
614 Tibetan Plateau, *J. Hydrol.*, 603, 127101, <https://doi.org/10.1016/j.jhydrol.2021.127101>, 2021.
- 615 O'Mara, F.P.: The role of grasslands in food security and climate change, *Ann. Bot-London* 110(6),  
616 1263–1270, <https://doi.org/10.1093/aob/mcs209>, 2012.
- 617 Qiu, D.X., Xu, R.R., Wu, C.X., Mu, X.M., Zhao, G.J., and Gao P.: Vegetation restoration improves  
618 soil hydrological properties by regulating soil physicochemical properties in the Loess Plateau,  
619 China, *J. Hydrol.*, 609, 127730, <https://doi.org/10.1016/j.jhydrol.2022.127730>, 2022.

620 Ren, F., Zhou, H.K., Zhao, X.Q., Han, F., Shi, L.N., Duan, J.C., and Zhao, J.Z.: Influence of simulated  
621 warming using OTC on physiological–biochemical characteristics of *Elymus nutans* in alpine  
622 meadow on Qinghai-Tibetan plateau, *Acta Ecol. Sinica*, 30(3), 166–171, <https://doi.org/10.1016/j.chnaes.2010.04.007>, 2010.

624 Robinson, M., Cognard-Plancq, A.L., Cosandey, C., David, J., Durand, P., Führer, H.W., Hall, R.,  
625 Hendriques, M.O., Marc, V., McCarthy, R., McDonnell, M., Martin, C., Nisbet, T., O’Dea, P.,  
626 Rodgers, M., and Zollner, A.: Studies of the impact of forests on peak flows and baseflows: a  
627 European perspective, *For. Ecol. Manag.*, 186, 85–97. [https://doi.org/10.1016/s0378-1127\(03\)00238-x](https://doi.org/10.1016/s0378-1127(03)00238-x), 2003.

629 Rodrigo-Comino, J., Brevik, E.C., and Cerdà, A.: The age of vines as a controlling factor of soil  
630 erosion processes in mediterranean vineyards, *Sci. Total Environ.*, 616–617, 1163–1173,  
631 <https://doi.org/10.1016/j.scitotenv.2017.10.204>, 2018.

632 Rumynin, V.G.: *Surface Runoff Generation, Vertical Infiltration and Subsurface Lateral Flow*. In:  
633 Overland flow dynamics and solute transport. Theory and Applications of Transport in Porous  
634 Media, vol 26. Springer, Cham, [https://doi.org/10.1007/978-3-319-21801-4\\_1](https://doi.org/10.1007/978-3-319-21801-4_1), 2015.

635 Saxton, K.E., and Rawls, W.J.: Soil water characteristic estimates by texture and organic matter for  
636 hydrologic solutions, *Soil Sci. Soc. Am. J.*, 70(5), 1569–1578,  
637 <https://doi.org/10.2136/sssaj2005.0117>, 2006.

638 Schwarz, M., Rist, A., Cohen, D., Giadrossich, F., Egorov, P., Büttner, D., Stolz, M., and Thormann,  
639 J.J.: Root reinforcement of soils under compression, *J. Geophys. Res-Earth*, 120(10), 2103–2120,  
640 <https://doi.org/10.1002/2015JF003632>, 2015.

641 Shang, Z.H., Ma, Y.S., Long, R.J., and Ding, L.M.: Effect of fencing, artificial seeding and  
642 abandonment on vegetation composition and dynamics of ‘black soil land’ in the headwaters of

- 643 the Yangtze and the Yellow Rivers of the Qinghai-Tibetan Plateau, *Land Degrad. Dev.*, 19(5),  
644 554–563, <https://doi.org/10.1002/ldr.861>, 2008.
- 645 Schmidt, S., Alewell, C., and Meusburger, K.: Monthly RUSLE soil erosion risk of Swiss grasslands, *J. Maps*, 15, 247–256, <https://doi.org/10.1080/17445647.2019.1585980>, 2019
- 646
- 647 Török, P., Brudvig, L.A., Kollmann, J., Price, J., and Tóthmérész, B.: The present and future of  
648 grassland restoration, *Restor. Ecol.*, 29(S1), e13378, <https://doi.org/10.1111/rec.13378>, 2021.
- 649 Wick, A.F., Geaumont, B.A., Sedivec, K.K., and Hendrickson, J.: *Grassland degradation*. In:  
650 Shroder, J.F., Sivanpillai, R. (Eds.), *Biological and environmental hazards, risks, and disasters*.  
651 Elsevier, pp. 257–276, 10.1016/B978-0-12-394847-2.00016-4, 2016.
- 652 Wang, B., Zhang, G., Yang, Y., Li, P., and Liu, J.: Response of soil detachment capacity to plant root  
653 and soil properties in typical grasslands on the Loess Plateau, *Agr. Ecosyst. Environ.*, 266, 68–  
654 75, <https://doi.org/10.1016/j.agee.2018.07.016>, 2018.
- 655 Wang, C.T., Wang, G.X., Liu, W., Wang, Y., Hu, L., and Ma, L.: Effects of establishing an artificial  
656 grassland on vegetation characteristics and soil quality in a degraded meadow, *Isr. Ecol. Evol.*,  
657 59(3), 141–153, <http://dx.doi.org/10.1080/15659801.2013.863669>, 2013.
- 658 Wang, L., Liang, T., and Zhang, Q.: Laboratory experiments of phosphorus loss with surface runoff  
659 during simulated rainfall, *Environ. Earth Sci.*, 70(6), 2839–2846, <http://dx.doi.org/10.1007/s12665-013-2344-9>, 2013.
- 660
- 661 Wang, Z.Q., Zhang, Y.Z., Yang, Y., Zhou, W., Gang, C.C., Zhang, Y., Li, J.L., An, R., Wang, K.,  
662 Odeh, I., and Qi, J.G.: Quantitative assess the driving forces on the grassland degradation in the  
663 Qinghai–Tibet Plateau, in China, *Ecol. Inform.*, 33, 32–44, [http://dx.doi.org/10.1016/j.ecoinf.](http://dx.doi.org/10.1016/j.ecoinf.2016.03.006)  
664 2016.03.006, 2016.
- 665 Wen, Y.R., Liu, B., Lin, L.T., Hu, M.M., Wen, X., Li, T.Y., Rong, J.D., and Yao, S.H.: Shelterbelt

- 666 [effects on soil redistribution on an arable slope by wind and water, \*Catena\*, 241, 108044.](#)
- 667 <https://doi.org/10.1016/j.catena.2024.108044>, 2024.
- 668 Wu, G. L., Huang, Z., Liu, Y.F., Cui, Z., Chang, X.F., Tian, F.P., López-Vicented, M., and Shi, Z.H.:
- 669 Soil water response of plant functional groups along an artificial legume grassland succession
- 670 under semi-arid conditions, *Agr. Forest Meteorol.*, 278, 107670.
- 671 <https://doi.org/10.1016/j.agrformet.2019.107670>, 2019.
- 672 Wu, G.L., Liu, Y.F., Cui, Z., Liu, Y., Shi, Z.H., Yin, R., and Kardol, P.: Trade-off between vegetation
- 673 type, soil erosion control and surface water in global semi-arid regions: A meta-analysis, *J. Appl.*
- 674 *Ecol.*, 57, 875–885. <https://doi.org/10.1111/1365-2664.13597>, 2020.
- 675 Wu, G.L., Liu, Z.H., Zhang, L., Hu, T., and Chen, J.: Effects of artificial grassland establishment on
- 676 soil nutrients and carbon properties in a black-soil-type degraded grassland, *Plant Soil*, 333(1–
- 677 2), 469–479, <https://doi.org/10.1007/s11104-010-0363-9>, 2010.
- 678 [Vanacker, V., Molina, A., Rosas, M. A., Bonnesoeur, V., Román-Dañobeytia, F., Ochoa-Tocachi, B.](#)
- 679 [F., and Buytaert, W.: The effect of natural infrastructure on water erosion mitigation in the Andes,](#)
- 680 [Soil](#), 8(1), 133–147, <https://doi.org/10.5194/soil-8-133-2022>, 2022.
- 681 Xia, L., Song, X.Y., Fu, N., Cui, S.Y., Li, L.J., Li, H.Y., and Li, Y.L.: Effects of forest litter cover on
- 682 hydrological response of hillslopes in the Loess Plateau of China, *Catena*, 181, 104076,
- 683 <https://doi.org/10.1016/j.catena.2019.104076>, 2019.
- 684 Xu, J.: A cave  $\delta^{18}\text{O}$  based 1800-year reconstruction of sediment load and streamflow: The Yellow
- 685 River source area, *Catena*, 161, 137–147, <http://dx.doi.org/10.1016/j.catena.2017.09.028>, 2018.
- 686 Zhang, G.H., Tang, K.M., Ren, Z.P., and Zhang, X.C.: Impact of grass root mass density on soil
- 687 detachment capacity by concentrated flow on steep slopes, *T. ASABE*, (56), 927–934, 2013.
- 688 Zhao, X., Huang, J., Wu, P., Gao, X.: The dynamic effects of pastures and crop on runoff and

689 sediments reduction at loess slopes under simulated rainfall conditions. *Catena*, 119, 1–7,  
690 <https://doi.org/10.1016/j.catena.2014.03.001>, 2014.

691 Zhao, Y.T., Pu, Y.F., Lin, H.L., and Tang, R.: Examining soil erosion responses to grassland  
692 conversation policy in Three-River Headwaters, China, *Sustainability*, 13(5), 2702,  
693 <https://doi.org/10.3390/su13052702>, 2021.

694 Zhou, J., Fu, B.J., Gao, G.Y., Lü, Y.H., Liu, Y., Lü, N., and Wang, S.: Effects of precipitation and  
695 restoration vegetation on soil erosion in a semi-arid environment in the Loess Plateau, China,  
696 *Catena*, 137, 1–11, <https://doi.org/10.1016/j.catena.2015.08.015>, 2016.

697 Zhu, P.Z., Zhang, G.H., Wang, H.X., Yang, H.Y., Zhang, B.J., and Wang, L.L.: Effectiveness of  
698 typical plant communities in controlling runoff and soil erosion on steep gully slopes on the  
699 Loess Plateau of China, *J. Hydrol.*, 602, 126714. <https://doi.org/10.1016/j.jhydrol.2021.126714>,  
700 2021.



701 **Table 1.** Results of path analysis of the factors affecting runoff depth.

Influence factor	Direct path coefficient	Indirect path coefficient						Sum of path coefficient	
		<i>RI</i> <sub>60</sub>	<i>ARI</i>	<i>RD</i>	<i>P</i>	<i>VC</i>	<i>LB</i>		Total
<i>RI</i> <sub>60</sub>	0.24*		0.25	-0.09	-0.11	0.02	0.00	0.07	0.31
<i>ARI</i>	0.37**	0.16		-0.34	-0.05	0.02	0.02	-0.19	0.18
<i>RD</i>	0.67**	-0.03	-0.18		-0.08	0.03	-0.03	-0.31	0.36
<i>P</i>	-0.18**	0.14	0.10	0.31		0.02	0.00	0.57	0.40
<i>VC</i>	0.29**	0.01	0.03	0.06	-0.01		-0.06	0.03	0.32
<i>LB</i>	-0.12	0.01	-0.09	0.18	0.00	0.15		0.25	0.13


702 Note: *RI*<sub>60</sub> is maximum 60-minute intensity (mm h<sup>-1</sup>), *ARI* is average rainfall intensity (mm h<sup>-1</sup>), *RD*  
703 is rainfall duration (h), *P* is rainfall amount (mm), *VC* is vegetation coverage (%), *LB* is litter biomass  
704 (g m<sup>-2</sup>). \* means the correlation is significant at 0.05 significance level, and \*\* means the correlation  
705 is significant at 0.01 significance level.

706


707 **Table 2.** Results of path analysis of the factors affecting soil loss per unit area.

Influence factor	Direct path coefficient	Indirect path coefficient								Sum of path coefficient
		<i>R</i>	<i>RI</i> <sub>60</sub>	<i>ARI</i>	<i>RD</i>	<i>P</i>	<i>VC</i>	<i>LB</i>	Total	
<i>R</i>	0.60**		-0.12	0.01	-0.10	0.11	0.01	0.01	-0.08	0.52
<i>RI</i> <sub>60</sub>	-0.29**	0.24		0.02	0.07	0.16	0.00	0.00	0.49	0.20
<i>ARI</i>	0.04	0.13	-0.19		0.21	0.07	0.01	0.02	0.25	0.28
<i>RD</i>	-0.41**	0.15	0.05	-0.02		0.13	0.00	-0.04	0.27	-0.13
<i>P</i>	0.28**	0.24	-0.17	0.01	-0.19		0.00	-0.01	-0.11	0.17
<i>VC</i>	0.03	-0.04	-0.04	0.01	-0.03	0.03		-0.06	-0.12	-0.10
<i>LB</i>	-0.10	-0.01	-0.01	-0.01	-0.16	0.03	0.02		-0.15	-0.25

708 Note: *R* is surface runoff (mm), *RI*<sub>60</sub> is maximum 60-minute intensity (mm h<sup>-1</sup>), *ARI* is average rainfall  
709 intensity (mm h<sup>-1</sup>), *RD* is rainfall duration (h), *P* is rainfall amount (mm), *VC* is vegetation coverage  
710 (%), *LB* is litter biomass (g m<sup>-2</sup>). \*\* means the correlation is significant at 0.01 significance level.



711 **Figure 1.** The location of the study area on the Qinghai-Tibetan Plateau, and the location of runoff  
 712 plots in the study area. (a) The location of the study area, (b) the fragmenting matic epipedon on the  
 713 alpine hillslope and (c) severely degraded meadows formed by the disappearance of matic epipedon  
 714 and (d) four runoff plots of severely degraded meadows (*SDM*) and mixed-cultivated grasslands. A  
 715 typical severely degraded meadow with a slope of 20° was selected to plant mixed grasses. Runoff  
 716 plots were photographed with a drone in the early stages of the 2022 growing season. *DE*,  
 717 *Deschampsia cespitosa* and *Elymus nutans*; *PE*, *Poa pratensis* L.cv. Qinghai and *Elymus nutans*; and  
 718 *PD*, *Poa pratensis* L.cv. Qinghai and *Deschampsia cespitosa*.



719


720 **Figure 2.** Changes in vegetation cover under various mixed-cultivated grasslands from 2019 to 2022.

721 Different capital letters mean that differences were significant in different years for the same

722 grassland community, and different lowercase letters mean that differences were significant between


723 different communities in the same year.

724




725 **Figure 3.** Changes in soil erosion and runoff under various mixed-cultivated grasslands from 2019 to  
 726 2022. (a) Runoff depth and (b) soil loss per unit area. Note: For the four treatment runoff plots, runoff  
 727 and sediment were measured 14, 18, and 10 times, respectively, during the growing season of 2019,  
 728 2020, and 2022. Different capital letters mean that differences were significant in different years for  
 729 the same grassland community, and different lowercase letters mean that differences were significant  
 730 between different communities in the same year. *SDM*, severely degraded meadows, *DE*,  
 731 *Deschampsia cespitosa* and *Elymus nutans*; *PE*, *Poa pratensis* L.cv. Qinghai and *Elymus nutans*; and  
 732 *PD*, *Poa pratensis* L.cv. Qinghai and *Deschampsia cespitosa*. The lines in the middle of the box  
 733 represent the median values. The squares in the box represent the average value.

734



735 **Figure 4.** Runoff, soil loss and sediment concentration reduction ratio under different mixed-  
736 cultivated grasslands from 2019 to 2022. (a) Runoff reduction ratio (*RRB*), (b) soil loss reduction  
737 ratio (*SRB*), (c) sediment concentration reduction ratio (*CRB*) and (d) the percent of runoff reduction  
738 ratio to soil loss reduction ratio (*RRSR*). Note: Different capital letters mean that differences were  
739 significant in different years for the same grassland community, and different lowercase letters mean  
740 that differences were significant between different communities in the same year. The lines in the  
741 middle of the box represent the median values. The squares in the box represent the average value.  
742



743 **Figure 5.** Changes in bulk density, root mass density, field capacity, capillary capacity, non-capillary  
744 porosity and soil cohesion in 0-10 cm soil layer when severely degraded meadow (*SDM*) were  
745 restored to mixed-cultivated grassland for 4 years. *DE*, *Deschampsia cespitosa* and *Elymus nutans*;  
746 *PE*, *Poa pratensis L.cv. Qinghai* and *Elymus nutans*; and *PD*, *Poa pratensis L.cv. Qinghai* and  
747 *Deschampsia cespitosa*. Percentages represent the increased rate of soil properties (increased rate =  
748  $(V_{DE}$  or  $V_{PE}$  or  $V_{PD} - V_{SDM})/V_{SDM}$ ), where  $V_{SDM}$ ,  $V_{DE}$ ,  $V_{PE}$  and  $V_{PD}$  are the mean values of soil  
749 characteristics of *SDM*, *DE*, *PE* and *PD*. Different lowercase letters mean that differences were  
750 significant between different communities. The lines in the middle of the box represent the median  
751 values. The squares in the box represent the average value.