

1 ~~Regulating effects of m~~Mixed-cultivated grasslands ~~promote~~ surface runoff generation and reduce
2 soil loss over time in ~~erosion reduction along with~~ restoration of alpine degraded hillsides

3 Yulei Ma¹, Yifan Liu^{1,2}, Jesús Rodrigo-Comino³, Manuel López-Vicente⁴, Gao-Lin Wu^{1,2,5}

4 ¹ *State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, College of Soil*
5 *and Water Conservation Science and Engineering (Institute of Soil and Water Conservation),*
6 *Northwest A & F University, Yangling, Shaanxi 712100, China*

7 ² *Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water*
8 *Resource, Yangling, Shaanxi 712100, China*

9 ³ *Departamento de Análisis Geográfico Regional y Geografía Física, Facultad de Filosofía y Letras,*
10 *Campus Universitario de Cartuja, University of Granada, Granada, Spain*

11 ⁴ *AQUATERRA Research Group, CICA-UDC, Universidade da Coruña. As Carballeiras s/n, Campus*
12 *de Elviña, A Coruña 15071, Spain*

13 ⁵ *CAS Center for Excellence in Quaternary Science and Global Change, Xi'an, 710061, China*

14 **Correspondence:** wugaolin@nwsuaf.edu.cn (G.L. Wu).

15 **Correspondence address:** State Key Laboratory of Soil Erosion and Dryland Farming on the Loess
16 Plateau, Northwest A & F University, No 26, Xinong Road, Yangling, Shaanxi 712100, P.R. China

17 Phone: +86- (29) 87012884 Fax: +86- (29) 87016082

18 **ABSTRACT**

19 Vegetation restoration is among the most effective measures for controlling runoff and soil erosion
20 resulting from human activity. Nevertheless, few studies have been undertaken to analyze the effects
21 of ~~grassland plant~~ restoration on maintaining the stability of the local runoff amount, especially, in
22 alpine degraded hillsides where mixed-cultivated grasslands predominate ~~in the landscape~~. In this
23 research, we conducted ~~in-situ monitoring using~~ runoff plots to investigate the impact of three
24 mixed-cultivated grasslands strategies, each ~~combining~~ two grass species per plot: (~~*Deschampsia*~~
25 ~~*cespitosa* and *Elymus nutans* (DE), *Poa pratensis* L.cv. Qinghai and *Elymus nutans* (PE), and *Poa*~~
26 ~~*pratensis* L.cv. Qinghai and *Deschampsia cespitosa* (PD);), on a 20-degree slope, assessing the
27 activation and volume of surface runoff and the magnitude of soil loss in alpine degraded hillsides
28 over three years: (~~2019, 2020 and 2022~~). A severely degraded meadow (*SDM*) plot was used as control.
29 The findings indicated that mixed-cultivated grasslands can effectively maintain runoff and reduce
30 soil loss as planting ages increases. Between 2019 and 2022, the values of the average runoff depth
31 for *DE*, *PE*, *PD* and *SDM* were 0.47, 0.55, 0.45 and 0.27 mm, respectively. Despite the increase in
32 runoff, On the contrary, when mixed-cultivated grassland restoration favored soil conservation played
33 a beneficial role in soil consolidation; the net soil loss per unit area of *SDM* was 1.4, 1.3 and 1.9
34 times greater than that in *DE*, *PE* and *PD*, respectively. This implies that protective measures should
35 be prioritized during the initial planting stage of cultivated grasslands in alpine degraded hillsides.
36 The key factors affecting soil loss and runoff were rainfall amount, duration and intensity (60-min
37 intensity). We conclude that the results of this study can serve as scientific guides to design efficient
38 policy decisions for planning the most effective vegetation restoration in the severely degraded
39 hillside alpine meadows. To boost the benefits of grassland restoration, we suggest that protective
40 measures should be prioritized during the initial planting stage of cultivated grasslands.~~

41 **Keywords:** Alpine meadow; Degraded hillside; Mixed-cultivated grassland; land management; runoff;
42 soil erosion.

43 1 Introduction

44 Grasslands are an essential component of terrestrial ecosystems and ~~natural~~ habitats for the
45 development of animal livestock (O'Mara, 2012). They make significant contributions to biodiversity
46 conservation, climate mitigation, carbon sequestration, and water supply and regulation (Bardgett et
47 al., 2021). Despite the importance of grasslands, about half of them are degraded globally, ~~with~~ and
48 5% of them undergoing severe degradation (based on net primary productivity), ~~and this issue~~ which
49 has become a major ~~concern~~ issue for ~~landscape conservation~~ humanity to overcome (Gang et al., 2014;
50 Török et al., 2021). Global grassland net primary productivity (NPP) has declined by 58.84 Tg C per
51 year. To date, numerous studies have been conducted to analyze the threatening drivers of degradation,
52 its negative impacts, and management and restoration methods for grassland degradation (Gang et al.,
53 2014; Grman et al., 2021; Han et al., 2020). Water and soil are critical for human survival and
54 development, as well as irreplaceable basic natural resources that maintain the function of natural
55 ecosystems and the development of socioeconomic systems. Grassland degradation causes the loss
56 of up to 90% of the soil structure that facilitates water movement (infiltration) and retention (water-
57 holding capacity) in soils (Wick et al., 2016), reduces carbon storage potential (Liebig et al., 2009),
58 and impedes soil functioning. Moreover, degraded grasslands are prone to severe soil erosion,
59 especially in mountainous areas. For example, in the Swiss alpine uplands, water erosion ranges from
60 0.14 to 1.25 t $\text{ha}^{-1} \text{month}^{-1}$ according to the phenological stage of the grasses (Schmidt et al., 2019);
61 and in the gully slope of the Loess Plateau, the average amount of soil erosion was 331.26 t km^{-2}
62 during the 2018–2020 grass growing season (Zhu et al., 2021).

UWIT
Can you use the same units -> t/km²/month or year?

63 Precipitation is the main water source of soil moisture supply in semi-arid areas and the
64 conversion of precipitation to runoff is one of the major contributors to river streamflow (Leung et al.
65 2015). In some previous experiences it was observed that ~~v~~Vegetation restoration ~~reduces~~ surface

and decreased (?)

66 ~~runoff while decreasing sediment production, which leading to lower river levels, threatening~~
67 ~~in turn influences~~ the health of river ecosystems (Dijk et al., 2007). A recent study ~~conducted by Wu~~
68 ~~et al. (2020) has proposed sustainable management strategies for semi-arid areas with a positive~~
69 ~~trade-off between surface runoff maintenance and erosion control. influenced by vegetation~~
70 ~~restoration, which in turn influences river flow recharge.~~ However, very few studies have addressed
71 to date there have been limited studies focused on how the effects of effectively restored grasslands
72 ~~can in regulate maintaining surface runoff water supply~~ and preventing soil erosion (Minea et al.,
73 2022). This topic is particularly important for alpine grasslands, which play a vital role in the supply
74 of fresh water and the development of livestock husbandry (Cui et al., 2022). Therefore, it is necessary
75 to assess the impacts of grassland restoration on runoff generation and soil protection. _

76 Vegetation restoration is widely considered as one of the most effective methods for controlling
77 runoff and soil erosion worldwide (Anache et al., 2018). The effects of vegetation cover properties
78 on runoff and soil loss reduction are strongly connected to plant species, leaf and branch coverage,
79 above-ground biomass, litter biomass, and root systems (Liu et al., 2022; Freschet and Roumet, 2017;
80 Gyssels et al., 2005; Zhu et al., 2021). Furthermore, the processes of runoff and soil loss are
81 significantly influenced by the improvement of soil characteristics with vegetation restoration
82 (Schwarz et al., 2015; Gyssels et al., 2005). The interaction between vegetation and soil could
83 stabilize the topsoil and alter soil properties (Saxton and Rawls, 2006; Ma et al., 2023a). Vegetation
84 restoration promotes the formation of soil aggregates, decreases soil bulk density, enhances organic
85 matter and nutrients and improves soil porosity, resulting in high soil hydraulic conductivity and field
86 capacity (Qiu et al., 2022; Saxton and Rawls, 2006). The above-interlinked soil properties alter soil
87 hydrological properties and ultimately influence hillslope and watershed hydrology, such as runoff
88 and soil erosion (Lu et al., 2020; Qiu et al., 2022). While vegetation restoration holds the potential to

89 be a key method of environmental restoration under human management, the inappropriate selection
90 of species can negatively impact the sustainability of local economic and environmental development
91 (Huang et al., 2017; 2019). For example, cultivated grasslands were already advocated as a sensible
92 solution for the conservation of soil and water, as well as the regrowth of vegetation in semi-arid
93 mountain areas (Liu et al., 2022; Wu et al., 2010). Grass communities with multiple stratified
94 structures are better at maintaining surface runoff and decreasing soil loss than those with a single
95 composition and structure (Mohammad and Adam, 2010).

96 Surface runoff ~~—also known as stormwater runoff or overland flow~~ ~~rainwater that moves over the~~
97 ~~ground surface~~— reaches the stream in the form of sheet, rill and gully flow (Rumynin, 2015). The
98 conversion of rainfall to overland flow depends on the rainfall intensity rate, the soil hydrological
99 properties, such as (non-)saturated hydraulic conductivity, matrix flux potential and field capacity,
100 and initial soil water content (López-Vicente and Navas, 2012; Gyssels et al., 2005; De Baets et al.,
101 2007). Because runoff is the primary driver of water erosion on hillslopes and serves as the main
102 agent for sediment transport ~~carrier of sediment transport~~, reducing the conversion of rainfall to
103 runoff is regarded as an effective way to control water erosion, ~~such as~~ through vegetation restoration
104 (Zhou et al., 2016; Zhu et al., 2021). On the other hand ~~However~~, ~~in for the~~ arid and semi-arid regions
105 ~~and headstreams (Qinghai-Tibetan Plateau)~~, surface runoff is the major water supply source to the
106 river streamflow, thereby it is vital for ensuring the sustainability of ecosystems and human activities
107 (Liu et al., 2020; Robinson et al., 2003). Therefore, ~~the fundamental objective of~~ restoration efforts
108 in areas with low rainfall should be oriented ~~is~~ to maintain runoff while reducing its level of sediment
109 concentration.

110 Soil erosion can be reduced by various factors, including the above- and below-ground biomass
111 of grasses, litter cover, and root systems (De Baets et al., 2007). Grasslands can control water erosion

See also work by :

6

[10.1051/agro:2007062](https://doi.org/10.1051/agro:2007062)

[10.1002/esp.447](https://doi.org/10.1002/esp.447)

[10.1016/j.catena.2012.12.012](https://doi.org/10.1016/j.catena.2012.12.012)

112 relying on the ~~the~~ aboveground biomass in dissipating flow energy (Bochet and García-
113 Fayos, 2004), living roots in decreasing soil detachment capacity (Zhang et al., 2013), grass plant
114 cover in intercepting rainfall (Liu et al., 2019), and litter cover in enhancing rainwater infiltration
115 (Liu et al., 2022). Moreover, the ~~reciprocal cementation and~~ interweaving of plant roots can
116 remarkably alter the physical properties of the topsoil, enhancing its resistance to erosion (Schwarz
117 et al., 2015; Wang et al., 2018). The impact of grassroots on the soil characteristics can be summarized
118 as follows: i) increasing the stability of soil aggregates through aggregating fine soil particles into
119 macroaggregates; ii) enhancing soil cohesion through interweaving with the soil; and iii) decreasing
120 soil bulk density through increasing soil porosity (Wu et al., 2019; Gyssels et al., 2005). For example,
121 numerous recent studies have confirmed that a grass with shallow yet dense fibrous root system
122 appears to be more effective at controlling water erosion than grass with good ground cover but low
123 root density (De Baets et al., 2007; Bochet et al., 2006).

124 Alpine meadows, especially in the Qinghai-Tibetan Plateau, constitute the predominant
125 ecosystem in China and the world, accounting for 44% and 6% of total grassland areas, ~~in China and~~
126 ~~the world~~, respectively (Wang et al., 2016). Over 50% of the alpine meadows have been subject to an
127 increasing degree of degradation (Bardgett et al., 2021), with the extent of degradation depending on
128 the meadow patch coverage resulting from the fragmentation of alpine meadows (Fig. 1b). Severely
129 degraded meadow^s (also known as “black beach” and “black soil-type degraded meadow”) formed
130 after the matic epipedon, typically 10 to 15 cm deep, was fully removed by overgrazing and rodent
131 activity exposing the sub-soil (Fig. 1c; Ma et al., 2023a; Shang et al., 2008). Severely degraded
132 meadows amounted to about 30% of the total area of alpine meadows on the Qinghai-Tibetan Plateau
133 (Shang et al., 2008). Recent studies by Niu et al. (2021) and Ma et al. (2023b) have observed that
134 fragmentation of alpine meadows and severely degraded meadows could reduce surface runoff and

135 ~~enhance soil erosion in alpine meadow hillslopes. Alpine meadows are fragile ecosystems when rapid~~
136 ~~changes are involved and due to climate change and overgrazing have suffered substantial~~
137 ~~degradation in recent decades (Fig. 1b and c).~~

138 ~~This situation is leading to a drop in vegetation cover and an increase in severely degraded~~
139 ~~meadows, especially for hillside grassland, ultimately posing a great hazard to the plateau from water~~
140 ~~and soil loss (Liu et al., 2022).~~

141 The Qinghai-Tibetan Plateau serves as the headwaters for many of Asia's major rivers (Xu, 2018).
142 ~~The eastern and southern parts of the Qinghai-Tibet Plateau, are influenced by the monsoon,~~
143 ~~and where rainfall is the primary source of streamflow (Cuo et al., 2014).~~ The long-term and
144 widespread degradation of hillside alpine meadows has disrupted the soil water balance, reducing
145 ~~surface runoff (Niu et al., 2019; Ma et al., 2023b).~~ This, in turn, diminishes river streamflow,
146 ultimately constraining the sustainable development of both local and downstream regions. The
147 importance of artificial grassland in restoring alpine degraded meadow is widely accepted (Wen et
148 al., 2018; Wu et al., 2010). ~~Artificial grassland –also known as tamed grassland, sowed grassland~~
149 ~~and cultivated grassland– refers to fields that have been broken up and replanted with exotic grasses~~
150 ~~and forbs and utilized for hay crop production or cattle grazing (Fisher et al., 2018).~~ The establishment
151 of artificial grassland on severely degraded areas provides a dual benefit by boosting productivity and
152 improving the ecological environment of alpine grasslands (Shang et al., 2008; Liu et al., 2022).

153 While previous ~~studies reports~~ have often focused on carbon sequestration capacity, vegetation
154 characteristics, soil quality and productivity of cultivated grassland (Wang et al., 2013; [Bai and](#)
155 [Cotrufo et al., 2022](#)), there has been a limited examination of the impacts of mixed-cultivated
156 grasslands on the provision of runoff and prevention of soil erosion on the alpine hillsides.
157 ~~Recently, Only~~ [Liu et al. \(2022\)](#) ~~evaluated the effects of plant morphological characteristics on runoff~~

158 [and soil erosion in different mixed-cultivated grassland under natural rainfall events](#). Here, we present
159 novel research to examine the ability of ~~alpine hillsides~~ cultivated grasslands to regulate runoff and
160 soil loss ~~through~~ and evaluated the effect of three different mixed-cultivated grasslands: (~~*Deschampsia cespitosa* and *Elymus*~~
161 ~~*nutans* (DE), *Poa pratensis* L.cv. Qinghai and *Elymus nutans* (PE), and *Poa pratensis* L.cv. Qinghai~~
162 ~~and *Deschampsia cespitosa* (PD),~~) compared to [a severely degraded meadows \(SDM\)](#) ~~in alpine~~
163 ~~degraded hillsides~~ by a three-year field experiment. In particular, this study aimed to (1) assess the
164 temporal variations in soil and water loss of DE, PE and PD grasslands during the growing season
165 and under natural rainfall; and (2) determine the key factors influencing the mixed-cultivated
166 grasslands in controlling runoff and soil erosion. ~~In this vein,~~ This study has realistic implications
167 for understanding the contribution of mixed-cultivated grasslands restoration on soil erosion control
168 in ~~the~~ degraded alpine hillsides.

169

170 2 Materials and methods

171 2.1 Study area

172 This study was carried out in the representative area of Zhique Village (33°40'01" N and 99°43'06" E,
173 elevation over 4200 m a.s.l), Dari County, Qinghai province, which served as a field experimental
174 site and model area for the restoration of severely degraded alpine meadow on the Qinghai-Tibetan
175 Plateau (Fig. 1a). The climate conditions correspond to a typical highland one with low temperatures
176 throughout the year, i.e., not showing [the typical four-season pattern \(spring, summer, autumn,](#)
177 ~~winter)~~ [distinct seasons, but rather just two main seasons: cold and warm.](#) ~~just cold and warm ones.~~ In
178 the study region, the average annual temperature is ~~-03.1~~ -3.1 °C, with monthly variations from -
179 ~~18.34.7~~ 18.34.7 °C in January to ~~12.47.5~~ 12.47.5 °C in July ([Li et al., 2018](#)) (~~values corresponded to the period between~~

180 ~~1981 and 2018; according to data source: d from the~~ [European Centre for Medium-Range Weather](#)
181 [Forecasts](#)). The average annual precipitation is 416 mm, with the majority of it falling from July to
182 September, [based on Climate Hazards Group InfraRed Precipitation with Station data \(CHIRPS\)](#).
183 ~~Nevertheless, t~~The majority of the precipitation and the warm season falls during the vegetation
184 growth period (from May to September), favoring optimal conditions for the development of
185 ~~plants~~vegetation. The soil type in the study area is classified as [Mat Cryi-gelic Cambisolsalpine](#)
186 ~~meadow soil~~ (IUSS-WRB, 2015). Currently, the remnant vegetation in this site is composed of an
187 alpine shrub (*Salix cupularis* and *Potentilla fruticosa*), alpine meadow (*Kobresia pygmaea*, *Kobresia*
188 *humilis* and *Kobresia capillifoli*) and swamp meadow (*Carex atrofusca*, *Poa annua* and *Carex parva*).

189 Soil erosion in the degraded alpine meadows ~~is~~was severe, ~~becoming~~which was the primary
190 source of sediment delivered to streams in the study area (Liu et al., 2022). The matic ~~epipedon~~
191 ~~epipedon~~ of alpine meadow has experienced fragmentation and even disappearance (Fig. 1b),
192 eventually forming a severely degraded meadow [\(Fig. 1c\)](#). ~~Before implementing the grassland~~
193 ~~restoration project, i.e., Subsidy and Incentive System for Grassland Conservation, t~~The average soil
194 erosion rate and the total erosion in the study area were 13.63 t ha⁻¹ y⁻¹ and 323.58 × 10⁶ t y⁻¹,
195 respectively, ~~before the implementation of the grassland restoration project, i.e., Subsidy and~~
196 ~~Incentive System for Grassland Conservation~~ (Zhao et al., 2021). Severely degraded meadows were
197 restored via mixed-cultivated grasslands ~~—~~fields were ploughed and replanted with two grass species
198 ~~—~~— and moderately degraded meadows were restored by broadcast sowing on the hillslopes during
199 the implementation of the grassland restoration project. The grass species used for the projects have
200 excellent characteristics like strong trampling tolerance, good palatability, abundant leaf quantity and
201 developed rhizome, such as *Poa pratensis* L. cv. Qinghai, *Deschampsia cespitosa* and *Elymus nutans*
202 (Shang et al., [2008](#)).

203

204 2.2 Experimental design and measurement

205 The degraded hillslopes are the main component of runoff generation and confluence areas on the
206 Qinghai-Tibetan Plateau. Hence, the grass species chosen for mixed-cultivated grasslands ~~should not~~
207 only must it be grazing-tolerant and good forage, but also prevent soil loss and maintain surface runoff.

208 Potential grass species should also be fully acclimated to harsh alpine climate and have
209 complementary morphological characteristics and living habits (Liu et al., 2022). The community
210 established by matching of grasses morphological characteristics and habits has a hierarchical vertical
211 cover structure and little inter- or intraspecific competition. Following the above-mentioned
212 guidelines for choosing grass species, we ultimately decided on three species (*Deschampsia cespitosa*,
213 *Poa pratensis* L. cv. Qinghai and *Elymus nutans*) from the most widely utilized grass species.
214 *Deschampsia cespitosa* is a cool-season bunching grass native to alpine environments. It typically
215 forms a low, dense tussock (to 30–50 cm tall) of very thin (0.5 cm wide), arching, flat to inrolled,
216 dark green grass blades (to 5 cm long). *Deschampsia cespitosa*, a common bottom grass, has 70% of
217 its grass stems growing between 0 and 30 cm tall. *Elymus nutans* is a common and important plant
218 species in the alpine meadows of the Qinghai-Tibetan pPlateau (Chen et al., 2009). It is a valuable
219 fodder grass in alpine locations that has been extensively employed for animal production, disturbed
220 grassland restoration, and artificial grassland construction due to its resilience to cold, drought and
221 pests (Ren et al., 2010). *Elymus nutans* is a herbaceous perennial species with sparsely tufted culms
222 that can grow to heights of 70 to 100 cm (Liu et al., 2022). *Poa pratensis* L. cv. *Qinghai* is the common
223 and dominant species native to the Qinghai-Tibetan Plateau. It is an excellent species that has been
224 selected and cultivated to restore degraded alpine meadows. Also, *Poa pratensis* L. cv. Qinghai is an
225 herbaceous perennial species with erect or geniculate base culms that grow 20–60 cm tall.

226 To reveal the effectiveness of mixed-cultivated grasslands in controlling runoff and soil loss on
227 hillsides, field observation of mixed grass plots designed by us was conducted from the 2019 to 2022
228 growing seasons. Therefore, one plot with severely degraded meadow (*SDM-*) as a control and three
229 plots with two mixed grass seeds per plot of *Deschampsia cespitosa* and *Elymus nutans* (*DE*), *Poa*
230 *pratensis* L.cv. Qinghai and *Elymus nutans* (*PE*), and *Poa pratensis* L.cv. Qinghai and *Deschampsia*
231 *cespitosa* (*PD*) were selected as the testing site (Fig. 1d). All four runoff plots were spaced 1m apart
232 and were located on the same hillside with the same elevation and soil texture. All plots were bounded
233 by steel plates (30 cm high and 2 mm thick sheet) and built during May 2019, with an area of 10 m²
234 (2 m wide and 5 m long parallel to the maximum slope gradient). To collect only runoff and soil loss
235 from the runoff plot, the steel plate was put vertically into the soil to a depth of about 10 cm, with the
236 remainder sticking out from the soil surface. At the outlet of each plot, a steel runoff collection and
237 calibrated tank (75 L) were set up to gather sediment and runoff. To prevent the collected runoff from
238 being lost to evaporation, the calibrated tank was set inside a sealed vat (Fig. 1d).

239 In addition, the grass seeding for each runoff plot was completed in May 2019. For the runoff
240 plots, grass seeds were distributed to a depth of less than 1 cm in strips at 20 cm intervals following
241 plowing. The seeding rate was set at 6.0 g m⁻² for *Poa pratensis* L.cv. Qinghai and *Deschampsia*
242 *cespitosa* and 4.5 g m⁻² for *Elymus nutans* to ensure a constant number of plants based on germination
243 and seedling emergence rates. None of the runoff plots experienced any human disturbance during
244 the observation period (2019–2022), including grazing, harvesting, and excavation.

245

246 **2.3 Rainfall, runoff and soil loss measurement**

247 A Vantage pro 2TM weather station (Davis Instruments Corp., USA) with a measurement accuracy of
248 4% *iwas* positioned next to the experimental plots to monitor precipitation intensity and duration (Fig.

249 1). A precipitation event was defined by the occurrence of a no-rain interval lasting more than 3 h
250 between them. A total of 42 precipitation events were recorded from 2019 to 2022 throughout the
251 growing season. Snow was not collected, and only rainfall was recorded during the growing season
252 (from June 15 to August 25). Precipitation characteristics of each event, including amount (P),
253 duration (RD), and maximum intensities of 60 minutes (RI_{60}) were recorded, and average
254 rainfall intensity (ARI) was calculated by dividing the total rainfall amount by the duration of the
255 rainfall event. were recorded. After each rainfall-runoff event, both runoff and sediment were
256 collected right away. The water level in the calibrated tank was first measured to calculate the runoff
257 volume. Then, runoff was fully mixed inside the calibrated tank using a stirring bar to thoroughly
258 whirl, and two 500 ml bottles were used to obtain mixture samples of sediment and runoff. When the
259 calibrated tank had less than 1000 ml of runoff sample, all runoff was collected. Lastly, the calibrated
260 tank was cleaned in order to collect sediment and runoff for the subsequent rainfall-runoff event. The
261 mixture samples in the bottle were transported back to the lab to be filtered on quantitative analysis
262 filter paper (30–50 μ m). The filter paper with sediment was oven-dried to a consistent weight at
263 105 °C. The ratio of soil loss amount to runoff volume in the mixed samples was applied to calculate
264 the sediment concentration. Finally, runoff volume and sediment concentration were multiplied to
265 calculate soil loss in each plot.

266 We collected runoff and soil erosion data during the growing season for the years 2019 to 2022.
267 Data for 2021 could not be collected due to the prevention and control strategies for coronavirus
268 (COVID-19). Soil erosion and runoff were portrayed in this work by soil erosion per unit area (g m^{-2})
269 and runoff depth (mm). The runoff depth (R) and soil erosion per unit area (S) could be calculated
270 using the following formulas:

$$R = \frac{V_R}{A} \times 10^3 \quad (1)$$

$$S = \frac{S_T}{A} \quad (2)$$

271 where V_R is the volume of runoff from runoff plots (m^3), S_T is the total amount of soil erosion from
 272 runoff plots (g), and A is the area of runoff plot (m^2).

273

274 2.4 Vegetation and soil properties measurement

275 Vegetation cover (VC), [including dead \(standing litter\) and living vegetation](#), was measured monthly
 276 from 2019 to 2022 growing seasons using a steel wire frame ($50 \text{ cm} \times 50 \text{ cm}$) subdivided into 25
 277 plots of $10 \text{ cm} \times 10 \text{ cm}$. Fig. 2 exhibited the change in vegetation coverage for all runoff plots from
 278 2019 to 2022. After collecting runoff samples each year, the quadrats ($50 \text{ cm} \times 50 \text{ cm}$) were positioned
 279 in the up-, mid-, and down-slope areas. Litter in each quadrant was collected and oven-dried to
 280 determine litter biomass (LB) (Zhu et al., 2021). The litter collection for 2019 was not completed due
 281 to the seeding of mixed-cultivated grasslands in May 2019, and the litter collection for 2020 and 2021
 282 was collected at the end of the runoff collection for the current year. Undisturbed soil samples were
 283 taken in the 0–10 cm soil layers using steel rings in 2022. All soil samples were saturated and then
 284 weighed (W_{sat}). Then the saturated soil samples were placed on the dry sand layer to drain water for
 285 about 2 h and 8 h, and weighed (W_{2h} and W_{8h}). Finally, soil samples were dried in an oven at $105 \text{ }^\circ\text{C}$
 286 for 24 h and then weighed (W_{dr}). Based on the above measurement, soil bulk density (BD , g cm^{-3}),
 287 total porosity (TP , %), capillary porosity (CP , %), [non-capillary porosity \(\$NCP\$, %\)](#), and [soil water](#)
 288 [content at field capacity \(\$FC\$, %\)](#) were determined as follows:

$$FC = \frac{(W_{8h} - W_{dr})}{(W_{dr} - W_{sr})} \quad (3)$$

$$BD = \frac{(W_{dr} - W_{sr})}{V} \quad (4)$$

$$TP = \left(1 - \frac{BD}{ds}\right) \times 100 \quad (5)$$

$$CP = \frac{(W_{2h} - W_{dr})}{V} \quad (6)$$

$$NCP = TP - CP \quad (7)$$

289 where W_{sr} is the weight of the ~~cut~~ steel ring (g), ds is the soil particle density (generally being 2.65
290 g cm⁻³), and V is the volume of the ~~cutting~~ ring (100 cm³).

291 In addition, root mass density (*RMD*) was obtained using a root drill, followed by washing with
292 water and drying in the oven. Four undisturbed samples were collected in each quadrat using a steel
293 ring (6.18 cm diameter and 2.0 cm height), and they were applied to a direct shear (ZJ type). The soil
294 cohesion ~~cohesion~~ was obtained by the Mohr–Coulomb theory (Fattel et al., 2011).

295

296 2.5 Calculating the reduction effect of runoff and soil loss

297 Four metrics were employed to assess the efficiencies of the mixed-cultivated grasslands in regulating
298 runoff and soil loss, which were: The runoff reduction ~~benefitratio~~ (*RR~~EB~~*, %), sediment
299 concentration reduction ~~benefitratio~~ (*CR~~EB~~*, %), soil erosion reduction ~~benefitratio~~ (*SR~~EB~~*, %), and
300 the percentage of runoff reduction ratio to soil loss reduction ratio (*RRSR*) (Zhao et al., 2014). High
301 values of *RR~~EB~~*, *SR~~EB~~* or *CR~~EB~~* indicated that vegetation was able to reduce runoff, soil erosion or
302 sediment concentration compared to the rates observed in the control plot (severely degraded
303 meadow). In addition, a low *RRSR* implied that vegetation was more beneficial in minimizing soil
304 erosion than in minimizing runoff (Liu et al., 2020). These indices were calculated as follows:

$$RR\del{EB} = \frac{R_c - R_v}{R_c} \times 100 \quad (8)$$

$$SR\del{BE} = \frac{S_c - S_v}{S_c} \times 100 \quad (9)$$

$$CR\del{EB} = \frac{C_c - C_v}{C_c} \times 100 \quad (10)$$

$$RRSR = \frac{RRBE}{SRBE} \times 100 \quad (11)$$

305 where R_c and R_v are the runoff depths of the degraded meadow plot and plots covered by mixed-
306 cultivated grasslands; S_c and S_v are the soil loss per unit area of the degraded meadow plot and
307 plots covered by mixed-cultivated grasslands; C_c and C_v are the sediment concentrations of the
308 degraded meadow plot and plots covered by mixed-cultivated grasslands, respectively.

309

310 2.6 Statistical analyses

311 ~~All data were analyzed u~~Using SPSS statistics software (IBM, USA, version 26.0), ~~all data were~~
312 ~~analyzed~~. The Kolmogorov–Smirnov test was used to test the normality of data. Duncan’s multiple
313 range tests of one-way analysis of variance (ANOVA) were applied to test for significant differences
314 between soil and vegetation characteristics, runoff depth, soil erosion amount, and runoff and soil
315 loss reduction ratio under various mixed-cultivated grasslands at 0.05 significance levels. Also, path
316 analysis is a form of multiple regression statistical analysis that is used to evaluate causal models by
317 examining the relationships between runoff, soil loss and soil and vegetation properties. By using this
318 method, one can identify the major factors influencing runoff and soil loss and determine the direct
319 and indirect effects of soil and vegetation properties on runoff and soil loss.

320

321 3 Results

322 3.1 ~~Mixed-cultivated grasslands modified r~~Runoff ~~yield~~ and soil loss ~~under various mixed-~~ 323 ~~cultivated grasslands~~

324 Mixed-cultivated grasslands ~~dramatically~~ increased runoff and reduced soil erosion. One-way
325 analysis of variance (ANOVA) revealed that runoff significantly ($P < 0.05$) increased after the

326 severely alpine degraded hillside was restored by the mixed-cultivated grassland (Fig. 3). During the
327 ~~three evaluated~~ growing seasons (~~of~~ 2019, 2020, and 2022), ~~the values of the average runoff depth~~
328 ~~for DE, PE, PD and SDM were~~ 0.47, 0.55, 0.45 and 0.27 mm, respectively. The average runoff
329 depths of *SDM* ~~in 2019, 2020, and 2022~~ were 0.23, 0.34 and 0.25 mm, respectively, all significantly
330 ($p < 0.05$) lower~~less~~ than (except for 2020) the average runoff of mixed-cultivated grassland *DE*, *PE*
331 and PD, which measured 0.44, 0.59 and 0.50 mm in 2019, 0.55, 0.51 and 0.38 mm in 2020, 0.43,
332 0.54 and 0.40 mm in 2022 (Fig. 3a). ~~However~~Regarding soil conservation, the amount of soil loss in
333 grasslands was significantly influenced by ~~the age of the~~ planting age. As depicted in Fig. 3b, soil
334 loss in DE, PE and PD (except for DE in 2019) were significantly ($p < 0.05$) higher than in 2019
335 and 2020 (the first and second years of planting) than those in the fourth year of planting (2022). In
336 the year 2020, soil loss produced by DE, PE, and PD was significantly higher ($p < 0.05$) greater than
337 that of SDM. ~~Satisfactorily~~However, the three mixed-cultivated grasslands did exhibit a clear
338 reduction in soil loss compared to SDM in 2022 (albeit not significantly), with soil loss per unit area
339 for SDM being 1.4, 1.3, and 1.9 times higher than those for DE, PE, and PD, respectively. No
340 significant differences ($p > 0.05$) was observed~~ere detected~~ in runoff depth and soil loss between *DE*
341 *PE* and *PD* in 2019, 2020 and 2022. The results showed that any of the three mixed-cultivated
342 grasslands (*DE*, *PE*, and *PD*) could be effective in controlling soil loss and maintaining runoff.

343

344 **3.2 Specific rRunoff and soil loss reduction ratios of the under various mixed-cultivated** 345 **grasslands**

346 Fig. 4 illustrates the runoff, soil loss and sediment concentration reduction ratio after planting various
347 mixed-cultivated grasslands. Lower *RR_{EB}* values indicated a better ability to maintain runoff for
348 mixed-cultivated grasslands, while higher *SR_{EB}* and *CR_{EB}* values indicated better effectiveness of

349 grasslands in soil loss reduction. The mean $RR\bar{E}B$ values of the grass community DE , PE_7 and PD
350 were -79.3%, -130.4% and -48.5% in 2019, -36.9%, -53.5% and -21.5% in 2020, and -115.4%, -156.1%
351 and -87.6% in 2022, respectively (Fig. 4a). Regardless of the combination of the above-mentioned
352 grass species, the average increase ratio of runoff in 2022 (the fourth years of planting) was
353 significantly ($p < 0.05$) higher than that in 2019 and 2020 (the first and second years of planting). The
354 $SREB$ of the three mixed-cultivated grasslands (DE , PE_7 and PD) increased with increasing planting
355 age. It is worth noting that the average $SREB$ values in the grassland communities of DE , PE and PD
356 were 18.0%, 24.3%₇ and 31.9% in 2022, respectively (Fig. 4b). The SRE values of DE , PE and PD in
357 2022 were significantly ($p < 0.05$) higher than those of 2019, whereas SRE values between 2020 and
358 2022 was significant ($p < 0.05$) for DE but not ($p > 0.05$) for PE and PD . Additionally, $CREB$ for all
359 mixed-cultivated grasslands in 2022 was significantly ($p < 0.05$) higher than that in 2019 and 2020.
360 The mean $CREB$ values of the cultivated-grassland communities DE , PE_7 and PD increased from -
361 120.9% to 55.8%, from -112.4% to 59.7%, and from -94.3% to 62.1% from 2019 to 2022, respectively
362 (Fig. 4c). Regardless of the age of the grasslands, the value of $RRSR$ was less than 1, suggesting that
363 the soil erosion reduction effect of the grasslands was higher than its runoff reduction effect (Fig. 4d).
364 No significant differences ($p > 0.05$) ~~appeared~~observed in RRB , SRB , CRB and $RRSR$ between DE
365 PE and PD in 2019, 2020 and 2022.

366

367 3.3 Key factors affecting runoff and soil loss

368 Precipitation characteristics and vegetation features played a significant role in influencing the
369 hydrological response of the soil. In this study, path analysis was applied to identify the key factors
370 affecting soil loss. The results of this analysis indicated that the sum of path coefficients of RI_{60} , RD ,
371 P and VC were 0.31, 0.36, 0.40 and 0.32, respectively (Table 1). This suggests that P , RD , VC and

372 RI_{60} had positive effects on runoff ~~yield~~amount, with P being the most influential factor. Direct
373 influences on runoff were primarily attributed to ARI and RD , with direct path coefficients of 0.37
374 and 0.67, respectively. Meanwhile, the influences of P and LB on runoff were mainly indirect, with
375 indirect path coefficients of 0.57 and 0.25, respectively. For instance, P , in combination with other
376 factors, particularly RI_{60} and RD , contributed significantly to runoff.

377 Soil loss was significantly influenced by R , RI_{60} , ARI and LB , ~~with R being the most relevant~~. The
378 sum of path coefficients of R , RI_{60} , ARI and LB were 0.52, 0.20, 0.28 and -0.25, respectively (Table
379 2). These results show that R , RI_{60} and ARI had a promotional effect, whereas LB had an inhibitory
380 effect on soil loss. Meanwhile, R and P had a direct positive influence on soil erosion, with direct
381 path coefficients of 0.60 and 0.28, whereas RI_{60} and RD had a direct negative influence on soil erosion,
382 with direct path coefficients of -0.29 and -0.41 (Table 2). In addition, the direct and indirect path
383 coefficients both indicated that LB had an inhibitory influence on the soil loss per unit area, with
384 values of -0.10 and -0.25, respectively.

385

386 4 Discussion

387 4.1 Benefits~~Contribution~~ of mixed-cultivated grasslands on soil conservation and runoff

388 maintenance

389 The mixed-cultivated grasslands (DE , PE_7 and PD) effectively maintained runoff and minimized soil
390 loss (Fig. 4). This finding is similar to those of studies conducted checking different grassland
391 communities (Liu et al., 2019; Liu et al., 2022). In this study, the mixed-cultivated grasslands
392 significantly increased surface runoff compared to the SDM . The difference in runoff between mixed-
393 cultivated grasslands and SDM may be attributed to the soil infiltration rate. Mixed-cultivated

394 grasslands had more abundance of fibrous roots in the topsoil compared with *SDM* (Fig. 5), and those
395 fine roots reduced infiltration by occupying the soil pore (Leung et al., 2015). In comparison to *SDM*,
396 soil non-capillary porosity (*NCP*) and field capacity (*FC*) of *DE*, *PE* and *PD* significantly decreased
397 by 46%, 32% and 48%, and increased by 55%, 59% and 48%, respectively (Fig. 5). This implied that
398 *SDM* was restored to mixed-cultivated grasslands with lower permeability and better water retention.
399 This was further evidence that infiltration was responsible for the difference in runoff between the
400 mixed-cultivated grasslands and *SDM*. Association between variables is not necessarily causal evidence

401 Soil loss in all three mixed-cultivated grassland communities (*DE*, *PE* and *PD*) was higher than
402 that in the *SDM* during the first- and second years following planting. However, by the fourth year,
403 the *SDM* exhibited higher soil loss than the three mixed-cultivated grasslands (Fig. 3). These changes
404 in soil erosion were dominantly attributed to the developing of the root system and improvement of
405 soil structure (Zhu et al., 2021). The loosening of the soil structure caused by the seeding method of
406 plowing resulted in a greater soil loss of the three mixed-cultivated grasslands than the *SDM* at the
407 beginning of the planting. We confirmed that the age of plantation was a key factor in understanding
408 the inter-annual changes of soil erosion. This idea was also demonstrated in other types of primary
409 land uses such as woody crops or young forests (Rodrigo-Comino, et al., 2018). Nevertheless, we
410 hypothesize that grassland topsoil demonstrated a stronger resilience to erosion as its root system
411 grew, which had a reinforcement impact on the soil and led to lower soil loss in the fourth year of
412 planting than that of the *SDM*. The topsoil (0-10 cm) of the grasslands had significantly different soil
413 properties from the *SDM* in the fourth year after planting, as detailed in Table 3. In comparison to
414 *SDM*, the root mass density and soil cohesion of grasslands *DE*, *PE* and *PD* increased by 672%, 890%
415 and 589%, and by 379%, 282% and 315%, respectively.

416

417 4.2 Effect of rainfall and ~~vegetation~~grassland community characteristics on runoff and soil 418 loss

419 Surface runoff and erosion process is influenced and constrained by rainfall depth, intensity and
420 duration, and by vegetation cover (*VC*) as well (Mohamadi and Kavian, 2015b; Bochet et al., 2006).
421 In this study, the *VC* had a directly promoted effect on surface runoff. Moreover, this result was in
422 line with the finding of Niu et al. (2021), who reported that the surface runoff increased with the
423 grassland coverage. Our results also indicated that rainfall amount (*P*) could have an indirect effect
424 on surface runoff through rainfall duration (*RD*) and maximum intensities of 60 minutes (*RI*₆₀). This
425 implies that heavier and longer-lasting rainfall events were more likely to lead to surface runoff
426 generation (Dos Santos et al., 2017). The findings demonstrated that runoff depth (*R*) and the average
427 rainfall intensity (*ARI*) were the most and second most influential factors in promoting soil erosion
428 (Table 2). The primary cause for this is that runoff velocity increases with higher precipitation
429 intensity (Wang et al., 2013), which likely enhances the capacity of soil detachment and transport by
430 surface runoff (Zhu et al., 2021). Furthermore, litter biomass (*LB*) had a direct and negative impact
431 on soil loss (Table 2), indicating that the effectiveness of grasslands in reducing soil loss increased as
432 litter biomass increased. Liu et al. (2022) found that the soil loss rate decreased with increasing litter
433 biomass in ~~the~~ grassland. Plant litter can intercept precipitation, reducing rainfall kinetic energy and
434 splash erosion, while also increasing surface roughness (Liu et al., 2017; Xia et al., 2019) All these
435 processes favor a reduction in runoff yield and soil loss rates.

436 The matching of morphological characteristics of plants can effectively reduce soil loss (Liu et
437 al., 2022). In this study, the reduction in soil loss in the early stages of mixed-cultivated grassland
438 planting (2019 and 2020) was attributed to grassland cover and ~~matching of plant morphological~~
439 characteristics of grass community. The morphological characteristics of *Deschampsia cespitosa*, *Poa*

440 *pratensis* L.cv. Qinghai and *Elymus nutans* are dense clump type, rhizomatic-sparse clump type, and
441 sparse clump perennial grasses, respectively. In addition, *Deschampsia cespitosa* and *Poa pratensis*
442 L.cv. Qinghai are bottom grasses, while *Elymus nutans* belongs to the top grass. The mix of dense and
443 sparse grasses (*DE* and *PD*), and mix of top and bottom grasses (*DE* and *PE*) can complement each
444 other morphologically and structurally, thereby more effectively reducing the kinetic energy of
445 raindrops (Liu et al., 2022). *Poa pratensis* L.cv. Qinghai, a rhizomatic grass, also has abundant root
446 systems intertwined with the soil, increasing soil cohesion and consequently reducing soil detachment
447 capacity (Wang et al., 2018). Overall, in this study, the morphological and root characteristics of
448 mixed-cultivated grasslands reduced runoff velocity, influenced water infiltration process and
449 decreased soil erodibility.

450

451 **4.3 Implications for ~~mixed-cultivated~~ grasslands restoration on ~~the~~ degraded alpine hillsides**

452 Our findings demonstrated that mixed-cultivated grasslands with complementing morphological
453 features and habits can be ~~more~~ effective at maintaining runoff and reducing soil erosion. Three
454 mixed-cultivated grasslands (*DE*, *PE*, and *PD*) exhibited an effective role in controlling soil loss on
455 the degraded alpine hillside. However, at the start of planting, the mixed planted grassland had a
456 greater soil erosion than ~~the~~ severely degraded meadow, whereas ~~the function of reducing~~ soil loss
457 was ~~reached~~ in the 4th year of planting (Figs. 2 and 3). This suggested that protection measures, such
458 as mesh covering and anti-trampling, may be taken into account to reduce soil loss in the initial
459 planting stage of cultivated grassland in alpine hillsides (Liu et al., 2022). Moreover, grass may also
460 be planted with a no-till system to avoid the initial increase of soil erosion at the initial phases of
461 cultivated grassland by destroying soil structure (Karayel and Sarauskis, 2019). In addition, spring
462 meltwater is the main driver of soil erosion in degraded alpine meadows in alpine regions, which

463 greatly increases turbidity of~~turbidizes~~ rivers (Zheng et al., 2022; Shi et al., 2020). The restoration of
464 severely degraded hillslope meadows increased vegetation cover and soil ability, both of which could
465 have an inhibitory impact against meltwater erosion (Liu et al., 2022). To better understand the effects
466 of cultivated grassland on meltwater erosion, future experiments under natural freezing and thawing
467 conditions need to be monitored.

468 Cultivated grasslands, considered a crucial component of vegetation restoration, have been
469 extensively utilized in the rehabilitation of degraded alpine hillsides (Shang et al., 2008). Nevertheless,
470 plant restoration is not necessarily beneficial to the long-term viability of on- and off-site ecosystems'
471 functions, including natural succession and river ecosystems. Therefore, the selected vegetation types
472 ought to be advantageous for the ecosystem's sustainability, both on- and off-site, such as maintaining
473 river streamflow and unrestricted natural succession. The seed prices of cultivated grass communities
474 of *Deschampsia cespitosa* and *Elymus nutans*, *Poa pratensis* L.cv. Qinghai and *Elymus nutans*, and
475 *Poa pratensis* L.cv. Qinghai and *Deschampsia cespitosa* were about \$690, \$750 and \$480 per ha.
476 Planting properly mixed-cultivated grassland on the alpine degraded hillsides can achieve both
477 environmental and economic benefits. This study proved that mixed-cultivated grasslands could
478 maintain runoff and decrease soil loss.

479 5 Conclusions

480 Based on the measured data during the 2019, 2020 and 2022 growing seasons, the planting of mixed-
481 cultivated grassland on severely degraded hillside alpine meadow ~~could~~ effectively maintained
482 surface runoff and decreased soil loss, especially after the mixed-cultivated grassland played a
483 positive role in consolidating the surface soil. The benefits were statistically significant compared
484 with the control plot, but differences between the three types of cultivated grasslands were not

485 ~~significant. The mean *CREB* values of the mixed-cultivated grasslands *DE*, *PE*, and *PD* increased~~
486 ~~from 120.9% to 55.8%, from 112.4% to 59.7%, and from 94.3% to 62.1% from 2019 to 2022,~~
487 ~~respectively.~~ Planting the mixed-cultivated grasslands after ploughing loosened the soil structure and
488 thus increased sediment concentration in runoff during the first stage after planting. Subsequently,
489 sediment concentration decreased with the growth of the root system of the mixed-cultivated
490 grasslands, ~~strengthening the sloping soils~~ ^{improving root and soil cohesion?} due to the root architecture. To guarantee that they can
491 perform the aforementioned functions, mixed-cultivated grasslands need protection measures in the
492 initial planting stage. Our results also suggested that mixed-cultivated grasslands ~~with complementary~~ ^{different but}
493 morphology and structure and abundant fine root systems were effective in maintaining surface runoff
494 and reducing soil erosion. Precipitation amount, duration, ~~vegetation coverage~~ ^{and composition} and maximum 60-
495 minute intensity, ~~and vegetation coverage~~ were the predominant factors affecting surface runoff and
496 soil loss. The erosion resistance contribution of the above-ground community characteristics and
497 below-ground roots along the cultivated time could maintain a relatively high surface runoff and
498 decrease sediment concentration. These findings have potential implications for understanding the
499 contribution of mixed-cultivated grasslands restoration on soil erosion control in the degraded
500 hillsides of alpine areas.

501
502 *Data availability.* [The data that support the findings of this study are available on request from the](#)
503 [corresponding author. All data needed to evaluate the conclusions in the paper are present in the paper.](#)

504
505 *Author contributions.* Yulei Ma: Investigation, Formal analysis, Methodology, Software, Writing -
506 original draft. Yifan Liu: Investigation, Formal analysis, Writing - review & editing. Jesús Rodrigo-
507 Comino: Interpretation of data, Writing - review & editing. Manuel López-Vicente: Interpretation of

508 data, Writing - review & editing. Gao-Lin Wu: Conceptualization, Funding acquisition, Supervision,
509 Writing - original draft, review & editing.

510

511 *Competing interests.* The authors declare that they have no known competing financial interests or
512 personal relationships that could have appeared to influence the work reported in this paper.

513

514 *Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional
515 claims in published maps and institutional affiliations

516 *Acknowledgments.* We thank Gall Corinna, Vanacker Veerle and Qianjin Liu for their constructive
517 comments and suggestions on this manuscript.

518 *Financial support.* This research was funded by the National Natural Science Foundation of China
519 (NSFC41930755, NSFC32230068), the Strategic Priority Research Program of the Chinese Academy
520 of Sciences (XDB40000000), and the Second Stage's Research and Technique Extending Project of
521 Sanjiangyuan Ecological Protection and Building in Qinghai (2019-S-1).

522

523 **References**

524 Anache, J.A.A., Flanagan, D.C., Srivastava, A., and Wendland, E.C.: Land use and climate change
525 impacts on runoff and soil erosion at the hillslope scale in the Brazilian Cerrado, *Sci. Total*
526 *Environ.*, 622–623, 140–151, <https://doi.org/10.1016/j.scitotenv.2017.11.257>, 2018.

527 [Bai, Y., Cotrufo, M.F.: Grassland soil carbon sequestration: Current understanding, challenges, and](#)

- 528 [solutions, Science, 377\(6606\), 603–608. https://doi.org/10.1126/science.abo2380, 2022.](https://doi.org/10.1126/science.abo2380)
- 529 Bardgett, R.D., Bullock, J.M., Lavorel, S., Manning, P., Schaffner, U., Ostle, N., Chomel, M.,
530 Durigan, G., Fry, E.L., Johnson, D., Lavallee, J.M., Le Provost, G., Luo, S., Png, K., Sankaran,
531 M., Hou, X.Y., Zhou, H.K., Ma, L., Ren, W.B., Li, X.L., Ding, Y., Li, Y.H., and Shi, H.X.:
532 Combatting global grassland degradation, *Nat. Rev. Earth Env.*, 2(10), 720–735,
533 <https://doi.org/10.1038/s43017-021-00207-2>, 2021.
- 534 Bochet, E., and García-Fayos, P.: Factors controlling vegetation establishment and water erosion on
535 motorway slopes in Valencia, Spain, *Restor. Ecol.*, 12(2), 166–174, [https://doi.org/](https://doi.org/10.1111/j.1061-2971.2004.0325.x)
536 [10.1111/j.1061-2971.2004.0325.x](https://doi.org/10.1111/j.1061-2971.2004.0325.x), 2004.
- 537 Bochet, E., Poesen, J., and Rubio, J.L.: Runoff and soil loss under individual plants of a semi-arid
538 Mediterranean shrubland: influence of plant morphology and rainfall intensity, *Earth Surf. Proc.*
539 *Land.*, 31(5), 536–549, <https://doi.org/10.1002/esp.1351>, 2006.
- 540 Chen, S., Ma, X., Zhang, X., and Chen, Z.: Genetic variation and geographical divergence in *Elymus*
541 *nutans* Griseb. (Poaceae: Triticeae) from West China, *Biochem. Syst. Ecol.*, 37(6), 716–722,
542 <https://doi.org/10.1016/j.bse.2009.12.005>, 2009.
- 543 Cui, Z., Liu, Y.F., Liu, Y., Leite, P.A.M., Shi, J.J., Shi, Z.H., and Wu, G.L.: Fragmentation alters the
544 soil water conservation capacity of hillside alpine meadows on the Qinghai-Tibetan Plateau,
545 *Geoderma*, 428, 116133, <https://doi.org/10.1016/j.geoderma.2022.116133>, 2022.
- 546 [Cuo, L., Zhang, Y.X., Zhu, F.X., and Liang, L.Q.: Characteristics and changes of streamflow on the](https://doi.org/10.1016/j.ejrh.2014.08.004)
547 [Tibetan Plateau: A review, *J. Hydrol-Reg. Stud.*, 2, 2014, 49–68,](https://doi.org/10.1016/j.ejrh.2014.08.004)
548 [https://doi.org/10.1016/j.ejrh.2014.08.004, 2014](https://doi.org/10.1016/j.ejrh.2014.08.004)
- 549 De Baets, S., Poesen, J., Knapen, A., Barberá, G.G., and Navarro, J.A.: Root characteristics of
550 representative Mediterranean plant species and their erosion-reducing potential during

551 concentrated runoff, *Plant Soil*, 294(1–2), 169–183, [https://doi.org/10.1007/s11104-007-9244-](https://doi.org/10.1007/s11104-007-9244-2)
552 2, 2007.

553 [Dijk, A. I. J. M., and Keenan, R. J: Planted forests and water in per spective. *Forest Ecol. Manag.*,](#)
554 [251, 1–9, <https://doi.org/10.1016/j.foreco.2007.06.010>, 2007.](#)

555 Dos Santos, J.C.N., de Andrade, E.M., Medeiros, P.H.A., Guerreiro, M.J.S., and de Queiroz Palácio,
556 H.A.: Effect of Rainfall Characteristics on Runoff and Water Erosion for Different Land Uses
557 in a Tropical Semiarid Region, *Water Resour. Manag.*, 31(1), 173–185, [https://doi.org/](https://doi.org/10.1007/s11269-016-1517-1)
558 10.1007/s11269-016-1517-1, 2017.

559 [Fisher, R.J., Sawa, B., and Prieto, B.: A novel technique using LiDAR to identify native-dominated](#)
560 [and tame-dominated grasslands in Canada, *Remote Sens. Environ.*, 218, 201-206,](#)
561 <https://doi.org/10.1016/j.rse.2018.10.003>, 2018.

562 Freschet, G.T., and Roumet, C.: Sampling roots to capture plant and soil functions, *Funct. Ecol.*, 31(8),
563 1506–1518, <https://doi.org/10.1111/1365-2435.12883>, 2017.

564 Gang, C.C., Zhou, W., Chen, Y.Z., Wang, Z.Q., Sun, Z.G., Li, J.L., Qi, J.G., and Odeh, I.:
565 Quantitative assessment of the contributions of climate change and human activities on global
566 grassland degradation, *Environ. Earth Sci.*, 72(11), 4273–4282, [https://doi.org/10.1007/s12665-](https://doi.org/10.1007/s12665-014-3322-6)
567 014-3322-6, 2014.

568 [Grman, E. Zirbel, C.R., Bauer, J.T., Groves, A.M., Bassett, T., and Brudvig, L.A.: Super - abundant](#)
569 [C₄ grasses are a mixed blessing in restored prairies, *Restor. Ecol.*, 29\(S1\), e13281.1–e13281.8,](#)
570 <https://doi.org/10.1111/rec.13281>, 2021.

571 Gysse, G., Poesen, J., Bochet, E., and Li, Y., Impact of plant roots on the resistance of soils to
572 erosion by water: a review, *Progr. Phys. Geogr.*, 29(2), 189–217, [https://doi.org/10.1191/](https://doi.org/10.1191/0309133305pp443ra)
573 0309133305pp443ra, 2005.

- 574 ~~Han, X., Li, Y.H., Du, X.F., Li, Y.B., Wang, Z.W., 1, Jiang S.W., and Li, Q.: Effect of grassland~~
575 ~~degradation on soil quality and soil biotic community in a semi arid temperate steppe, Ecol.~~
576 ~~Process., 9, 63, <https://doi.org/10.1186/s13717-020-00256-3>, 2020.~~
- 577 Huang, Z., Tian, F.P., Wu, G.L., Liu, Y., and Dang, Z.Q.: Legume grasslands promote precipitation
578 infiltration better than gramineous grasslands in arid regions, Land Degrad. Dev., 28(1), 309–
579 316, <https://doi.org/10.1002/ldr.2635>, 2017.
- 580 Huang, Z., Liu, Y.F., Cui, Z., Liu, Y., Wang, D., Tian, F.P., and Wu, G.L.: Natural grasslands
581 maintain soil water sustainability better than planted grasslands in arid areas, Agr. Ecosyst.
582 Environ., 286(1), 106683, <https://doi.org/10.1016/j.agee.2019.106683>, 2019.
- 583 Karayel, D., and Šarauskis, E.: Environmental impact of no-tillage farming, Environ. Res. Eng.
584 Manag., 75(1), 7–12, <http://dx.doi.org/10.5755/j01.arem.75.1.20861>, 2019.
- 585 ~~Li, W., Wang, J.L., Zhang, X.J., Shangli, S., and Wenxia, C.: Effect of degradation and rebuilding of~~
586 ~~artificial grasslands on soil respiration and carbon and nitrogen pools on an alpine meadow of~~
587 ~~the Qinghai Tibetan Plateau, Ecol. Eng., 111, 134–142, [https://doi.org/10.1016/j.ecoleng.](https://doi.org/10.1016/j.ecoleng.2017.10.013)~~
588 ~~2017.10.013, 2018.~~
- 589 Leung, A.K., Garg, A., Coe, J.L., Ng, C.W.W., and Hau, B.C.H.: Effects of the roots of *Cynodon*
590 *dactylon* and *Schefflera heptaphylla* on water infiltration rate and soil hydraulic conductivity,
591 Hydrol. Process., 29(15), 3342–3354. <https://doi.org/10.1002/hyp.1045>, 2015.
- 592 Liu, W.J., Luo, Q.P., Lu, H.J., Wu, J.E., and Duan, W.P.: The effect of litter layer on controlling
593 surface runoff and erosion in rubber plantations on tropical mountain slopes, SW China, Catena,
594 149, 167–175, <https://doi.org/10.1016/j.catena.2016.09.013>, 2017.
- 595 Liu, Y., Li, S.Y., Niu, Y.L., Cui, Z., Zhang, Z.C., Wang, Y.L., Ma, Y.S., L'opez-Vicente, M., and
596 Wu, G.L.: Effectiveness of mixed cultivated grasslands to reduce sediment concentration in

597 runoff on hillslopes in the Qinghai-Tibetan Plateau, *Geoderma*, 422, 115933, [https://doi.org/](https://doi.org/10.1016/j.geoderma.2022.115933)
598 [10.1016/j.geoderma.2022.115933](https://doi.org/10.1016/j.geoderma.2022.115933), 2022.

599 Liu, Y.F., Dunkerley, D., López-Vicente, M., Shi, Z.H., and Wu, G.L.: Trade-off between surface
600 runoff and soil erosion during the implementation of ecological restoration programs in semiarid
601 regions: A meta-analysis, *Sci. Total Environ.*, 712, 136477, [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.scitotenv.2019.136477)
602 [scitotenv.2019.136477](https://doi.org/10.1016/j.scitotenv.2019.136477), 2020.

603 Liu, Y.F., Liu, Y., Wu, G.L., and Shi, Z.H.: Runoff maintenance and sediment reduction of different
604 grasslands based on simulated rainfall experiments., *J. Hydrol.*, 572, 329–335, [https://doi.org/10.](https://doi.org/10.1016/j.jhydrol.2019.03.008)
605 [1016/j.jhydrol.2019.03.008](https://doi.org/10.1016/j.jhydrol.2019.03.008), 2019.

606 Liu, Y., Zhao, L.R., Liu, Y.F., Huang, Z., Shi, J.J., Wang, Y.L., Ma, Y.S., Lucas-Borja, M.E., López-
607 Vicente, M., and Wu, G.L.: Restoration of a hillslope grassland with an ecological grass species
608 (*Elymus tangutorum*) favors rainfall interception and water infiltration and reduces soil loss on
609 the Qinghai-Tibetan Plateau, *Catena*, 219, 106632, [https://doi.org/10.1016/j.catena.2022.](https://doi.org/10.1016/j.catena.2022.106632)
610 [106632](https://doi.org/10.1016/j.catena.2022.106632), 2022.

611 [López-Vicente, M., and Navas, A.: A new distributed rainfall-runoff \(DR2\) model based on soil](https://doi.org/10.1016/j.agwat.2011.12.007)
612 [saturation and runoff cumulative processes, *Agricultural Water Management*, 104, 128–141.](https://doi.org/10.1016/j.agwat.2011.12.007)
613 <https://doi.org/10.1016/j.agwat.2011.12.007>, 2012.

614 Lu, J.R., Zhang, Q., Werner, A.D., Li, Y.L., Jiang, S.Y., and Ta, Z.Q.: Root-induced changes of soil
615 hydraulic properties – a review. *J. Hydrol.*, 589, 125203 [https://doi.](https://doi.org/10.1016/j.jhydrol.2020.125203)
616 [org/10.1016/j.jhydrol.2020.125203](https://doi.org/10.1016/j.jhydrol.2020.125203), 2020.

617 Ma, Y.L., Liu, Y.F., López-Vicente, M., and Wu G.L.: Divergent shift of normal alpine meadow
618 towards shrub and degraded meadows reduces soil water retention and storage capacity, *J.*
619 *Hydrol.*, 625, 130109, <https://doi.org/10.1016/j.jhydrol.2023.130109>, 2023^a.

620 [Ma, Y.L., Liu, Y.F., López-Vicente, M., and Wu, G.L.: Divergent shift of normal alpine meadow](#)
621 [exacerbated soil loss of hillslope alpine meadows based on field experiments. *Int. Soil Water*](#)
622 [Conse. Res., In Press, <https://doi.org/10.1016/j.iswcr.2023.11.007>, 2023b.](#)

623 Minea, G., Mititelu-Ionuș, O., Gyasi-Agyei, Y., Ciobotaru, N., and Rodrigo-Comino, J.: Impacts of
624 grazing by small ruminants on hillslope hydrological processes: A review of European current
625 understanding, *Water Resour. Res.*, 58, e2021WR030716, <https://doi.org/10.1029/2021WR>,
626 2022.

627 Mohamadi, M.A., and Kavian, A.: Effects of rainfall patterns on runoff and soil erosion in field plots,
628 *Int. Soil Water Conse.*, 3(4), 273–281, <http://dx.doi.org/10.1016/j.iswcr.2015.10.001>, 2015.

629 Mohammad, A.G., and Adam, M.A.: The impact of vegetative cover type on runoff and soil erosion
630 under different land uses, *Catena*, 81(2), 97–103, <http://dx.doi.org/10.1016/j.catena.2010.01.008>,
631 2010.

632 Niu, Y.L., Li, S.Y., Liu, Y., Shi, J.J., Wang, Y.L., Ma, Y.S., and Wu, G.L.: Regulation of alpine
633 meadow patch coverage on runoff and sediment under natural rainfall on the eastern Qinghai-
634 Tibetan Plateau, *J. Hydrol.*, 603, 127101, <https://doi.org/10.1016/j.jhydrol.2021.127101>, 2021.

635 O'Mara, F.P.: The role of grasslands in food security and climate change, *Ann. Bot-London* 110(6),
636 1263–1270, <https://doi.org/10.1093/aob/mcs209>, 2012.

637 Qiu, D.X., Xu, R.R., Wu, C.X., Mu, X.M., Zhao, G.J., and Gao P.: Vegetation restoration improves
638 soil hydrological properties by regulating soil physicochemical properties in the Loess Plateau,
639 China, *J. Hydrol.*, 609, 127730, <https://doi.org/10.1016/j.jhydrol.2022.127730>, 2022.

640 Ren, F., Zhou, H.K., Zhao, X.Q., Han, F., Shi, L.N., Duan, J.C., and Zhao, J.Z.: Influence of simulated
641 warming using OTC on physiological–biochemical characteristics of *Elymus nutans* in alpine
642 meadow on Qinghai-Tibetan plateau, *Acta Ecol. Sinica*, 30(3), 166–171, <https://doi.org/>

643 10.1016/j.chnaes.2010.04.007, 2010.

644 Robinson, M., Cognard-Plancq, A.L., Cosandey, C., David, J., Durand, P., Führer, H.W., Hall, R.,
645 Hendriques, M.O., Marc, V., McCarthy, R., McDonnell, M., Martin, C., Nisbet, T., O’Dea, P.,
646 Rodgers, M., and Zollner, A.: Studies of the impact of forests on peak flows and baseflows: a
647 European perspective, *For. Ecol. Manag.*, 186, 85–97. [https://doi.org/10.1016/s0378-](https://doi.org/10.1016/s0378-1127(03)00238-x)
648 1127(03)00238-x, 2003.

649 Rodrigo-Comino, J., Brevik, E.C., and Cerdà, A.: The age of vines as a controlling factor of soil
650 erosion processes in mediterranean vineyards, *Sci. Total Environ.*, 616–617, 1163–1173,
651 <https://doi.org/10.1016/j.scitotenv.2017.10.204>, 2018.

652 Rumynin, V.G.: *Surface Runoff Generation, Vertical Infiltration and Subsurface Lateral Flow*. In:
653 Overland flow dynamics and solute transport. Theory and Applications of Transport in Porous
654 Media, vol 26. Springer, Cham, https://doi.org/10.1007/978-3-319-21801-4_1, 2015.

655 Saxton, K.E., and Rawls, W.J.: Soil water characteristic estimates by texture and organic matter for
656 hydrologic solutions, *Soil Sci. Soc. Am. J.*, 70(5), 1569–1578,
657 <https://doi.org/10.2136/sssaj2005.0117>, 2006.

658 Schwarz, M., Rist, A., Cohen, D., Giadrossich, F., Egorov, P., Büttner, D., Stolz, M., and Thormann,
659 J.J.: Root reinforcement of soils under compression, *J. Geophys. Res-Earth*, 120(10), 2103–2120,
660 <https://doi.org/10.1002/2015JF003632>, 2015.

661 Shang, Z.H., Ma, Y.S., Long, R.J., and Ding, L.M.: Effect of fencing, artificial seeding and
662 abandonment on vegetation composition and dynamics of ‘black soil land’ in the headwaters of
663 the Yangtze and the Yellow Rivers of the Qinghai-Tibetan Plateau, *Land Degrad. Dev.*, 19(5),
664 554–563, <https://doi.org/10.1002/ldr.861>, 2008.

665 [Schmidt, S., Alewell, C., and Meusburger, K.: Monthly RUSLE soil erosion risk of Swiss grasslands.](#)

666 [J. Maps, 15, 247–256, https://doi.org/10.1080/ 17445647.2019.1585980, 2019](https://doi.org/10.1080/17445647.2019.1585980)

667 Török, P., Brudvig, L.A., Kollmann, J., Price, J., and Tóthmérész, B.: The present and future of
668 grassland restoration, *Restor. Ecol.*, 29(S1), e13378, <https://doi.org/10.1111/rec.13378>, 2021.

669 [Wick, A.F., Geaumont, B.A., Sedivec, K.K., and Hendrickson, J.: *Grassland degradation*. In:
670 \[Shroder, J.F., Sivanpillai, R. \\(Eds.\\), *Biological and environmental hazards, risks, and disasters*.
671 \\[Elsevier, pp. 257–276, 10.1016/B978-0-12-394847-2.00016-4, 2016.\\]\\(#\\)\]\(#\)](#)

672 Wang, B., Zhang, G., Yang, Y., Li, P., and Liu, J.: Response of soil detachment capacity to plant root
673 and soil properties in typical grasslands on the Loess Plateau, *Agr. Ecosyst. Environ.*, 266, 68–
674 75, <https://doi.org/10.1016/j.agee.2018.07.016>, 2018.

675 Wang, C.T., Wang, G.X., Liu, W., Wang, Y., Hu, L., and Ma, L.: Effects of establishing an artificial
676 grassland on vegetation characteristics and soil quality in a degraded meadow, *Isr. Ecol. Evol.*,
677 59(3), 141–153, <http://dx.doi.org/10.1080/15659801.2013.863669>, 2013.

678 Wang, L., Liang, T., and Zhang, Q.: Laboratory experiments of phosphorus loss with surface runoff
679 during simulated rainfall, *Environ. Earth Sci.*, 70(6), 2839–2846, [http://dx.doi.org/10.1007/
680 s12665-013-2344-9](http://dx.doi.org/10.1007/s12665-013-2344-9), 2013.

681 Wang, Z.Q., Zhang, Y.Z., Yang, Y., Zhou, W., Gang, C.C., Zhang, Y., Li, J.L., An, R., Wang, K.,
682 Odeh, I., and Qi, J.G.: Quantitative assess the driving forces on the grassland degradation in the
683 Qinghai–Tibet Plateau, in China, *Ecol. Inform.*, 33, 32–44, [http://dx.doi.org/10.1016/j.ecoinf.
684 2016.03.006](http://dx.doi.org/10.1016/j.ecoinf.2016.03.006), 2016.

685 Wu, G. L., Huang, Z., Liu, Y.F., Cui, Z., Chang, X.F., Tian, F.P., López-Vicented, M., and Shi, Z.H.:
686 Soil water response of plant functional groups along an artificial legume grassland succession
687 under semi-arid conditions, *Agr. Forest Meteorol.*, 278, 107670.
688 <https://doi.org/10.1016/j.agrformet.2019.107670>, 2019.

- 689 [Wu, G.L., Liu, Y.F., Cui, Z., Liu, Y., Shi, Z.H., Yin, R., and Kardol, P.: Trade-off between vegetation](#)
690 [type, soil erosion control and surface water in global semi-arid regions: A meta-analysis, J. Appl.](#)
691 [Ecol., 57, 875–885.https://doi.org/10.1111/1365-2664.13597, 2020.](#)
- 692 Wu, G.L., Liu, Z.H., Zhang, L., Hu, T., and Chen, J.: Effects of artificial grassland establishment on
693 soil nutrients and carbon properties in a black-soil-type degraded grassland, *Plant Soil*, 333(1–
694 2), 469–479, <https://doi.org/10.1007/s11104-010-0363-9>, 2010.
- 695 Xia, L., Song, X.Y., Fu, N., Cui, S.Y., Li, L.J., Li, H.Y., and Li, Y.L.: Effects of forest litter cover on
696 hydrological response of hillslopes in the Loess Plateau of China, *Catena*, 181, 104076,
697 <https://doi.org/10.1016/j.catena.2019.104076>, 2019.
- 698 Xu, J.: A cave $\delta^{18}\text{O}$ based 1800-year reconstruction of sediment load and streamflow: The Yellow
699 River source area, *Catena*, 161, 137–147, <http://dx.doi.org/10.1016/j.catena.2017.09.028>, 2018.
- 700 Zhang, G.H., Tang, K.M., Ren, Z.P., and Zhang, X.C.: Impact of grass root mass density on soil
701 detachment capacity by concentrated flow on steep slopes, *T. ASABE*, (56), 927–934, 2013.
- 702 [Zhao, X., Huang, J., Wu, P., Gao, X.: The dynamic effects of pastures and crop on runoff and](#)
703 [sediments reduction at loess slopes under simulated rainfall conditions. *Catena* 119, 1–7.](#)
704 <https://doi.org/10.1016/j.catena.2014.03.001>, 2014.
- 705 [Zhao, Y.T., Pu, Y.F., Lin, H.L., and Tang, R.: Examining soil erosion responses to grassland](#)
706 [conversation policy in Three-River Headwaters, China, *Sustainability*, 13\(5\), 2702,](#)
707 <https://doi.org/10.3390/su13052702>, 2021.
- 708 Zhou, J., Fu, B.J., Gao, G.Y., Lü, Y.H., Liu, Y., Lü, N., and Wang, S.: Effects of precipitation and
709 restoration vegetation on soil erosion in a semi-arid environment in the Loess Plateau, China,
710 *Catena*, 137, 1–11, <https://doi.org/10.1016/j.catena.2015.08.015>, 2016.
- 711 Zhu, P.Z., Zhang, G.H., Wang, H.X., Yang, H.Y., Zhang, B.J., and Wang, L.L.: Effectiveness of

712 typical plant communities in controlling runoff and soil erosion on steep gully slopes on the
713 Loess Plateau of China, *J. Hydrol.*, 602, 126714. <https://doi.org/10.1016/j.jhydrol.2021.126714>,
714 2021.

715 **Table 1.** Results of path analysis of the factors affecting runoff depth.

Influence factor	Direct path coefficient	Indirect path coefficient						Sum of path coefficient	
		<i>RI</i> ₆₀	<i>ARI</i>	<i>RD</i>	<i>P</i>	<i>VC</i>	<i>LB</i>		Total
<i>RI</i> ₆₀	0.24*		0.25	-0.09	-0.11	0.02	0.00	0.07	0.31
<i>ARI</i>	0.37**	0.16		-0.34	-0.05	0.02	0.02	-0.19	0.18
<i>RD</i>	0.67**	-0.03	-0.18		-0.08	0.03	-0.03	-0.31	0.36
<i>P</i>	-0.18**	0.14	0.10	0.31		0.02	0.00	0.57	0.40
<i>VC</i>	0.29**	0.01	0.03	0.06	-0.01		-0.06	0.03	0.32
<i>LB</i>	-0.12	0.01	-0.09	0.18	0.00	0.15		0.25	0.13


716 Note: *RI*₆₀ is maximum 60-minute intensity (mm h⁻¹), *ARI* is average rainfall intensity (mm h⁻¹), *RD*
717 is rainfall duration (h), *P* is rainfall amount (mm), *VC* is vegetation coverage (%), *LB* is litter biomass
718 (g m⁻²). * means the correlation is significant at 0.05 significance level, and ** means the correlation
719 is significant at 0.01 significance level.

720


721 **Table 2.** Results of path analysis of the factors affecting soil loss per unit area.

Influence factor	Direct path coefficient	Indirect path coefficient								Sum of path coefficient
		<i>R</i>	<i>RI</i> ₆₀	<i>ARI</i>	<i>RD</i>	<i>P</i>	<i>VC</i>	<i>LB</i>	Total	
<i>R</i>	0.60**		-0.12	0.01	-0.10	0.11	0.01	0.01	-0.08	0.52
<i>RI</i> ₆₀	-0.29**	0.24		0.02	0.07	0.16	0.00	0.00	0.49	0.20
<i>ARI</i>	0.04	0.13	-0.19		0.21	0.07	0.01	0.02	0.25	0.28
<i>RD</i>	-0.41**	0.15	0.05	-0.02		0.13	0.00	-0.04	0.27	-0.13
<i>P</i>	0.28**	0.24	-0.17	0.01	-0.19		0.00	-0.01	-0.11	0.17
<i>VC</i>	0.03	-0.04	-0.04	0.01	-0.03	0.03		-0.06	-0.12	-0.10
<i>LB</i>	-0.10	-0.01	-0.01	-0.01	-0.16	0.03	0.02		-0.15	-0.25

722 Note: *R* is surface runoff (mm), *RI*₆₀ is maximum 60-minute intensity (mm h⁻¹), *ARI* is average rainfall
723 intensity (mm h⁻¹), *RD* is rainfall duration (h), *P* is rainfall amount (mm), *VC* is vegetation coverage
724 (%), *LB* is litter biomass (g m⁻²). ** means the correlation is significant at 0.01 significance level.



725 **Figure 1.** The location of the study area on the Qinghai-Tibetan Plateau, and the location of runoff
 726 plots in the study area. (a) The location of the study area, (b) the fragmenting mattic epipedon
 727 epipedon on the alpine hillslope and (c) severely degraded meadows formed by the disappearance of
 728 mattic epipedon-epipedon and (d) four runoff plots of severely degraded meadows (SEMSDM)
 729 and mixed-cultivated grasslands. A typical severely degraded meadow with a slope of 20° was
 730 selected to plant mixed grasses. Runoff plots were photographed with a drone in the early stages of
 731 the 2022 growing season. *DE*, *Deschampsia cespitosa* and *Elymus nutans*; *PE*, *Poa pratensis L.cv.*
 732 Qinghai and *Elymus nutans*; and *PD*, *Poa pratensis L.cv.* Qinghai and *Deschampsia cespitosa*.



733


734 **Figure 2.** Changes in vegetation cover under various mixed-cultivated grasslands from 2019 to 2022.

735 Different capital letters mean that differences were significant in different years for the same

736 grassland community, and different lowercase letters mean that differences were significant between


737 different communities in the same year.

738




739 **Figure 3.** Changes in soil erosion and runoff under various mixed-cultivated grasslands from 2019 to
740 2022. (a) Runoff depth and (b) soil loss per unit area. Note: For the four treatment runoff plots, runoff
741 and sediment were measured 14, 18, and 10 times, respectively, during the growing season of 2019,
742 2020, and 2022. Different capital letters mean that differences were significant in different years for
743 the same grassland community, and different lowercase letters mean that differences were significant
744 between different communities in the same year. *SDM*, severely degraded meadows, *DE*,
745 *Deschampsia cespitosa* and *Elymus nutans*; *PE*, *Poa pratensis* L.cv. Qinghai and *Elymus nutans*; and
746 *PD*, *Poa pratensis* L.cv. Qinghai and *Deschampsia cespitosa*. The lines in the middle of the box
747 represent the median values. The squares in the box represent the average value.

748



749 **Figure 4.** Runoff, soil loss and sediment concentration reduction ratio under different mixed-
 750 cultivated grasslands from 2019 to 2022. (a) Runoff reduction ratio ($RR\bar{E}B$), (b) soil loss reduction
 751 ratio ($SR\bar{E}B$), (c) sediment concentration reduction ratio ($CR\bar{E}B$) and (d) the percent of runoff
 752 reduction ratio to soil loss reduction ratio ($RRSR$). Note: Different capital letters mean that differences
 753 were significant in different years for the same grassland community, and different lowercase letters
 754 mean that differences were significant between different communities in the same year. [The lines in](#)
 755 [the middle of the box represent the median values. The squares in the box represent the average value.](#)
 756



757 **Figure 5.** Changes in bulk density, root mass density, field capacity, capillary capacity, non-capillary
 758 porosity and soil cohesion in 0-10 cm soil layer when severely degraded meadow (*SDM*) were
 759 restored to mixed-cultivated grassland for 4 years. *DE*, *Deschampsia cespitosa* and *Elymus nutans*;
 760 *PE*, *Poa pratensis L.cv. Qinghai* and *Elymus nutans*; and *PD*, *Poa pratensis L.cv. Qinghai* and
 761 *Deschampsia cespitosa*. Percentages represent the increased rate of soil properties (increased rate =
 762 $(V_{DE}$ or V_{PE} or $V_{PD} - V_{SDM})/V_{SDM}$), where V_{SDM} , V_{DE} , V_{PE} and V_{PD} are the mean values of soil
 763 characteristics of *SDM*, *DE*, *PE* and *PD*. Different lowercase letters mean that differences were
 764 significant between different communities. The lines in the middle of the box represent the median
 765 values. The squares in the box represent the average value.