

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

## **Changes in Mediterranean flood processes and seasonality**

Yves Tramblay 1\*

Patrick Arnaud 2

Guillaume Artigue 1

Michel Lang 3

Emmanuel Paquet 4

Luc Neppel 1

Eric Sauquet 3

1 HSM, Univ. Montpellier, CNRS, IRD, IMT-Mines Alès, Montpellier, France

2 RECOVER, INRAE, Aix Marseille Université, Aix-en-Provence, France

3 RiverLy, INRAE, Villeurbanne, France

4 EDF-DTG, Saint Martin le Vinoux, France

\* corresponding author, HydroSciences-Montpellier, Hydropolis, Faculté de Pharmacie, 15 Avenue Charles Flahaut, 34000 Montpellier. [yves.tramblay@ird.fr](mailto:yves.tramblay@ird.fr)

REVISED MANUSCRIPT WITH MARKS

33

## 34 Abstract

35

36 Floods are a major natural hazard in the Mediterranean region, causing deaths and  
37 extensive damages. Recent studies have shown that intense rainfall events are becoming  
38 more extreme in this region, but paradoxically without leading to an increase in the severity  
39 of floods. Consequently, it is important to understand how flood events are changing to  
40 explain this absence of trends in flood magnitude despite increased rainfall extremes. A  
41 database of 98 stations in Southern France with an average record of 50 years of daily river  
42 discharge data between ~~1958-1959~~ and 2021 was considered, together with a high-  
43 resolution reanalysis product providing precipitation and simulated soil moisture. Flood  
44 events, corresponding to an average occurrence of one event per year (5317 events in  
45 total), were extracted and classified into excess rainfall, short rainfall and long rainfall event  
46 types. ~~Further, s~~Several flood event characteristics have been also analyzed/computed:  
47 flood event durations, base flow contribution to floods/the peak, runoff coefficient, total and  
48 maximum event rainfall and antecedent soil moisture. The evolution through time of the  
49 flood event characteristics and seasonality were analyzed. Results indicated that, in most  
50 basins, floods tend to occur earlier during the year, the mean flood date being on average  
51 advanced by one month between 1959-1990 and 1991-2021. This seasonal shift ~~can~~could  
52 be attributed to the increased frequency of southern-circulation weather types during spring  
53 and summer. An increase in total and extreme event precipitation has been observed,  
54 associated with a decrease of antecedent soil moisture before rainfall events, linked to a  
55 smaller contribution of base flow during floods. The majority of flood events are associated  
56 with excess rainfall on saturated soils, but their relative proportion is decreasing over time  
57 notably in spring with a concurrent increased frequency of short rain floods. Therefore, this  
58 study shows that even in the absence of trends, flood properties may change over time and  
59 these changes need to be accounted for when analyzing the long-term evolution of flood  
60 hazards.

61

62

63

64

65

66

67

68

69

70

# 1. Introduction

There is a growing interest in understanding the evolution of floods occurring in different regions in the context of climate change. The recent sixth report of the Intergovernmental Panel on Climate Change (Ali et al., 2022), reported a mixture of observed trends in different Mediterranean countries, with both increasing and decreasing river floods and overall, a low confidence in their attribution to climate change. Several large-scale studies on changes in flood risk (Slater et al., 2021a; Blöschl et al., 2017, 2019) have indicated for the Mediterranean region a possible decrease over the last decades. This difficulty in detecting possible changes in flood hazard is doubtless linked to the fact that there are different types of floods (Tarasova et al., 2019; Berghuijs et al., 2019; Stein et al., 2020; Trambly et al., 2022). Indeed, a distinction can be made between floods associated with soil saturation excess, soil infiltration excess or snowmelt-driven floods and the relative frequency of these different types of floods may change over time (Zhang et al., 2022). Furthermore, these changes can occur at local to regional scales, given complex combinations of climatic and physiographic triggers, making global generalization of changes in flood risk hazardous, if not irrelevant (Whitfield, 2012; Blöschl et al., 2015).

Only a few studies are focusing on changes in flood types over time, while it is important to understand the long-term changes in flood processes to evaluate how flood risk can evolve, in order notably to better adapt the flood mitigation strategies (Merz et al., 2014). The main limitation to perform such studies is the need for long time-series of river discharge, to have large samples of flood events to evaluate how their properties may change over time. For many regions of the world, the lack of observed river discharge data prevents this type of analysis. ~~Only a few~~Some studies attempted to analyze the changes in different types of floods for different regions (Berghuijs et al., 2019; Mao et al., 2019; Kemter et al., 2020; Bertola et al., 2021; Liu et al., 2022; Trambly et al., 2022, Tarasova et al., 2023). Most of these studies rely on flood classification schemes, with various complexity depending on the type of data available, allowing a data-based separation of floods into their distinct generation mechanisms (Tarasova et al., 2019; Berghuijs et al., 2019; Tarasova et al., 2020; Stein et al., 2020, 2021; Trambly et al., 2022). For basins under a Mediterranean climate, several studies reported complex interactions between precipitation increases and lower antecedent soil moisture, leading to thresholds effects (Brunner et al., 201) in the catchment response to changing hydro-climatic conditions (Wasko and Nathan, 2019; Cao et al., 2020; Bertola et al., 2021). Recent large-scale studies (Jiang et al., 2022, Tarasova et al., 2023) suggested a reduction of the frequency of floods driven by soil saturation in Europe, including basins in the Mediterranean area.

In French Mediterranean basins, several studies reported an increase in precipitation extremes (Trambly et al., 2013; Blanchet and Creutin, 2022; Ribes et al., 2019) that did not translate into increased floods (Trambly et al., 2019). It is hypothesized that, as many regions of the world, a decrease in soil moisture linked with a greater aridity can potentially offset the increase in precipitation extremes and thus not increase flood severity (Sharma et al., 2018; Trambly et al., 2019; Wasko and Nathan, 2019; Wasko et al., 2021; Huang et al., 2022). Excess soil moisture was previously identified as an important flood driver notably in the Mediterranean (Kemter et al., 2020; Bertola et al., 2021), indicating that they can play an important role. Yet, beside ~~the~~ trend detection or changes in flood types, no study has

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

118 provided an in-depth analysis of the joint long-term evolution of flood processes in  
119 Mediterranean basins~~these regions~~, in relation to their drivers such as precipitation, soil  
120 moisture and the evolution of synoptic weather patterns associated with floods. Therefore,  
121 the objective of the study is to evaluate how the characteristics of Mediterranean floods are  
122 evolving in time, not only in terms of flood event types, but also the seasonality of events,  
123 their duration, runoff coefficients, and antecedent and antecedent soil moisture. A recent  
124 study (Tramblay et al., 2019) indicated no significant trends on flood hazards for a large  
125 ensemble of basins located in southern France. This database is used herein to further  
126 analyze the possible changes in flood generating processes and in the seasonality of flood  
127 events.

Code de champ modifié

128  
129 In section 2 are presented the different datasets used for this analysis, including river  
130 discharge, precipitation, soil moisture data and weather types classification. In section 3 the  
131 methods are presented, for event extraction, analysis of the seasonality and changes in the  
132 different flood drivers. Results are presented in section 4.

## 134 2. Data

135  
136 We consider 98 catchments in southern France (Figure 1) where the time series of daily  
137 discharge exceeds 30 years of complete data between 1959 and 2021 (Tramblay et al.,  
138 2019). Among them, 48 basins have more than 50 years of data and the basins selected are  
139 not influenced by reservoir or dam regulation. The catchment sizes are ranging from 14 km<sup>2</sup>  
140 to 3195 km<sup>2</sup>, with a mean size equal to 480 km<sup>2</sup> (see Table t1 in supplementary materials).  
141 Basins with a nival regime were removed, identified from the river discharge hydrographs  
142 and removing basins with more than 20% of precipitation falling as snow. In addition to river  
143 discharge data, the precipitation and soil moisture for each basin has been retrieved from  
144 the SAFRAN-ISBA-MODCOU (SIM) reanalysis covering the whole France territory at 8 km  
145 spatial resolution (Vidal et al., 2010; Habets et al., 2008). Precipitation and soil moisture data  
146 have been extracted and averaged at the catchment scale. The soil moisture data extracted  
147 from SIM is a soil wetness index obtained from the normalization of the volumetric soil  
148 moisture content with the wilting point and field capacity, that ranges between 0 and 1. Land  
149 cover classes (forest, agricultural, urban) corresponding to 2018 have been extracted from  
150 the CORINE land cover inventory (Büttner, 2014). In addition, we used the weather type  
151 classification from *Electricité de France* (EDF), corresponding to a daily classification into 8  
152 synoptic situations associated with rainfall events over France (Garavaglia et al., 2010). This  
153 classification is built on geopotential heights at 700 and 1000 hPa pressure levels associated  
154 with rainy days over France.

Code de champ modifié

Code de champ modifié

Code de champ modifié

## 156 3. Methods

### 158 3.1 Extraction of flood events

159  
160 We extracted a sample of flood events with a mean occurrence of 1 event per year using a  
161 peaks-over-threshold approach. This type of sampling is chosen since low annual maximum  
162 runoff could be observed during dry years (Farquharson et al., 1992). A de-clustering  
163 algorithm is applied to identify single events to avoid introducing autocorrelation in the  
164 analysis and ensuring that flood events are independent, using two rules (Lang et al., 1999):

Code de champ modifié

Code de champ modifié

165 first a minimum of  $n$  days between events, with  $n = 5 + \log(\text{catchment area})$  and second,  
166 between two consecutive peaks, runoff must drop below  $\frac{2}{3}$  of the smallest peak. The  
167 maximum daily runoff of each event is kept. This means that for an event lasting several  
168 days, only the maximum daily discharge, and the corresponding date, are kept. ~~Then, f~~For  
169 each flood event, we ~~computed~~extracted the total rainfall and maximum rainfall. The  $n$ -day  
170 previous precipitation is extracted. Total rainfall for each event is estimated by a cumulative  
171 sum of precipitation ~~starting the day of the flood and before a flood and~~ this aggregation  
172 stops if there are two consecutive days with precipitation close to zero (1 mm) to account for  
173 rainfall intermittency within events. The maximum daily precipitation is extracted from the  
174 same time interval used to compute total event precipitation. The soil moisture at the  
175 beginning of the events is extracted from the previous day of the start of the rainfall event. A  
176 base flow filter has been used to separate direct runoff and base flow for each ~~event~~time  
177 series, using the Lyne Hollick Filter (Lyne and Hollick, 1979), with its default parameters. For  
178 each flood event, the base flow corresponding to the peak has been extracted to estimate  
179 the direct runoff, corresponding to the event rainfall contribution, in addition to base flow.  
180 Different metrics characterizing each flood event have been computed: total rainfall (mm),  
181 event maximum rainfall (mm), duration of the rainfall event (days), duration of the flood event  
182 (days), antecedent soil moisture (0-1) and runoff coefficient (0-1). The runoff coefficient was  
183 computed for each event as the ratio of direct runoff depth and total event precipitation

### 185 3.2 Analysis of the mean date of occurrence

186  
187 Circular statistics (Burn, 1997; Berens, 2009) are used to analyze flood timing. The dates are  
188 first converted into an angular value, then from this sample of angular values, the mean date  
189 of occurrence ( $\theta$ ) can be computed, together with the concentration index ( $r$ ) which  
190 measures the variability of the flood occurrences around the mean date. Using the dates of  
191 flood events, considering hydrological years starting September 1,  $\theta$  and  $r$  are computed  
192 from the sample of dates. The first step in the analysis of seasonality is to test against  
193 circular uniformity. Circular uniformity refers to the case in which all angular values of flood  
194 dates around the circle are equally likely, indicative of the absence of flood seasonality. In  
195 that case, the computation of the mean date would have little relevance. The Rayleigh  
196 (Fisher, 1993) and the Hermans-Rasson (Landler et al., 2019) tests are used to test against  
197 uniformity for unimodal distributions, to verify the presence of flood seasonality (ie. meaning  
198 that floods do not occur randomly throughout the year). To associate flood events and  
199 weather types, for each rainy day corresponding to flood events, the weather type has been  
200 extracted from the weather type's classification.

### 202 3.3 Classification of flood generating processes

203  
204 A classification is applied to the flood events, adapted from a previously implemented  
205 classification at the global scale (Stein et al., 2020), in the United States (Stein et al., 2021)  
206 and Africa (Tramblay et al., 2022). This approach relates the occurrence of rainfall amounts  
207 above various thresholds to the occurrence of floods. Flood events in each catchment are  
208 classified according to three hydrometeorological generating processes, namely, the excess  
209 rainfall, short rainfall, long rainfall using a decision tree. Excess rainfall is defined as a flood  
210 event triggered by rainfall higher than average occurring over wet soils (i.e. soil moisture  
211 above than 50% saturation), short rainfall as a single daily rainfall event above high  
212 thresholds (the 95th percentile computed over the whole time series of rainfall) and long

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

213 rainfall as several consecutive days (> 2 days) with rainfall above the 95th percentile of  
214 rainfall summed over 7 days. The classification first evaluates if a larger than average multi-  
215 day rainfall fell on wet soil to determine if the flood event was an excess rainfall type of flood.  
216 If that was not the case, it evaluates whether the thresholds for long rainfall and then short  
217 rainfall are exceeded. If no process could be identified, the class “other” is assigned.  
218

### 219 3.4 Changes in flood characteristics

220  
221 To assess the changes over time in flood dates and generating mechanisms, we split the  
222 records of each station into two periods of equal length. Given that most stations have  
223 records starting after the 1960s, on average the first period is ranging from 1959 to 1990 and  
224 the second one from 1991 to 2021, with a pivot year within +/- 5 years around 1991, allowing  
225 the comparison of the two time slices across the different stations. To assess the relative  
226 changes in the flood drivers, the frequency of each driver for each time period has been  
227 computed, and then we computed for each station the relative change (%) in each driver  
228 contribution (Berghuijs et al., 2019). In addition, to detect trends on the long-term frequency  
229 of event types per year pooled at the regional scale, we used the Mann-Kendall test for  
230 trends, modified to account for autocorrelation in the time series (Hamed and Ramachandra  
231 Rao, 1998).  
232

233 We use the same approach to estimate changes in the different flood events characteristics,  
234 and we applied the two-tailed Wilcoxon test to check the difference in medians. In addition,  
235 to assess the regional significance of the trends, we also computed the Mann-Kendall test  
236 on flood events characteristics pooled at the regional scale. For flood dates, we computed  
237 the mean dates of occurrence for the two time periods and assessed the significance of the  
238 difference using the Watson and Williams test, which is a circular analogue to the two  
239 sample t-test (Watson and Williams, 1956). Finally, to estimate potential relationships  
240 between different flood characteristics, the Spearman correlation coefficient ( $\rho$ ) is computed.  
241

## 242 4. Results and discussion

### 243 4.1 Change in flood event characteristics

244  
245 There are several changes in flood event characteristics as seen in Figure 2 between the  
246 two sub-periods, 1959-1990 and 1991-2021. On average, total event precipitation is  
247 increasing in 65 basins (66% of basins), maximum event rainfall is increasing in 76 basins,  
248 consistent with previous studies in this area (Ribes et al., 2019; Trambly et al., 2019;  
249 Blanchet and Creutin, 2022), while on the opposite antecedent soil moisture is decreasing in  
250 71 basins, baseflow contribution is decreasing in 75 basins and runoff coefficient is  
251 decreasing in 68 basins. These changes in soil moisture, base flow and runoff coefficients  
252 are consistent with an overall increase of aridity in southern Europe mostly driven by higher  
253 evapotranspiration (Trambly et al., 2020) and have been also observed in other regions  
254 with a similar climate (Ho et al., 2022). ~~The number of local statistically significant changes~~  
255 ~~for each flood event characteristic are~~ number of local statistically significant changes for  
256 each flood event characteristic is given in Table 1. These numbers remain small but it should  
257 be reminded that sample sizes are quite short for a robust statistical assessment in a context  
258 of high interannual variability. To overcome this issue, we also assessed the regional  
259

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

260 [significance of these changes in flood event characteristics. We performed a regional](#)  
261 [pooling of the events and applied the Mann-Kendall test to detect trends in the regional](#)  
262 [series of event characteristics. As shown in table 1, all the detected changes are regionally](#)  
263 [significant except the decrease in base flow contribution to peak discharge during floods.](#)

264 Overall, an increase in total event rainfall can be observed, mostly caused by the increase of  
265 maximum rainfall during the events (the changes in the two variables are correlated, with  $\rho =$   
266 0.52), while the flood event durations are on average decreasing, consistent with studies at  
267 the global scale (Wasko et al., 2021).

268  
269 These changes in precipitation are associated with a decrease of antecedent soil moisture,  
270 before the beginning of the rainfall events. This decrease is also related to a smaller  
271 contribution of base flow during floods [in some basins](#). There is indeed a significant  
272 correlation between the relative changes in the base flow contribution to peak runoff and soil  
273 moisture ( $\rho = 0.56$ ), indicating that the soil moisture decrease is likely the main driver of  
274 these changes. There is also for most basins a significant nonlinear relationship (exponential  
275 type) between the flood events antecedent soil moisture and runoff coefficients, as reported  
276 in many studies (Penna et al., 2011; Rogger et al., 2013; Raynaud et al., 2015; Tarasova et  
277 al., 2020). Indeed, for the first time period, 1959-1990 the median Spearman correlation  
278 between antecedent soil moisture and flood runoff coefficients [\(see supplementary figure](#)  
279 [S1\)](#) is equal to 0.43 and is significant at the 5% level in 56 basins (67 basins at the 10%  
280 significance level). For the second time period, 1991-2021, the median correlation is  
281 increasing to 0.45 and is significant in 64 basins at the 5% significance level (68 at the 10%  
282 significance level). These results show, contrary to popular belief, that at the catchment  
283 scale drier soils produce less runoff, and this characteristic is even slightly accentuating over  
284 time. Indeed, increased runoff coefficients induced either by hydrophobic soil conditions  
285 following droughts (Burch et al., 1989), soil crusting and sealing (Bissonnais and Singer,  
286 1993) or compaction (Alaoui et al., 2018), are well documented processes that mostly occur  
287 at the local plot scale that do not produce discernible effects at the catchment scale. This  
288 observation is reinforced by the fact that no negative correlation between runoff coefficients  
289 and initial soil moisture was detected. About the explanatory factors of the association  
290 between soil moisture and runoff coefficients, we found stronger correlations (significant at  
291 the 5% level) between these two variables in catchments with higher percentage of urban or  
292 agricultural areas, and on the opposite weaker correlations along with increased percentage  
293 of forests or mean catchment altitude. There is only a very small increase of the correlations  
294 for larger basins (no significant correlation with basin sizes), indicating that this relationship  
295 between soil moisture and runoff coefficient remains valid for all basins scales considered in  
296 the present study. This relationship between runoff coefficients and antecedent soil moisture  
297 remained stable between 1959-1990 and 1991-2021 (Figure 3).

#### 298 **4.2 Changes in flood dates**

299  
300 Floods in southern France tend to occur mainly during November or December for basins  
301 close to the Mediterranean, East of the Cévennes mountainous range, while for basins  
302 located on the western part of the region, they tend to occur later during winter months,  
303 centered in January or February (Figure 4). Both the Rayleigh and Hermans-Rasson tests  
304 reject the null hypothesis of uniformity at the 5% level, indicating that floods do not occur  
305 randomly throughout the year. In most cases, the seasonal distribution is unimodal, except  
306 for a few cases; in about 15 stations the maximum occurrence of floods is observed in late  
307

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

308 autumn or winter and a secondary minor peak of occurrence is observed, usually centered  
309 around the month of March or April. These floods are associated with rainfall events rather  
310 than snowmelt, since for only 3 basins the snowfall contribution reaches 19% of total  
311 precipitation whereas the snowfall contribution is much lower for the remaining 12 basins  
312 (less than 5%).

313  
314 For 79 basins, floods tend to occur earlier during the year, on average by -22 days between  
315 1958-1959-1990 and 1991-2021 (Figure 5, left panel). On the opposite, for 19 basins the  
316 mean flood date occurs later in the second period with an average of +12 days. These  
317 changes in the mean date are significant in 26 basins at the 0.1 level according to the  
318 Watson and Williams test (18 basins at the 0.05 significance level). There are two  
319 differentiated spatial patterns: all basins where floods tend to occur earlier are located  
320 widespread in the center of the Mediterranean region, and basins where floods tend to occur  
321 later are found only in the northwestern margin of the domain. In these basins, the mean  
322 floods occur in late winter, until February and March. The same spatial patterns of changes  
323 in mean flood dates have been observed by Blöschl et al., (2017), but without providing the  
324 possible causes of these seasonal shifts. For the concentration index (i.e., the variability  
325 around the mean date) similarly two different patterns are found: for basins where floods  
326 tend to occur earlier, the concentration index is increasing, meaning more floods are  
327 clustered around the mean date, while for western basins where floods tend to occur later,  
328 the concentration index is decreasing, meaning a larger variability in flood dates (Figure 5,  
329 right panel).

330  
331 To assess the regional changes in flood dates, we first separated in two regional samples  
332 the stations where floods tend to occur earlier (sample 1) or later (sample 2). Then we used  
333 the Watson-William test, previously used to assess changes in flood dates in each station, to  
334 compare these two regional samples. The test results indicate that for the 19 stations where  
335 floods tend to occur later, the change in flood dates are not significant at the 5% level (p  
336 value = 0.0821), on the opposite, for the 79 stations where floods are occurring earlier, the  
337 change is significant (p value = 5.34.10^-8).

### 339 4.3 Associations between flood occurrence and weather patterns

340  
341 The seasonal patterns observed for the floods are closely related to the occurrence of  
342 different weather types in different sub-regions. As shown in figure 6, most basins located  
343 east of the Cévennes mountainous range have floods associated with WT4, Southern  
344 Circulation, and western basins with WT2, ~~Atlantic circulation~~ Steady Oceanic. The most  
345 frequent pattern associated with 37% of floods, WT4, is known to be triggering intense  
346 rainfall events in this region (Ducrocq et al., 2008; Trambly et al., 2013). Interestingly, the  
347 WT6, Eastern circulation, and WT7, Southwestern circulation, are both associated to a  
348 lesser extent with floods across the whole region, but without notable spatial differences in  
349 the relative frequency of floods associated with these weather types. Change in flood  
350 seasonality could ~~seasonality could~~ an be ascribed to changes in the seasonal occurrence of  
351 the weather types (Figure 7): WT4 tends to occur more frequently from March to August  
352 during 1991-2021 compared to 1959-1990, and these changes are statistically significant  
353 (see supplementary figure S2). When looking at the actual count of WT4 days, this change  
354 represents an increase of 69 events during that 6-month period for 1991-2021, so an  
355 average of +2.2 days per year. Associated with a warmer Mediterranean Sea over the last

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié



356 decades notably during summer (Pastor et al., 2020), the combination of these two factors  
357 could explain the earlier occurrence of floods east of the Cévennes mountainous range.  
358 Similarly, there is an ~~an notable~~ increased frequency of WT2 in January, February and March  
359 between 1991-2021 and 1959-1990, ~~that is also significant (supplementary figure S2),~~ that  
360 could be possibly related to the later occurrence of floods west of the Cévennes range.  
361 Although this change in weather types seasonality leading to heavy rainfalls is a plausible  
362 cause of the observed changes in the flood seasonality, more research is needed to better  
363 understand these relationships and attribute changes in flood seasonality. Notably, to  
364 analyze in more detail the moisture supply from the Mediterranean or Atlantic seas, the  
365 interaction with the atmospheric thermodynamics, the duration, localization and the spatial  
366 dependence of the rainfall episodes inducing floods.

Code de champ modifié

#### 368 4.4 Changes in flood generating processes

369  
370 When first applying the classification of flood-generating processes on all floods, we find a  
371 predominance of excess rainfall events (Figure 8), followed by long rain and short rains, that  
372 is consistent with the known flood-generating processes in this region (Mediterranean  
373 episodes) and, in particular, the strong influence of saturated soils on runoff generation with  
374 floods mostly occurring during the autumn (Tramblay et al., 2010, 2019). The category  
375 'other' regroups only 0.97% of ~~floods, floods~~ and it represents mostly events with very low  
376 precipitation amounts, likely due to an underestimation of rainfall in the SAFRAN database  
377 for some events. It is worth noticing that despite the large sample of basins considered, the  
378 patterns are remarkably consistent and homogeneous across different basin sizes and  
379 locations. There is a significant, yet low, correlation ( $\rho = 0.26$ , p-value = 0.008) between the  
380 ratio of excess rain floods and catchment size, with a larger proportion of excess rain in  
381 larger basins, while on the opposite there is an even weaker and negative correlation ( $\rho = -$   
382  $0.16$ , p-value = 0.09) between the ratio of short rain and basin size. It should be noted that  
383 floods driven by short rain episodes are potentially affecting smaller regions than floods  
384 driven by excess floods (Brunner and Dougherty, 2022). For 30 basins (not necessarily the  
385 largest ones), the proportion of excess rain exceeds 80% of the total number of flood events  
386 (see supplementary figure S23). For short rain and long rain, the maximum contributions  
387 observed much lower, are 36% and 32%, respectively, but these maximum values are only  
388 found in small basins. Indeed, basins when short rain or long rain exceed 30% of episodes  
389 are only found in basins smaller than 100 km<sup>2</sup>.

Code de champ modifié

390  
391 The mean date of occurrence is significantly different between the three flood types drivers,  
392 according to the Watson and Williams test. As shown on figure 9, the highest proportion of  
393 floods induced by short rain is observed during September to November, while the floods  
394 induced by long rain are mostly occurring during October to December, and excess rain  
395 floods are observed in late autumn and winter, with a peak in ~~January~~ February. This is  
396 consistent with the annual soil moisture cycle in this region: at the end of the summer the  
397 soils are dry and it takes several months to replenish the soil moisture level, which is at  
398 highest during winter. If examining the long-term changes in this ~~seasonal-monthly~~  
399 repartition of flood ~~types drivers~~ (see supplementary figure S34), the frequency of excess  
400 rain is decreasing from February to ~~October-April, and also in October,~~  
401 while increasing during winter months. This implies that the season during which excess rain floods are  
402 occurring is reducing in length and concentrated during wet winter months. On the opposite

Code de champ modifié

403 [the frequency of short or long rain floods is increasing in June and September, months that](#)  
404 [are getting drier over time in this region.-](#)

405  
406 The noticeable changes in flood processes over time are a reduction of Excess rainfall in 71  
407 basins and an increased frequency of short rains in 53 basins and Long rains in 63 basins  
408 (Figure 10), while short rain and long rain floods are decreasing for 19 and 22 basins,  
409 respectively. For excess rain, there are only 16 basins where their relative proportion is  
410 increasing; they are mostly located on the margins of the Alps and Pyrenees mountains. For  
411 more frequent events (ie. if considering an average of 3 episodes per year instead of one),  
412 the number of basins with a change is larger, with a reduction of Excess rainfall in 82 basins  
413 out of 98 (results not shown). This indicates that the soil moisture depletion has more  
414 impacts on small to moderate flood events, as previously observed by Bertola et al. (2021).  
415 There is no relationship between the rate of change in the different flood generating  
416 processes and catchment sizes indicating a clear regional pattern. The average magnitude  
417 [across all basins](#) of these changes remains low, on average -4.1% for excess rain, +1.2% for  
418 short rain and + 2.1% long rain. Yet, the magnitude of these changes is ranging from +15%  
419 to -21% for excess rain, +11% to -20% for short rain and +12% to -11% for long rain,  
420 depending on the catchment, indicating that local catchment characteristics could strongly  
421 modulate the regional signal. In addition, the average values over the whole domain are  
422 hiding some local changes: for instance, short rainfall floods are increasing in the  
423 southeastern part of the Cévennes while decreasing for the northwestern part as seen in  
424 Figure 10.

#### 425 426 **4.5 Regional changes**

427  
428 To assess whether the changes in the relative influence of the three different flood types are  
429 significant at the regional scale, we computed for each year the relative frequency of the  
430 different flood types, all basins together. It is indeed not possible to do this analysis for each  
431 station independently, due to the small size of the samples over the two periods. These  
432 changes in the occurrence of flood types are significant at the regional scale according to the  
433 Mann-Kendall test (Figure 11), for the frequency of excess rain floods and short rain floods,  
434 at the 5% significance level, but not for the long rain floods. [All events pooled regionally, the](#)  
435 [decrease in excess rain floods is equal to -13% between 1959-1990 and 1991-2021, and the](#)  
436 [increase of short rain floods is equal to +36%. In addition, to assess whether these results](#)  
437 [are robust to the thresholds used in the classification of flood events, a Monte Carlo](#)  
438 [experiment has been also conducted. Results show \(see supplementary figure S5\) that](#)  
439 [regional changes in excess rain and short rain floods are not dependent on classification](#)  
440 [thresholds, while it is not the case for long rain floods.](#) In terms of flood severity for the  
441 different flood types, the median flood computed for each basin is strongly correlated to  
442 basin size ( $\rho = 0.78$ ) for floods caused by excess rain, short rain ( $\rho = 0.80$ ) and long rain ( $\rho =$   
443  $0.75$ ); and very similar results are found for the maximum flood. On the contrary, the specific  
444 discharge of flood peaks is non linearly related to basin sizes, with a clear threshold effect  
445 for basins smaller than 500 km<sup>2</sup> that have a much larger specific discharge than larger  
446 basins.

447  
448 Given that there are different flood sample sizes in the different basins corresponding to  
449 different flood-generating processes, we pooled regionally the [flood events](#). [To do so, we](#)  
450 [computed the specific discharge for each event \(i.e. the flood magnitude divided by](#)

451 [catchment area](#)) to analyze the distributions of specific discharge for all [the](#) events  
452 associated with excess rain, long rain or short rain. [Specific discharge is used herein since it](#)  
453 [is a good indicator of flash floods severity, notably in this Mediterranean region \(Delrieu et](#)  
454 [al., 2005, Ruin et al., 2008\).](#) -Figure 12 shows that the short rain floods are more severe, in  
455 terms of specific discharge, than excess rain or long rain floods at the regional level ([as](#)  
456 [shown also by Tarasova et al., 2023](#)). The regional distributions [for the different event types](#)  
457 are different according to the Kolmogorov-Smirnov test. It must be noted that for a given  
458 basin the magnitude of the different types of floods may not be very different, showing the  
459 strong variability from one event to another that is not solely linked to the flood trigger. When  
460 comparing [the different flood distributions between](#) the time periods 1959-1990 and 1991-  
461 2021, the differences in flood magnitudes between excess rain, long and short rain are  
462 reduced. This is mainly due to a [slight](#) decrease in the specific discharge of short rain floods,  
463 notably for flood events with a return level higher than 10 years, while the excess rain floods  
464 show very little changes in intensity over time.

465

## 466 5. Conclusions

467

468 The aim of this study was to document the evolution of the characteristics of Mediterranean  
469 floods, using a large database of long river discharge records in southern France. In most  
470 basins, floods tend to occur earlier during the year, the mean flood date being on average  
471 advanced by one month. This seasonal shift can be attributed to the increased frequency of  
472 southern-circulation weather types during spring and [summer, that summer that](#) are strongly  
473 associated with the occurrence of floods in this region. Over time, floods also tend to be  
474 more clustered in time over the different basins, as reflected by a decreasing variability in  
475 flood occurrence throughout the year. On the contrary, for the westernmost basins  
476 influenced by Atlantic circulation patterns, floods tend to occur later during the year, also due  
477 to a seasonal shift of the flood-generating circulation patterns that are occurring more  
478 frequently in late winter. During floods, an increase in total and extreme event precipitation  
479 has been observed, associated with a decrease of antecedent soil moisture before rainfall  
480 events, linked to a smaller contribution of base flow during floods. It can be concluded that it  
481 is the depletion of soil water content, due to increased aridity in south France notably related  
482 to higher evapotranspiration rates ([Tramblay et al., 2020](#)), that is the likely cause of the  
483 absence of flood trends despite the increase in extreme rainfall. It should be also noted that  
484 over all basins, dryer soils are associated with lower runoff coefficients, and this relationship  
485 remains valid over time. The majority of flood events are associated with excess rainfall on  
486 saturated soils, but that proportion is decreasing over time with a concurrent increased  
487 frequency of short rain, potentially leading to more severe floods. At the regional scale,  
488 floods induced by short rains are indeed of higher magnitude, but due to a lower runoff  
489 coefficient induced by drier antecedent soil moisture, the specific discharge associated with  
490 short rain flood is decreasing over time. These results are consistent with those obtained in  
491 other regions, showing that floods do not necessarily increase with the increase in extreme  
492 precipitation, and that soil moisture seems to play a key role in explaining these changes  
493 and the lack of trends ultimately on flood hazard ([Wasko and Nathan, 2019; Bertola et al.,](#)  
494 [2021; Wasko et al., 2021](#)). The results of the present study are rather homogeneous given  
495 the different catchment sizes and land use types, indicating that changes in flood types are  
496 mainly resulting from regional climate change and not only local changes, such as land  
497 cover or agricultural practice changes, [n](#)or the increase of urban and peri-urban areas.

Code de champ modifié

498 Nonetheless, if the observed trend in increased short rain floods is persisting in the  
499 upcoming decades, the severity of floods, particularly the most important ones, could  
500 increase along with the rise in rainfall extremes particularly in areas where the soil infiltration  
501 potential is low, such as in mountainous or urbanized areas, that have expanded a lot in  
502 recent years. This aspect could be further investigated using climate scenarios.

503 One of the main perspectives of this work would be to perform a similar analysis at sub-daily  
504 time steps, that would be more adapted to analyze changes in flash floods characteristics,  
505 notably in terms of the flashiness response of the catchments (Baker et al., 2004; Li et al.,  
506 2022). Indeed, the daily time step prevents a thorough analysis of changes in rainfall  
507 patterns, notably at shorter time steps. Yet, there is no gridded dataset of hourly precipitation  
508 before the 2000s in Southern France, that does not allow to repeat such a similar study over  
509 62 years. However, given the availability of radar rainfall over France, it would be possible to  
510 analyze, at least for the recent years after 2000, the evolution of several characteristics,  
511 such as the evolution of storm hydrographs, concentration time and the flashiness response  
512 of the basins. Another relevant prospective work would be to analyze the spatial extent of  
513 floods. Given the future evolution of weather types associated with floods in combination  
514 with more local to regional characteristics, such as soil moisture state, these types of events  
515 may impact simultaneously wider, or smaller, parts of the region considered, and this could  
516 have serious implications on risk management (Brunner et al., 2021; Brunner and  
517 Dougherty, 2022). Therefore, the joint analysis of flood occurrence in nearby basins would  
518 be highly relevant. Finally, there is also a need for new approaches to incorporate these  
519 changes in flood generating process into engineering practice (Slater et al., 2021b), notably  
520 to estimate the return levels for different types of infrastructure design.

Code de champ modifié

Code de champ modifié

Code de champ modifié

521

## 522 **Acknowledgements**

523 This study is a contribution to the EXPLORE2 project funded by the French Ministry of  
524 Ecological Transition. The access to the Météo France and Banque Hydro databases is  
525 acknowledged. [The authors wish to thanks 3 anonymous reviewers for their comments.](#)

## 526 **Data availability**

527 The computed catchment-based indicators are available upon request to the corresponding  
528 author.

## 529 **Author contributions**

530 Y.T.: Conceptualization, investigation, data curation, formal analysis, writing—original draft  
531 preparation. YT designed the experiments, performed the analyses, and wrote the paper.  
532 PA, GA, ML, EP, LN and ES Conceptualization, data curation, writing—reviewing and  
533 editing.

## 534 **Competing interests**

535 The authors declare that they have no conflict of interest.

536

## INDEX OF FIGURES

Figure 1: Map of the selected catchments

Figure 2: Relative changes in different flood event characteristics between 1959-1990 and 1991-2021

Figure 3: Relationship between the flood event runoff coefficients and antecedent soil moisture for the two time periods considered: 1959-1990 and 1991-2021. For each box, the central line indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points. The numbers at the top of the figure indicate the number of events in each category.

Figure 4: Mean date of flood occurrence

Figure 5: Changes in mean flood date (left) and changes in the concentration index (right) between 1959-1990 and 1991-2021

Figure 6: Frequency of the different weather types associated with flood events, the names of the different weather types are from Garavaglia et al. (2010)

Figure 7: Change in monthly frequency of weather types 1, 2, 4, 6 and 7 between 1959-1990 and 1991-2021

Figure 8: Flood event classification into four categories: excess rain, long rain, short rain and others

Figure 9: Mean monthly frequency of occurrence for the three flood drivers: excess rain, short rain, and long rain

Figure 10: Relative changes in the frequency of excess rain, short rain and long rain between 1959-1990 and 1991-2021

Figure 11: Regional frequency of excess rain, short rain, and long rain floods between 1959 and 2021. The gray lines denote a least-square linear fit to represent the long-term tendency

Figure 12: Distribution of regionally sampled floods corresponding to excess rain, short rain and long rain types of floods for the two time periods 1959-1990 and 1991-2021

585  
586  
587  
588  
589  
590

Table 1: Number of local significant changes in the median of flood events characteristics detected by the Wilcoxon test and results of the regional Mann-Kendall test on flood event characteristics

Indicator	Number of significant <u>local</u> changes (Wilcoxon, 10%)	<u>pvalue of the regional MK test</u>	<u>Regional changes between 1959 and 2022 (%)</u>
Flood event duration	17	<a href="#">0.0046178</a>	<a href="#">-0.40%</a>
Base flow contribution to peak	15	<a href="#">0.5687962</a>	<a href="#">-8.62%</a>
Runoff coefficient	19	<a href="#">0.0000002</a>	<a href="#">-14.62%</a>
Total event rainfall	16	<a href="#">0.0011851</a>	<a href="#">9.01%</a>
Maximum event rainfall	27	<a href="#">0.0000000</a>	<a href="#">13.47%</a>
Antecedent soil moisture	12	<a href="#">0.0000008</a>	<a href="#">-9.80%</a>

591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613

614

615 **References**

616

617 Alaoui, A., Rogger, M., Peth, S., and Blöschl, G.: Does soil compaction increase floods? A  
618 review, *Journal of Hydrology*, 557, 631–642, <https://doi.org/10.1016/j.jhydrol.2017.12.052>,  
619 2018.

620

621 Ali, E., Cramer, W., Carnicer, J., Georgopoulou, E., Hilmi, N. J. M., Cozannet, G. L., and  
622 Lionello, P.: Cross-Chapter Paper 4: Mediterranean Region, edited by: Pörtner, H. O.,  
623 Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M.,  
624 Langsdorf, S., Lösckke, S., Möller, V., Okem, A., and Rama, B., *Climate Change 2022:*  
625 *Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth*  
626 *Assessment Report of the Intergovernmental Panel on Climate Change*, 2233–2272,  
627 <https://doi.org/10.1017/9781009325844.021.2233>, 2022.

628

629 Baker, D. B., Richards, R. P., Loftus, T. T., and Kramer, J. W.: A new flashiness index:  
630 characteristics and applications to midwestern rivers and streams, *J Am Water Resources*  
631 *Assoc*, 40, 503–522, <https://doi.org/10.1111/j.1752-1688.2004.tb01046.x>, 2004.

632

633 Berens, P.: *CircStat: A MATLAB Toolbox for Circular Statistics*, *J. Stat. Soft.*, 31,  
634 <https://doi.org/10.18637/jss.v031.i10>, 2009.

635

636 Berghuijs, W. R., Harrigan, S., Molnar, P., Slater, L. J., and Kirchner, J. W.: The Relative  
637 Importance of Different Flood-Generating Mechanisms Across Europe, *Water Resour. Res.*,  
638 2019WR024841, <https://doi.org/10.1029/2019WR024841>, 2019.

639

640 Bertola, M., Viglione, A., Vorogushyn, S., Lun, D., Merz, B., and Blöschl, G.: Do small and  
641 large floods have the same drivers of change? A regional attribution analysis in Europe,  
642 *Hydrol. Earth Syst. Sci.*, 25, 1347–1364, <https://doi.org/10.5194/hess-25-1347-2021>, 2021.

643

644 Bissonnais, Y. L. and Singer, M. J.: Seal Formation, Runoff, and Interrill Erosion from  
645 Seventeen California Soils, *Soil Science Society of America Journal*, 57, 224–229,  
646 <https://doi.org/10.2136/sssaj1993.03615995005700010039x>, 1993.

647

648 Blanchet, J. and Creutin, J.-D.: Instrumental agreement and retrospective analysis of trends  
649 in precipitation extremes in the French Mediterranean Region, *Environ. Res. Lett.*, 17,  
650 074011, <https://doi.org/10.1088/1748-9326/ac7734>, 2022.

651

652 Blöschl, G., Gaál, L., Hall, J., Kiss, A., Komma, J., Nester, T., Parajka, J., Perdigão, R. A. P.,  
653 Plavcová, L., Rogger, M., Salinas, J. L., and Viglione, A.: Increasing river floods: fiction or  
654 reality?: Increasing river floods, *WIREs Water*, 2, 329–344,  
655 <https://doi.org/10.1002/wat2.1079>, 2015.

656

657 Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A. P., Merz, B., Arheimer, B., Aronica, G. T.,  
658 Bilibashi, A., Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G. B., Claps, P.,  
659 Fiala, K., Frolova, N., Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss,  
660 A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-  
661 Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M.,

662 Ovcharuk, V., Radevski, I., Rogger, M., Salinas, J. L., Sauquet, E., Šraj, M., Szolgay, J.,  
663 Viglione, A., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate shifts timing  
664 of European floods, *Science*, 357, 588–590, <https://doi.org/10.1126/science.aan2506>, 2017.  
665  
666 Blöschl, G., Hall, J., Viglione, A., Perdigão, R. A. P., Parajka, J., Merz, B., Lun, D., Arheimer,  
667 B., Aronica, G. T., Bilibashi, A., Boháč, M., Bonacci, O., Borga, M., Čanjevac, I., Castellarin,  
668 A., Chirico, G. B., Claps, P., Frolova, N., Ganora, D., Gorbachova, L., Gül, A., Hannaford, J.,  
669 Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O.,  
670 Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A.,  
671 Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Salinas, J. L., Sauquet, E., Šraj, M.,  
672 Szolgay, J., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate both  
673 increases and decreases European river floods, *Nature*, 573, 108–111,  
674 <https://doi.org/10.1038/s41586-019-1495-6>, 2019.  
675  
676 Brunner, M. I. and Dougherty, E. M.: Varying Importance of Storm Types and Antecedent  
677 Conditions for Local and Regional Floods, *Water Resources Research*, 58,  
678 <https://doi.org/10.1029/2022WR033249>, 2022.  
679  
680 Brunner, M. I., Slater, L., Tallaksen, L. M., and Clark, M.: Challenges in modeling and  
681 predicting floods and droughts: A review, *WIREs Water*, 8,  
682 <https://doi.org/10.1002/wat2.1520>, 2021.  
683  
684 [Brunner, M. I., Swain, D. L., Wood, R. R., Willkofer, F., Done, J. M., Gilleland, E., and](https://doi.org/10.1038/s43247-021-00248-x)  
685 [Ludwig, R.: An extremeness threshold determines the regional response of floods to](https://doi.org/10.1038/s43247-021-00248-x)  
686 [changes in rainfall extremes. \*Communications Earth & Environment\*, 2:1,](https://doi.org/10.1038/s43247-021-00248-x)  
687 <https://doi.org/10.1038/s43247-021-00248-x>, 2021  
688  
689  
690 Burch, G. J., Moore, I. D., and Burns, J.: Soil hydrophobic effects on infiltration and  
691 catchment runoff, *Hydrol. Process.*, 3, 211–222, <https://doi.org/10.1002/hyp.3360030302>,  
692 1989.  
693  
694 Burn, D. H.: Catchment similarity for regional flood frequency analysis using seasonality  
695 measures, *Journal of Hydrology*, 202, 212–230, [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-1694(97)00068-1)  
696 [1694\(97\)00068-1](https://doi.org/10.1016/S0022-1694(97)00068-1), 1997.  
697  
698 Büttner, G.: CORINE Land Cover and Land Cover Change Products, in: *Land Use and Land*  
699 *Cover Mapping in Europe*, vol. 18, edited by: Manakos, I. and Braun, M., Springer  
700 Netherlands, Dordrecht, 55–74, [https://doi.org/10.1007/978-94-007-7969-3\\_5](https://doi.org/10.1007/978-94-007-7969-3_5), 2014.  
701  
702 Cao, Q., Gershunov, A., Shulgina, T., Ralph, F. M., Sun, N., and Lettenmaier, D. P.: Floods  
703 due to Atmospheric Rivers along the U.S. West Coast: The Role of Antecedent Soil Moisture  
704 in a Warming Climate, *Journal of Hydrometeorology*, 21, 1827–1845,  
705 <https://doi.org/10.1175/JHM-D-19-0242.1>, 2020.  
706  
707 [Delrieu, G., Nicol, J., Yates, E., Kirstetter, P.-E., Creutin, J.-D., Anquetin, S., Obled, C.,](https://doi.org/10.1038/s43247-021-00248-x)  
708 [Saulnier, G.-M., Ducrocq, V., Gaume, E., Payrastre, O., Andrieu, H., Ayrat, P.-A., Bouvier,](https://doi.org/10.1038/s43247-021-00248-x)  
709 [C., Neppel, L., Livet, M., Lang, M., du-Châtelet, J. P., Walpersdorf, A., and Wobrock, W.:](https://doi.org/10.1038/s43247-021-00248-x)



710 [The Catastrophic Flash-Flood Event of 8–9 September 2002 in the Gard Region, France: A](#)  
711 [First Case Study for the Cévennes–Vivarais Mediterranean Hydrometeorological](#)  
712 [Observatory. Journal of Hydrometeorology, 6:1, 34–52, https://doi.org/10.1175/jhm-400.1,](#)  
713 [2005.](#)

714

715 Ducrocq, V., Nuissier, O., Ricard, D., Lebeauupin, C., and Thouvenin, T.: A numerical study of  
716 three catastrophic precipitating events over southern France. II: Mesoscale triggering and  
717 stationarity factors, Q.J.R. Meteorol. Soc., 134, 131–145, <https://doi.org/10.1002/qj.199>,  
718 2008.

719

720 Farquharson, F. A. K., Meigh, J. R., and Sutcliffe, J. V.: Regional flood frequency analysis in  
721 arid and semi-arid areas, Journal of Hydrology, 138, 487–501, [https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-1694(92)90132-F)  
722 [1694\(92\)90132-F](#), 1992.

723

724 Fisher, N. I.: Statistical analysis of circular data, Cambridge University Press, Cambridge  
725 [England]; New York, NY, USA, 277 pp., 1993.

726

727 Garavaglia, F., Gailhard, J., Paquet, E., Lang, M., Garçon, R., and Bernardara, P.:  
728 Introducing a rainfall compound distribution model based on weather patterns sub-sampling,  
729 Hydrol. Earth Syst. Sci., 14, 951–964, <https://doi.org/10.5194/hess-14-951-2010>, 2010.

730

731 Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M., Soubeyroux, J.-M. :  
732 A 50-year high-resolution atmospheric reanalysis over France with the Safran system.  
733 International Journal of Climatology, 2010, 30 (11), 1627-1644.  
734 <https://doi.org/10.1002/joc.2003>, 2010.

735

736 Hamed, K. H. and Ramachandra Rao, A.: A modified Mann-Kendall trend test for  
737 autocorrelated data, Journal of Hydrology, 204, 182–196, [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-1694(97)00125-X)  
738 [1694\(97\)00125-X](#), 1998.

739

740 Ho, M., Nathan, R., Wasko, C., Vogel, E., and Sharma, A.: Projecting changes in flood event  
741 runoff coefficients under climate change, Journal of Hydrology, 615, 128689,  
742 <https://doi.org/10.1016/j.jhydrol.2022.128689>, 2022.

743

744 Huang, H., Fischella, M. R., Liu, Y., Ban, Z., Fayne, J. V., Li, D., Cavanaugh, K. C., and  
745 Lettenmaier, D. P.: Changes in Mechanisms and Characteristics of Western U.S. Floods  
746 Over the Last Sixty Years, Geophysical Research Letters, 49,  
747 <https://doi.org/10.1029/2021GL097022>, 2022.

748

749 [Jiang, S., Bevacqua, E., and Zscheischler, J.: River flooding mechanisms and their changes](#)  
750 [in Europe revealed by explainable machine learning, Hydrol. Earth Syst. Sci., 26, 6339–](#)  
751 [6359, https://doi.org/10.5194/hess-26-6339-2022, 2022.](#)

752

753 Kemter, M., Merz, B., Marwan, N., Vorogushyn, S., and Blöschl, G.: Joint Trends in Flood  
754 Magnitudes and Spatial Extents Across Europe, Geophys. Res. Lett., 47,  
755 <https://doi.org/10.1029/2020GL087464>, 2020.

756

a mis en forme : Anglais (États-Unis)

757 Landler, L., Ruxton, G. D., and Malkemper, E. P.: The Hermans–Rasson test as a powerful  
758 alternative to the Rayleigh test for circular statistics in biology, *BMC Ecol*, 19, 30,  
759 <https://doi.org/10.1186/s12898-019-0246-8>, 2019.

760

761 Lang, M., Ouarda, T. B. M. J., and Bobée, B.: Towards operational guidelines for over-  
762 threshold modeling, *Journal of Hydrology*, 225, 103–117, [https://doi.org/10.1016/S0022-1694\(99\)00167-5](https://doi.org/10.1016/S0022-1694(99)00167-5), 1999.

763

764

765 Li, Z., Gao, S., Chen, M., Gourley, J. J., Liu, C., Prein, A. F., and Hong, Y.: The  
766 conterminous United States are projected to become more prone to flash floods in a high-  
767 end emissions scenario, *Commun Earth Environ*, 3, 86, <https://doi.org/10.1038/s43247-022-00409-6>, 2022.

768

769

770 Liu, J., Feng, S., Gu, X., Zhang, Y., Beck, H. E., Zhang, J., and Yan, S.: Global changes in  
771 floods and their drivers, *Journal of Hydrology*, 614, 128553,  
772 <https://doi.org/10.1016/j.jhydrol.2022.128553>, 2022.

773

774 Lyne, V. D. and Hollick, M.: Stochastic time-variable rainfall runoff modelling, *Hydrology and*  
775 *Water Resources Symposium*, Institution of Engineers, Australia, Perth (1979), Barton,  
776 Australia, 1979.

777

778 Mao, Y., Zhou, T., Leung, L. R., Tesfa, T. K., Li, H., Wang, K., Tan, Z., and Getirana, A.:  
779 Flood Inundation Generation Mechanisms and Their Changes in 1953–2004 in Global Major  
780 River Basins, *J. Geophys. Res. Atmos.*, 124, 11672–11692,  
781 <https://doi.org/10.1029/2019JD031381>, 2019.

782

783 Merz, B., Aerts, J., Arnbjerg-Nielsen, K., Baldi, M., Becker, A., Bichet, A., Blöschl, G.,  
784 Bouwer, L. M., Brauer, A., Cioffi, F., Delgado, J. M., Gocht, M., Guzzetti, F., Harrigan, S.,  
785 Hirschboeck, K., Kilsby, C., Kron, W., Kwon, H.-H., Lall, U., Merz, R., Nissen, K., Salvatti, P.,  
786 Swierczynski, T., Ulbrich, U., Viglione, A., Ward, P. J., Weiler, M., Wilhelm, B., and Nied, M.:  
787 Floods and climate: emerging perspectives for flood risk assessment and management, *Nat.*  
788 *Hazards Earth Syst. Sci.*, 14, 1921–1942, <https://doi.org/10.5194/nhess-14-1921-2014>,  
789 2014.

790

791 Pastor, F., Valiente, J. A., and Khodayar, S.: A Warming Mediterranean: 38 Years of  
792 Increasing Sea Surface Temperature, *Remote Sensing*, 12, 2687,  
793 <https://doi.org/10.3390/rs12172687>, 2020.

794

795 Penna, D., Tromp-van Meerveld, H. J., Gobbi, A., Borga, M., and Dalla Fontana, G.: The  
796 influence of soil moisture on threshold runoff generation processes in an alpine headwater  
797 catchment, *Hydrol. Earth Syst. Sci.*, 15, 689–702, <https://doi.org/10.5194/hess-15-689-2011>,  
798 2011.

799

800 Raynaud, D., Thielen, J., Salamon, P., Burek, P., Anquetin, S., and Alfieri, L.: A dynamic  
801 runoff co-efficient to improve flash flood early warning in Europe: evaluation on the 2013  
802 central European floods in Germany: Dynamic runoff co-efficient for flash flood early warning  
803 in Europe, *Met. Apps*, 22, 410–418, <https://doi.org/10.1002/met.1469>, 2015.

804

805 Ribes, A., Thao, S., Vautard, R., Dubuisson, B., Somot, S., Colin, J., Planton, S., and  
806 Soubeyroux, J.-M.: Observed increase in extreme daily rainfall in the French Mediterranean,  
807 *Clim Dyn*, 52, 1095–1114, <https://doi.org/10.1007/s00382-018-4179-2>, 2019.

808  
809 Rogger, M., Viglione, A., Derx, J., and Blöschl, G.: Quantifying effects of catchments storage  
810 thresholds on step changes in the flood frequency curve: Step Changes in the Flood  
811 Frequency Curve, *Water Resour. Res.*, 49, 6946–6958, <https://doi.org/10.1002/wrcr.20553>,  
812 2013.

813  
814 [Ruin, I., Creutin, J.-D., Anquetin, S., & Lutoff, C.: Human exposure to flash floods – Relation  
815 between flood parameters and human vulnerability during a storm of September 2002 in  
816 Southern France. \*Journal of Hydrology\*, 361:1-2, 199–213,  
817 <https://doi.org/10.1016/j.jhydrol.2008.07.044>, 2008.](https://doi.org/10.1016/j.jhydrol.2008.07.044)

818  
819 Sharma, A., Wasko, C., and Lettenmaier, D. P.: If Precipitation Extremes Are Increasing,  
820 Why Aren't Floods?, *Water Resour. Res.*, 54, 8545–8551,  
821 <https://doi.org/10.1029/2018WR023749>, 2018.

822  
823 Slater, L., Villarini, G., Archfield, S., Faulkner, D., Lamb, R., Khouakhi, A., and Yin, J.: Global  
824 Changes in 20-Year, 50-Year, and 100-Year River Floods, *Geophysical Research Letters*,  
825 48, <https://doi.org/10.1029/2020GL091824>, 2021a.

826  
827 Slater, L. J., Anderson, B., Buechel, M., Dadson, S., Han, S., Harrigan, S., Kelder, T., Kowal,  
828 K., Lees, T., Matthews, T., Murphy, C., and Wilby, R. L.: Nonstationary weather and water  
829 extremes: a review of methods for their detection, attribution, and management, *Hydrol.*  
830 *Earth Syst. Sci.*, 25, 3897–3935, <https://doi.org/10.5194/hess-25-3897-2021>, 2021b.

831  
832 Stein, L., Pianosi, F., and Woods, R.: Event-based classification for global study of river  
833 flood generating processes, *Hydrological Processes*, 34, 1514–1529,  
834 <https://doi.org/10.1002/hyp.13678>, 2020.

835  
836 Stein, L., Clark, M. P., Knoben, W. J. M., Pianosi, F., and Woods, R. A.: How Do Climate  
837 and Catchment Attributes Influence Flood Generating Processes? A Large-Sample Study for  
838 671 Catchments Across the Contiguous USA, *Water Res*, 57,  
839 <https://doi.org/10.1029/2020WR028300>, 2021.

840  
841 Tarasova, L., Merz, R., Kiss, A., Basso, S., Blöschl, G., Merz, B., Viglione, A., Plötner, S.,  
842 Guse, B., Schumann, A., Fischer, S., Ahrens, B., Anwar, F., Bárdossy, A., Bühler, P.,  
843 Haberlandt, U., Kreibich, H., Krug, A., Lun, D., Müller-Thomy, H., Pidoto, R., Primo, C.,  
844 Seidel, J., Vorogushyn, S., and Wietzke, L.: Causative classification of river flood events,  
845 *WIREs Water*, 6, <https://doi.org/10.1002/wat2.1353>, 2019.

846  
847 Tarasova, L., Basso, S., Wendi, D., Viglione, A., Kumar, R., and Merz, R.: A Process-Based  
848 Framework to Characterize and Classify Runoff Events: The Event Typology of Germany,  
849 *Water Resour. Res.*, 56, <https://doi.org/10.1029/2019WR026951>, 2020.

850

851 [Tarasova, L., Lun, D., Merz, R. Blösch, G., Basso, S., Bertola, M., Miniussi, A., Radovec, O.,](#)  
852 [Samaniego, L., Thober, S., Kumar, R.: Shifts in flood generation processes exacerbate](#)  
853 [regional flood anomalies in Europe. Commun Earth Environ 4, 49,](#)  
854 <https://doi.org/10.1038/s43247-023-00714-8>, 2023.

855  
856 Trambly, Y., Bouvier, C., Martin, C., Didon-Lescot, J.-F., Todorovik, D., and Domergue, J.-  
857 M.: Assessment of initial soil moisture conditions for event-based rainfall–runoff modelling,  
858 Journal of Hydrology, 387, 176–187, <https://doi.org/10.1016/j.jhydrol.2010.04.006>, 2010.

859  
860 Trambly, Y., Neppel, L., Carreau, J., and Najib, K.: Non-stationary frequency analysis of  
861 heavy rainfall events in southern France, Hydrological Sciences Journal, 58, 280–294,  
862 <https://doi.org/10.1080/02626667.2012.754988>, 2013.

863  
864 Trambly, Y., Mimeau, L., Neppel, L., Vinet, F., and Sauquet, E.: Detection and attribution of  
865 flood trends in Mediterranean basins, Hydrol. Earth Syst. Sci., 23, 4419–4431,  
866 <https://doi.org/10.5194/hess-23-4419-2019>, 2019.

867  
868 Trambly, Y., Koutroulis, A., Samaniego, L., Vicente-Serrano, S. M., Volaire, F., Boone, A.,  
869 Le Page, M., Llasat, M. C., Albergel, C., Burak, S., Cailleret, M., Kalin, K. C., Davi, H.,  
870 Dupuy, J.-L., Greve, P., Grillakis, M., Hanich, L., Jarlan, L., Martin-StPaul, N., Martínez-  
871 Vilalta, J., Mouillot, F., Pulido-Velazquez, D., Quintana-Seguí, P., Renard, D., Turco, M.,  
872 Türkeş, M., Trigo, R., Vidal, J.-P., Vilagrosa, A., Zribi, M., and Polcher, J.: Challenges for  
873 drought assessment in the Mediterranean region under future climate scenarios, Earth-  
874 Science Reviews, 210, 103348, <https://doi.org/10.1016/j.earscirev.2020.103348>, 2020.

875  
876 Trambly, Y., Villarini, G., Saidi, M. E., Massari, C., and Stein, L.: Classification of flood-  
877 generating processes in Africa, Sci Rep, 12, 18920, [https://doi.org/10.1038/s41598-022-](https://doi.org/10.1038/s41598-022-23725-5)  
878 [23725-5](https://doi.org/10.1038/s41598-022-23725-5), 2022.

879  
880 Wasko, C. and Nathan, R.: Influence of changes in rainfall and soil moisture on trends in  
881 flooding, Journal of Hydrology, 575, 432–441, <https://doi.org/10.1016/j.jhydrol.2019.05.054>,  
882 2019.

883  
884 Wasko, C., Nathan, R., Stein, L., and O’Shea, D.: Evidence of shorter more extreme rainfalls  
885 and increased flood variability under climate change, Journal of Hydrology, 603, 126994,  
886 <https://doi.org/10.1016/j.jhydrol.2021.126994>, 2021.

887  
888 Watson, G. S. and Williams, E. J.: On the Construction of Significance Tests on the Circle  
889 and the Sphere, Biometrika, 43, 344, <https://doi.org/10.2307/2332913>, 1956.

890  
891 Whitfield, P. H.: Floods in future climates: a review: Changing floods in future climates, J.  
892 Flood Risk Manage, 5, 336–365, <https://doi.org/10.1111/j.1753-318X.2012.01150.x>, 2012.

893  
894 Zhang, S., Zhou, L., Zhang, L., Yang, Y., Wei, Z., Zhou, S., Yang, D., Yang, X., Wu, X.,  
895 Zhang, Y., Li, X., and Dai, Y.: Reconciling disagreement on global river flood changes in a  
896 warming climate, Nat. Clim. Chang., 12, 1160–1167, [https://doi.org/10.1038/s41558-022-](https://doi.org/10.1038/s41558-022-01539-7)  
897 [01539-7](https://doi.org/10.1038/s41558-022-01539-7), 2022.

898

899

900