

1 **Changes in Mediterranean flood processes and seasonality**

2

3 Yves Tramblay 1\*

4 Patrick Arnaud 2

5 Guillaume Artigue 1

6 Michel Lang 3

7 Emmanuel Paquet 4

8 Luc Neppel 1

9 Eric Sauquet 3

10

11 1 HSM, Univ. Montpellier, CNRS, IRD, IMT-Mines Alès, Montpellier, France

12 2 RECOVER, INRAE, Aix Marseille Université, Aix-en-Provence, France

13 3 RiverLy, INRAE, Villeurbanne, France

14 4 EDF-DTG, Saint Martin le Vinoux, France

15

16

17 \* corresponding author, HydroSciences-Montpellier, Hydropolis, Faculté de  
18 Pharmacie, 15 Avenue Charles Flahaut, 34000 Montpellier. [yves.tramblay@ird.fr](mailto:yves.tramblay@ird.fr)

19

20

21

22

23

24

25

26

27 REVISÉD MANUSCRIPT V2

28

29

30

31

32

33 **Abstract**

34

35 Floods are a major natural hazard in the Mediterranean region, causing deaths and extensive  
36 damages. Recent studies have shown that intense rainfall events are becoming more extreme  
37 in this region, but paradoxically without leading to an increase in the severity of floods.  
38 Consequently, it is important to understand how flood events are changing to explain this  
39 absence of trends in flood magnitude despite increased rainfall extremes. A database of 98  
40 stations in Southern France with an average record of 50 years of daily river discharge data  
41 between 1959 and 2021 was considered, together with a high-resolution reanalysis product  
42 providing precipitation and simulated soil moisture and a classification of weather patterns  
43 associated with rainfall events over France. Flood events, corresponding to an average  
44 occurrence of one event per year (5317 events in total), were extracted and classified into  
45 excess rainfall, short rainfall and long rainfall event types. Several flood events characteristics  
46 have been also analyzed: flood event durations, base flow contribution to floods, runoff  
47 coefficient, total and maximum event rainfall and antecedent soil moisture. The evolution  
48 through time of these flood event characteristics and seasonality were analyzed. Results  
49 indicated that, in most basins, floods tend to occur earlier during the year, the mean flood date  
50 being on average advanced by one month between 1959-1990 and 1991-2021. This seasonal  
51 shift could be attributed to the increased frequency of southern-circulation weather types  
52 during spring and summer. An increase in total and extreme event precipitation has been  
53 observed, associated with a decrease of antecedent soil moisture before rainfall events. The  
54 majority of flood events are associated with excess rainfall on saturated soils, but their relative  
55 proportion is decreasing over time notably in spring with a concurrent increased frequency of  
56 short rain floods. For most basins there is a positive correlation between antecedent soil  
57 moisture and flood event runoff coefficients that is remaining stable over time, with dryer soils  
58 producing less runoff and a lower contribution of base flow to floods. In a context of an  
59 increasing aridity, this relationship is the likely cause of the absence of trends in flood  
60 magnitudes observed in this region and the change of event types. These changes in flood  
61 characteristics are quite homogeneous over the domain studied, suggesting that they are  
62 rather linked to the evolution of the regional climate than to catchments characteristics.  
63 Consequently, this study shows that even in the absence of trends, flood properties may  
64 change over time and these changes need to be accounted for when analyzing the long-term  
65 evolution of flood hazards.

66

67

68

69

70

71

72

73  
74  
75  
76

## 77 1. Introduction

78

79 There is a growing interest in understanding the evolution of floods occurring in different  
80 regions in the context of climate change. The recent sixth report of the Intergovernmental  
81 Panel on Climate Change (Ali et al., 2022), reported a mixture of observed trends in different  
82 Mediterranean countries, with both increasing and decreasing river floods and overall, a low  
83 confidence in their attribution to climate change. Several large-scale studies on changes in  
84 flood risk (Slater et al., 2021a; Blöschl et al., 2017, 2019) have indicated for the Mediterranean  
85 region a possible decrease over the last decades. This difficulty in detecting possible changes  
86 in flood hazard is doubtless linked to the fact that there are different types of floods (Tarasova  
87 et al., 2019; Berghuijs et al., 2019; Stein et al., 2020; Trambly et al., 2022). Indeed, a  
88 distinction can be made between floods associated with soil saturation excess, soil infiltration  
89 excess or snowmelt-driven floods and the relative frequency of these different types of floods  
90 may change over time (Zhang et al., 2022). Furthermore, these changes can occur at local to  
91 regional scales, given complex combinations of climatic and physiographic triggers, making  
92 global generalization of changes in flood risk hazardous, if not irrelevant (Whitfield, 2012;  
93 Blöschl et al., 2015).

94

95 Only a few studies are focusing on changes in flood types over time, while it is important to  
96 understand the long-term changes in flood processes to evaluate how flood risk can evolve,  
97 in order notably to better adapt the flood mitigation strategies (Merz et al., 2014). The main  
98 limitation to perform such studies is the need for long time-series of river discharge, to have  
99 large samples of flood events to evaluate how their properties may change over time. For  
100 many regions of the world, the lack of observed river discharge data prevents this type of  
101 analysis. Some studies attempted to analyze the changes in different types of floods for  
102 different regions (Berghuijs et al., 2019; Mao et al., 2019; Kemter et al., 2020; Bertola et al.,  
103 2021; Liu et al., 2022; Trambly et al., 2022, Tarasova et al., 2023). Most of these studies rely  
104 on flood classification schemes, with various complexity depending on the type of data  
105 available, allowing a data-based separation of floods into their distinct generation mechanisms  
106 (Tarasova et al., 2019; Berghuijs et al., 2019; Tarasova et al., 2020; Stein et al., 2020, 2021;  
107 Trambly et al., 2022). For basins under a Mediterranean climate, several studies reported  
108 complex interactions between precipitation increases and lower antecedent soil moisture,  
109 leading to thresholds effects (Brunner et al., 201) in the catchment response to changing  
110 hydro-climatic conditions (Wasko and Nathan, 2019; Cao et al., 2020; Bertola et al., 2021).  
111 Recent large-scale studies (Jiang et al., 2022, Tarasova et al., 2023) suggested a reduction  
112 of the frequency of floods driven by soil saturation in Europe, including basins in the  
113 Mediterranean area. Yet, these studies rely on coarse resolution climate forcings provided by  
114 E-OBS and there is a need to assess whether these findings are also valid in smaller basins  
115 with high-resolution datasets, notably with a better estimation of precipitation extremes. For  
116 instance, in Bertola et al. (2021) a decrease in 7-days maximum precipitation is found with E-

117 OBS for basins in Southern France, while Ribes et al. (2019) observed an opposite increasing  
118 trend using a network of about 700 rain gauges in that area. In addition, recent studies at the  
119 European scale focused on annual maxima only, when to attribute changes in flood event  
120 types it is also relevant to consider other metrics such as the runoff coefficient and the flood  
121 event duration, that require data about flood events and not only their maxima.

122  
123 In French Mediterranean basins, several studies reported an increase in precipitation  
124 extremes (Tramblay et al., 2013; Blanchet and Creutin, 2022; Ribes et al., 2019) that did not  
125 translate into increased floods (Tramblay et al., 2019). It is hypothesized that, as many regions  
126 of the world, a decrease in soil moisture linked with a greater aridity can potentially offset the  
127 increase in precipitation extremes and thus not increase flood severity (Sharma et al., 2018;  
128 Tramblay et al., 2019; Wasko and Nathan, 2019; Wasko et al., 2021; Huang et al., 2022).  
129 Excess soil moisture was previously identified as an important flood driver notably in the  
130 Mediterranean (Kemter et al., 2020; Bertola et al., 2021), indicating that they can play an  
131 important role. Yet, beside trend detection or changes in flood types, no study has provided  
132 an in-depth analysis of the joint long-term evolution of flood processes in Mediterranean  
133 basins, in relation to their drivers such as precipitation, soil moisture and the evolution of  
134 synoptic weather patterns associated with floods. In particular, there is a need to assess how  
135 these changes may be modulated in different catchment sizes with different land use, notably  
136 in the Mediterranean context where floods are generated by rainfall events characterized by  
137 a strong spatial and temporal variability. Therefore, the objective of the study is to evaluate  
138 how the characteristics of Mediterranean floods are changing. A recent study (Tramblay et al.,  
139 2019) indicated no significant trends on flood hazards for a large ensemble of basins located  
140 in southern France. This database is used herein and expanded with a weather types  
141 classification linked to floods to further analyze the possible changes in flood event  
142 characteristics. This analysis provides a comprehensive view at the regional scale of the  
143 evolution in time of flood-events characteristics (runoff coefficients, soil moisture, base flow  
144 contribution...), flood event seasonality in relation to large scale atmospheric patterns, flood-  
145 events types (excess rain floods, short rain floods...) and their respective magnitude.

146  
147 In section 2 are presented the different datasets used for this analysis, including river  
148 discharge, precipitation, soil moisture data and weather types classification. In section 3 the  
149 methods are presented, for event extraction, analysis of the seasonality and changes in the  
150 different flood drivers. Results are presented in section 4.

## 151 152 **2. Data**

153  
154 We consider 98 catchments in southern France (Figure 1) where the time series of daily  
155 discharge exceeds 30 years of complete data between 1959 and 2021 (Tramblay et al., 2019).  
156 Among them, 48 basins have more than 50 years of data and the basins selected are not  
157 influenced by reservoir or dam regulation. The catchment sizes are ranging from 14 km<sup>2</sup> to  
158 3195 km<sup>2</sup>, with a mean size equal to 480 km<sup>2</sup> (see Table t1 in supplementary materials).  
159 Basins with a nival regime were removed, identified from the river discharge hydrographs and  
160 removing basins with more than 20% of precipitation falling as snow. In addition to river  
161 discharge data, the precipitation and soil moisture for each basin has been retrieved from the  
162 SAFRAN-ISBA-MODCOU (SIM) reanalysis covering the whole France territory at 8 km spatial  
163 resolution (Vidal et al., 2010). Precipitation and soil moisture data have been extracted and

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

164 averaged at the catchment scale. The soil moisture data extracted from SIM is a soil wetness  
165 index obtained from the normalization of the volumetric soil moisture content with the wilting  
166 point and field capacity, that ranges between 0 and 1. Land cover classes (forest, agricultural,  
167 urban) corresponding to 2018 have been extracted from the CORINE land cover inventory  
168 (Büttner, 2014). In addition, we used the weather type classification from *Electricité de France*  
169 (EDF), corresponding to a daily classification into 8 synoptic situations associated with rainfall  
170 events over France (Garavaglia et al., 2010). This classification is built on geopotential heights  
171 at 700 and 1000 hPa pressure levels associated with rainy days over France.  
172

### 173 **3. Methods**

#### 174 **3.1 Extraction of flood events**

175  
176  
177 We extracted a sample of flood events with a mean occurrence of 1 event per year using a  
178 peaks-over-threshold approach. This type of sampling is chosen since low annual maximum  
179 runoff could be observed during dry years (Farquharson et al., 1992). A de-clustering algorithm  
180 is applied to identify single events to avoid introducing autocorrelation in the analysis and  
181 ensuring that flood events are independent, using two rules (Lang et al., 1999): first a minimum  
182 of  $n$  days between events, with  $n = 5 + \log(\text{catchment area})$  and second, between two  
183 consecutive peaks, runoff must drop below  $\frac{2}{3}$  of the smallest peak. The maximum daily runoff  
184 of each event is kept. This means that for an event lasting several days, only the maximum  
185 daily discharge, and the corresponding date, are kept. Then, for each flood event, we  
186 computed the total rainfall and maximum rainfall. The  $n$ -day previous precipitation is extracted.  
187 Total rainfall for each event is estimated by a cumulative sum of precipitation starting the day  
188 of the flood and this aggregation stops if there are two consecutive days with precipitation  
189 close to zero (1 mm) to account for rainfall intermittency within events. The maximum daily  
190 precipitation is extracted from the same time interval used to compute total event precipitation.  
191 The soil moisture at the beginning of the events is extracted from the previous day of the start  
192 of the rainfall event. A base flow filter has been used to separate direct runoff and base flow  
193 for each time series, using the Lyne Hollick Filter (Lyne and Hollick, 1979), with its default  
194 parameters. For each flood event, the base flow corresponding to the peak has been extracted  
195 to estimate the direct runoff, corresponding to the event rainfall contribution, in addition to base  
196 flow. Different metrics characterizing each flood event have been computed: total rainfall  
197 (mm), event maximum rainfall (mm), duration of the rainfall event (days), duration of the flood  
198 event (days), antecedent soil moisture (0-1) and runoff coefficient (0-1). The runoff coefficient  
199 was computed for each event as the ratio of direct runoff depth and total event precipitation  
200

#### 201 **3.2 Analysis of the mean date of occurrence**

202  
203 Circular statistics (Burn, 1997; Berens, 2009) are used to analyze flood timing. The dates are  
204 first converted into an angular value, then from this sample of angular values, the mean date  
205 of occurrence ( $\theta$ ) can be computed, together with the concentration index ( $r$ ) which measures  
206 the variability of the flood occurrences around the mean date. Using the dates of flood events,  
207 considering hydrological years starting September 1,  $\theta$  and  $r$  are computed from the sample  
208 of dates. The first step in the analysis of seasonality is to test against circular uniformity.  
209 Circular uniformity refers to the case in which all angular values of flood dates around the  
210 circle are equally likely, indicative of the absence of flood seasonality. In that case, the

211 computation of the mean date would have little relevance. The Rayleigh (Fisher, 1993) and  
212 the Hermans-Rasson (Landler et al., 2019) tests are used to test against uniformity for  
213 unimodal distributions, to verify the presence of flood seasonality (ie. meaning that floods do  
214 not occur randomly throughout the year). To associate flood events and weather types, for  
215 each rainy day corresponding to flood events, the weather type has been extracted from the  
216 weather type's classification.

217

### 218 **3.3 Classification of flood generating processes**

219

220 A classification is applied to the flood events, adapted from a previously implemented  
221 classification at the global scale (Stein et al., 2020), in the United States (Stein et al., 2021)  
222 and Africa (Tramblay et al., 2022). This approach relates the occurrence of rainfall amounts  
223 above various thresholds to the occurrence of floods. Flood events in each catchment are  
224 classified according to three hydrometeorological generating processes, namely, the excess  
225 rainfall, short rainfall, long rainfall using a decision tree. Excess rainfall is defined as a flood  
226 event triggered by rainfall higher than average occurring over wet soils (i.e. soil moisture above  
227 than 50% saturation), short rainfall as a single daily rainfall event above high thresholds (the  
228 95th percentile computed over the whole time series of rainfall) and long rainfall as several  
229 consecutive days (> 2 days) with rainfall above the 95th percentile of rainfall summed over 7  
230 days. The classification first evaluates if a larger than average multi-day rainfall fell on wet soil  
231 to determine if the flood event was an excess rainfall type of flood. If that was not the case, it  
232 evaluates whether the thresholds for long rainfall and then short rainfall are exceeded. If no  
233 process could be identified, the class "other" is assigned.

234

### 235 **3.4 Changes in flood characteristics**

236

237 To assess the changes over time in flood dates and generating mechanisms, we split the  
238 records of each station into two periods of equal length. Given that most stations have records  
239 starting after the 1960s, on average the first period is ranging from 1959 to 1990 and the  
240 second one from 1991 to 2021, with a pivot year within +/- 5 years around 1991, allowing the  
241 comparison of the two time slices across the different stations. To assess the relative changes  
242 in the flood drivers, the frequency of each driver for each time period has been computed, and  
243 then we computed for each station the relative change (%) in each driver contribution  
244 (Berghuijs et al., 2019). In addition, to detect trends on the long-term frequency of event types  
245 per year pooled at the regional scale, we used the Mann-Kendall test for trends, modified to  
246 account for autocorrelation in the time series (Hamed and Ramachandra Rao, 1998).

247

248 We use the same approach to estimate changes in the different flood events characteristics,  
249 and we applied the two-tailed Wilcoxon test to check the difference in medians. In addition, to  
250 assess the regional significance of the trends, we also computed the Mann-Kendall test on  
251 flood events characteristics pooled at the regional scale. For flood dates, we computed the  
252 mean dates of occurrence for the two time periods and assessed the significance of the  
253 difference using the Watson and Williams test, which is a circular analogue to the two sample  
254 t-test (Watson and Williams, 1956). Finally, to estimate potential relationships between  
255 different flood characteristics, the Spearman correlation coefficient ( $\rho$ ) is computed.

256

## 257 **4. Results and discussion**

258

#### 259 **4.1 Change in flood event characteristics**

260

261 There are several changes in flood event characteristics as seen in Figure 2 between the two  
262 sub-periods, 1959-1990 and 1991-2021. On average, total event precipitation is increasing in  
263 65 basins (66% of basins), maximum event rainfall is increasing in 76 basins, consistent with  
264 previous studies in this area (Ribes et al., 2019; Trambly et al., 2019; Blanchet and Creutin,  
265 2022), while on the opposite antecedent soil moisture is decreasing in 71 basins, baseflow  
266 contribution is decreasing in 75 basins and runoff coefficient is decreasing in 68 basins. These  
267 changes in soil moisture, base flow and runoff coefficients are consistent with an overall  
268 increase of aridity in southern Europe mostly driven by higher evapotranspiration (Trambly  
269 et al., 2020) and have been also observed in other regions with a similar climate (Ho et al.,  
270 2022). The number of local statistically significant changes for each flood event characteristic  
271 is given in Table 1. These numbers remain small but it should be reminded that sample sizes  
272 are quite short for a robust statistical assessment in a context of high interannual variability.  
273 To overcome this issue, we also assessed the regional significance of these changes in flood  
274 event characteristics. We performed a regional pooling of the events and applied the Mann-  
275 Kendall test to detect trends in the regional series of event characteristics. As shown in table  
276 1, all the detected changes are regionally significant except the decrease in base flow  
277 contribution to peak discharge during floods. Overall, an increase in total event rainfall can be  
278 observed, mostly caused by the increase of maximum rainfall during the events (the changes  
279 in the two variables are correlated, with  $\rho = 0.52$ ), while the flood event durations are on  
280 average decreasing, consistent with studies at the global scale (Wasko et al., 2021).

281

282 These changes in precipitation are associated with a decrease of antecedent soil moisture,  
283 before the beginning of the rainfall events. This decrease is also related to a smaller  
284 contribution of base flow during floods in some basins. There is indeed a significant correlation  
285 between the relative changes in the base flow contribution to peak runoff and soil moisture ( $\rho$   
286 = 0.56), indicating that the soil moisture decrease is likely the main driver of these changes.  
287 There is also for most basins a significant nonlinear relationship (exponential type) between  
288 the flood events antecedent soil moisture and runoff coefficients, as reported in many studies  
289 (Penna et al., 2011; Rogger et al., 2013; Raynaud et al., 2015; Tarasova et al., 2020). Indeed,  
290 for the first time period, 1959-1990 the median Spearman correlation between antecedent soil  
291 moisture and flood runoff coefficients (see supplementary figure S1) is equal to 0.43 and is  
292 significant at the 5% level in 56 basins (67 basins at the 10% significance level). For the  
293 second time period, 1991-2021, the median correlation is increasing to 0.45 and is significant  
294 in 64 basins at the 5% significance level (68 at the 10% significance level). These results  
295 show, contrary to popular belief, that at the catchment scale drier soils produce less runoff,  
296 and this characteristic is even slightly accentuating over time. Indeed, increased runoff  
297 coefficients induced either by hydrophobic soil conditions following droughts (Burch et al.,  
298 1989), soil crusting and sealing (Bissonnais and Singer, 1993) or compaction (Alaoui et al.,  
299 2018), are well documented processes that mostly occur at the local plot scale that do not  
300 produce discernible effects at the catchment scale. This observation is reinforced by the fact  
301 that no negative correlation between runoff coefficients and initial soil moisture was detected.  
302 About the explanatory factors of the association between soil moisture and runoff coefficients,  
303 we found stronger correlations (significant at the 5% level) between these two variables in  
304 catchments with higher percentage of urban or agricultural areas, and on the opposite weaker

305 correlations along with increased percentage of forests or mean catchment altitude. There is  
306 only a very small increase of the correlations for larger basins (no significant correlation with  
307 basin sizes), indicating that this relationship between soil moisture and runoff coefficient  
308 remains valid for all basins scales considered in the present study. This relationship between  
309 runoff coefficients and antecedent soil moisture remained stable between 1959-1990 and  
310 1991-2021 (Figure 3).

311

#### 312 **4.2 Changes in flood dates**

313

314 Floods in southern France tend to occur mainly during November or December for basins  
315 close to the Mediterranean, East of the Cévennes mountainous range, while for basins located  
316 on the western part of the region, they tend to occur later during winter months, centered in  
317 January or February (Figure 4). Both the Rayleigh and Hermans-Rasson tests reject the null  
318 hypothesis of uniformity at the 5% level, indicating that floods do not occur randomly  
319 throughout the year. In most cases, the seasonal distribution is unimodal, except for a few  
320 cases; in about 15 stations the maximum occurrence of floods is observed in late autumn or  
321 winter and a secondary minor peak of occurrence is observed, usually centered around the  
322 month of March or April. These floods are associated with rainfall events rather than snowmelt,  
323 since for only 3 basins the snowfall contribution reaches 19% of total precipitation whereas  
324 the snowfall contribution is much lower for the remaining 12 basins (less than 5%).

325

326 For 79 basins, floods tend to occur earlier during the year, on average by -22 days between  
327 1959-1990 and 1991-2021 (Figure 5, left panel). On the opposite, for 19 basins the mean flood  
328 date occurs later in the second period with an average of +12 days. These changes in the  
329 mean date are significant in 26 basins at the 0.1 level according to the Watson and Williams  
330 test (18 basins at the 0.05 significance level). There are two differentiated spatial patterns: all  
331 basins where floods tend to occur earlier are located widespread in the center of the  
332 Mediterranean region, and basins where floods tend to occur later are found only in the  
333 northwestern margin of the domain. In these basins, the mean floods occur in late winter, until  
334 February and March. The same spatial patterns of changes in mean flood dates have been  
335 observed by Blöschl et al. (2017), but without providing the possible causes of these seasonal  
336 shifts. For the concentration index (i.e., the variability around the mean date) similarly two  
337 different patterns are found: for basins where floods tend to occur earlier, the concentration  
338 index is increasing, meaning more floods are clustered around the mean date, while for  
339 western basins where floods tend to occur later, the concentration index is decreasing,  
340 meaning a larger variability in flood dates (Figure 5, right panel).

341

342 To assess the regional changes in flood dates, we first separated in two regional samples the  
343 stations where floods occur earlier (sample 1) or later (sample 2). Then we used the Watson-  
344 William test, previously used to assess changes in flood dates in each station, to compare  
345 these two regional samples. The test results indicate that for the 19 stations where floods tend  
346 to occur later, the change in flood dates are not significant at the 5% level ( $p$  value = 0.0821),  
347 on the opposite, for the 79 stations where floods are occurring earlier, the change is significant  
348 ( $p$  value =  $5.34 \cdot 10^{-8}$ ).

349

#### 350 **4.3 Associations between flood occurrence and weather patterns**

351



352 The seasonal patterns observed for the floods are closely related to the occurrence of different  
353 weather types in different sub-regions. As shown in figure 6, most basins located east of the  
354 Cévennes mountainous range have floods associated with WT4, Southern Circulation, and  
355 western basins with WT2, Steady Oceanic. The most frequent pattern associated with 37% of  
356 floods, WT4, is known to be triggering intense rainfall events in this region (Ducrocq et al.,  
357 2008; Tramblay et al., 2013). Interestingly, the WT6, Eastern circulation, and WT7,  
358 Southwestern circulation, are both associated to a lesser extent with floods across the whole  
359 region, but without notable spatial differences in the relative frequency of floods associated  
360 with these weather types. Change in flood seasonality could be ascribed to changes in the  
361 seasonal occurrence of the weather types (Figure 7): WT4 tends to occur more frequently  
362 from March to August during 1991-2021 compared to 1959-1990, and these changes are  
363 statistically significant (see supplementary figure S2). When looking at the actual count of WT4  
364 days, this change represents an increase of 69 events during that 6-month period for 1991-  
365 2021, so an average of +2.2 days per year. Associated with a warmer Mediterranean Sea over  
366 the last decades notably during summer (Pastor et al., 2020), the combination of these two  
367 factors could explain the earlier occurrence of floods east of the Cévennes mountainous  
368 range. Similarly, there is an increased frequency of WT2 in January, February and March  
369 between 1991-2021 and 1959-1990, that is also significant (supplementary figure S2) that  
370 could be possibly related to the later occurrence of floods west of the Cévennes range.  
371 Although this change in weather types seasonality leading to heavy rainfalls is a plausible  
372 cause of the observed changes in the flood seasonality, more research is needed to better  
373 understand these relationships and attribute changes in flood seasonality. Notably, to analyze  
374 in more detail the moisture supply from the Mediterranean or Atlantic seas, the interaction with  
375 the atmospheric thermodynamics, the duration, localization and the spatial dependence of the  
376 rainfall episodes inducing floods.

#### 377 378 **4.4 Changes in flood generating processes**

379  
380 When first applying the classification of flood-generating processes on all floods, we find a  
381 predominance of excess rainfall events (Figure 8), followed by long rain and short rains, that  
382 is consistent with the known flood-generating processes in this region (Mediterranean  
383 episodes) and, in particular, the strong influence of saturated soils on runoff generation with  
384 floods mostly occurring during the autumn (Tramblay et al., 2010, 2019). The category 'other'  
385 regroups only 0.97% of floods and it represents mostly events with very low precipitation  
386 amounts, likely due to an underestimation of rainfall in the SAFRAN database for some events.  
387 It is worth noticing that despite the large sample of basins considered, the patterns are  
388 remarkably consistent and homogeneous across different basin sizes and locations. There is  
389 a significant, yet low, correlation ( $\rho = 0.26$ ,  $p$ -value = 0.008) between the ratio of excess rain  
390 floods and catchment size, with a larger proportion of excess rain in larger basins, while on  
391 the opposite there is an even weaker and negative correlation ( $\rho = -0.16$ ,  $p$ -value = 0.09)  
392 between the ratio of short rain and basin size. It should be noted that floods driven by short  
393 rain episodes are potentially affecting smaller regions than floods driven by excess floods  
394 (Brunner and Dougherty, 2022). For 30 basins (not necessarily the largest ones), the  
395 proportion of excess rain exceeds 80% of the total number of flood events (see supplementary  
396 figure S3). For short rain and long rain, the maximum contributions observed much lower, 36%  
397 and 32%, respectively, but these maximum values are only found in small basins. Indeed,

398 basins when short rain or long rain exceed 30% of episodes are only found in basins smaller  
399 than 100 km<sup>2</sup>.

400

401 The mean date of occurrence is significantly different between the three flood types, according  
402 to the Watson and Williams test. As shown on figure 9, the highest proportion of floods induced  
403 by short rain is observed during September to November, while the floods induced by long  
404 rain are mostly occurring during October to December, and excess rain floods are observed  
405 in late autumn and winter, with a peak in February. This is consistent with the annual soil  
406 moisture cycle in this region: at the end of the summer the soils are dry and it takes several  
407 months to replenish the soil moisture level, which is at highest during winter. If examining the  
408 long-term changes in this monthly repartition of flood types (see supplementary figure S4), the  
409 frequency of excess rain is decreasing from February to April, and also in October, while  
410 increasing during winter months. This implies that the season during which excess rain floods  
411 are occurring is reducing in length and concentrated during wet winter months. On the  
412 opposite the frequency of short or long rain floods is increasing in June and September,  
413 months that are getting drier over time in this region.

414

415 The noticeable changes in flood processes over time are a reduction of Excess rainfall in 71  
416 basins and an increased frequency of short rains in 53 basins and Long rains in 63 basins  
417 (Figure 10), while short rain and long rain floods are decreasing for 19 and 22 basins,  
418 respectively. For excess rain, there are only 16 basins where their relative proportion is  
419 increasing; they are mostly located on the margins of the Alps and Pyrenees mountains. For  
420 more frequent events (ie. if considering an average of 3 episodes per year instead of one), the  
421 number of basins with a change is larger, with a reduction of Excess rainfall in 82 basins out  
422 of 98 (results not shown). This indicates that the soil moisture depletion has more impacts on  
423 small to moderate flood events, as previously observed by Bertola et al. (2021). There is no  
424 relationship between the rate of change in the different flood generating processes and  
425 catchment sizes indicating a clear regional pattern. The average magnitude across all basins  
426 of these changes remains low, on average -4.1% for excess rain, +1.2% for short rain and +  
427 2.1% long rain. Yet, the magnitude of these changes is ranging from +15% to -21% for excess  
428 rain, +11% to -20% for short rain and +12% to -11% for long rain, depending on the catchment,  
429 indicating that local catchment characteristics could strongly modulate the regional signal. In  
430 addition, the average values over the whole domain are hiding some local changes: for  
431 instance, short rainfall floods are increasing in the southeastern part of the Cévennes while  
432 decreasing for the northwestern part as seen in Figure 10.

433

#### 434 **4.5 Regional changes**

435

436 To assess whether the changes in the relative influence of the three different flood types are  
437 significant at the regional scale, we computed for each year the relative frequency of the  
438 different flood types, all basins together. It is indeed not possible to do this analysis for each  
439 station independently, due to the small size of the samples over the two periods. These  
440 changes in the occurrence of flood types are significant at the regional scale according to the  
441 Mann-Kendall test (Figure 11), for the frequency of excess rain floods and short rain floods, at  
442 the 5% significance level, but not for the long rain floods. All events pooled regionally, the  
443 decrease in excess rain floods is equal to -13% between 1959-1990 and 1991-2021, and the  
444 increase of short rain floods is equal to +36%. In addition, to assess whether these results are  
445 robust to the thresholds used in the classification of flood events, a Monte Carlo experiment

446 has been also conducted. Results show (see supplementary figure S5) that regional changes  
447 in excess rain and short rain floods are not dependent on classification thresholds, while it is  
448 not the case for long rain floods. In terms of flood severity for the different flood types, the  
449 median flood computed for each basin is strongly correlated to basin size ( $\rho = 0.78$ ) for floods  
450 caused by excess rain, short rain ( $\rho = 0.80$ ) and long rain ( $\rho = 0.75$ ); and very similar results  
451 are found for the maximum flood. On the contrary, the specific discharge of flood peaks is non  
452 linearly related to basin sizes, with a clear threshold effect for basins smaller than 500 km<sup>2</sup>  
453 that have a much larger specific discharge than larger basins.

454  
455 Given that there are different flood sample sizes in the different basins corresponding to  
456 different flood-generating processes, we pooled regionally the flood events. To do so, we  
457 computed the specific discharge for each event (i.e. the flood magnitude divided by catchment  
458 area) to analyze the distributions of specific discharge for all the events associated with excess  
459 rain, long rain or short rain. Specific discharge is used herein since it is a good indicator of  
460 flash floods severity, notably in this Mediterranean region (Delrieu et al., 2005, Ruin et al.,  
461 2008). Figure 12 shows that the short rain floods are more severe, in terms of specific  
462 discharge, than excess rain or long rain floods at the regional level (as shown also by Tarasova  
463 et al., 2023). The regional distributions for the different event types are different according to  
464 the Kolmogorov-Smirnov test. It must be noted that for a given basin the magnitude of the  
465 different types of floods may not be very different, showing the strong variability from one event  
466 to another that is not solely linked to the flood trigger. When comparing the different flood  
467 distributions between the time periods 1959-1990 and 1991-2021, the differences in flood  
468 magnitudes between excess rain, long and short rain are reduced. This is mainly due to a  
469 slight decrease in the specific discharge of short rain floods, notably for flood events with a  
470 return level higher than 10 years, while the excess rain floods show very little changes in  
471 intensity over time.

## 472 473 **5. Conclusions**

474  
475 The aim of this study was to document the evolution of the characteristics of Mediterranean  
476 floods, using long river discharge records in southern France, [a high-resolution climate and  
477 surface reanalysis that is used operationally to monitor water resources and extremes in  
478 France, and a classification of weather patterns. This large regional database with an average  
479 of 50 years of records allowed to detect regional significant changes on several flood events  
480 characteristics](#). In most basins, floods tend to occur earlier during the year, the mean flood  
481 date being on average advanced by one month. This seasonal shift can be attributed to the  
482 increased frequency of southern-circulation weather types during spring and summer that are  
483 strongly associated with the occurrence of floods in this region. Over time, floods also tend to  
484 be more clustered in time over the different basins, as reflected by a decreasing variability in  
485 flood occurrence throughout the year. On the contrary, for the westernmost basins influenced  
486 by Atlantic circulation patterns, floods tend to occur later during the year, also due to a  
487 seasonal shift of the flood-generating circulation patterns that are occurring more frequently in  
488 late winter. [These results indicate that changes in large-scale atmospheric circulation  
489 described by weather types is the likely driver of the seasonal shift of flood dates previously  
490 observed for French Mediterranean regions \(Blöschl et al., 2017\).](#)

491 During floods, an increase in total and extreme event precipitation has been observed,  
492 associated with a decrease of antecedent soil moisture before rainfall events, linked to a  
493 smaller contribution of base flow during floods. It can be concluded that it is the depletion of  
494 soil water content, due to increased aridity in south France notably related to higher  
495 evapotranspiration rates (Tramblay et al., 2020), that is the likely cause of the absence of flood  
496 trends in this region despite the increase in extreme rainfall. It should be also noted that over  
497 all basins, dryer soils are associated with lower runoff coefficients, and this relationship  
498 remains valid over time. This result contrasts sharply with the generally accepted idea that in  
499 a drier climate we observe stronger runoff. While this statement could be valid at the plot scale,  
500 we show herein that it is an opposite relationship found for the whole range of catchment sizes  
501 considered in the present work. The majority of flood events are associated with excess rainfall  
502 on saturated soils, but that proportion is decreasing over time with a concurrent increased  
503 frequency of short rain, potentially leading to more severe floods, as previously shown at the  
504 European scale by Tarasova et al. (2023). At the regional scale, we show that floods induced  
505 by short rains are indeed of higher magnitude, but due to a lower runoff coefficient induced by  
506 drier antecedent soil moisture, the specific discharge associated with short rain flood is also  
507 decreasing over time. These results are consistent with those obtained in other regions of the  
508 world, showing that floods do not necessarily increase with the increase in extreme  
509 precipitation, and that soil moisture seems to play a key role in explaining these changes and  
510 the lack of trends ultimately on flood hazard (Wasko and Nathan, 2019; Bertola et al., 2021;  
511 Wasko et al., 2021). The results of the present study are rather homogeneous given the  
512 different catchment sizes and land use types, indicating that changes in flood types are mainly  
513 resulting from regional climate change and not only local changes, such as land cover or  
514 agricultural practice changes, nor the increase of urban and peri-urban areas. Nonetheless, if  
515 the observed trend in increased short rain floods is persisting in the upcoming decades, the  
516 severity of floods, particularly the most important ones, could increase along with the rise in  
517 rainfall extremes particularly in areas where the soil infiltration potential is low, such as in  
518 mountainous or urbanized areas, that have expanded a lot in recent years in Mediterranean  
519 regions. For other types of basins, notably in lowland areas with agricultural or natural  
520 landscapes, caution should be exerted before extrapolating such hazards in the future, since  
521 we show herein a potential reduction over time of the specific runoff even for short rain floods.  
522 This aspect could be further investigated using climate scenarios.

523 One of the main perspectives of this work would be to perform a similar analysis at sub-daily  
524 time steps, that would be more adapted to analyze changes in flash floods characteristics,  
525 notably in terms of the flashiness response of the catchments (Baker et al., 2004; Li et al.,  
526 2022). Indeed, the daily time step prevents a thorough analysis of changes in rainfall patterns,  
527 notably at shorter time steps. Yet, there is no gridded dataset of hourly precipitation before the  
528 2000s in Southern France, that does not allow to repeat such a similar study over 62 years.  
529 However, given the availability of radar rainfall over France, it would be possible to analyze,  
530 at least for the recent years after 2000, the evolution of several characteristics, such as the  
531 evolution of storm hydrographs, concentration time and the flashiness response of the basins.  
532 Another relevant prospective work would be to analyze the spatial extent of floods. Given the  
533 future evolution of weather types associated with floods in combination with more local to  
534 regional characteristics, such as soil moisture state, these types of events may impact  
535 simultaneously wider, or smaller, parts of the region considered, and this could have serious  
536 implications on risk management (Brunner et al., 2021; Brunner and Dougherty, 2022).  
537 Therefore, the joint analysis of flood occurrence in nearby basins would be highly relevant.

538 Finally, there is also a need for new approaches to incorporate these changes in flood  
539 generating process into engineering practice (Slater et al., 2021b), notably to estimate the  
540 return levels for different types of infrastructure design.

541

#### 542 **Acknowledgements**

543 This study is a contribution to the EXPLORE2 project funded by the French Ministry of  
544 Ecological Transition. The access to the Météo France and Banque Hydro databases is  
545 acknowledged. The authors wish to thanks 3 anonymous reviewers and the associated Editor  
546 Manuela Brunner for their comments.

#### 547 **Data availability**

548 The computed catchment-based indicators for each flood events in all basins are accessible  
549 from the online repository: <https://doi.org/10.5281/zenodo.8075639>  
550 are available upon request to the corresponding author.

551 The complete time series of river runoff can be accesses freely here:  
552 <https://hydro.eaufrance.fr/>

553 The SAFRAN/ISBA outputs can be ordered freely for scientific purpose here:  
554 <https://donneespubliques.meteofrance.fr>

555

#### 556 **Author contributions**

557 Y.T.: Conceptualization, investigation, data curation, formal analysis, writing—original draft  
558 preparation. YT designed the experiments, performed the analyses, and wrote the paper. PA,  
559 GA, ML, EP, LN and ES Conceptualization, data curation, writing—reviewing and editing.

#### 560 **Competing interests**

561 The authors declare that they have no conflict of interest.

562

563

564

565

566

567

568

569

a mis en forme : Anglais (États-Unis)

a mis en forme : Anglais (États-Unis)

Code de champ modifié

570 Table 1: Number of local significant changes in the median of flood events characteristics  
 571 detected by the Wilcoxon test and results of the regional Mann-Kendall test on flood event  
 572 characteristics  
 573

<b>Indicator</b>	<b>Number of significant local changes (Wilcoxon, 10%)</b>	<b>pvalue of the regional MK test</b>	<b>Regional changes between 1959 and 2022 (%)</b>
Flood event duration	17	<b>0.0046178</b>	-0.40%
Base flow contribution to peak	15	<b>0.5687962</b>	-8.62%
Runoff coefficient	19	<b>0.0000002</b>	-14.62%
Total event rainfall	16	<b>0.0011851</b>	9.01%
Maximum event rainfall	27	<b>0.0000000</b>	13.47%
Antecedent soil moisture	12	<b>0.0000008</b>	-9.80%

574  
 575  
 576  
 577  
 578  
 579  
 580  
 581  
 582  
 583  
 584  
 585  
 586  
 587  
 588  
 589  
 590  
 591  
 592  
 593  
 594  
 595  
 596  
 597  
 598

**References**

599  
600 Alaoui, A., Rogger, M., Peth, S., and Blöschl, G.: Does soil compaction increase floods? A  
601 review, *Journal of Hydrology*, 557, 631–642, <https://doi.org/10.1016/j.jhydrol.2017.12.052>,  
602 2018.  
603  
604 Ali, E., Cramer, W., Carnicer, J., Georgopoulou, E., Hilmi, N. J. M., Cozannet, G. L., and  
605 Lionello, P.: Cross-Chapter Paper 4: Mediterranean Region, edited by: Pörtner, H. O.,  
606 Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M.,  
607 Langsdorf, S., Lösckke, S., Möller, V., Okem, A., and Rama, B., *Climate Change 2022:*  
608 *Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth*  
609 *Assessment Report of the Intergovernmental Panel on Climate Change*, 2233–2272,  
610 <https://doi.org/10.1017/9781009325844.021.2233>, 2022.  
611  
612 Baker, D. B., Richards, R. P., Loftus, T. T., and Kramer, J. W.: A new flashiness index:  
613 characteristics and applications to midwestern rivers and streams, *J Am Water Resources*  
614 *Assoc*, 40, 503–522, <https://doi.org/10.1111/j.1752-1688.2004.tb01046.x>, 2004.  
615  
616 Berens, P.: *CircStat: A MATLAB Toolbox for Circular Statistics*, *J. Stat. Soft.*, 31,  
617 <https://doi.org/10.18637/jss.v031.i10>, 2009.  
618  
619 Berghuijs, W. R., Harrigan, S., Molnar, P., Slater, L. J., and Kirchner, J. W.: The Relative  
620 Importance of Different Flood-Generating Mechanisms Across Europe, *Water Resour. Res.*,  
621 2019WR024841, <https://doi.org/10.1029/2019WR024841>, 2019.  
622  
623 Bertola, M., Viglione, A., Vorogushyn, S., Lun, D., Merz, B., and Blöschl, G.: Do small and  
624 large floods have the same drivers of change? A regional attribution analysis in Europe,  
625 *Hydrol. Earth Syst. Sci.*, 25, 1347–1364, <https://doi.org/10.5194/hess-25-1347-2021>, 2021.  
626  
627 Bissonnais, Y. L. and Singer, M. J.: Seal Formation, Runoff, and Interrill Erosion from  
628 Seventeen California Soils, *Soil Science Society of America Journal*, 57, 224–229,  
629 <https://doi.org/10.2136/sssaj1993.03615995005700010039x>, 1993.  
630  
631 Blanchet, J. and Creutin, J.-D.: Instrumental agreement and retrospective analysis of trends  
632 in precipitation extremes in the French Mediterranean Region, *Environ. Res. Lett.*, 17,  
633 074011, <https://doi.org/10.1088/1748-9326/ac7734>, 2022.  
634  
635 Blöschl, G., Gaál, L., Hall, J., Kiss, A., Komma, J., Nester, T., Parajka, J., Perdigão, R. A. P.,  
636 Plavcová, L., Rogger, M., Salinas, J. L., and Viglione, A.: Increasing river floods: fiction or  
637 reality?: Increasing river floods, *WIREs Water*, 2, 329–344,  
638 <https://doi.org/10.1002/wat2.1079>, 2015.  
639  
640 Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A. P., Merz, B., Arheimer, B., Aronica, G. T.,  
641 Bilibashi, A., Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G. B., Claps, P.,  
642 Fiala, K., Frolova, N., Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss,  
643 A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-  
644 Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M.,  
645 Ovcharuk, V., Radevski, I., Rogger, M., Salinas, J. L., Sauquet, E., Šraj, M., Szolgay, J.,

646 Viglione, A., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate shifts timing  
647 of European floods, *Science*, 357, 588–590, <https://doi.org/10.1126/science.aan2506>, 2017.  
648

649 Blöschl, G., Hall, J., Viglione, A., Perdigão, R. A. P., Parajka, J., Merz, B., Lun, D., Arheimer,  
650 B., Aronica, G. T., Bilibashi, A., Boháč, M., Bonacci, O., Borga, M., Čanjevac, I., Castellarin,  
651 A., Chirico, G. B., Claps, P., Frolova, N., Ganora, D., Gorbachova, L., Gül, A., Hannaford, J.,  
652 Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O.,  
653 Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A.,  
654 Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Salinas, J. L., Sauquet, E., Šraj, M.,  
655 Szolgay, J., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate both  
656 increases and decreases European river floods, *Nature*, 573, 108–111,  
657 <https://doi.org/10.1038/s41586-019-1495-6>, 2019.  
658

659 Brunner, M. I. and Dougherty, E. M.: Varying Importance of Storm Types and Antecedent  
660 Conditions for Local and Regional Floods, *Water Resources Research*, 58,  
661 <https://doi.org/10.1029/2022WR033249>, 2022.  
662

663 Brunner, M. I., Slater, L., Tallaksen, L. M., and Clark, M.: Challenges in modeling and  
664 predicting floods and droughts: A review, *WIREs Water*, 8,  
665 <https://doi.org/10.1002/wat2.1520>, 2021.  
666

667 Brunner, M. I., Swain, D. L., Wood, R. R., Willkofer, F., Done, J. M., Gilleland, E., and  
668 Ludwig, R.: An extremeness threshold determines the regional response of floods to  
669 changes in rainfall extremes. *Communications Earth & Environment*, 2:1,  
670 <https://doi.org/10.1038/s43247-021-00248-x>, 2021  
671

672 Burch, G. J., Moore, I. D., and Burns, J.: Soil hydrophobic effects on infiltration and  
673 catchment runoff, *Hydrol. Process.*, 3, 211–222, <https://doi.org/10.1002/hyp.3360030302>,  
674 1989.  
675

676 Burn, D. H.: Catchment similarity for regional flood frequency analysis using seasonality  
677 measures, *Journal of Hydrology*, 202, 212–230, [https://doi.org/10.1016/S0022-1694\(97\)00068-1](https://doi.org/10.1016/S0022-1694(97)00068-1), 1997.  
678

679 Büttner, G.: CORINE Land Cover and Land Cover Change Products, in: *Land Use and Land  
680 Cover Mapping in Europe*, vol. 18, edited by: Manakos, I. and Braun, M., Springer  
681 Netherlands, Dordrecht, 55–74, [https://doi.org/10.1007/978-94-007-7969-3\\_5](https://doi.org/10.1007/978-94-007-7969-3_5), 2014.  
682

683 Cao, Q., Gershunov, A., Shulgina, T., Ralph, F. M., Sun, N., and Lettenmaier, D. P.: Floods  
684 due to Atmospheric Rivers along the U.S. West Coast: The Role of Antecedent Soil Moisture  
685 in a Warming Climate, *Journal of Hydrometeorology*, 21, 1827–1845,  
686 <https://doi.org/10.1175/JHM-D-19-0242.1>, 2020.  
687

688 Delrieu, G., Nicol, J., Yates, E., Kirstetter, P.-E., Creutin, J.-D., Anquetin, S., Obled, C.,  
689 Saulnier, G.-M., Ducrocq, V., Gaume, E., Payrastré, O., Andrieu, H., Ayrat, P.-A., Bouvier,  
690 C., Neppel, L., Livet, M., Lang, M., du-Châtelet, J. P., Walpersdorf, A., and Wobrock, W.:  
691 The Catastrophic Flash-Flood Event of 8–9 September 2002 in the Gard Region, France: A  
692 First Case Study for the Cévennes–Vivarais Mediterranean Hydrometeorological  
693



694 Observatory. *Journal of Hydrometeorology*, 6:1, 34–52, <https://doi.org/10.1175/jhm-400.1>,  
695 2005.  
696  
697 Ducrocq, V., Nuissier, O., Ricard, D., Lebeau-pin, C., and Thouvenin, T.: A numerical study of  
698 three catastrophic precipitating events over southern France. II: Mesoscale triggering and  
699 stationarity factors, *Q.J.R. Meteorol. Soc.*, 134, 131–145, <https://doi.org/10.1002/qj.199>,  
700 2008.  
701  
702 Farquharson, F. A. K., Meigh, J. R., and Sutcliffe, J. V.: Regional flood frequency analysis in  
703 arid and semi-arid areas, *Journal of Hydrology*, 138, 487–501, [https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-1694(92)90132-F)  
704 1694(92)90132-F, 1992.  
705  
706 Fisher, N. I.: *Statistical analysis of circular data*, Cambridge University Press, Cambridge  
707 [England]; New York, NY, USA, 277 pp., 1993.  
708  
709 Garavaglia, F., Gailhard, J., Paquet, E., Lang, M., Garçon, R., and Bernardara, P.:  
710 Introducing a rainfall compound distribution model based on weather patterns sub-sampling,  
711 *Hydrol. Earth Syst. Sci.*, 14, 951–964, <https://doi.org/10.5194/hess-14-951-2010>, 2010.  
712  
713 Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M., Soubeyroux, J.-M. :  
714 A 50-year high-resolution atmospheric reanalysis over France with the Safran system.  
715 *International Journal of Climatology*, 2010, 30 (11), 1627-1644.  
716 <https://doi.org/10.1002/joc.2003>, 2010.  
717  
718 Hamed, K. H. and Ramachandra Rao, A.: A modified Mann-Kendall trend test for  
719 autocorrelated data, *Journal of Hydrology*, 204, 182–196, [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-1694(97)00125-X)  
720 1694(97)00125-X, 1998.  
721  
722 Ho, M., Nathan, R., Wasko, C., Vogel, E., and Sharma, A.: Projecting changes in flood event  
723 runoff coefficients under climate change, *Journal of Hydrology*, 615, 128689,  
724 <https://doi.org/10.1016/j.jhydrol.2022.128689>, 2022.  
725  
726 Huang, H., Fischella, M. R., Liu, Y., Ban, Z., Fayne, J. V., Li, D., Cavanaugh, K. C., and  
727 Lettenmaier, D. P.: Changes in Mechanisms and Characteristics of Western U.S. Floods  
728 Over the Last Sixty Years, *Geophysical Research Letters*, 49,  
729 <https://doi.org/10.1029/2021GL097022>, 2022.  
730  
731 Jiang, S., Bevacqua, E., and Zscheischler, J.: River flooding mechanisms and their changes  
732 in Europe revealed by explainable machine learning, *Hydrol. Earth Syst. Sci.*, 26, 6339–  
733 6359, <https://doi.org/10.5194/hess-26-6339-2022>, 2022.  
734  
735 Kemter, M., Merz, B., Marwan, N., Vorogushyn, S., and Blöschl, G.: Joint Trends in Flood  
736 Magnitudes and Spatial Extents Across Europe, *Geophys. Res. Lett.*, 47,  
737 <https://doi.org/10.1029/2020GL087464>, 2020.  
738  
739 Landler, L., Ruxton, G. D., and Malkemper, E. P.: The Hermans–Rasson test as a powerful  
740 alternative to the Rayleigh test for circular statistics in biology, *BMC Ecol*, 19, 30,  
741 <https://doi.org/10.1186/s12898-019-0246-8>, 2019.

a mis en forme : Français (France)

742  
743 Lang, M., Ouarda, T. B. M. J., and Bobée, B.: Towards operational guidelines for over-  
744 threshold modeling, *Journal of Hydrology*, 225, 103–117, [https://doi.org/10.1016/S0022-](https://doi.org/10.1016/S0022-1694(99)00167-5)  
745 1694(99)00167-5, 1999.  
746  
747 Li, Z., Gao, S., Chen, M., Gourley, J. J., Liu, C., Prein, A. F., and Hong, Y.: The  
748 conterminous United States are projected to become more prone to flash floods in a high-  
749 end emissions scenario, *Commun Earth Environ*, 3, 86, [https://doi.org/10.1038/s43247-022-](https://doi.org/10.1038/s43247-022-00409-6)  
750 00409-6, 2022.  
751  
752 Liu, J., Feng, S., Gu, X., Zhang, Y., Beck, H. E., Zhang, J., and Yan, S.: Global changes in  
753 floods and their drivers, *Journal of Hydrology*, 614, 128553,  
754 <https://doi.org/10.1016/j.jhydrol.2022.128553>, 2022.  
755  
756 Lyne, V. D. and Hollick, M.: Stochastic time-variable rainfall runoff modelling, *Hydrology and*  
757 *Water Resources Symposium*, Institution of Engineers, Australia, Perth (1979), Barton,  
758 Australia, 1979.  
759  
760 Mao, Y., Zhou, T., Leung, L. R., Tesfa, T. K., Li, H., Wang, K., Tan, Z., and Getirana, A.:  
761 Flood Inundation Generation Mechanisms and Their Changes in 1953–2004 in Global Major  
762 River Basins, *J. Geophys. Res. Atmos.*, 124, 11672–11692,  
763 <https://doi.org/10.1029/2019JD031381>, 2019.  
764  
765 Merz, B., Aerts, J., Arnbjerg-Nielsen, K., Baldi, M., Becker, A., Bichet, A., Blöschl, G.,  
766 Bouwer, L. M., Brauer, A., Cioffi, F., Delgado, J. M., Gocht, M., Guzzetti, F., Harrigan, S.,  
767 Hirschboeck, K., Kilsby, C., Kron, W., Kwon, H.-H., Lall, U., Merz, R., Nissen, K., Salvatti, P.,  
768 Swierczynski, T., Ulbrich, U., Viglione, A., Ward, P. J., Weiler, M., Wilhelm, B., and Nied, M.:  
769 Floods and climate: emerging perspectives for flood risk assessment and management, *Nat.*  
770 *Hazards Earth Syst. Sci.*, 14, 1921–1942, <https://doi.org/10.5194/nhess-14-1921-2014>,  
771 2014.  
772  
773 Pastor, F., Valiente, J. A., and Khodayar, S.: A Warming Mediterranean: 38 Years of  
774 Increasing Sea Surface Temperature, *Remote Sensing*, 12, 2687,  
775 <https://doi.org/10.3390/rs12172687>, 2020.  
776  
777 Penna, D., Tromp-van Meerveld, H. J., Gobbi, A., Borga, M., and Dalla Fontana, G.: The  
778 influence of soil moisture on threshold runoff generation processes in an alpine headwater  
779 catchment, *Hydrol. Earth Syst. Sci.*, 15, 689–702, <https://doi.org/10.5194/hess-15-689-2011>,  
780 2011.  
781  
782 Raynaud, D., Thielen, J., Salamon, P., Burek, P., Anquetin, S., and Alfieri, L.: A dynamic  
783 runoff co-efficient to improve flash flood early warning in Europe: evaluation on the 2013  
784 central European floods in Germany: Dynamic runoff co-efficient for flash flood early warning  
785 in Europe, *Met. Apps*, 22, 410–418, <https://doi.org/10.1002/met.1469>, 2015.  
786  
787 Ribes, A., Thao, S., Vautard, R., Dubuisson, B., Somot, S., Colin, J., Planton, S., and  
788 Soubeyroux, J.-M.: Observed increase in extreme daily rainfall in the French Mediterranean,  
789 *Clim Dyn*, 52, 1095–1114, <https://doi.org/10.1007/s00382-018-4179-2>, 2019.

790  
791 Rogger, M., Viglione, A., Derx, J., and Blöschl, G.: Quantifying effects of catchments storage  
792 thresholds on step changes in the flood frequency curve: Step Changes in the Flood  
793 Frequency Curve, *Water Resour. Res.*, 49, 6946–6958, <https://doi.org/10.1002/wrcr.20553>,  
794 2013.  
795  
796 Ruin, I., Creutin, J.-D., Anquetin, S., & Lutoff, C.: Human exposure to flash floods – Relation  
797 between flood parameters and human vulnerability during a storm of September 2002 in  
798 Southern France. *Journal of Hydrology*, 361:1-2, 199–213,  
799 <https://doi.org/10.1016/j.jhydrol.2008.07.044>, 2008.  
800  
801 Sharma, A., Wasko, C., and Lettenmaier, D. P.: If Precipitation Extremes Are Increasing,  
802 Why Aren't Floods?, *Water Resour. Res.*, 54, 8545–8551,  
803 <https://doi.org/10.1029/2018WR023749>, 2018.  
804  
805 Slater, L., Villarini, G., Archfield, S., Faulkner, D., Lamb, R., Khouakhi, A., and Yin, J.: Global  
806 Changes in 20-Year, 50-Year, and 100-Year River Floods, *Geophysical Research Letters*,  
807 48, <https://doi.org/10.1029/2020GL091824>, 2021a.  
808  
809 Slater, L. J., Anderson, B., Buechel, M., Dadson, S., Han, S., Harrigan, S., Kelder, T., Kowal,  
810 K., Lees, T., Matthews, T., Murphy, C., and Wilby, R. L.: Nonstationary weather and water  
811 extremes: a review of methods for their detection, attribution, and management, *Hydrol.*  
812 *Earth Syst. Sci.*, 25, 3897–3935, <https://doi.org/10.5194/hess-25-3897-2021>, 2021b.  
813  
814 Stein, L., Pianosi, F., and Woods, R.: Event-based classification for global study of river  
815 flood generating processes, *Hydrological Processes*, 34, 1514–1529,  
816 <https://doi.org/10.1002/hyp.13678>, 2020.  
817  
818 Stein, L., Clark, M. P., Knoben, W. J. M., Pianosi, F., and Woods, R. A.: How Do Climate  
819 and Catchment Attributes Influence Flood Generating Processes? A Large-Sample Study for  
820 671 Catchments Across the Contiguous USA, *Water Res*, 57,  
821 <https://doi.org/10.1029/2020WR028300>, 2021.  
822  
823 Tarasova, L., Merz, R., Kiss, A., Basso, S., Blöschl, G., Merz, B., Viglione, A., Plötner, S.,  
824 Guse, B., Schumann, A., Fischer, S., Ahrens, B., Anwar, F., Bárdossy, A., Bühler, P.,  
825 Haberlandt, U., Kreibich, H., Krug, A., Lun, D., Müller-Thomy, H., Pidoto, R., Primo, C.,  
826 Seidel, J., Vorogushyn, S., and Wietzke, L.: Causative classification of river flood events,  
827 *WIREs Water*, 6, <https://doi.org/10.1002/wat2.1353>, 2019.  
828  
829 Tarasova, L., Basso, S., Wendi, D., Viglione, A., Kumar, R., and Merz, R.: A Process-Based  
830 Framework to Characterize and Classify Runoff Events: The Event Typology of Germany,  
831 *Water Resour. Res.*, 56, <https://doi.org/10.1029/2019WR026951>, 2020.  
832  
833 Tarasova, L., Lun, D., Merz, R., Blöschl, G., Basso, S., Bertola, M., Miniussi, A., Radovec, O.,  
834 Samaniego, L., Thober, S., Kumar, R.: Shifts in flood generation processes exacerbate  
835 regional flood anomalies in Europe. *Commun Earth Environ* 4, 49,  
836 <https://doi.org/10.1038/s43247-023-00714-8>, 2023.

837  
838 Tramblay, Y., Bouvier, C., Martin, C., Didon-Lescot, J.-F., Todorovik, D., and Domergue, J.-  
839 M.: Assessment of initial soil moisture conditions for event-based rainfall–runoff modelling,  
840 *Journal of Hydrology*, 387, 176–187, <https://doi.org/10.1016/j.jhydrol.2010.04.006>, 2010.  
841  
842 Tramblay, Y., Neppel, L., Carreau, J., and Najib, K.: Non-stationary frequency analysis of  
843 heavy rainfall events in southern France, *Hydrological Sciences Journal*, 58, 280–294,  
844 <https://doi.org/10.1080/02626667.2012.754988>, 2013.  
845  
846 Tramblay, Y., Mimeau, L., Neppel, L., Vinet, F., and Sauquet, E.: Detection and attribution of  
847 flood trends in Mediterranean basins, *Hydrol. Earth Syst. Sci.*, 23, 4419–4431,  
848 <https://doi.org/10.5194/hess-23-4419-2019>, 2019.  
849  
850 Tramblay, Y., Koutroulis, A., Samaniego, L., Vicente-Serrano, S. M., Volaire, F., Boone, A.,  
851 Le Page, M., Llasat, M. C., Albergel, C., Burak, S., Cailleret, M., Kalin, K. C., Davi, H.,  
852 Dupuy, J.-L., Greve, P., Grillakis, M., Hanich, L., Jarlan, L., Martin-StPaul, N., Martinez-  
853 Vilalta, J., Mouillot, F., Pulido-Velazquez, D., Quintana-Seguí, P., Renard, D., Turco, M.,  
854 Türkeş, M., Trigo, R., Vidal, J.-P., Vilagrosa, A., Zribi, M., and Polcher, J.: Challenges for  
855 drought assessment in the Mediterranean region under future climate scenarios, *Earth-*  
856 *Science Reviews*, 210, 103348, <https://doi.org/10.1016/j.earscirev.2020.103348>, 2020.  
857  
858 Tramblay, Y., Villarini, G., Saidi, M. E., Massari, C., and Stein, L.: Classification of flood-  
859 generating processes in Africa, *Sci Rep*, 12, 18920, [https://doi.org/10.1038/s41598-022-](https://doi.org/10.1038/s41598-022-23725-5)  
860 [23725-5](https://doi.org/10.1038/s41598-022-23725-5), 2022.  
861  
862 Wasko, C. and Nathan, R.: Influence of changes in rainfall and soil moisture on trends in  
863 flooding, *Journal of Hydrology*, 575, 432–441, <https://doi.org/10.1016/j.jhydrol.2019.05.054>,  
864 2019.  
865  
866 Wasko, C., Nathan, R., Stein, L., and O’Shea, D.: Evidence of shorter more extreme rainfalls  
867 and increased flood variability under climate change, *Journal of Hydrology*, 603, 126994,  
868 <https://doi.org/10.1016/j.jhydrol.2021.126994>, 2021.  
869  
870 Watson, G. S. and Williams, E. J.: On the Construction of Significance Tests on the Circle  
871 and the Sphere, *Biometrika*, 43, 344, <https://doi.org/10.2307/2332913>, 1956.  
872  
873 Whitfield, P. H.: Floods in future climates: a review: Changing floods in future climates, *J.*  
874 *Flood Risk Manage*, 5, 336–365, <https://doi.org/10.1111/j.1753-318X.2012.01150.x>, 2012.  
875  
876 Zhang, S., Zhou, L., Zhang, L., Yang, Y., Wei, Z., Zhou, S., Yang, D., Yang, X., Wu, X.,  
877 Zhang, Y., Li, X., and Dai, Y.: Reconciling disagreement on global river flood changes in a  
878 warming climate, *Nat. Clim. Chang.*, 12, 1160–1167, [https://doi.org/10.1038/s41558-022-](https://doi.org/10.1038/s41558-022-01539-7)  
879 [01539-7](https://doi.org/10.1038/s41558-022-01539-7), 2022.  
880  
881  
882  
883

884  
885  
886  
887  
888  
889  
890  
891

# INDEX OF FIGURES




Figure 1: Map of the selected catchments




Figure 2: Relative changes in different flood event characteristics between 1959-1990 and 1991-2021




Figure 3: Relationship between the flood event runoff coefficients and antecedent soil moisture for the two time periods considered: 1959-1990 and 1991-2021. For each box, the central line indicates the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The whiskers extend to the most extreme data points. The numbers at the top of the figure indicate the number of events in each category.




Figure 4: Mean date of flood occurrence




Figure 5: Changes in mean flood date (left) and changes in the concentration index (right) between 1959-1990 and 1991-2021






Figure 6: Frequency of the different weather types associated with flood events, the names of the different weather types are from Garavaglia et al. (2010)




Figure 7: Change in monthly frequency of weather types 1, 2, 4, 6 and 7 between 1959-1990 and 1991-2021




Figure 8: Flood event classification into four categories: excess rain, long rain, short rain and others




Figure 9: Mean monthly frequency of occurrence for the three flood drivers: excess rain, short rain, and long rain




Figure 10: Relative changes in the frequency of excess rain, short rain and long rain between 1959-1990 and 1991-2021




Figure 11: Regional frequency of excess rain, short rain, and long rain floods between 1959 and 2021. The gray lines denote a least-square linear fit to represent the long-term tendency




Figure 12: Distribution of regionally sampled floods corresponding to excess rain, short rain and long rain types of floods for the two time periods 1959-1990 and 1991-2021