

1 **Spatiotemporal responses of runoff to climate change on the southern Tibetan Plateau**

2 He Sun<sup>1\*</sup>, Tandong Yao<sup>1,2</sup>, Fengge Su<sup>1,2\*</sup>, Wei Yang<sup>1,2</sup>, Deliang Chen<sup>3</sup>

3

4 <sup>1</sup>State Key Laboratory of Tibetan Plateau Earth System, Environment and Resources  
5 (TPESER), Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing  
6 100101, China

7 <sup>2</sup>University of Chinese Academy of Sciences, Beijing 100101, China

8 <sup>3</sup>Regional Climate Group, Department of Earth Sciences, University of Gothenburg,  
9 Gothenburg 405 30, Sweden

10

11

12

13

14

15

16

17

18 \*Corresponding author: He Sun, Fengge Su

19 State Key Laboratory of Tibetan Plateau Earth System, Resources and Environment, Institute  
20 of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

21 Email: [sunhe@itpcas.ac.cn](mailto:sunhe@itpcas.ac.cn), [fgsu@itpcas.ac.cn](mailto:fgsu@itpcas.ac.cn)

## 22 **Abstract**

23 A comprehensive understanding of spatiotemporal runoff changes at a sub-basin scale of the  
24 Yarlung Zangbo (YZ) basin on the southern Tibetan Plateau (TP), amidst varying climatic and  
25 cryospheric conditions, is imperative for effective water resources management. However,  
26 spatiotemporal differences of runoff composition, change and the attribution within the YZ  
27 basin have not been extensively explored, primarily due to the lack of hydrometeorological  
28 observations, especially in the downstream region. In this study, we investigated historical and  
29 future evolution of annual and seasonal total water availability, as well as glacier runoff and  
30 snowmelt contributions across six sub-basins of the YZ with a particular focus on the  
31 comparison between the upstream Nuxia (NX) basin and the downstream Nuxia-Pasighat  
32 (NX-BXK) basin, based on a newly generated precipitation dataset and a well-validated  
33 model with streamflow, glacier mass, and snow cover observations. Our findings revealed  
34 large spatiotemporal differences in changes exist within the YZ basin for 1971–2020. Firstly,  
35 runoff generation was dominated by rainfall runoff throughout the YZ basin, with glacier  
36 runoff playing more important role in the annual total runoff (19%) in the NX-BXK sub-basin  
37 compared to other sub-basins. Notably, glacier runoff contributed 52% of the total runoff at  
38 the Pasighat outlet of the YZ basin. Secondly, annual runoff exhibited an increasing trend in  
39 the NX basin but a decreasing trend in the NX-BXK, primarily attributed to rainfall runoff  
40 changes influenced by atmospheric moisture. Glacier runoff enhanced water supply, by  
41 offsetting the decreasing contribution from rainfall. Total runoff will consistently increase

42 (27–100 mm/10yr) across the sub-basins through the 21st century, resulting from increased  
43 rainfall runoff and a minor effect of increased snowmelt and glacier runoff.

44

45 **Keywords**

46 Runoff Composition; Runoff Changes; VIC-Glacier Hydrological Model; Yarlung Zangbo;  
47 Tibetan Plateau

48

49 **Highlights**

50 ● Runoff generation is dominated by rainfall runoff (59%–72%) in the YZ, and the largest  
51 glacier runoff contribution is in the downstream sub-basin (16%–19%).

52 ● Annual runoff trends indicate an increase in the NX but a decrease in the NX-BXK for  
53 1971–2020, due to contrasting precipitation changes.

54 ● Total runoff across the sub-basins will consistently increase (27–100 mm/10yr) through  
55 the 21st century, mostly resulting from increased rainfall runoff.

56

57

58

59

60

61

## 62 **1 Introduction**

63 Climatic and cryospheric changes have profoundly affected hydrological processes in high-  
64 mountain regions. The Tibetan Plateau (TP), known as the Asian Water Tower, supplying  
65 freshwater to nearly 2 billion people. Marked atmospheric warming since the 1980s has  
66 changed the balance between liquid and solid states of water, leading to shifts in river runoff,  
67 glacier, and snow melt dynamics (Yao et al. 2022). These drastic changes in the upper  
68 mountains of the TP pose a threat to the sustainability of the downstream water supply.

69

70 The Yarlung Zangbo (YZ, Figure 1) river basin, located in the southern TP is the largest river  
71 basin of the TP and a vital freshwater source for the Tibet Autonomous Region (TAR). It  
72 constitutes the main agricultural region in the TAR (Yang et al., 1989; Zhong et al., 2014).  
73 Like elsewhere on TP, a rapid ongoing temperature rise (0.3–0.4°C per decade) since the mid-  
74 1960s potentially influences runoff processes and water resources availability in the YZ basin  
75 (Yao et al., 2012; Li et al., 2018). The YZ basin, spanning approximately 250,000 km<sup>2</sup>,  
76 exhibits diverse climatic systems, including the Indian summer monsoon and the westerly  
77 system, and varying glacier and snow conditions (Zhang et al., 2013). These factors contribute  
78 to spatiotemporal differences in runoff changes within the YZ basin. Therefore, gaining a  
79 comprehensive understanding of runoff regimes and flow changes at the sub-basin scale is  
80 crucial for informed decision-making in water resources management and social development.

81

82 While numerous studies have investigated runoff regimes and changes using hydrological  
83 models in the YZ basin, most have focused solely on the region upstream of the Nuxia (NX)  
84 hydrological station (Figure 1, Table 1) (Chen et al., 2017; Cuo et al., 2019; Su et al., 2016;  
85 Zhang et al., 2013; Zhao et al., 2019; Cui et al., 2023; Gu et al., 2023). This focus is due to the  
86 nearest national hydrological station's proximity to the mainstream outlet, providing long-  
87 term daily records (> 50years). Conversely, the glacierized downstream region (about 65% of  
88 the total glacier area in the YZ), particularly between NX and Pasighat outlet (NX-BXK,  
89 Figure 1) has received less attention. This lack of focus is attributed to limited  
90 hydrometeorological and glacier observations in this sub-basin. Remarkably, this region  
91 exhibits the largest glacier retreat in the TP, with a length reduction rate of  $48.2\text{m yr}^{-1}$  and an  
92 area decrease of  $0.57\% \text{ yr}^{-1}$  during the 1970s–2000s (Yang et al., 2013; Yao et al., 2012).  
93 These changes have the potential to significantly alter the runoff regime, influencing the  
94 quantity, timing, and variability of flows across space and time. However, the characteristics  
95 and changes in runoff, along with the effect of glacier melt on water supply, remain unclear in  
96 the NX-BXK sub-basin.

97

98 The NX basin, with an area of approximately  $201,548 \text{ km}^2$ , presents divergent glacier and  
99 snow conditions (Table 2). For example, the region upstream of the Lhatse hydrological  
100 station (Figure 1), the source region of the YZ river, is influenced by both monsoon and  
101 westerlies, experiencing higher precipitation in spring and winter compared to other NX sub-

102 basins (Figure S1 in Supporting Information). The Lhasa (LS) and Rikaze (RKZ) sub-basins,  
103 vital crop centers for the central Tibet Autonomous Region, play a crucial role in irrigation  
104 water resources. The LS sub-basin, with about 23% snow cover contrasts with the RKZ sub-  
105 basin, which has little glacier and snow coverage (Table 2). This difference suggests that the  
106 water supply in the RKZ sub-basin is more sensitive to climate change. Moreover, runoff in  
107 the region between Yangcun and NX hydrological stations (YC-NX, Figure 1) contributes 51%  
108 to the total runoff at the NX hydrological station (Sun and Su, 2020), making runoff regimes  
109 and changes in this sub-basin influential for the entire NX basin. Therefore, a comprehensive  
110 investigation into runoff regimes and changes in different sub-basins is essential for a nuanced  
111 understanding of the mechanisms underlying runoff changes in response to climate change.

112

113 While many hydrological studies focus on the region upstream of the NX hydrological station  
114 (Zhang et al., 2013; Lutz et al., 2014; Zhao et al., 2019; Sun and Su, 2020; Khanal et al., 2021;  
115 Nan et al., 2021; Wang et al., 2021), considerable differences in runoff regimes and change  
116 studies exist in the NX basin (Table 1). These differences may arise from variations in forcing  
117 inputs for hydrological model simulations. Accurate precipitation inputs play an important  
118 role in reliable hydrological model simulations. However, high mountain precipitation in the  
119 YZ basin is still inadequately represented in gauge-based, satellite-based, and reanalysis-  
120 based estimates, or outputs of regional climate models (Wang and Zeng, 2012; Liu et al., 2020;  
121 Sun et al., 2021). The mean annual precipitation ranges from 360–1236 mm in the YZ basin

122 (Qi et al. 2018; Sun; Su 2020; Tong et al. 2014), resulting in significant uncertainties in  
123 hydrological simulations. This inconsistency in gridded datasets is often underestimated in  
124 hydro-climate studies, especially in glacierized basins with limited data coverage. Even when  
125 realistic runoff simulations are achieved at the catchment outlet, they cannot guarantee  
126 reasonable results (Zhao et al., 2019) due to the compensation between precipitation-induced  
127 runoff and snow/glacier melting. For example, Lutz et al. (2014) simulated glacier runoff in  
128 the NX basin with the Spatial Processes in HYdrology (SPHY) model driven by the Asian  
129 Precipitation-Highly-Resolved Observational Data Integration Towards Evaluation  
130 (APHRODITE) precipitation estimates, suggesting that glacier runoff contributed about 16%  
131 to total runoff. In contrast, Khanal et al. (2021) proposed that glacier runoff contributed about  
132 1.8% to total runoff with the same model driven by the newly released fifth-generation  
133 reanalysis (ERA5) precipitation of the European Centre for Medium-Range Weather  
134 Forecasts (Table 1), primarily due to the overestimation of the ERA5 precipitation estimate.  
135 Sun and Su (2020) indicated that the contribution of glacier runoff would increase by 7–10%  
136 with a unit decrease in mean annual precipitation. Therefore, an accurate precipitation  
137 estimate is crucial as a model input to simulate runoff regimes, and further quantify the effect  
138 of glaciers and snowmelt on runoff in the NX basin.

139

140 These uncertainties in the hydrological simulation will be introduced and enlarged on the  
141 uncertainty in future projections (Lutz et al. 2016). Existing studies of hydrological responses

142 to future climate changes have been a debate in the NX sub-basin of the YZ basin. For  
143 example, Lutz et al. (2014) forced the Spatial Processes in Hydrology (SPHY) model using  
144 outputs from 4 global climate models (GCMs) and showed that runoff would be increased by  
145 3–13% relative to the reference period 1998–2007 until at least 2050s due to the increasing  
146 precipitation in the YZ basin. Zhao et al. (2019) projected the future runoff changes with 5  
147 GCM outputs using an extended Variable Infiltration Capacity (VIC) macroscale hydrological  
148 model (VIC-CAS), suggesting that the total runoff will increase by 16–31% by the end of this  
149 century relative to the reference period 1970–2010 because of increased rainfall-induced  
150 runoff in the YZ basin. Cui et al. (2023) also suggested that total runoff in the YZ basin will  
151 increase of  $7.3\pm 11\%$  by 2070s relative to the 1985–2014 resulted from rainfall runoff.  
152 Meanwhile, Su et al. (2016) projected a runoff increase of 6.7–14.4% in the 2041–2070  
153 relative to the reference period 1971–2000 in the YZ forced by the VIC-Glacier model with  
154 ensemble outputs of 20 GCMs, and attributed the runoff increases to the rising glacier melt  
155 runoff. In addition, future flow evolution and the effect of different runoff compositions on  
156 total runoff in the NX-BXK are also unclear.

157

158 To address these issues, this study divided the YZ into six sub-basins and collected  
159 streamflow observations at three hydrological stations (Yigong, Bomi, and Motuo) and glacier  
160 mass balance observations (Parlung No.94) in the NX-BXK sub-basin, filling in the gap of  
161 scarce data coverage. Additionally, streamflow observations at five national hydrological



162 stations and glacier mass balance observations at a site (Gurenhekou) in the five sub-basins of  
163 the NX (Figure 1), together with hydrological stations in the NX-BXK, constitute a unique  
164 observation basis. This basis allows us to validate the glacier-hydrology model and reveal  
165 runoff regimes, and changes at the sub-basin scale. Precipitation observations at 280 gauges,  
166 were collected, and a high spatiotemporal resolution (10 km; daily) precipitation dataset was  
167 generated using a machine learning algorithm based on these gauges (Sun et al., 2022).

168

169 Leveraging this basin-wide observation dataset, this study comprehensively investigates  
170 runoff compositions, changes, and attributions across six sub-basins in the YZ for 1971–2020,  
171 with a particular focus on the comparison between the NX and NX-BXK. This investigation  
172 employs the process-based and well-established Variable Infiltration Capacity (VIC)-Glacier  
173 hydrological model. Furthermore, the study assesses the future evolution of annual and  
174 seasonal total water availability, glacier runoff, and snowmelt, using an ensemble of multiple  
175 global climate models (GCMs) from the latest release of the Coupled Model Intercomparison  
176 Project Phase 6 (CMIP6). The objectives are to: (1) use the model framework to identify  
177 spatiotemporal characteristics in runoff compositions and changes at the sub-basin scale under  
178 heterogeneous climate and glacier/snow conditions. (2) quantify the contributions of three  
179 major runoff compositions (glacier, rainfall, and snowmelt runoff) to total runoff among  
180 different sub-basins, and investigate their responses to climate changes. (3) assess future  
181 water availability under 21st-century climate-cryosphere change, assisting policy-makers and

182 water managers in adopting strategies. These findings are anticipated to provide a basic  
183 framework for studying cryospheric basin hydrological cycles in the TP and provide  
184 adaptation strategies for rational water resource management, and social, and economic  
185 development grounded in a robust scientific understanding.

186

## 187 **2 Study Area**

188 In this study, the YZ basin is divided into six sub-basins based on flow direction and locations  
189 of hydrological stations (Figure 1; Table 2). There are five sub-basins located upstream of the  
190 Nuxia (NX) hydrological station, collectively termed the NX basin, with an additional sub-  
191 basin lying between Nuxia and Pasighat (NX-BXK) hydrological stations. The NX basin  
192 comprises the upstream sub-basins of Lhatse (LZ), Shigatse (RKZ), and Lhasa (LS)  
193 hydrological stations, along with the sub-basins between Lhatse and Yangcun (LZ-YC)  
194 hydrological station and between Yangcun and Nuxia (YC-NX) hydrological station.

195

196 The climate in the YZ is characterized by a wet and warm summer and a cool, dry winter,  
197 with precipitation mostly dominated by the summer monsoon, contributing 70–90% of annual  
198 totals during June–September (Figure S1). Furthermore, mean annual precipitation increases  
199 downstream in the YZ basin, ranging from 283 mm upstream to 1465 mm downstream,  
200 averaging about 774 mm for the entire YZ basin (Table 2). All sub-basins exhibit similar  
201 seasonal temperature patterns, with peaks mainly occurring in July–August (Figure S1).

202 Glacier coverage varies from 0.9% (LZ-YC) to 10.2% (NX-BXK), averaging 3.3% for the  
203 entire YZ basin. The YC-NX (2.8%) and NX-BXK (10.2%) sub-basins host the most  
204 extensive glacier coverage (Table 1). The mean annual snow cover fraction (SCF) ranges from  
205 7% (RKZ) to 32% (NX-BXK), with an average of 19% across the YZ basin.

206

## 207 **3 Data and Method**

### 208 **3.1 Data**

209 Daily precipitation, maximum and minimum temperature, and wind speed estimates with a  
210 spatial resolution of 10×10 km were used as the VIC-Glacier model forcing inputs in this  
211 work. Historical meteorological data during 1971–2100 was adopted from Sun et al. (2022).  
212 The daily precipitation data with a spatial resolution of 10×10 km for 1961–2020 was  
213 reconstructed by correcting gridded estimates from the ERA5 precipitation of the European  
214 Centre for Medium-Range Weather Forecasts (ECMWF) based on 580 rain gauges in the  
215 monsoon-dominated TP region (290 rain gauges in the YZ basin, Figure 1) and the Random  
216 Forest-based (RF) machine learning algorithm (Sun et al., 2022). Inputs of the RF algorithm  
217 selected in this study include: 1) geographical features (e.g., longitude, latitude, elevation,  
218 slope gradient and aspect), which influence precipitation distribution, and 2) climatic features  
219 derived from the ERA5 (e.g., convective available potential energy, lifting condensation level,  
220 and total column water vapor), which represent the potential for the generation and  
221 development of precipitation. The corrected precipitation data set was evaluated at a point

222 scale by comparing it with gauge observations, and has been inversely evaluated by the  
223 hydrological model, which demonstrates its suitability for hydrological simulation (Sun et al.,  
224 2022).It was downloaded from the National Tibetan Plateau/Third Pole Environment Data  
225 Center (TPDC, <https://doi.org/10.11888/Atmos.tpdc.272885>).

226

227 Using the newly generated daily meteorological data for 1961–2020, Sun et al. (2024) applied  
228 the Bias Corrected Spatial Disaggregation (BCSD) statistical downscaling approach (Wood  
229 2002; Wood et al. 2004) to downscale and bias-correct daily transient meteorological data,  
230 including precipitation, maximum and minimum temperature, and wind speed, with a spatial  
231 resolution of 10×10 km through the 21st century from 10 GCMs from the CMIP6 under the  
232 Shared Socioeconomic Pathways (SSPs)2-4.5 and SSP5-8.5 scenario. Daily transient climate  
233 estimates, at a spatial resolution of 10×10 km for 1971–2100 under 20 scenarios (10 GCMs ×  
234 2 SSPs) from Sun et al. (2024) were directly employed to drive the VIC-Glacier model to  
235 continuous runoff production in the YZ basin.

236

237 Observed streamflow, observed glacier mass balance, satellite-based glacier area and snow  
238 cover fraction estimates were applied to calibrate and validate the model in this study.  
239 Monthly streamflow since 1971 were collected at eight hydrological stations (Lhatse, Shigatse,  
240 Lhasa, Yangcun, Nuxia, Yigong, Bomi, and Motuo from the Ministry of Water Resources,  
241 China (Figure 1, Table S1).

242

243 Two shapefiles of glacier inventory, the first Glacier Inventory of China (CGI) from the  
244 “Environment & Ecological Science Data Center for west China”  
245 (<http://westdc.westgis.ac.cn/glacier>) and Randolph Glacier Inventory (RGI) 6.0  
246 ([https://www.glims.org/RGI/rgi60\\_dl.html](https://www.glims.org/RGI/rgi60_dl.html)), were used to describe the glacier information of  
247 the YZ in the VIC-Glacier model. Observed annual glacier mass balance data from  
248 Gurenhekou and Parlung No.94 glacier sites since 2005 were used to validate the performance  
249 of glacier model (<http://www.tpdc.ac.cn>, Figure 1, Table S1). The Moderate Resolution  
250 Imaging Spectroradiometer (MODIS) 10CM reporting the maximum percentage of snow  
251 cover during an 8-day period in 0.05° resolution grid (<https://nsidc.org/> data) during 2006–  
252 2018 was used to calculate the snow cover fraction (SCF) and to compare with VIC-Glacier  
253 model simulations in the YZ basin.

254

255 The model required land surface characteristics, including soil texture and vegetation types,  
256 were adopted from Sun and Su (2020). These data were used as initial model inputs, and  
257 remained unchanged in simulation period.

258

### 259 **3.2 VIC-Glacier Hydrological Model**

260 The present study employed the physically-based and distributed VIC hydrological model  
261 (Liang et al., 1994; Liang et al., 1996) linked with a simple degree-day glacier melt algorithm

262 (Hock, 2003), referred to as VIC-Glacier. This modeling framework facilitates the  
263 comprehensive simulation of the physical exchange of water and energy within a grid mesh  
264 encompassing soil, vegetation, and the atmosphere. The VIC-Glacier effectively models  
265 surface water balance compositions, including evapotranspiration, surface runoff, baseflow  
266 (subsurface drainage into the local stream channel network, as opposed to groundwater  
267 recharge), and total soil moisture, including liquid and ice content in each soil layer. The  
268 model integrates a two-layer energy-balance snow model (Cherkauer and Lettenmaier, 1999)  
269 and a frozen soil/permafrost algorithm (Cherkauer and Lettenmaier, 1999; 2003). These  
270 components account for ground snowpack, snow within the vegetation canopy, snow atop  
271 lake ice, and sublimation of snow. In each time step, the model calculates the rain or snow  
272 fraction contributing to the snowpack. Subsequently, all energy fluxes are computed,  
273 triggering melt if the energy balance is positive. The VIC-Glacier model has demonstrated its  
274 effectiveness in hydrological simulations for various high-mountainous TP basins (Meng et al.,  
275 2019; Su et al., 2016; Sun and Su, 2020; Tong et al., 2016; Zhang et al., 2013; Zhao et al.,  
276 2019).

277

278 Here, the modeling framework at a 10 km  $\times$  10 km spatial resolution and a three-hourly time  
279 step was adopted from Sun and Su (2020). To categorize the total runoff sources in this study,  
280 we partitioned it into three components: rainfall runoff, snowmelt runoff, and glacier runoff.  
281 Glacier runoff was defined as all water generated in the glacierized area, including rainfall,

282 snow melt, and ice melt in the glacierized area.

283

$$284 \quad \text{Total runoff (TR)} = \text{Rainfall runoff (RR)} + \text{Snowmelt runoff (SR)} + \\ 285 \quad \text{Glacier runoff (GR)} \quad (1)$$

$$286 \quad \text{Rainfall runoff contribution} = \frac{RR}{TR} \times 100\% \quad (2)$$

$$287 \quad \text{Snowmelt runoff contribution} = \frac{SR}{TR} \times 100\% \quad (3)$$

288

289

290 Recognizing the influence of glacier melt at different elevations, each glacierized grid cell

291 underwent division into various elevation bands with an interval of 100 m (Kan et al., 2018).

292 The simulated total runoff of each grid is from both the glacierized and non-glacierized areas,

293 that is,

$$294 \quad R_i = f \times R_{glac} + (1 - f) \times R_{vic} \quad (4)$$

295 Where,  $R_i$  is the total runoff (mm) in grid  $I$ ,  $f$  is the percentage of glacier area, and glacier area

296 is updated every year,  $R_{glac}$  is the runoff (mm) from the glacier area calculated by the glacier

297 model, and  $R_{vic}$  is the sum of surface runoff and baseflow runoff (mm) for non-glacierized

298 areas calculated by the VIC model, including both rainfall and seasonal snowmelt runoff.

299  $R_{glac}$  can be calculated as:

$$300 \quad M_i = \begin{cases} DDF \times (T - T_{base}); & T > T_{base} \\ 0; & T \leq T_{base} \end{cases} \quad (5)$$

$$301 \quad R_{glac} = M_1 + M_2 + \dots + M_i; i = 1, 2, 3, \dots, n \quad (6)$$

302 where,  $M_i$  is the meltwater (mm) from elevation band  $j$  and  $n$  is the total number of elevation  
303 bands in grid  $i$ ;  $DDF$  is the degree-day factors of glacier or snow melt ( $\text{mm } ^\circ\text{C}^{-1} \text{ day}^{-1}$ );  $T$  ( $^\circ\text{C}$ )  
304 is the daily average air temperature above the glacier surface;  $T_{base}$  ( $^\circ\text{C}$ ) is the temperature  
305 threshold for glacier and snow melt ( $0^\circ\text{C}$ ). In a precipitation event, rainfall occurs when the  
306 air temperature is above  $0^\circ\text{C}$ ; otherwise, it snows. In the presence of a snowpack on the  
307 glacier, the snow melts first before glacier melting starts, following the same degree-day  
308 approach but different degree-day factors.

309

310 The calculated glacier area and volume were updated every year in the model by the volume-  
311 area scaling approach (Bahr et al., 1997). An exponential form (equation 7), derived from  
312 glacier observations in western China (Liu et al., 2003), converts glacier area to volume for a  
313 basin:

$$314 \quad V = 0.04S^{1.35} \quad (7)$$

315 Where,  $V$  is glacier volume ( $\text{km}^3$ ) and  $S$  is glacier area ( $\text{km}^2$ ). Initial glacier volume was  
316 determined using glacier area data from the first Glacier Inventory of China (CGI V1.0,  
317 <http://westdc.westgis.ac.cn/glacier>) dataset, which presented glacier area for period  
318 1970s–1990s. The Randolph Glacier Inventory (RGI V6.0) dataset presented glacier area for  
319 period 2000s–2010s. Therefore, glacier area was simulated by glacier model since 1971, and  
320 then it was updated every year with the snowfall accumulation and simulated ice melt from all  
321 the glacier cells in the glacier model based on volume-area scaling approach. The simulated  
322 mean annual glacier area during 2000–2010 was compared with the Randolph Glacier  
323 Inventory (RGI V6.0) dataset in the YZ basin and its six sub-basins.



324

### 325 **3.3 Model Calibration and Validation**

326 The VIC-Glacier model requires the calibration of two sets of parameters: (1) the degree day  
327 factor (DDF), related to glacier runoff simulation; and (2) VIC model parameters related to  
328 runoff simulation in non-glacierized regions. The latter includes parameters (Table 3) such as  
329 the depth of the first and second soil layers (D1 and D2), the infiltration shape parameter  
330 ( $B_{inf}$ ), and three base flow parameters, including the maximum velocity of baseflow  
331 ( $D_{smax}$ ), a fraction of  $D_{smax}$  where non-linear baseflow begins ( $D_s$ ), and a fraction of  
332 maximum soil moisture where non-linear baseflow occurs ( $W_s$ ).

333

334 To adjust the internal stores of energy and water from the initial state to equilibrium, the VIC-  
335 Glacier model underwent a spin-up from 1961–1970, with subsequent simulation for the years  
336 1971–2020. In addition, 1971–2000 was selected as the calibration period and 2001–2015 the  
337 validation period based on the observed monthly streamflow for 1971–2015. Calibration and  
338 validation of the VIC-Glacier hydrological model followed a systematic two-step approach,  
339 employing observed streamflow, glacier mass balance, and satellite-based estimates of glacier  
340 area and snow cover fraction (Table S1). Model performance was assessed using metrics such  
341 as Nash-Sutcliffe efficiency (NSE), relative bias (RB, %), and correlation coefficient (CC).  
342 The optimization process utilized a trial-and-error method to minimize bias against predefined  
343 criteria. Linear regression was employed to calculate annual and seasonal trends of

344 precipitation, temperature, and runoff.

345

346 The model calibration and validation were conducted using a two-step approach to overcome  
347 equifinality problems. First, initial values of DDF parameters in the glacier model related to  
348 glacier and snowmelt were adopted from Sun and Su (2020). The glacier model was  
349 calibrated to match the glacier area observations from RGI V6 for 2000s–2010s in the YZ and  
350 six sub-basins, and validated by observed mass balance data from the Gurenhekou site in the  
351 NX basin and the Parlung No.94 glacier site in the NX-BXK sub-basin (Figure S2). Given the  
352 good performance in simulating glacier area (with RB of mostly < 7%, Figure 2c) and good  
353 consistency (CCs of 0.65–0.96) in annual variations between observed glacier mass balance  
354 and simulation, final DDF values ( $6.5\text{--}11.0 \text{ mm}^\circ\text{C}^{-1} \text{ day}^{-1}$ ) were determined across six sub-  
355 basins (Table 3).

356

357 Second, the VIC-related parameters were validated against observed streamflow and satellite-  
358 based snow cover fraction (SCF) data. The infiltration parameter ( $B_{\text{inf}}$ ) and the second soil  
359 layer depth (D2) have been identified as the most sensitive parameters (Zhang et al., 2013).  
360 The  $B_{\text{inf}}$  which defines the shape of the variable infiltration capacity curve has a common  
361 range of 0–0.4, while the D2 mainly determines the moisture storage capacity of the VIC  
362 model, with a range of 0.5–1.0 (Liang et al., 1996; Shi et al., 2008). The simulated monthly  
363 streamflow captured well the magnitudes and patterns of observation at eight hydrological

364 stations, with NSEs of 0.71 to 0.86 and RBs of within  $\pm 8\%$  for the calibration and validation  
365 period across the sub-basins (Table 3, Figure S3). To further validate the model, monthly  
366 satellite-based SCF data for the years 2001–2019 in the YZ basin were compared with the  
367 model simulations (Figure S4). The simulated SCF closely mirrors the monthly variations  
368 observed in the satellite-based data, exhibiting a CC of 0.60–0.82 ( $p < 0.05$ ) and RB within  
369  $\pm 12\%$  across sub-basins. This alignment suggests the VIC-Glacier model's satisfactory  
370 performance in simulating snow cover dynamics in the YZ basin.

371

## 372 **4 Result**

### 373 **4.1 Hydrological Response to Historical Climate Changes**

#### 374 **4.1.1 Runoff Composition**

375 The credibility of our model allows for a reasonable interpretation of the current runoff  
376 composition and change, and their responses to climate change. Examining simulated  
377 streamflow across sub-basins reveals significant differences in each sub-basin contribution to  
378 the total runoff at the Pasighat outlet of the YZ basin (Figure 1b). The NX-BXK emerges as  
379 the most critical runoff-generating area, contributing approximately 52% to the total runoff in  
380 the YZ basin, followed by YC-NX (25%), LS (10%), and other sub-basins with contributions  
381 ranging from 3% to 6%.

382

383 According to the source of runoff generation, total runoff is partitioned into three  
384 compositions in this study: rainfall runoff, snowmelt runoff, and glacier runoff. In this study,

385 glacier runoff is defined as all water generated in the glacierized area, including rainfall, snow  
386 melt, and ice melt in the glacierized area. Rainfall and snowmelt runoff are originating from  
387 the non-glacierized area. Different runoff regimes of rainfall runoff, snowmelt, and glacier  
388 runoff influence their contributions to total runoff across the six sub-basins of the YZ with  
389 heterogeneous surface characteristics (Figure 2). Rainfall runoff dominates the mean annual  
390 total runoff in the YZ and all its sub-basins from 1971 to 2020, contributing 59%–78% to  
391 annual total runoff, with an average of 62% in the entire YZ basin. Snowmelt contributes 22%  
392 to annual total runoff in the YZ basin, varying from 14% to 36% across six sub-basins, with  
393 the LS sub-basin having the largest contribution at 36%. Glacier runoff contributes 16% to the  
394 annual total runoff in the YZ basin, ranging from 5% to 19% across six sub-basins for 1971–  
395 2020. The highest contributions are in the NX-BXK (19%) and YC-NX (16%) sub-basins,  
396 which have the largest glacier coverage in the YZ basin (Table 2).

397

398 Figure 3 shows the spatial pattern of average annual rainfall runoff, snowmelt, and glacier  
399 runoff for 1971–2020, along with their percentages at different elevation bands in the YZ  
400 basin. The spatial pattern of average annual rainfall runoff (Figure 3a) is similar to that of  
401 total runoff (Figure 1b) and precipitation (Figure 3b), decreasing from east to west, with the  
402 NX-BXK sub-basin exhibiting the largest runoff. Similarly, the largest snowmelt and glacier  
403 runoff occur in the NX-BXK sub-basin, consistent with the spatial distribution of glacier and  
404 snow cover area, constituting about 65% of total glacier area and 34% of total snow coverage

405 in the YZ (Figure 3c–f). Approximately 84% of the YZ basin runoff originates from middle  
406 altitudes (3500–5500 m), with 62% from 4500–5500 m and 22% from above 3500–4500 m,  
407 primarily contributed by rainfall and seasonal snow (80%–83%, Figure 3g). About 8% of the  
408 basin runoff is generated from high altitudes (>5500 m), where 29% of the flow is from  
409 glacier runoff, and the remainder is from rainfall (50%) and snowmelt (21%). In low altitudes  
410 (<3500 m), 8% of the basin runoff is primarily from rainfall (82%) and snowmelt (11%), with  
411 only 7% attributed to glacier runoff.

412

413 The seasonal pattern of total runoff remains consistent across the six sub-basins within the YZ  
414 for 1971–2020, with more than 60% of the annual total runoff occurring in June–September  
415 and 10%–15% in November–February (Figure 2a–h). This seasonal pattern aligns with the  
416 rainfall runoff, which peaks in July–August, reflecting the peak in total runoff in the YZ and  
417 its sub-basins (Figure 2 a–h). Snowmelt predominantly takes place from April–October, with  
418 peak months varying across sub-basins. In LZ, RKZ, LS, and LZ-YC sub-basins, the peak is  
419 in July–September (Figure 2a–d), attributed to the melting of fresh snowfall in the warm  
420 season. Conversely, in the YC-NX (Figure 2e) and NX-BXK (Figure 2f) sub-basins, the peak  
421 is in May–June, possibly due to snowfall accumulation during October–March. Simulated  
422 glacier runoff occurs mainly from June to September for all basins, peaking in July–August,  
423 coinciding with the co-occurrences of peak precipitation and temperature.

424

## 425 **4.1.2 Runoff Changes and the Response to Climate Changes**

### 426 **(a) Annual Scales**

427 Figure 4 illustrates annual trends in precipitation, temperature, total runoff, and three runoff  
428 compositions (rainfall, glacier, and snowmelt runoff) across the six sub-basins for 1971–2020,  
429 respectively. Annual variations for precipitation, temperature, and simulated runoff in each  
430 sub-basin are presented in Figure S5–S10. All sub-basins exhibit significant warming trends  
431 ( $0.3\text{--}0.5\text{ }^{\circ}\text{C}/10\text{yr}$ ,  $p<0.05$ ), with precipitation tending to increase ( $6\text{--}15\text{ mm}/10\text{yr}$ ) in the LZ,  
432 LZ-YC, LS, RKZ, and YC-NX sub-basins (Figure 4a–e) upstream of the NX hydrological  
433 station (NX basin). Conversely, the NX-BXK sub-basin experiences a significant decrease in  
434 precipitation ( $-35\text{ mm}/10\text{yr}$ ,  $p<0.05$ , Figure 4f).

435

436 Simulated annual total runoff demonstrates increasing trends of  $8.1\text{--}18.8\text{ mm}/10\text{yr}$  for 1971–  
437 2020 across all sub-basins within the NX basin, except for the RKZ sub-basin with an  
438 insignificant change ( $-1.1\text{ mm}/10\text{yr}$ ), resulting in a significantly increasing trend of  $9.4$   
439  $\text{mm}/10\text{yr}$  ( $p<0.05$ ) over the entire NX basin (Table 4). Strong correlations between annual  
440 variation of total runoff, precipitation, and rainfall runoff exist in these sub-basins (CC of  
441  $0.90\text{--}0.99$ ,  $p<0.05$ ), while total runoff shows weak relationships with temperature and glacier  
442 runoff. This suggests the predominant role of rainfall runoff from nonglacierized areas, with  
443 minor impacts from glacier runoff on annual runoff, along with significant increases in  
444 precipitation and temperature (Figure 4a). In contrast, the NX-BXK sub-basin exhibits a

445 significantly decreasing trend of 9.4 mm/10yr ( $p < 0.05$ ) for 1971–2020 (Figure 4f), resulting  
446 from significant decreases in rainfall runoff (-22 mm/10yr) and seasonal snowmelt (-5.5  
447 mm/10yr) from non-glacierized areas. Glacier runoff, however, exhibits a significantly  
448 increasing trend (6.0 mm/10yr,  $p < 0.05$ , Table 3) in NX-BXK during the same period, partially  
449 compensating for the decreasing trend of total runoff in this sub-basin. The integrated result is  
450 a weakly increasing trend of 3.1 mm/10yr in total runoff for the entire YZ basin (Table 4),  
451 primarily attributed to increases in rainfall (3.0 mm/10yr) and glacier runoff (2.1 mm/10yr).  
452 Snowmelt tends to decrease (-1 to -6 mm/10yr) in the YZ and its sub-basins during 1971–  
453 2020, associated with a reduction in solid precipitation and an increase in liquid precipitation  
454 (Figure S12), along with significant temperature increases.

455

456 Cuo et al. (2019) investigated precipitation and streamflow mutations in the YZ basin using  
457 Mann-Kendall analysis, identifying a streamflow mutation in 1997 at the NX hydrological  
458 station. This abrupt change is consistent with our long-term runoff observations. This abrupt  
459 change is consistent with our long-term runoff observations. Total runoff trends are opposite  
460 before and after the year 1998 in the YZ, and its NX and NX-BXK sub-basins (Table 4).  
461 During 1971–1997, annual total runoff shows increasing trends (8.9–48.1 mm/10yr) in the  
462 basins during 1971–1997, mainly due to an increasing trend in rainfall and glacier runoff  
463 (Table 4). However, during 1998–2020, total runoff showed insignificant decreasing trends (-  
464 0.3 to -3.3 mm/10yr), attributed to a decreasing trend in rain runoff induced by the weakening

465 Indian monsoon from 1998–2000 (Table 4). It is noteworthy that the rate of decrease in  
466 precipitation is faster in NX-BXK (-16.0 mm/10yr) than in NX (-7.0 mm/10yr, Table 4).  
467 However, the decline in total runoff is less pronounced in NX-BXK (-0.3 mm/10yr) compared  
468 to NX (-3.3 mm/10yr, Table 3) during 1998–2000. This discrepancy arises from different  
469 influences of glacier runoff on total runoff between NX and NX-BXK sub-basins. A more  
470 rapid increase in glacier runoff in NX-BXK (16 mm/10yr) than in NX (0.7 mm/10yr, Table 4)  
471 partly compensates for the quicker decline in rainfall-runoff, resulting in a slower overall  
472 decrease in total runoff in NX-BXK.

473

474 Figure 5 illustrates the mean monthly vertical integral of atmospheric moisture budget in June,  
475 July, August, and September from ERA5 data across the YZ basin for 1971–2020. It  
476 demonstrates an increasing trend in the NX basin but a decreasing trend in the NX-BXK. This  
477 pattern corresponds with precipitation trends in the NX and NX-BXK sub-basins, influencing  
478 rainfall runoff in these areas. Additionally, teleconnection indices can modulate circulation  
479 patterns over a region, thereby affecting precipitation and its induced runoff. Among the 10  
480 teleconnection indices (Text S1), Pacific Decadal Oscillation (PDO) and El Niño/Southern  
481 Oscillation (ENSO) exhibit significantly negative consistency with precipitation, while  
482 Atlantic Multidecadal Oscillation (AMO) shows significantly positive consistency (CC=0.43,  
483  $p<0.05$ ) with precipitation for 1971–2020 in the NX basin (Figure S13a, d). The change in  
484 runoff induced by precipitation is mostly influenced by EASM with significantly positive



485 consistency (CC=0.38,  $p<0.05$ ) for 1971–2020 in the NX-BXK sub-basin (Figure S13a, e).  
486 The streamflow mutation in 1997, associated with the precipitation mutation, is also  
487 influenced by NAO and ENSO in the NX and EASM in the NX-BXK.

488

#### 489 **(b) Seasonal Scales**

490 Because of the similarity in annual runoff regimes and changes across five sub-basins within  
491 the NX basin, here, we particularly focus on comparing the NX and NX-BXK sub-basins at  
492 seasonal scales for 1971–2020. The high-altitude NX basin exhibits faster warming trends  
493 (0.2–0.5 °C/10yr) in each season compared to the low-altitude NX-BXK basin (0.16–  
494 0.23 °C/10yr, Figure 6d) for 1971–2020. Seasonal precipitation trends increase (2–9 mm/10yr)  
495 for 1971–2020 in the NX basin (Figure 6a), particularly in summer, influenced mainly by the  
496 AMO and PDO (Figure S14). Conversely, in the NX-BXK, precipitation decreased (-18 to -2  
497 mm/10yr) for 1971–2020, influenced by the EASM in summer and the AMO in autumn.  
498 Consequently, total runoff during 1971–2020 reflects similar trends to precipitation, affected  
499 by increased rainfall (1–6 mm/10yr) and glacier runoff (1 mm/10yr) in the NX (Figure 6a),  
500 and decreased rainfall (-10 to -3 mm/10yr) and snowmelt (-2 mm/10yr), along with increased  
501 glacier runoff (1–5 mm/10yr) in the NX-BXK (Figure 6b). Due to these different trends in the  
502 two sub-basins, total runoff shows an increasing trend in summer (5 mm/10yr) but decreasing  
503 trends (-1 mm/10yr) in other seasons for 1971–2020 in the YZ basin (Figure 6c), attributed to  
504 the dominance of rainfall runoff.

505

506 Relative to the period 1971–1997, divergent seasonal changes in total runoff are apparent in  
507 the YZ basin during 1998–2020 (Figure 7). In the NX basin, total runoff tends to increase by  
508 about 5%–22% in all seasons, with the largest increases during May–August (11%–22%),  
509 mainly due to increases in rain-induced and glacier runoff. The smallest increases occur  
510 during December–February (5%–6%), mostly due to increased rainfall runoff (3%–5%) in the  
511 NX (Figure 7a, d; Table S2). Snowmelt significantly increases during March–May (24%–50%)  
512 due to early snow melting (Figure 7d; Table S2), potentially benefiting agricultural water  
513 supplies. Conversely, total runoff in the NX-BXK sub-basin decreases by about 3%–20% in  
514 all seasons (Figure 7b; Table S2) due to declines in rainfall runoff (3%–23%) and seasonal  
515 snowmelt (4%–28%). This indicates a trend toward drier conditions, although increased  
516 glacier runoff (2%–12%) somewhat compensates for the loss of total runoff in July–August  
517 (Figure 7e). The integrated result of seasonal runoff changes in NX and NX-BXK shows total  
518 runoff in the YZ increases by 2%–4% in June–September, mostly due to increases in rain-  
519 induced (3%–7%) and glacier runoff (2%–6%), while it decreases in other months due to  
520 decreased rain-induced runoff (2%–8%) and seasonal snowmelt (3%–10%, Figure 7c, f, Table  
521 S2).

522

523 The distinct seasonal changes in rainfall, snowmelt, and glacier runoff largely play a crucial  
524 role in determining the seasonal shifts in their contributions to total runoff across the entire

525 YZ basin and its NX and NX-BXK sub-basins. Compared to the period 1971–1997, the  
526 contribution of rainfall increases by 5%–8% from May to October in the NX for 1998–2020,  
527 whereas glacier and snowmelt contributions decline by -0.3% to -2% and -5% to -7%,  
528 respectively (Figure 7g, Table S2). Conversely, in the NX-BXK, contributions from rainfall  
529 runoff and snowmelt decrease by -2% to -6% during May–October, while glacier contribution  
530 increases by 2%–7% in these months (Figure 7h, Table S2), underscoring the growing  
531 significance of this season in sustaining summer water supplies in the NX-BXK. Taken  
532 together, for the entire YZ basin (Figure 7i), glacier contribution increases by 0.5%–2%  
533 (Table S2) during June–October, and the seasonal changes in rainfall and snowmelt  
534 contributions to total runoff closely mirror those observed in the NX basin.

535

#### 536 **4.2 Hydrological Response to Future Climate Changes**

537 Historical differences in total runoff changes in the NX and NX-BXK sub-basins are  
538 projected to weaken in the future. The YZ basin is projected to experience increased  
539 precipitation (7–33 mm/10yr) and higher temperatures (0.3–0.8 °C/10yr) under the SSP2-4.5  
540 and SSP5-8.5 scenarios throughout the 21st century (Table 5). Predictions indicate an increase  
541 in total runoff for the NX (7–27 mm/10yr) and BXK (34–100 mm/10yr) for 2021–2100 under  
542 both SSPs, with significant increases (36–142 mm/10yr) anticipated in the latter half of the  
543 century (2071–2100) under SSP5-8.5 (Table 5). The changes in total runoff are projected to be  
544 primarily influenced by increased rainfall runoff, with minor contributions from increased

545 snowmelt and glacier runoff under both SSPs scenario through the 21st century (Table 5).  
546 However, in comparison to the 1971–2000 mean, a reduction of approximately -6% to -14%  
547 is projected in the first half of the 21st century (2021–2050) in the YZ and its NX and NX-  
548 B XK sub-basins under the SSP2-4.5 and SSP5-8.5 scenario (Figure 8). This reduction is  
549 attributed to decreased rainfall (-9% to -19%) and snowmelt (-5% to -6%), which may result  
550 in the decline of freshwater supply. Conversely, there is a broadly consistent increase (6%–  
551 32%) in total runoff in the second half of the 21st century (2071–2100), mainly driven by  
552 increased rainfall (4%–52%) and glacier runoff (9%–78%), suggesting that the YZ basin will  
553 not face a water supply crisis in the end of 21st century.

554

555 Changes in meltwater from glaciers and seasonal snow significantly impact total runoff,  
556 influencing both quantity and timing and are particularly important for water availability  
557 during warm and dry seasons (Barnett et al., 2005). Relative to the 1971–2000 mean, the  
558 future annual hydrograph appears relatively stable across all sub-basins (Figure 9), with 60%–  
559 80% of mean annual runoff occurring from June to September. However, a decline of about -  
560 18% to -3% is projected in each month during 2021–2050 under the two SSPs due to  
561 decreased monthly precipitation and precipitation-induced rainfall and snowmelt runoff  
562 (Table S3 and S4). In contrast, there is an anticipated increase of about 6%–40% in each  
563 month, particularly in summer (25%–40%), during 2071–2100 under the two SSPs. The  
564 increased total runoff in the NX basin is primarily attributed to increased rainfall runoff and

565 spring snowmelt, indicating an earlier spring snow melt and delayed fall freeze-up (Figure 9a,  
566 b). Similarly, the increased total runoff in the NX-BXK basin is mostly a result of increased  
567 rainfall and glacier runoff, coupled with decreased snowmelt (Figure 9c, d), primarily due to  
568 reduced snowfall with ongoing warming in each month (Figure S4 and S5). Future changes in  
569 seasonal runoff across the entire YZ basin closely align with those in the NX-BXK sub-basin  
570 (Figure 9e, f) due to its significant contribution to the overall runoff of the YZ basin.

571

## 572 **5 Discussion**

573 Forcing inputs, parameters, and representation of physical processes are major sources of  
574 uncertainty in hydrological model simulations.

575

576 Precipitation is the most important atmospheric input for land surface hydrology models, but  
577 none of the multiple precipitation datasets proves equally suitable for all basins in the TP due  
578 to the high spatiotemporal variability in their performance at the sub-basin scale (Dahri et al.  
579 2021). The variation in precipitation datasets for high mountains can lead to significant  
580 differences in meltwater contribution (Lutz et al., 2014; Zhao et al., 2019; Sun and Su, 2020;  
581 Khanal et al., 2021; Nan et al., 2021; Wang et al., 2021). Duethmann et al. (2014) applied a  
582 multi-objective genetic algorithm to characterize the trade-off curve between model  
583 performance in terms of discharge and snow cover area in Central Asia, suggesting that good  
584 discharge simulations at the catchment outlet cannot guarantee good internal functioning of

585 the model, as different forcing inputs may result in error compensation among different runoff  
586 compositions. Jost et al. (2012) simulated glacier runoff of 25 large glacierized basins  
587 (>50,000 km<sup>2</sup>) in North and South America, Europe, Asia, and New Zealand, suggesting that  
588 the runoff differences ranged from 0.07 % for weakly glacier-influenced basins to 252 % for  
589 strongly glacier-influenced basins. They also suggested that hydrologic model calibration in  
590 glacier-fed catchments was difficult, because errors in modelling snow accumulation can be  
591 offset by compensating errors in glacier melt. Zhang et al. (2013) simulated glacier runoff by  
592 the VIC-Glacier model with the APHRODITE precipitation estimates in the upper Indus (UI)  
593 river basin of the TP during 1961–2009, and suggested that contribution of glacier runoff to  
594 total runoff was about 48.2. However, Meng et al. (2023) simulated glacier runoff by the  
595 VIC-Glacier model with the corrected MERRA-2 precipitation estimates in the UI basin,  
596 suggested that glacier runoff contributed of 24% to total runoff. The difference between  
597 Zhang et al. (2013) and Mengand (2023) mostly resulted from the higher amount of corrected  
598 MERRA-2 than APHRODITE precipitation estimates in the UI basin, because the  
599 underestimation of precipitation-induced runoff would be compensated by glacier runoff.

600

601 Like elsewhere on earth, the aforementioned issues are typical of the YZ basin. In the case of  
602 the NX basin, glacier melt contributed approximately 2–18% to the total runoff in existing  
603 research (Table 1), mostly resulting from differences in forcing inputs used in hydrological  
604 models. The YZ basin received less attention regarding glacier runoff contributions in the

605 NX-BXK, with significant inconsistencies in glacier contributions evident in these studies  
606 (Table 1). Sun and Su (2020) suggested that mean annual glacier runoff contributed about 45%  
607 to total runoff in the NX-BXK sub-basin for 1980–2000, using a hydrological model without  
608 calibration and validation due to a lack of hydrometeorological observations in the sub-basin.  
609 In this study, we utilized newly acquired rain gauge data, and streamflow, glacier mass  
610 balance, and glacier and snow cover observations in the NX-BXK sub-basin, glacier runoff  
611 was simulated using the well-validated VIC-Glacier model, forced by a comprehensively  
612 reconstructed long-term precipitation dataset in this study. The updated contribution of glacier  
613 runoff to total runoff during 1971–2020 in the NX-BXK sub-basin was determined to be 19%.  
614 Furthermore, accurate historical precipitation estimates have the potential to reduce  
615 uncertainty in future projections with the large spread in the GCMs, forming the basis for  
616 correcting future GCM estimates. Different study period also results in the difference of  
617 hydrological model simulation. For example, streamflow also mutates in 1997 at the RKZ  
618 sub-basin of the YZ (Figure S15). Increased precipitation and evaporation caused an  
619 insignificant runoff change during 1971–1997. However, due to significant decrease of  
620 precipitation and increase of evaporation, runoff decreased during 1998–2000, resulting in the  
621 insignificant decrease for 1971–2000 (Figure S15).

622

623

624 Hydrological model themselves have their own uncertainties, such as model parameters and

625 structure of physical processes, which are ideally all taken into account. Reliable parameters  
626 play a crucial role in accurate runoff simulation by hydrological models. The DDF emerges as  
627 the most sensitive parameter for the degree-day glacier model (Hock 2003; Radić; Hock  
628 2010). Zhang et al. (2013) examined the sensitivity of glacier melt runoff to DDF parameters,  
629 suggesting that average annual glacier runoff could change by about 10% with each one unit  
630 change in DDF ( $\text{mm } ^\circ\text{C}^{-1} \text{ day}^{-1}$ ). In this study, the DDF parameters are derived based on  
631 observed glacier mass balance data, with intensive validations on glacier melt, including  
632 observed glacier mass balance and satellite-based glacier area estimates. The uncertainty  
633 associated with VIC model parameters is generally lower than the uncertainties from  
634 precipitation inputs. Su et al. (2022) indicated that changes in the RB are within 8% when  
635  $B_{\text{inf}}$  ranges from 0.05 to 0.4,  $D_2$  ranges from 0.5 to 3.0 m, and the changes in NSE are  
636 generally within 0.1. Therefore, high-density hydrometeorological observations are expected  
637 to better constrain the model and further improve the description of hydrological responses to  
638 climate and spatiotemporal changes in glacier/snow.

639

640 Uncertainties are introduced by different representation of physical processes in hydrological  
641 model, especially the snow and glacier melt simulation in high-mountainous basins. Existing  
642 studies used different definitions of runoff composition. For example, Lutz et al. (2014) and  
643 Khanal et al. (2021) divided total runoff into four compositions: rainfall runoff, snow melt,  
644 glacier melt and baseflow. Some studies also further divided the glacier melt into ice melt and



645 supraglacial snowmelt (Armstrong et al. 2018; Wang et al. 2021). In this study, we divided  
646 total runoff into rainfall runoff, snow melt and glacier runoff. Baseflow is a relatively stable  
647 streamflow composition, and it plays an important role in sustaining surface water flow,  
648 especially for the winter half-year when surface water availability is limited. The VIC model  
649 accounts for baseflow (<https://vic.readthedocs.io/en/master/>), which is comprised of three soil  
650 layers to represent the rapid dynamics of soil moisture movement during storm events  
651 (surface runoff) and the slower deep inter-storm response in the bottom layer (baseflow).  
652 Figure S16 shows mean annual contribution and annual variation contribution of rainfall  
653 runoff, snowmelt, glacier runoff and baseflow to total runoff. The baseflow contribution was  
654 relatively stable, and it only contributed of 4% to total runoff in the NX basin since 1971.  
655 Wang et al. (2022) quantified the contribution of baseflow by the water and energy budget-  
656 based distributed hydrological model (WEB-DHM), and suggested mean annual baseflow  
657 contributed of 3.3% to total runoff. However, different model structures to represent baseflow  
658 processes may also result in uncertainties. In addition, the effect of climate change on the  
659 baseflow in the YZ basin remains uncertain mainly due to the generally poor understanding of  
660 mountain aquifers. Detailed study of infiltration and recharge processes, aquifer  
661 characteristics, and flow pathways needs to be a focus of future research to predict how  
662 baseflow will respond to the changes in climate and cryosphere.

663

664 The representation of glacier melting processes introduces substantial uncertainties in model

665 simulations. The accuracy of distinguishing between debris-free and debris-covered glacier  
666 extents at the basin scale critically influences the simulated contribution of glacier runoff.  
667 Currently, the differentiation between these two glacier surface types relies on elevation  
668 constraints. However, due to the observation in these two glacier surface types, the DDFs  
669 were set to the same value in the debris-free and debris-covered glacier. To address this  
670 pivotal issue, additional glacier observations encompassing both surface types, coupled with  
671 high-quality remote sensing mapping, would solve this key issue. This approach holds the  
672 potential to refine distinctions between debris-free and debris-covered glaciers, thereby  
673 enhancing the precision of model simulations concerning glacier melting processes.

674

675 Another key issue is the restricted comprehension of the effect of snow and ice sublimation on  
676 glacier runoff. Sublimation can potentially be an important component of the high-altitude  
677 water balance in the Himalayan region (Lutz et al., 2016). Sublimation was mostly calculated  
678 based on gauge measurement and estimated using an elevation-dependent potential  
679 sublimation function (Lutz et al., 2016; Khanal et al., 2021; Stigter et al. 2018). Stigter et al.  
680 (2018) suggested that the fraction of snowfall sublimation may be much higher than 21% at  
681 wind-exposed locations in the Himalayan region. Lutz et al. (2016) and Khanal et al. (2021)  
682 proposed that snow sublimation accounts for approximately 10% in the UI basin and 2%–3%  
683 in the YZ basin. Furthermore, the impact of snow sublimation diminished as a result of a  
684 smaller fraction of precipitation falling as snow with ongoing warming (Khanal et al., 2021).

685 Yang et al. (2013) investigated mass balance of a maritime glacier on the YZ basin of the  
686 southeast TP during 2005–2010, and indicated that the mass loss by way of  
687 sublimation/evaporation was quite negligible (about -0.07 m/yr).

688

689 Runoff change is also influenced by land cover and land use. Liu et al. (2023) studied the  
690 effect of vegetation growth induced by climate change to runoff variation during 1981–2010  
691 in the YZ basin with the Variable Infiltration Capacity (VIC) model, suggesting that  
692 implanting grassland effectively reduces flash flood runoff in the short term and balances  
693 groundwater runoff in the long term. Broad-leaved and coniferous forests, with their longer  
694 growth cycles, also play a key role in adjusting soil moisture. Ji et al. (2023) explored the  
695 effect of vegetation growth on runoff changes in the YZ basin by computing the functional  
696 equation for the Normalized Difference Vegetation Index (NDVI) and Budyko parameter,  
697 suggesting that the NDVI and discharge both presented an increasing trend, and the  
698 contributions of NDVI on streamflow change in the 1998–2015 were about 43.04%.

699

700

## 701 **6 Conclusions**

702 This study comprehensively investigates runoff composition, flow changes, and their  
703 attribution across six sub-basins in the YZ for 1971–2020, with a particular focus on the  
704 comparison between the NX and NX-BXK using a newly generated precipitation dataset and

705 a well-validated large-scale VIC-Glacier model with observed streamflow at eight  
706 hydrological stations, glacier mass balance data at two sites, and satellite-based glacier and  
707 snow cover estimates. The study also assesses the future evolution of annual and seasonal  
708 total water availability, as well as glacier runoff and snowmelt contributions, using an  
709 ensemble of multiple GCMs from CMIP6 under two SSPs. The key findings are summarized  
710 as follows:

711

712 1. Large regional differences in runoff regimes were observed in the YZ basin for 1971–2020.  
713 The NX-BXK contributed 52% to total runoff at the Pasighat outlet of the YZ basin, followed  
714 by the YC-NX (25%), LS (10%), and other sub-basins. While rain-induced runoff dominated  
715 the entire YZ (59%–72%), glacier runoff played a more important role in annual total runoff  
716 in downstream sub-basins (16%–19%), particularly in summer (23%–35%).

717

718 2. Regional differences in runoff changes were identified in the YZ basin. Annual runoff  
719 generally increased (8–19 mm/10yr) during 1971–2020 in all sub-basins of the NX basin, but  
720 a significant decrease is noted in the NX-BXK sub-basin (-9.4 mm/10yr). Total runoff trends  
721 reversed after 1998 for all sub-basins of the YZ, with increasing trends during 1971–1997 and  
722 decreasing trends during 1998–2020, influenced by changes in summer rainfall runoff due to  
723 atmospheric moisture and teleconnection indices (PDO, ENSO, and AMO). Glacier runoff  
724 mitigated the decreasing contribution from rainfall since 1998, exhibiting an increased effect

725 on water supply.

726

727 3. Total runoff will consistently increase (27–100 mm/10yr) across the sub-basins through the  
728 21st century, with increases of 7–27 mm/10yr in NX and 34–100 mm/10yr in NX-BXK under  
729 two SSPs, resulting from increased rainfall runoff and minor effect of increased snowmelt and  
730 glacier runoff. Relative to the 1971–2000 mean, a decrease of about -6% to -14% is expected  
731 in the first half of the 21st century (2021–2050), followed by a consistent increase (6%–32%)  
732 in the second half (2071–2100).

733

#### 734 **Acknowledgments**

735 This study was financially supported by the National Natural Science Foundation of China  
736 (42201140), a project funded by the China Postdoctoral Science Foundation (2022M723256),  
737 and the Second Tibetan Plateau Scientific Expedition and Research (STEP) Program  
738 (2019QZKK0201). We extend our gratitude to Daqing Yang, Ying Li, Qikai Sun and Tinghai  
739 Ou for their constructive suggestions.

740

#### 741 **Data availability**

742 Daily precipitation, maximum and minimum temperature, and wind speed estimates with a  
743 spatial resolution of 10×10 km during 1971–2100 were adopted from Sun et al. (2022), and  
744 were downloaded from the National Tibetan Plateau/Third Pole Environment Data Center

745 (TPDC, <https://doi.org/10.11888/Atmos.tpdc.272885>). Daily transient climate estimates, at a  
746 spatial resolution of 10×10 km for 1971–2100 under 20 scenarios (10 GCMs × 2 SSPs) used  
747 in this study were from Sun et al. (2024). Observed streamflow was from the Ministry of  
748 Water Resources, China. Two shapefiles of glacier inventory were downloaded from the  
749 “Environment & Ecological Science Data Center for west China”  
750 (<http://westdc.westgis.ac.cn/glacier>) and Randolph Glacier Inventory (RGI) 6.0  
751 ([https://www.glims.org/RGI/rgi60\\_dl.html](https://www.glims.org/RGI/rgi60_dl.html)). Observed annual glacier mass balance data from  
752 Gurenhekou and Parlung No.94 glacier sites since 2005 were downloaded from the TPDC.  
753 The snow cover fraction (SCF) estimates during 2006–2018 were from the Moderate  
754 Resolution Imaging Spectroradiometer (MODIS) 10CM ([https://nsidc.org/ data](https://nsidc.org/data)).

#### 755 **Author Contributions**

756 He Sun: Conceptualization, Formal analysis, Investigation, Methodology, Resources,  
757 Visualization, Funding acquisition, Writing draft. Tandong Yao: Writing (review and editing).  
758 Fengge Su: Writing (review and editing). Wei Yang: Editing and provision of glacier mass  
759 balance data. Deliang Chen: Writing (review and editing).

760

#### 761 **Competing interests**

762 The authors declare that they have no known competing financial interests or personal  
763 relationships that could have appeared to influence the work reported in this paper.

764

765 **References**

- 766 An, B., Wang, W., Yang, W., Wu, G., Guo, Y., Zhu, H., et al., 2021. Process, mechanisms, and  
767 early warning of glacier collapse-induced river blocking disasters in the Yarlung Tsangpo  
768 Grand Canyon, southeastern Tibetan Plateau, *Sci. Total. Environ.* 151652.  
769 <https://doi.org/10.1016/j.scitotenv.2021.151652>.
- 770 Armstrong, R. L., Rittger, K., Brodzik, M. J., Racoviteanu, A., Barrett, A. P., Khalsa, S. J. S.,  
771 et al., 2019. Runoff from glacier ice and seasonal snow in High Asia: separating melt water  
772 sources in river flow. *Reg. Environ. Change.* 19, 1249-1261. [https://doi.org/10.1007/s10113-](https://doi.org/10.1007/s10113-018-1429-0)  
773 [018-1429-0](https://doi.org/10.1007/s10113-018-1429-0).
- 774 Bahr, D. B., Meier, M. F., Peckham, S. D., 1997. The physical basis of glacier volume-area  
775 scaling. *J. Geophys. Res. Solid. Earth.* 102, 20355-20362. <https://doi.org/10.1029/97jb01696>.
- 776 Barnett, T. P., Adam, J. C., Lettenmaier, D. P., 2005. Potential impacts of a warming climate  
777 on water availability in snow-dominated regions. *Nature.* 438, 303-309.  
778 <https://doi.org/10.1038/nature04141>.
- 779 Chen, C., Zhang, L., Xiao, T., He, J., 2020. Barrier lake bursting and flood routing in the  
780 Yarlung Tsangpo Grand Canyon in October 2018. *J. Hydrol.* 583.  
781 <https://doi.org/10.1016/j.jhydrol.2020.124603>.
- 782 Chen, X., Long, D., Hong, Y., Zeng, C., Yan, D., 2017. Improved modeling of snow and  
783 glacier melting by a progressive two-stage calibration strategy with GRACE and multisource  
784 data: How snow and glacier meltwater contributes to the runoff of the Upper Brahmaputra

785 River basin?, *Water Resour. Res.* 53, 2431–2466. <https://doi.org/10.1002/2016WR019656>.

786 Cui, T., Li, Y., Yang, L., Nan, Y., Li, K., Tudaji, M., et al., 2023. Non-monotonic changes in  
787 Asian Water Towers' streamflow at increasing warming levels. *Nat. Commun.* 14(1), 1176.  
788 <https://doi.org/10.1038/s41467-023-36804-6>.

789 Cuo, L., Zhang, Y., 2017. Spatial patterns of wet season precipitation vertical gradients on the  
790 Tibetan Plateau and the surroundings. *Sci. Rep.* 7, 5057. [https://doi.org/10.1038/s41598-017-](https://doi.org/10.1038/s41598-017-05345-6)  
791 [05345-6](https://doi.org/10.1038/s41598-017-05345-6).

792 Cuo, L., Li, N., Liu, Z., Ding, J., Liang, L., Zhang, Y., Gong, T., 2019. Warming and human  
793 activities induced changes in the Yarlung Tsangpo basin of the Tibetan plateau and their  
794 influences on streamflow. *J. Hydrol-Reg. Stud.* 25, 100625.  
795 <https://doi.org/10.1016/j.ejrh.2019.100625>.

796 Dahri, Z. H., Ludwig, F., Moors, E., Ahmad, S., Ahmad, B., Shoaib, M., et al., 2021. Spatio -  
797 temporal evaluation of gridded precipitation products for the high - altitude Indus basin. *Int. J.*  
798 *Climatol.* 41, 4283-4306. <https://doi.org/10.1002/joc.7073>.

799 Duethmann, D., Peters, J., Blume, T., Vorogushyn, S., & Güntner, A., 2014. The value of  
800 satellite-derived snow cover images for calibrating a hydrological model in snow-dominated  
801 catchments in Central Asia. *Water Resources Res.* 50, 2002-2021. [https://doi.org/](https://doi.org/10.1002/2013WR014382)  
802 [10.1002/2013WR014382](https://doi.org/10.1002/2013WR014382).

803 Gao, C., Liu, L., Ma, D., He, K., Xu, Y. P., 2019. Assessing responses of hydrological  
804 processes to climate change over the southeastern Tibetan Plateau based on resampling of



805 future climate scenarios. *Sci. Total. Environ.* 664, 737-752.  
806 <https://doi.org/10.1016/j.scitotenv.2019.02.013>.

807 Gu, H., Xu, Y. P., Liu, L., Xie, J., Wang, L., Pan, S., Guo, Y., 2023. Seasonal catchment  
808 memory of high mountain rivers in the Tibetan Plateau. *Nat. Commun.* 14(1), 3173.  
809 <https://doi.org/10.1038/s41467-023-38966-9>.

810 Hoang, L. P., Lauri, H., Kummu, M., Koponen, J., van Vliet, M. T. H., Supit, I., Leemans, R.,  
811 et al., 2016. Mekong River flow and hydrological extremes under climate change. *Hydrol.*  
812 *Earth Syst. Sci.* 20, 3027-3041. <https://doi.org/10.5194/hess-20-3027-2016>.

813 Hock, R., 2003. Temperature index melt modelling in mountain areas, *J. Hydrol.* 282, 104-115.  
814 [https://doi.org/10.1016/s0022-1694\(03\)00257-9](https://doi.org/10.1016/s0022-1694(03)00257-9).

815 Ji, G., Yue, S., Zhang, J., Huang, J., Guo, Y., Chen, W. 2023. Assessing the impact of  
816 vegetation variation, climate and human factors on the streamflow variation of yarlung  
817 zangbo river with the corrected budyko equation. *Forests*, 14(7), 1312.

818 Jost, G., Moore, R. D., Menounos, B., Wheate, R., 2012. Quantifying the contribution of  
819 glacier runoff to streamflow in the upper Columbia River Basin, Canada. *Hydrol. Earth. Syst.*  
820 *Sc.* 16, 849-860. <https://doi.org/10.5194/hess-16-849-2012>.

821 Kan, B., Su, F., Xu, B., Xie, Y., Li, J., Zhang, H., 2018. Generation of High Mountain  
822 Precipitation and Temperature Data for a Quantitative Assessment of Flow Regime in the  
823 Upper Yarkant Basin in the Karakoram J. *Geophys. Res. Atmos.* 123, 8462-8486.  
824 <https://doi.org/10.1029/2017jd028055>.

825 Khanal, S., Lutz, A. F., Kraaijenbrink, P. D., van den Hurk, B., Yao, T., & Immerzeel, W. W.,

826 2021. Variable 21st century climate change response for rivers in High Mountain Asia at  
827 seasonal to decadal time scales. *Water Resources Res.* e2020WR029266.  
828 <https://doi.org/10.1029/2020WR029266>.

829 Li, C., Su, F., Yang, D., Tong, K., Meng, F., Kan, B., 2018. Spatiotemporal variation of snow  
830 cover over the Tibetan Plateau based on MODIS snow product, 2001-2014. *Int. J. Climatol.*  
831 38, 708-728, <https://doi.org/10.1002/joc.5204>.

832 Liang, X., Lettenmaie, D. P., Wood, E. F., Burges, S. J., 1994. A simple hydrologically based  
833 model of land-surface water and energy fluxes. *J. Geophys. Res. Atmos.* 99, 14415–14428.  
834 <https://doi.org/10.1029/94jd00483>.

835 Liang, X., Lettenmaier, D. P., Wood, E. F., 1996. One-dimensional statistical dynamic  
836 representation of subgrid spatial variability of precipitation in the two-layer variable  
837 infiltration capacity model. *J. Geophys. Res. Atmos.* 101, 21403-21422.  
838 <https://doi.org/10.1029/96jd01448>.

839 Liu, L., Gu, H., Xie, J., Xu, Y. P., 2020. How well do the ERA-Interim, ERA-5, GLDAS-2.1  
840 and NCEP-R2 reanalysis datasets represent daily air temperature over the Tibetan Plateau? *Int.*  
841 *J. Climatol.* 41, 1484-1505. <https://doi.org/10.1002/joc.6867>.

842 Liu, L., Pan, S. L., Bai, Z. X., Xu, Y. P., 2018. Potential application of hydrological ensemble  
843 prediction in forecasting flood and its components over the Yarlung Zangbo River Basin,  
844 China. *Hydrol. Earth. Syst. Sci. Discuss.* 1-33, <https://doi.org/10.5194/hess-2018-179>.

845 Liu, S., Sun, W., Shen, Y., Li, G., 2003. Glacier changes since the Little Ice Age maximum in

846 the western Qilian Shan, northwest China, and consequences of glacier runoff for water  
847 supply, *J. Glaciol.* 49, 117-124, <https://doi.org/10.3189/172756503781830926>.

848 Liu, X., Lu, H., Yang, K., Xu, Z., Wang, J. 2023. Responses of runoff processes to vegetation  
849 dynamics during 1981–2010 in the Yarlung Zangbo River basin. *Journal of Hydrology:  
850 Regional Studies*, 50, 101553.

851 Liu, Y., Wu, G., Hong, J., Dong, B., Duan, A., Bao, Q., Zhou, L., 2012. Revisiting Asian  
852 monsoon formation and change associated with Tibetan Plateau forcing: II. *Chang. Clim.  
853 Dynam.*, 39, 1183-1195, <https://doi.org/10.1007/s00382-012-1335-y>.

854 Lutz, A. F., Immerzeel, W. W., Kraaijenbrink, P. D., Shrestha, A. B., Bierkens, M. F., 2016.  
855 Climate Change Impacts on the Upper Indus Hydrology: Sources, Shifts and Extremes, *PLoS  
856 One*, 11, e0165630. <https://doi.org/10.1371/journal.pone.0165630>.

857 Lutz, A. F., Immerzeel, W. W., Shrestha, A. B., Bierkens, M. F. P., 2014. Consistent increase  
858 in High Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Change.* 4,  
859 587-592. <https://doi.org/10.1038/nclimate2237>.

860 Meng, F. C., Su, F. G., Li, Y., Tong, K., 2019. Changes in Terrestrial Water Storage During  
861 2003-2014 and Possible Causes in Tibetan Plateau. *J. Geophys. Res. Atmos.* 124, 2909-2931.  
862 <https://doi.org/10.1029/2018jd029552>.

863 Meng, F., Su, F., Sun, H., Huang, J., Li, C., 2023. Divergent runoff regime revealed by  
864 hydrological simulations with corrected precipitation in the upper Indus. *J. Hydrol.* 626,  
865 130315. <https://doi.org/10.1016/j.jhydrol.2023.130315>.

866 Miao, L., Li, S., Zhang, F., Chen, T., Shan, Y., and Zhang, Y., 2020. Future Drought in the Dry  
867 Lands of Asia Under the 1.5 and 2.0 °C Warming Scenarios. *Earth's Future*. 8,  
868 e2019EF001337. <https://doi.org/10.1029/2019ef001337>.

869 Nan, Y., Tian, L., He, Z., Tian, F., Shao, L., 2021. The value of water isotope data on  
870 improving process understanding in a glacierized catchment on the Tibetan Plateau. *Hydrol.*  
871 *Earth Syst. Sci.*, 25(6), 3653-3673. <https://doi.org/10.5194/hess-25-3653-2021>.

872 Niu, Q., Liu, L., Heng, J., Li, H., Xu, Z., 2020. A Multi-Index Evaluation of Drought  
873 Characteristics in the Yarlung Zangbo River Basin of Tibetan Plateau, Southwest China. *Front.*  
874 *Earth. Sc.* 8, 213. <https://doi.org/10.3389/feart.2020.00213>.

875 Pradhan, N. S., Das, P. J., Gupta, N., Shrestha, A. B., 2021. Sustainable Management Options  
876 for Healthy Rivers in South Asia: The Case of Brahmaputra, *Sustainability*, 13, 1987.  
877 <https://doi.org/10.3390/su13031087>.

878 Pritchard, H. D., 2019. Asia's shrinking glaciers protect large populations from drought stress.  
879 *Nature*, 569, 649-654. <https://doi.org/10.1038/s41586-019-1240-1>.

880 Qi, W., Liu, J., Chen, D., 2018. Evaluations and Improvements of GLDAS2.0 and GLDAS2.1  
881 Forcing Data's Applicability for Basin Scale Hydrological Simulations in the Tibetan Plateau.  
882 *J. Geophys. Res. Atmos.* 123, 1128-113,148. <https://doi.org/10.1029/2018JD029116>.

883 Radić, V., Hock, R., 2010. Regional and global volumes of glaciers derived from statistical  
884 upscaling of glacier inventory data. *J. Geophys. Res-earth.* 115.  
885 <https://doi.org/10.1029/2009JF001373>.

886 Shi, X. G., Wood, A. W., Lettenmaier, D. P., 2008. How Essential is Hydrologic Model  
887 Calibration to Seasonal Streamflow Forecasting? *J. Hydrometeorol.* 9(6), 1350–1363.  
888 <https://doi.org/10.1175/2008JHM1001.1>.

889 Stigter, E. E., M. Litt, J. F. Steiner, P. N. J. Bonekamp, J. M. Shea, M. F. P. Bierkens, et al.,  
890 2018. The Importance of Snow Sublimation on a Himalayan Glacier. *Front. Earth. Sci.* 6, 108.  
891 <https://doi.org/10.3389/feart.2018.00108>.

892 Su, F., Hong, Y., Lettenmaier, D. P., 2008. Evaluation of TRMM Multisatellite Precipitation  
893 Analysis (TMPA) and Its Utility in Hydrologic Prediction in the La Plata Basin, *J.*  
894 *Hydrometeorol.* 9, 622-640, <https://doi.org/10.1175/2007jhm944.1>.

895 Su, F., Pritchard, H. D., Yao, T., Huang, J., Ou, T., Meng, F., et al., 2022. Contrasting Fate of  
896 Western Third Pole's Water Resources Under 21st Century Climate Change. *Earth's Future.* 10.  
897 <https://doi.org/10.1029/2022ef002776>.

898 Su, F., Zhang, L., Ou, T., Chen, D., Yao, T., Tong, K., et al., 2016. Hydrological response to  
899 future climate changes for the major upstream river basins in the Tibetan Plateau. *Glob. Planet.*  
900 *Chang.* 136, 82-95. <https://doi.org/10.1016/j.gloplacha.2015.10.012>.

901 Sun, H. Su, F., 2020. Precipitation correction and reconstruction for streamflow simulation  
902 based on 262 rain gauges in the upper Brahmaputra of southern Tibetan Plateau. *J. Hydrol.*  
903 590. <https://doi.org/10.1016/j.jhydrol.2020.125484>.

904 Sun, H., Su, F., He, Z., Ou, T., Chen, D., Li, Z., et al., 2021. Hydrological evaluation of high-  
905 resolution precipitation estimates from the WRF model in the Third Pole river basins. *J.*

906 Hydrometeorol., <https://doi.org/10.1175/jhm-d-20-0272.1>.

907 Sun, H., Yao, T., Su, F., He, Z., Tang, G., Li, N., et al., 2022. Corrected ERA5 precipitation by  
908 machine learning significantly improved flow simulations for the Third Pole basins. *J.*  
909 *Hydrometeorol.* 23. <https://doi.org/10.1175/JHM-D-22-0015.1>.

910 Sun, H., Yao, T., Su, F., Ou, T., He, Z., Tang, G., et al., 2024. Increased glacier melt enhances  
911 future extreme floods in the southern Tibetan Plateau. *Adv. Climate. Change. Res.*

912 Tang, Q., Lan, C., Su, F., Liu, X., Sun, H., Ding, J. et al., 2019. Streamflow change on the  
913 Qinghai-Tibet Plateau and its impacts. *Chin. Sci. Bull.* 64(27), 2807–2821.  
914 <https://doi.org/10.1360/TB-2019-0141>.

915 Tong, K., Su, F., Xu, B., 2016. Quantifying the contribution of glacier meltwater in the  
916 expansion of the largest lake in Tibet. *J. Geophys. Res. Atmos.* 121, 11158–11173.  
917 <https://doi.org/10.1002/2016jd025424>.

918 Tong, K., Su, F., Yang, D., Zhang, L., Hao, Z., 2014. Tibetan Plateau precipitation as depicted  
919 by gauge observations, reanalyses and satellite retrievals. *Int. J. Climatol.* 34, 265-285.  
920 <https://doi.org/10.1002/joc.3682>.

921 Wang, A., Zeng, X., 2012. Evaluation of multireanalysis products with in situ observations  
922 over the Tibetan Plateau. *J. Geophys. Res. Atmos.* 117(D5).  
923 <https://doi.org/10.1029/2011jd016553>.

924 Wang, A., Wang, Y., Su, B., Kundzewicz, Z. W., Tao, H., Wen, S., et al., 2020. Comparison of  
925 Changing Population Exposure to Droughts in River Basins of the Tarim and the Indus.

926 Earth's Future. 8. <https://doi.org/10.1029/2019ef001448>.

927 Wang, L., Cuo, L., Luo, D., Su, F., Ye, Q., Yao, T., et al., 2022: Observing multi-sphere  
928 hydrological changes in the largest river basin of the Tibetan Plateau. *Bull. Am. Meteorol. Soc.*  
929 103(6), E1595-E1620. <https://doi.org/10.1175/BAMS-D-21-0217.1>.

930 Wang, X., Luo, Y., Sun, L., & Shafeeque, M., 2021. Different climate factors contributing for  
931 runoff increases in the high glacierized tributaries of Tarim River Basin, China. *J. Hydrol-reg.*  
932 *stud.* 36, 100845. <https://doi.org/10.1016/j.ejrh.2021.100845>.

933 Wang, Y., Wang, L., Zhou, J., Yao, T., Yang, W., Zhong, X., et al., 2021. Vanishing glaciers at  
934 southeast Tibetan Plateau have not offset the declining runoff at Yarlung Zangbo. *Geophys.*  
935 *Res. Lett.* 48, e2021GL094651. <https://doi.org/10.1029/2021gl094651>.

936 Wu, G., Duan, A., 2008. Weakening Trend in the Atmospheric Heat Source over the Tibetan  
937 Plateau during Recent Decades. Part I: Observations. *J. Clim.* 21, 3149-3164.  
938 <https://doi.org/10.1175/2007jcli1912.1>.

939 Wood, A. W., 2002. Long-range experimental hydrologic forecasting for the eastern United  
940 States. *J. Geophys. Res. Atmos.* 107(D20), ACL-6. <https://doi.org/1029/2001JD000659>.

941 Wood, A. W., Leung, L. R., Sridhar, V., Lettenmaier, D. P., 2004. Hydrologic Implications of  
942 Dynamical and Statistical Approaches to Downscaling Climate Model Outputs. *Clim. Change.*  
943 62, 189-216. <https://doi.org/10.1023/B:CLIM.0000013685.99609.9e>.

944 Wu, G., Duan, A., 2009. Weakening Trend in the Atmospheric Heat Source over the Tibetan  
945 Plateau during Recent Decades. Part II: Connection with Climate Warming. *J. Clim.* 22, 4197-

946 4212. <https://doi.org/10.1175/2009jcli2699.1>.

947 Yang, W., Guo, X., Yao, T., Yang, K., Zhao, L., Li, S., et al., 2011. Summertime surface  
948 energy budget and ablation modeling in the ablation zone of a maritime Tibetan glacier. *J.*  
949 *Geophys. Res. Atmos.* 116. <https://doi.org/10.1029/2010jd015183>.

950 Yang, W., Yao, T., Guo, X., Zhu, M., Li, S., Kattel, D. B., 2013. Mass balance of a maritime  
951 glacier on the southeast Tibetan Plateau and its climatic sensitivity. *J. Geophys. Res. Atmos.*  
952 118, 9579-9594. <https://doi.org/10.1002/jgrd.50760>.

953 Yang, Y., Gao, D., Li, B., 1989. Study on the moisture passage on the lower reaches of the  
954 Yarlung Zangbo river, *Sci. China (Series B)* 32, 580-593.

955 Yang, Z., Zhuo, M., Lu, H., Ma, P., Zhou, K., 2014b. Characteristics of precipitation variation  
956 and its effects on runoff in the Yarlung Zangbo River basin during 1961–2010. *Journal of*  
957 *Glaciology and Geocryology*, 36, 166-172 (in chinese).

958 Yao, T., Bolch, T., Chen, D., Gao, J., Immerzeel, W., Piao, S., et al., 2022. The imbalance of  
959 the Asian water tower. *Nat. Rev. Earth. Environ.* 3(10), 618-632. [https://doi.org/s43017-022-](https://doi.org/s43017-022-00299-4)  
960 00299-4.

961 Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., et al., 2012. Different glacier  
962 status with atmospheric circulations in Tibetan Plateau and surroundings, *Nat. Clim. Chang.* 2,  
963 663-667. <https://doi.org/10.1038/nclimate1580>.

964 Yao, T. 2004. Recent glacial retreat in High Asia in China and its impact on water resource in  
965 Northwest China, *Sci. China (Series D)*. 47, 1065. <https://doi.org/10.1360/03yd0256>.



966 Zhang, L., Su, F., Yang, D., Hao, Z., Tong, K., 2013. Discharge regime and simulation for the  
967 upstream of major rivers over Tibetan Plateau, *J. Geophys. Res. Atmos.* 118, 8500-8518.  
968 <https://doi.org/10.1002/jgrd.50665>.

969 Zhao, C., Yang, W., Westoby, M., An, B., Wu, G., Wang, W., et al., 2022. Brief  
970 communication: An approximately 50 Mm<sup>3</sup> ice-rock avalanche on 22  
971 March 2021 in the Sedongpu valley, southeastern Tibetan Plateau. *The Cryosphere*. 16, 1333-  
972 1340, <https://doi.org/10.5194/tc-16-1333-2022>.

973 Zhao, Q., Ding, Y., Wang, J., Gao, H., Zhang, S., Zhao, C., et al., 2019. Projecting climate  
974 change impacts on hydrological processes on the Tibetan Plateau with model calibration  
975 against the Glacier Inventory Data and observed streamflow. *J. Hydrol.* 573, 60-81.  
976 <https://doi.org/10.1016/j.jhydrol.2019.03.043>.

977 Zhong, L., Ma, Y., Fu, Y., Pan, X., Hu, W., Su, Z., et al., 2014. Assessment of soil water  
978 deficit for the middle reaches of Yarlung-Zangbo River from optical and passive microwave  
979 images. *Remote. Sens. Environ.* 142, 1-8. <https://doi.org/10.1016/j.rse.2013.11.008>.

980