

Deducing Land-Atmosphere Coupling Regimes from SMAP Soil Moisture

Payal R. Makhasana^{1*}, Joseph A. Santanello², Patricia M. Lawston-Parker^{2,3}, Joshua K. Roundy¹

¹Department of Civil, Environmental, and Architectural Engineering, University of Kansas, Lawrence, Kansas

²Hydrological Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, Maryland

³Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland

Correspondence to: Payal R. Makhasana (prmakhasana@ku.edu)

Abstract. In recent years, there has been a growing recognition of the significance of Land-Atmosphere (L-A) interactions and feedback mechanisms in understanding and predicting Earth's water and energy cycles. Soil moisture plays a critical role in mediating the strength of L-A interactions, and is important for understanding the complex and governing processes across this interface. This study aims to identify the significance of soil moisture in identifying L-A coupling strength within the Convective Triggering Potential (CTP) and Humidity Index (HI) framework. To address this, a consistent and reliable dataset of atmospheric profiles is created by merging CTP and HI using Triple Collocation (TC) with three reanalysis datasets. The merged CTP and HI product demonstrates enhanced performance globally as compared to the individual datasets when validated with radiosonde and satellite observations. This merged product of CTP and HI is then used to compare the L-A coupling strength based on Soil Moisture Active Passive Level 3 (SMAPL3) and SMAP Level 4 (SMAPL4) over two decades (2003-2022) where L-A coupling strength is defined as the persistence probability within the dry and wet coupling regimes. Results indicate that the persistency-based coupling strength is related to the ability of soil moisture to predict future atmospheric humidity and dry vs. wet coupling state. The coupling strength in SMAPL4 is consistently stronger than in SMAPL3 and is likely due to its reliance on a land surface model and reduced susceptibility to random noise. The difference in coupling strength based on the same CTP-HI underscores the importance of soil moisture data in estimating coupling strength within the CTP-HI framework. These findings lay the groundwork for understanding the role of L-A interactions and drought evolution due to soil moisture variations, by providing insight into the quantification of coupling strength and its role in drought monitoring and forecast efforts.

Key Word. Land-atmosphere coupling, SMAP observation, Triple collocation, CTP-HI framework.

1 Introduction

Land-atmosphere (L-A) interactions are critical to Earth's complex climate processes and environmental sustainability (Seo and Dirmeyer, 2022; Seneviratne and Stöckli, 2008). These interactions are primarily driven by the two-way energy, momentum, and mass exchanges between the land surface and the overlying atmosphere (Hsu and Dirmeyer, 2023). Among

Deleted: on of

Deleted: and their importance for weather and climate prediction

Deleted: which

Deleted: are

Deleted: interactions between the land surface and the atmosphere

Deleted: is within the framework a d is created using Triple Collocation (TC) and three reanalysis datasets. To isolate the impact of soil moisture within this framework, is used toofCTP and HI globally as compared to the individual datasets when validated with radiosonde and satellite observationsThis merged product of CTP and HI is then used to compareL-A Coupling strength based on where L-A coupling strength is defined as the persistence probability within the dry and wet coupling regimesthe based is related to the ability of soil moisture to predict future atmospheric humidity as measured by the lag correlation between the two variables. This suggests trophoshericprovides a means of prediction forthe coupling stateThe coupling strength in SMAPL4 is consistently stronger as compared tothan in SMAPL3 eResults indicate that higher coupling strength, based on the persistence in the wet and dry coupling regime, is indicative of higher lag-correlation between soil moisture and the CTP and HI (i.e. soil moisture predicts future atmospheric states). This suggests that temporal persistence plays a crucial role in L-A coupling strength. ven withwhen accounting for the difference in sample sizeThe increased coupling strength in SMAPL4 is likely due to the assimilation process and its reliance on a land surface model which reduces susceptibility to random noise compared to SMAPL3. While it is not clear which of the two datasets better represents the "real world coupling", the difference in coupling strength based on the same CTP-HI, underscores the importance of soil moisture data onin estimating coupling strength within the CTP-HI framework.

Deleted: This is due to SMAPL4's advanced assimilation algorithms, which increase temporal autocorrelation by making it less susceptible to random noise.

Deleted: ; therefore, this research assesses the impact of different soil moisture datasets on the classification and distribution of L-A coupling regimes. Using SMAP Level 3 (SMAPL3) and SMAP Level 4 (SMAPL4) soil moisture data, we examine the persistence of dry and wet coupling regimes over two decades (2003-2022), exploring how soil moisture influences coupling classification. An inherent challenge in assessing the significance of soil moisture in L-A coupling classification lies in the need for consistent and u... [1]

Deleted: sensitivity

Deleted: of

Deleted: gaining

Deleted: by gaining insight into the quantification of coupling strength, thereby

Deleted: providing critical insights for

Deleted: future

Deleted: modelling

Deleted: and prediction

Formatted: Font color: Red

Deleted: system

the many components influencing L-A interactions, soil moisture is a critical element (Zhou et al., 2019; Santanello et al., 2018; Saini et al., 2016; Alexander et al., 2022; Wakefield et al., 2019; Findell et al., 2023). Soil moisture is not merely a passive participant but an active modifier of water and heat transfer between the land and the atmosphere (Qi et al., 2023). It serves as a vital component in the climate system and is considered an Essential Climate Variable (ECV) (Liu et al., 2020; Miranda Espinosa et al., 2020; Pratola et al., 2015).

Soil moisture is a critical intermediary in L-A feedbacks, affecting a wide range of atmospheric processes at local and regional scales (Seo and Ha, 2022). Its influence on partitioning energy at the land surface into sensible and latent heat fluxes directly impacts weather patterns, climate variability, and extreme meteorological events (Zhou et al., 2019). L-A interactions are often separated into different regimes (Bennet et al., 2023), such as dry and wet coupling regimes. Dry coupling refers to conditions where the land's dryness limits moisture availability to the atmosphere, typically resulting in less cloud formation and precipitation, leading to hotter, drier conditions that exacerbate droughts. Conversely, wet coupling describes a scenario where abundant soil moisture enhances evaporation and transpiration, increasing humidity, cloud formation, and potentially precipitation. Understanding these interactions is crucial to diagnosing and predicting how the land condition may affect weather and climate.

Classifying L-A coupling regimes requires characterizing both the land state and the atmosphere. An integrated approach, highlighted by Findell and Eltahir (2003), provides a framework for understanding the complex exchanges and feedback loops between the Earth's surface and the air above. Specifically, this framework uses the Convective Triggering Potential (CTP) and Humidity Index (HI) as metrics to classify L-A feedback based on the likelihood of convective precipitation. The CTP is critical in gauging the potential for convection by assessing atmospheric stability, while the HI quantifies the moisture content in the lower troposphere. The CTP-HI framework, was further developed by Roundy et al. (2013), which added the use of soil moisture within the CTP-HI framework. This eliminated the need for a model-based approach for linking the thermodynamic preconditioning of the lower troposphere to the triggering or suppression of convective activities in response to surface conditions. This method underscores the critical role of soil moisture as a crucial driver influencing convective dynamics and provides a means for understanding and predicting weather patterns and regimes based on L-A interactions (Roundy and Wood, 2015; Roundy et al., 2014).

Despite this, the sensitivity of the classification system to soil moisture remains largely unexplored, especially relative to satellite-based observations of soil moisture. Furthermore, quantifying the impact of soil moisture within the CTP-HI framework requires a consistent and reliable data set of atmospheric profiles (i.e., for CTP and HI). However, this is not straightforward, as global observations of lower tropospheric temperature and humidity are limited (Teixeira et al., 2021). Reanalysis data provide the best global estimate of atmospheric data for quantifying the CTP-HI, yet inherent uncertainties in the data cannot be overlooked. These uncertainties often arise from varied data sources, measurement approaches, and spatial and temporal resolution. These factors, along with diverse data assimilation techniques, can contribute to biases and uncertainties within climate reanalysis, thus complicating the quantification of land-atmosphere dynamics, as highlighted by both Jach et al. (2022) and Mukherjee and Mishra (2022).

Deleted:

Deleted: the foundation and ...ctive modifier of water and heat transfer between the land and the atmosphere (Qi et al., 2023). It serves as a vital component in the climate system and represe (... [2])

Deleted: characterizes as

Deleted: e...sential Ce...imate Vv (... [3])

Formatted: Italian

Field Code Changed

Formatted (... [4])

Formatted (... [5])

Field Code Changed

Field Code Changed

Deleted: ...loops (... [6])

Deleted: coupling

Deleted: more ...recipitation. Understanding these interactions is crucial tofor...comprehending ...iagnosing and predicting how the the land's...condition affects ...ay affect the atmosphere and vice versa to predictits impact on ...eather patterns, manage ecoo (... [7])

Deleted: incorporating

Deleted: measures...haracterizing of (... [8])

Deleted: to illuminate the L-A coupling

Deleted: offers insight...rovides a framework for understar (... [9])

Deleted: Findell and Eltahir

Deleted: they...his framework used (... [10])

Deleted: within a n observationally driven modelling framework

Deleted: Within this framework,

Deleted: formation

Deleted: measures

Deleted: low atmospheric moisture levels... The CTP-HI (... [11])

Deleted: advanced...eveloped by Roundy et al. (2013), w (... [12])

Deleted: and thus ...liminated the need for a model- (... [13])

Deleted: levels that could

Deleted: as an indicator of the surface conditions

Deleted: This method provides a sophisticated

Deleted:

Deleted: tool

Deleted:)... ((... [14])

Deleted: Roundy et al. (2013) introduced an approach to (... [15])

Deleted: the extent of soil moisture's sensitivity within thi (... [16])

Deleted: ¶

Deleted: sensitivity ...ithin the CTP-HI framework requir (... [17])

295 One way to address the challenge of uncertainties in reanalysis datasets, is to employ data merging techniques, as suggested by Sun and Fu (2021), Lu et al. (2021), and Feng and Wang (2018), to create a more reliable dataset of CTP and HI based on multiple reanalysis data. Many merging techniques exist, including M-kernel merging (Zhou et al., 2003), optimal interpolation (Lorenzo et al., 2017), Random Forest algorithms (Nguyen et al., 2021), and approaches rooted in Bayesian analysis (Wilson and Fronczyk, 2016) to name a few. However, the Triple Collocation (TC) method has emerged as an invaluable technique for estimating error variances within datasets, as evidenced by research from Gruber et al., (2017), Gruber et al., (2020), Stoffelen, (1998), and Saha et al. (2020). The widespread application of the TC method is primarily due to its effectiveness in utilizing the statistical attributes of multiple datasets to reduce the bias and quantify the uncertainties associated with each dataset. Consequently, the TC method is an ideal choice to create a more robust merged CTP and HI metric for analysis of L-A coupling strength.

305 This study aims to enhance our understanding of L-A coupling classification by developing and applying a merged CTP and HI dataset to isolate the impact of different soil moisture products on the CTP-HI framework and the associated coupling strength. Specifically, it focuses on the L-A coupling that results from using either the Soil Moisture Active Passive Level 3 (SMAPL3, Entekhabi et al., 2016) or SMAP Level 4 (SMAPL4, Rolf et al., 2017) soil moisture products in the CTP-HI framework. The goal of this comparative study is to uncover how soil moisture, as detected through direct satellite observations (SMAPL3) and assimilated data products (SMAPL4), influences L-A coupling strength across the globe. The anticipated insights into how soil moisture variability influences coupling strength will provide critical advancements for assessing hydrological extremes. The comprehensive analysis of the coupling series, detailed in the results and discussion section, underscores the contributions of this study towards improving predictive models for weather and climate applications.

2 Methodology

315 2.1 L-A Coupling Classification

To quantify the L-A coupling, the observational-based CTP-HI framework developed by Roundy et al. (2013) is used. This framework is similar to the Findell and Eltahir (2003) framework in that it utilizes measures of atmospheric stability (CTP), and humidity (HI), but goes a step further by using soil moisture (SM) to classify the CTP-HI into coupling regimes. The use of soil moisture to classify the CTP-HI space provides an observationally based approach to L-A classification, which is different from the Findell and Eltahir (2003) approach that utilized model simulations to classify the CTP-HI space. A brief overview of the classification methodology from Roundy et al. (2013) is given below, organized into three sections. First, the input variables are discussed. Next, a description of the methodology used to classify the CTP-HI space into coupling regimes is presented. Finally, the development of a daily timeseries of L-A coupling classification and its use in creating a measure of coupling strength is described. A conceptual diagram of this process is given in Fig 1.

Deleted: ...ets, is to employ data merging techniques, as suggested by Sun and Fu (2021), Lu et al. (2021), and Feng and Wang (2018), to create a more reliable data (... [18])

Deleted: (

Deleted: (

Deleted: (

Formatted: Danish

Field Code Changed

Field Code Changed

Deleted: Peng et al. (2021), Dash and Sinha (2022),

Deleted: Its widespread application ...s primarily due to its effectiveness in utilizing...the statistical attributes of multiple datasets to reduce the bias discern the accurate signal ...nd quantify the uncertainties associated with each dataset. Therefore...consequently, using the...the TC method to merge reanalysis data sets of CTP and HI based (... [20])

Formatted: Danish

Formatted (... [19])

Field Code Changed

Deleted: has a the potential to provide

Deleted: analysis of land-atmosphere interactions with which to assess the impact of soil moisture on the strength of the coupling regimes.

Deleted: s

Deleted: ting...the impact of different soil moisture datas (... [21])

Deleted: outcome ...rom using either the Soil Moisture Active Passive Level 3 (SMAPL3)...satellite remote sensing ... ADDIN ZOTERO_ITEM CSL_CITATION {"citationID":"UYi8QM0x","properties":{"formattedCitation":"(Entekhabi et al., 2016)","plainCitation":"(Entekhabi et al., 2016)","noteIndex":0},"citationItems":[{"id":390,"uris":["http://zote ro.org/users/11597444/items/8IG297GU"],"itemData":{"id":390,"type":"dataset","DOI":"10.5067/7KKNQ5UURM2W","publisher":"[object Object]","source":"DOI.org (Datacite)","title":"SMAP L3 Radar/Radiometer Global Daily 9 km EASE-Grid Soil Moisture, Version 3","URL":"http://nsidc.org/data/SPL3SMAP/versions/3","author":{"family":"Entekhabi","given":"Dara"}, {"family":"Das","given":"Narendra"}, {"family":"Njoku","given":"Eni"}, {"family":"Johnson","given":"Joel"}, {"family":"Shi","given":"Jiancheng"}],"accessed (... [22])

Deleted: we use ...he observational-based revised (... [23])

Deleted: developed...eveloped by Roundy et al. (2013) is (... [24])

Deleted: ...is similar to originally developed by (... [25])

Deleted: is based on the work of

Deleted: ,

Deleted: and was revised by Roundy et al. (2013) to be a (... [26])

Deleted: ...and measure of atmospheric (... [27])

Deleted: differs in that it utilizes...oes a step further by us (... [28])




Figure 1: A visual representation of the Soil Moisture (SM), Convective Triggering Potential (CTP), and Humidity Index (HI) on a thermodynamic diagram, along with a joint probability space of CTP-HI-SM at a specific grid location (38.89, -115.59). The CTP-HI framework is depicted after applying a two-sample Kolmogorov-Smirnov test on the joint probability space of CTP-HI-SM for the classification period from April 2015 to 2022.

2.1.1 Classification Input Variables

The left panel in Fig. 1 shows a conceptual diagram of the input variables required and includes an atmospheric thermodynamic diagram showing the CTP (atmospheric stability) and HI (atmospheric moisture) as well as a representation of SM (surface moisture). The CTP is the integrated area between the temperature profile and a moist adiabat between 100 and 300 hPa above the surface. It quantifies the stability of the atmosphere, where a more negative value indicates stable conditions, and a more positive value indicates unstable conditions. The HI quantifies moisture content in the lower atmosphere, explicitly reflecting the lower tropospheric dew point depression. The HI is calculated as the sum of the dew point depressions at 50 and 150 hPa above the surface. A large value of HI indicates a dry atmosphere as there is a significant difference between the temperature and the dew point temperature at the specified pressure levels. As the dew point temperature approaches the temperature, the HI decreases and eventually reaches zero when the atmosphere is saturated. SM is a measure of surface wetness and is usually taken as the top-level soil moisture, thus making satellite remote sensing estimates an ideal input. Daily values of each of the three input variables are typically taken before sunrise to capture the state of the land and atmosphere prior to the impact of solar radiation. It is not always possible to have estimates right before sunrise, especially for remote sensing variables which are available only at a fixed overpass time. For these cases, any observation obtained before sunrise may be used, but care must be taken when comparing datasets with inconsistent measurement times (Roundy and Santanello, 2017).

2.1.2 Classification of the CTP-HI Space

The classification process relies on daily values of the early morning estimates of CTP, HI and SM over a classification period. In this work, the classification period was selected as April 2015 to December 2022, to be consistent with the SMAP

Formatted: Centered

Formatted: Font: Bold

Formatted: Space Before: 12 pt, After: 12 pt

Moved (insertion) [1]

Deleted: a

Deleted: -level

Deleted: , suggesting low moisture content

Deleted: depend on the

Deleted:

Deleted: time

Deleted: can work

Formatted: Space Before: 12 pt, After: 12 pt

Deleted: While the revised framework explicitly incorporates and soil moisture to classify coupling regime in CTP-HI framework, the CTP-HI space into coupling regimes as shown in Fig. 1. The left panel in Fig. 1 shows the conceptual thermodynamic diagram soil moisture (SM), CTP, and HI

Deleted: .

observational record. An example of the joint probability space, with the CTP in the x-axis, the HI in the y-axis, and the SM averaged over bins in the CTP-HI space is given in the middle panel of Fig. 1. This joint probability space is then used to define L-A coupling regimes within the 2-dimensional CTP-HI space based on the distribution of soil moisture. This is done by comparing the soil moisture in each bin to the climatological soil moisture using the two-sample Kolmogorov-Smirnov test. Bins with soil moisture distributions significantly wetter than the climatological distribution are classified as a wet regime bin, while those with significantly drier soil moisture distributions are classified as a dry regime bin. This process is repeated using multiple bin sizes and statistical significance thresholds within the CTP-HI space to arrive at a probabilistic classification considering algorithmic uncertainty (see Roundy et. al 2013 for more details) as shown in the right panel. The fundamental underpinning of this approach is that consistent areas within the CTP-HI space with fundamentally different distributions of soil moisture represent a persistent coupling between the land and the atmosphere. Wet coupling regions in the CTP-HI framework are typified by a positive feedback loop through higher surface moisture which results in increased latent heat fluxes that can initiate convection through an increase in moist static energy and a lower lifting condensation level. On the other hand, the dry coupling regime is identified in regions of drier soil moisture within the CTP-HI framework and is characteristic of high sensible heat fluxes leading to strong boundary layer growth and possibly convection, thus causing negative feedback. However, this regime is often associated with less frequent and smaller precipitation events, resulting in an overall drying effect. Locations with a non-dominant combination of wet and dry classification based on the uncertainty parameters are considered to be in a transitional L-A coupling regime. Lastly, areas in the CTP-HI framework that do not show a significantly different distribution of soil moisture relative to the climatology are classified as atmospherically controlled, signifying that the convective processes are predominantly driven by atmospheric factors rather than land surface feedback. To simplify the analysis and to emphasize the crucial role that soil moisture plays in defining the dry and wet regimes, the atmospherically controlled and transitional regimes are merged into a single category termed atmospherically controlled for this analysis. The classification process is done for each dataset of CTP-HI-SM at the grid scale thus allowing for the coupling classification to account for the difference in climatology across datasets and regions around the globe, overcoming a limitation of the original Findell and Eltahir (2003) framework (Ferguson and Wood, 2011).

2.1.3 L-A Coupling Strength

Once the CTP-HI space is classified based on estimates of morning observations of CTP, HI, and SM, a daily coupling timeseries can be generated. The daily coupling is determined by mapping the CTP and HI values for a day onto the classified CTP-HI space (right panel Fig. 1). For example, if the CTP and HI for a particular day map to a wet coupling regime, then that day is classified as a wet coupling regime day. This process is repeated for every day where there is an estimate of CTP and HI. Since the process for determining the daily coupling regime does not require the SM variable, the coupling timeseries can extend beyond the availability of SM data if there are CTP and HI data. Therefore, even though the CTP-HI space was classified on data from 2015-2022, the time series of daily coupling was extended to 2003 based on the availability of CTP and HI data

- Deleted:** s
- Deleted:** the
- Deleted:** which
- Deleted:** typified
- Deleted:** which result in
- Deleted:** that can lead to
- Deleted:** a negative feedback
- Deleted:** te
- Deleted:** s
- Moved (insertion) [4]**
- Deleted:** streamline
- Deleted:** simply
- Deleted:** the
- Deleted:** is
- Deleted:** global
- Deleted:** which was
- Deleted:** Soil moisture plays a crucial role in defining coupling space within CTP-HI framework.
- Moved up [1]:** The CTP is the integrated area between the moist adiabat between 100 and 300 hPa above the surface. It quantifies the stability of the atmosphere, where a more negative value indicates a stable condition, and a more positive value indicates unstable conditions. The HI quantifies moisture content in the lower atmosphere, explicitly reflecting the low-level dew point depression. The HI is calculated as the sum of the dew point depressions at 50 and 150 hPa above the surface. A large value of HI indicates a dry atmosphere as there is a significant difference between the temperature and the dew point temperature at the specified pressure levels, suggesting low moisture content. As the dew point temperature approaches the temperature, the HI decreases and reaches zero when the atmosphere is saturated. ¶
- Moved up [3]:** Soil moisture plays a crucial role in defining
- Deleted:** ¶ (... [29])
- Deleted:** framework
- Deleted:** established
- Deleted:** classification
- Deleted:** classification
- Deleted:** , morning observations of CTP and HI are used (... [30])
- Deleted:** classification
- Deleted:** is
- Deleted:** classification

700 from remote sensing. This extended period provides for a more robust analysis of L-A coupling strength. Within this context, coupling strength is defined as the persistency of a specific coupling regime over the derived timeseries. The coupling strength is calculated by applying a first order three-state (dry, wet, and atmospherically controlled) Markov Chain model to the timeseries. A first order three state Markov Chain describes the evolution of the coupling state through three persistence probabilities and six transitional probabilities based on a one-day lag (i.e. tomorrows coupling state is only dependent on the current coupling state). Of the nine probabilities calculated, only the two persistence probabilities for the dry and wet regimes are used to define the coupling strength (i.e. the probability that it remains in its current state). Since coupling strength is a probability expressed as a percentage (ranging from 0% to 100%, with 100% indicating strong coupling or persistence), higher percentages signify a stable interaction between the land and the atmosphere that can impact weather patterns and short-term climate variability. In contrast, low values of coupling strength indicates weaker L-A interactions.

710 2.2 Merged CTP and HI

Evaluating the sensitivity of the coupling strength to different soil moisture datasets requires a consistent dataset of CTP-HI. The only global CTP and HI datasets available are from reanalysis or satellite. Relying solely on a single reanalysis dataset for CTP-HI could introduce biases and limit the comparability when evaluated alongside multiple soil moisture products. Moreover, satellite-derived CTP-HI estimates face significant challenges, such as missing observations and lower vertical resolution (Roundy and Santanello, 2017), which can impact the quality and reliability of the data.

To address the limitations inherent in single-source CTP-HI estimates, a merged CTP-HI product is developed from three different reanalysis datasets. This approach aims to provide a more comprehensive and reliable benchmark for comparing the impact of soil moisture dynamics on L-A coupling strength. Previous research has shown that the Triple Collocation (TC) data merging methodology is reliable for combining hydrologic variables without requiring ground-based observational data (Yilmaz et al., 2012). This study uses a 30-day centered window (15 days on either side of the day) that removes the effect of seasonality. To reduce complexity due to leap years, the analysis only considers 28 days in February for each year. The TC technique employs a least-squares approach to calculate weight distributions for each dataset based on the root mean square error (RMSE). A core assumption of this method is the independence of each dataset, that is a lack of correlated errors between the datasets. This condition ensures the accuracy of the TC method, as otherwise the merged estimate is prone to biases.

725 Mathematically, the errors within each dataset can be articulated as a linear combination of mutually independent error terms, as mentioned in Equation 1 (Wu et al., 2020).

$$\theta_i = a_i + b_i\theta + \varepsilon_i \quad (1)$$

where, θ_i are collocated measurements of an arbitrary variable (here CTP and HI) for $i = 1, 2,$ and 3 related to the true value θ with ε_i as random errors, a_i and b_i correspond to the intercept and slope obtained through ordinary least squares regression.

730 Considering these assumptions and their potential implications, the resulting merged dataset provides a more comprehensive and accurate representation of the underlying physical phenomenon.

- Deleted:** time period
- Deleted:** of coupling classification
- Formatted:** Font: Not Bold
- Moved (insertion) [5]**
- Deleted:** fall within the CTP-HI framework and identify whether it belong to the wet, dry, or atmospherically controlled regime. The classification period used to develop CTP-HI framework, spanning from April 2015 to 2022, that aligns with the availability of SMAP data. Coupling time series can be generated if CTP and HI data are available. In this study, we used observations from 2003 to 2022 for CTP and HI, enabling us to generate a time series of coupling classifications for the entire period from 2003 to 2022.
- 2.2 L-A Coupling Strength [31]
- Deleted:** One way to define c
- Deleted:** in the context of L-A interactions ...s defined by ...s the persistency of a specific coupling regime over the time [32]
- Deleted:** classification...trength time series ...s then entered...calculated by applying into...a first order three-state (dry, wet, and atmospherically controlled) Markov Chain model to the timeseries. A first order three state Markov Chain describes the evolution of the coupling state through three The outcome of this model is the ...ersistence probabilities and six transitional probabilities based on a one-day lag (i.e. tomorrows coupling state is only dependent on the current coupling state), which is defined as coupling strength [33]
- Deleted:** Since the coupling strength is a probability, , This model calculates the transition probabilities between the three state [34]
- Deleted:** signifies...ignify a robust, [35]
- Deleted:**
- Moved up [4]:** To streamline the analysis and emphasize [37]
- Deleted:** a
- Deleted:** This quantification reflects the intensity and dur [36]
- Deleted:** 3
- Deleted:** different soil moisture data sets on ...he couplin [38]
- Deleted:** datasets
- Deleted:** . These factors
- Deleted:** using data ...rom three different reanalysis datas [39]
- Deleted:** is developed... This approach aims to provide a [40]
- Deleted:** Additionally, the
- Deleted:** comprises
- Deleted:** compound event
- Deleted:** the ...eap years and keep consistency [41]
- Deleted:** ,
- Deleted:** and
- Deleted:** tion...between ...rrors betweenacross [42]
- Deleted:** is crucial for ensuring...nsures the accuracy of th [43]

The TC methodology is employed to create a merged CTP and HI dataset from three reanalysis products: the Modern-Era Retrospective Analysis for Research and Application, version 2 (MERRA2), the Climate Forecast System Reanalysis (CFSR), and European Centre for Medium-Range Weather Forecast (ECMWF) Reanalysis v5 (ERA5). The TC method requires choosing a reference dataset (used to estimate the true values of the measured physical phenomenon). MERRA2 is selected to be the reference dataset, however, it is worth noting that previous studies have indicated that the choice of reference dataset does not impact the outcome of TC error variance (Anderson et al., 2012). The first step involves converting the CFSR and ERA5 estimates (θ) into the MERRA2 climatology (θ') using equations (2) and (3) to establish a common reference framework for the datasets.

$$\theta'_{CFSR} = \mu_{MERRA2} + (\theta_{CFSR} - \mu_{CFSR}) \left(\frac{\sigma_{MERRA2}}{\sigma_{CFSR}} \right) \quad (2)$$

$$\theta'_{ERA5} = \mu_{MERRA2} + (\theta_{ERA5} - \mu_{ERA5}) \left(\frac{\sigma_{MERRA2}}{\sigma_{ERA5}} \right) \quad (3)$$

In the above step, μ represents the mean and σ is the standard deviation of the respective datasets. The normalized composites are linearly scaled and used as input in TC analysis, as described below in equations (4) – (6). Each dataset was rescaled to a consistent grid resolution of $1^\circ \times 1^\circ$ before applying the TC method and was evaluated from 2003 to 2022 to calculate the TC error value (ε^2).

$$\varepsilon_{MERRA2}^2 = \{(\theta_{MERRA2} - \theta'_{CFSR})(\theta_{MERRA2} - \theta'_{ERA5})\} \quad (4)$$

$$\varepsilon_{CFSR}^2 = \{(\theta_{CFSR} - \theta'_{MERRA2})(\theta_{CFSR} - \theta'_{ERA5})\} \quad (5)$$

$$\varepsilon_{ERA5}^2 = \{(\theta_{ERA5} - \theta'_{CFSR})(\theta_{ERA5} - \theta'_{MERRA2})\} \quad (6)$$

The above equation brackets indicate the temporal average of differences between two datasets over the study area.

Mathematically, the ideal merger of a variable from numerous datasets requires information regarding errors, such that a highly accurate data source receives the larger weight for merging and vice versa (Chen et al., 2022). To generate an unbiased merged data product from the three datasets, the sum of individual weights at each grid cell should be one ($w_{MERRA2} + w_{CFSR} + w_{ERA5} = 1$) (Gruber et al., 2017). The merged outcome or cost function is calculated using equations (7) to (9), which minimizes the error variance in the merged outcome obtained from the least square approach and highly depends on the TC error (Lyu et al., 2021).

$$w_{MERRA2} = \frac{\varepsilon_{CFSR}^2 \varepsilon_{ERA5}^2}{\varepsilon_{MERRA2}^2 \varepsilon_{CFSR}^2 + \varepsilon_{MERRA2}^2 \varepsilon_{ERA5}^2 + \varepsilon_{CFSR}^2 \varepsilon_{ERA5}^2} \quad (7)$$

$$w_{CFSR} = \frac{\varepsilon_{MERRA2}^2 \varepsilon_{ERA5}^2}{\varepsilon_{MERRA2}^2 \varepsilon_{CFSR}^2 + \varepsilon_{MERRA2}^2 \varepsilon_{ERA5}^2 + \varepsilon_{CFSR}^2 \varepsilon_{ERA5}^2} \quad (8)$$

$$w_{ERA5} = \frac{\varepsilon_{MERRA2}^2 \varepsilon_{CFSR}^2}{\varepsilon_{MERRA2}^2 \varepsilon_{CFSR}^2 + \varepsilon_{MERRA2}^2 \varepsilon_{ERA5}^2 + \varepsilon_{CFSR}^2 \varepsilon_{ERA5}^2} \quad (9)$$

The merged CTP is then calculated using the weighted sum of the individual datasets:

$$CTP_{merged} = w_{MERRA2} \cdot CTP_{MERRA2} + w_{CFSR} \cdot CTP_{CFSR} + w_{ERA5} \cdot CTP_{ERA5} \quad (10a)$$

$$HI_{merged} = w_{MERRA2} \cdot HI_{MERRA2} + w_{CFSR} \cdot HI_{CFSR} + w_{ERA5} \cdot HI_{ERA5} \quad (10b)$$

Deleted: is

Deleted: triple collocation

Deleted:

Deleted: was

Deleted: t

Deleted: during the

Deleted: period

Deleted: variables

Deleted: Besides,

Deleted: t

Deleted: the

Deleted: ($w_x + w_y + w_z = 1$)

Deleted: In the context of this analysis, the variables x, y, and z correspond to the datasets MERRA2, CFSR, and ERA5, respectively. ...

Deleted: ing

Formatted: Font: (Default) +Body (Times New Roman)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Deleted: ¶

$$w_x = \frac{\varepsilon_y^2 \varepsilon_z^2}{\varepsilon_x^2 \varepsilon_y^2 + \varepsilon_x^2 \varepsilon_z^2 + \varepsilon_y^2 \varepsilon_z^2} \quad (7) ¶$$

$$w_y = \frac{\varepsilon_x^2 \varepsilon_z^2}{\varepsilon_x^2 \varepsilon_y^2 + \varepsilon_x^2 \varepsilon_z^2 + \varepsilon_y^2 \varepsilon_z^2} \quad (8) ¶$$

$$w_z = \frac{\varepsilon_x^2 \varepsilon_y^2}{\varepsilon_x^2 \varepsilon_y^2 + \varepsilon_x^2 \varepsilon_z^2 + \varepsilon_y^2 \varepsilon_z^2} \quad (9) ¶$$

The cost function to merge three datasets is, ¶

$$J = \varepsilon_m^2 = w_x \varepsilon_x^2 + w_y \varepsilon_y^2 + (1 - w_x - w_y) \varepsilon_z^2 \quad (10)$$

The weights remain consistent with the three datasets. The resulting merged CTP and HI, as well as the individual reanalysis products (MERRA2, CFSR, and ERA5) are evaluated against estimates of CTP and HI from in-situ radiosondes and satellite remote sensing based on several summary metrics, including the Mean Absolute Error (MAE), bias, and the correlation coefficient (CC) as discussed in section 4.1.

3. Dataset Description

This study utilizes datasets that include soil moisture products derived from satellite remote sensing, as well as atmospheric profiles for calculating the CTP and HI metrics. The computation of CTP and HI requires surface pressure, 2-meter temperature (T2m) and dew point temperature (DP2M), and vertical profiles of humidity (q) and temperature (T). A summary of these datasets, including their horizontal and temporal resolutions and coverage, is presented in Table 1.

Satellite remote sensing and in-situ data are used to assess the performance of the merged CTP-HI dataset. Specifically, CTP and HI are calculated using data from the Atmospheric Infrared Sounder Version 7 (AIRSv7) as well as radiosonde observations from Integrated Global Radiosonde Archive Version 2 (IGRA2). To ensure spatial consistency in the study, all datasets were standardized to a uniform spatial resolution of 1°x1°, which aligns the analysis with the spatial and temporal coverage of the AIRSv7 from 2003 to 2022.

The merged product is validated using the AIRS overpass time (~1:30 AM local time) to leverage the benefits of remote sensing data (i.e., global coverage). However, the theoretical basis for the CTP-HI framework relies on early morning observations of CTP and HI (Findell and Eltair, 2003; Roundy et al. 2013), which more closely align with the SMAP overpass time (~6 AM local time). Estimates of CTP and HI calculated from reanalysis at 1:30 AM and 6:30 AM reveal variations that suggest that the timing of data acquisition may influence these measurements. Therefore, the merged product is created at two different times, the AIRS overpass time (~1:30AM) and at sunrise. The validation of the merged product is based on CTP and HI calculated at the AIRS overpass time so that it can be directly compared to AIRS, while the merged sunrise CTP and HI is used for the analysis on coupling strength to be consistent with previous L-A coupling work.

Table 1: Description of the dataset for Convective Triggering Potential (CTP), Humidity Index (HI), and Soil Moisture (SM) used in this study.

Dataset	Type	Variable	Horizontal Resolution	Vertical Resolution	Temporal Resolution	Temporal range
MERRA2	Reanalysis	CTP, HI	0.5°x0.625°	72 Levels	6 hours	2003 to 2022
CFSR	Reanalysis	CTP, HI	0.5° x 0.5°	64 Levels	6 hours	2003 to 2022
ERA5	Reanalysis	CTP, HI	31 km	137 Levels	1 hour	2003 to 2022
IGRA2	In-situ	CTP, HI	-	N/A	6-12 hours	2003 to 2022
AIRSv7	Remote Sensing	CTP, HI	1°x1°	24 Levels	12 hours	2003 to 2022

Moved up [2]: Additionally, the study comprises a 30-day centered window (15 days on either side of the compound event) that removes the effect of seasonality. To reduce complexity due to the leap year and keep consistency, the analysis only considers 28 days in February for each year.

- Deleted:** e
- Deleted:** analysis approach
- Deleted:** and an assimilation approach
- Deleted:** frameworks
- Deleted:** ,
- Deleted:** crucial
- Deleted:** and
- Deleted:** ,
- Deleted:** , in addition to surface-level data including 2-meter temperature (T2m) and dew point temperature (DP2M)
- Deleted:** To
- Deleted:** , the analysis
- Deleted:** also
- Deleted:** includes
- Deleted:** satellite remote sensing observation and
- Deleted:** utilized
- Deleted:** i
- Deleted:** for validation
- Deleted:** Comparisons between data from AIRS nighttime overpasses reanalysis and sunrise reanalysis have revealed variations in the estimates of CTP and HI, suggesting
- Deleted:** while
- Deleted:** ed
- Deleted:** using the
- Deleted:** overpass of
- Deleted:** coupling classification

SMAPL3	Remote Sensing	SM	9 km	N/A	12 hours	April 2015 to 2022
SMAPL4	Assimilated Soil Moisture	SM	9 km	N/A	3 hours	April 2015 to 2022

3.1 CTP and HI Datasets

The CTP and HI is calculated using three reanalysis datasets (MERRA2, CFSR and ERA5), satellite estimates, and in-situ observations. When performing TC analysis, it is crucial to consider the presence of biases and errors in the datasets across different variables and applications. For instance, Park et al. (2020), Dong et al. (2020), Arshad et al. (2021), and Kozubek (2020) have observed variations among these reanalysis datasets in their respective studies. Yingshan et al. (2022) found seasonal trend variations in all three datasets and concluded that ERA5 demonstrated superior performance in short-wave and long-wave radiation compared to MERRA2. Zhang et al. (2021) evaluated the surface air temperature in China and reported significant interannual variability in the MERRA2, CFSR, and ERA5 datasets. Hassler and Lauer (2021) noted that performance in tropical areas varies depending on the subset of data used, such as land-only, ocean-only, or land-atmosphere-ocean. Santanello et al. (2015) reported a dry bias in CFSR and a wet bias in MERRA in the overall surface planetary boundary layer (PBL) based on local land-atmosphere coupling (LoCo) analyses over the U.S. Southern Great Plains. A description of each of the CTP-HI datasets is given below.

Deleted: triple collocation

3.1.1 The Modern-Era Retrospective Analysis for Research and Application, version 2 (MERRA2)

NASA's Global Modelling and Assimilation Office (GMAO) developed MERRA2 as an atmospheric reanalysis dataset, employing the Goddard Earth Observing System (GEOS) [Atmospheric General Circulation Model \(AGCM\)](#). [The AGCM is a sophisticated numerical model that simulates the Earth's atmospheric processes, providing a comprehensive framework for understanding climate dynamics and variability](#). MERRA2 provides 6-hourly observations with an approximate spatial resolution of 0.5°x0.625° and includes 72 hybrid pressure levels ranging from the surface to 0.01hPa (Gelaro et al. 2017). The data assimilation system of MERRA2 utilizes the 3D-var algorithm and spans from 1980 to the present. Gelaro et al. (2017) describe [how the dataset incorporates observation-based precipitation to force the land model, ensuring realistic precipitation inputs](#), along with advancements and improvements made in the system.

Deleted: i

Deleted: Catchment Land Surface model.

Deleted: (Ochege et al., 2021)

Deleted: the assimilation process of MERRA2

3.1.2 The Climate Forecast System Reanalysis (CFSR)

The Climate Forecast System Reanalysis (CFSR) is developed by the National Center for Environmental Prediction (NCEP). It covers the period from 1979 to the present. It provides six-hourly variable estimations, including 64 atmospheric levels at a 0.5° x 0.5° horizontal resolution (Saha et al. 2010). Operating as a global coupled atmosphere-ocean-land surface-sea ice system, CFSR incorporates satellite radiance data and employs the Integrated Forecasting System (IFS) Cycle 41r2 with the 3D-var data assimilation system. Observations are carefully considered for each component during the assimilation process of the CFSR dataset, [however CFSR uses observation-based precipitation to force the land model, enhancing precipitation accuracy](#), as highlighted in Saha et al. (2010).

Deleted: s

Deleted:

Deleted: (Centella-Artola et al., 2020)

Deleted: CFSR is a homogeneous reanalysis product suited for long-term climate studies, as observations play a crucial role in the assimilation process.

3.1.3 European Centre for Medium-Range Weather Forecast (ECMWF) Reanalysis v5 (ERA5)

ERA5, the fifth ECMWF reanalysis data of global climate, is accessible from January 1959 to the present and produced by the Copernicus Climate Change Service (C3S). ERA5 provides hourly land and atmospheric climate variable estimations at approximately a 31 km spatial resolution and 137 levels from the surface to 80 km (Bell et al., 2021). It employs the Integrated Forecasting System (IFS) Cycle 41r2 and assimilates satellite and in-situ observations. ERA5 includes advanced screen-level assimilation for 2m temperature and relative humidity components, where the soil moisture is nudged to better match the 2-meter observations. ERA5 assimilates soil moisture from spaceborne scatterometers and integrates various precipitation data sources, improving soil moisture and precipitation estimates. Hersbach et al. (2020) compared ERA5 with radiosonde data and showed temperature, wind, and humidity improvements in the troposphere for the latest version.

Differences in land surface observations among these datasets can impact atmospheric variables and introduce biases. Soil moisture influences evaporation and humidity, while observation-based precipitation enhances land model accuracy, influencing atmospheric moisture and stability. ERA5 benefits from direct soil moisture assimilation, which potentially reduces bias. In contrast, MERRA2 and CFSR use observation-based precipitation to force their land models and rely on model-generated soil moisture. This approach can introduce bias in temperature and humidity profiles due to uncertainties in the modeled soil moisture.

3.1.4 The Integrated Global Radiosonde Archive (IGRA) Version 2

The Integrated Global Radiosonde Archive Version 2 (IGRA2) is a comprehensive dataset provided by the National Center for Environmental Information (NCEI) of the National Oceanic and Atmospheric Administration (NOAA) in the United States. It offers access to high-quality radiosonde observations over 1500 stations worldwide from 1905 to the present (Durre and Yin, 2008). The dataset has undergone quality control and adjustments to correct instrument biases across various regions. These procedures ensure that the IGRA records are homogeneous and robust, making them valuable for long-term climate studies.

Despite its comprehensive coverage, the IGRA2 dataset presents challenges, including non-uniform data distribution across the world due to variable observation frequencies. Although radiosonde observations are typically recorded twice daily at 00:00 and 12:00 Coordinated Universal Time (UTC), in some regions, additional observations are taken at 06:00 and 18:00 UTC. Therefore, the frequency and timing of these observations differ among stations and locations. In this study, calculations for CTP and HI are performed using radiosonde data that fall within a ± 3 -hour window of the night-time AIRS overpass (~01:30 AM local time). This targeted approach aligns CTP and HI calculations with the same observation time during the merging and validation process. Also, when multiple observations are available for a single grid cell, the selection criteria prioritizes the station offering the most frequent data. A map of the geographic location of the IGRA2 radiosonde observation sites, along with regional totals in North America (NAM), South America (SAM), Africa (AFR), Europe (EUR), Asia (ASA),

Deleted: (Centella-Artola et al., 2020).

Formatted: Font: 10 pt

Deleted: Understanding these differences is essential for interpreting reanalysis data.

Formatted: Font: 10 pt

Deleted: from

Deleted: Globe

Deleted: R

Deleted: ;

Deleted: vary


Deleted: C

Deleted: UTC

Deleted: on

Deleted: es across the Globe

and Australia (AUS) is given in Fig. 2. Across the world, 534 locations have data available from 2003 to 2022. Most observations are in Asia, followed by North America and Europe.



1155 Figure 2: IGRA2 observation stations across the Globe, along with regions selected for further analysis and the number of observation stations in each region.

3.1.5 Atmospheric Infrared Sounder Version 7 (AIRSv7)

AIRS was launched in 2002 on NASA's Aqua satellite. AIRS retrieved thermodynamic profiles (temperature and humidity) using passive radiance observations. This study focuses on the descending orbit, covering Northern to Southern Hemisphere movement with an equator crossing at ~1:30 am local time. Data has a 2x/daily temporal resolution, capturing half of the 8-day Aqua orbit repeat cycle. Level 3 files contain averaged quality and geophysical parameters in 1°x1° grid cells, including humidity and temperature profiles at 24 pressure levels from 1 to 1000 hPa (AIRS Project, 2020). In Version 7, short-wave exclusion in the retrieval algorithm reduces bias, and targeted channel selection focused on water vapor retrieval improves temperature/water vapor profile performance (Zhang et al., 2023a). It should be noted that for the processing and analysis of the AIRSv7 data, no alterations were made to the predefined quality control (QC) flags.

3.2 Soil Moisture Active Passive (SMAP)

NASA's Soil Moisture Active Passive (SMAP) mission provides global monitoring of soil moisture content. The Enhanced SMAP Level 3 (SMAPL3) product, derived from the foundational Level 1 and 2 data, provide standardized, gridded global soil moisture at 9 km resolution with the capability to observe the entirety of the Earth's surface every 2-3 days (O'Neill et al. 2021). While the Enhanced SMAP Level 3 is provided at a 9 km resolution, it should be noted that the native radiometer footprint is at ~36 km and the brightness temperatures are interpolated to the 9 km resolution using an optimally localized average method. SMAP Level 4 (SMAPL4) data are produced using a land surface modelling system that assimilates SMAP brightness temperatures. SMAPL4 integrates SMAP Level 2 brightness temperature measurements, along with initialization

Deleted: Working with AMSU-A,

Deleted: , geopotential height,

Formatted: French

Deleted: mapping

Deleted: s

Deleted: (O'Neill et al. 2021)

Deleted: ~30

Deleted: t 9

Deleted: global

Deleted:

Deleted:

Deleted: derived

Deleted: from

and forcing inputs from the Catchment Land Surface Model (CLSM) (Reichle et al. 2017), thus producing 3-hourly, comprehensive estimates of surface and root-zone soil moisture at a 9 km spatial resolution (Qiu et al., 2021). SMAPL3 provides direct satellite remote sensing retrievals of surface soil moisture, while SMAPL4 uses a sophisticated data assimilation system within a modelling framework to create a more comprehensive (i.e., in time and space) soil moisture record. This differentiation is the basis of the analysis, providing a contrasting comparison of the impact of different estimates of soil moisture on the L-A coupling framework and the resulting impact on L-A coupling strength. The morning overpass is used for SMAPL3, while the sunrise soil moisture is estimated through linear interpolation from the 3-hour data for SMAPL4. All available SMAPL3 data are used without filtering based on the quality flags in order to maintain a larger dataset for a comprehensive analysis. SMAPL3 also has missing observations due to the satellite's orbital constraints, which is not the case with the SMAPL4 dataset due to its reliance on the land surface model. This difference in data completeness may affect the comparative analysis. To quantify this impact, a dataset called SMAPL4 L3 is also included, which utilizes SMAPL4 but with the temporal coverage of SMAPL3 data, ensuring a comparison with equal samples of soil moisture estimates.

4. Results

4.1 Evaluation of Merged CTP-HI

The foundational concept of the TC method emphasizes that inaccuracies in individual datasets should remain uncorrelated to avoid errors in the merged product. However, in practice, it is likely that the three reanalysis datasets will exhibit correlated errors. In Fig. 3(a), the scatterplots show the correlated errors between different reanalysis datasets (MERRA2, CFSR, and ERA5) with respect to the IGRA2 observations at a location in Kansas, United States (coordinates 39.96, -95.26). For instance, MERRA2 vs CFSR represents the errors in MERRA2 plotted against the errors in CFSR, both with respect to the IGRA2 observations. These errors are analysed at the AIRS overpass (~1:30 AM local time). The analysis indicates a significant correlation of errors, with a correlation coefficient of -0.85 when assessed against IGRA2 data. This correlation appears more pronounced for the CTP than the HI, suggesting a stronger association of errors within the stability metric (CTP). This is further seen at the global scale in Fig. 3(b) showing boxplots for observed error correlation across the 534 IGRA2 locations. Most locations have a correlation exceeding 0.7. This high correlation points to similar error sources in the datasets for CTP and HI across the three datasets considered. This may stem from similar approaches and data used in assimilating temperature and humidity profiles or their radiances, while HI may be influenced more by differences in model physics and parameters, screen-level nudging (in ERA5), compounded by challenges in assimilating near-surface humidity data. Even though the datasets employ diverse methods of Data Assimilation (DA) and feature distinct physics, parameters, etc., the common input of in-situ and satellite data likely contributes to this correlation. These correlated errors may affect the accuracy of the resulting merged dataset, which is validated against observations in section 4.1.2.

Deleted: or

Deleted: . At the same time... while SMAPL4 data builds... [44]

Deleted: this with a

Deleted: vital for our...he basis of the analysis, enabling us to assess...roviding a contrasting comparison of the benefits ...mpact of incorporating the...ifferent observations ...imates of soil moisture into [45]

Deleted: modelling system in capturing soil moisture dynamics

Deleted: This study utilizes a

Deleted: points ...re used without filtering based on the quality flags in order . This decision is made ...o maintain a larger dataset for a comprehensive analysis and ensure analysis remain comprehensive [46]

Deleted: It is also essential to recognize that

Deleted: Also, ...MAPL3 also has missing observations due to the satellite's orbital constraints, which is not the case with the comprehensive [47]

Deleted: spanning from April 2015 to 2022

Deleted: variance ...ifference in data completeness could [48]

Deleted: will

Deleted: address ...uantify this impact, we introduce [49]

Deleted: synthetic ...ata [50]

Deleted: labelled

Deleted: an

Deleted: This study uses the morning observations, specifically from the sunrise overpass , of SMAP soil moisture data, which are crucial in capturing its interactions with the atmospheric processes. The different SMAP data products are used to classify develop the CTP-HI framework from 2015 to 2022 for all datasets to assure consistency with the analysis. Since soil moisture measurements are only needed for the classification period, the time series of coupling classification can still cover the entire time series of CTP-HI from 2003 to 2022.

Deleted: ¶

Deleted: the

Deleted: amplification of

Deleted: in the aggregated outcome from multiple sources...n the merged product. However, in practice, it is likely that theThe merged dataset, derived from...three reanalysis datasets , ... [51]

Deleted:


Deleted: were...analysed across the datasets MERRA2, CFSR, and ERA5 ...t the AIRS overpass time ...~1:30 AM local time), using observational data from IGRA2 at a location in Kansa... [52]

Deleted: This may stem from similar errors in assimilatin... [53]

Deleted: aving

Deleted: The reason for the similarities across the differe... [54]

Deleted: ing...distinct physics, parameters, etc., the comm... [55]



1370 **Figure 3: (a) Observed errors between IGRA2 and reanalysis datasets at a location in Kansas, United States (Latitude: 39.96, Longitude: -95.26) for CTP and HI from 2003 to 2022, along with the Pearson correlation displayed in the upper left corner. (b) Observed error correlation between reanalysis data sets at IGRA2 stations across the Globe (529 sites).**

The triple collocation-based error (TC error) using equations (4)-(6) given in the supplementary material (Fig. S1) reveals a remarkably high TC error for the CTP within the MERRA2 and CFSR datasets, most notably over the northern hemisphere and South America. The merged product is constructed based on the weights derived from the error variance calculated by equations (7)-(9). Locations with a higher error variance are assigned lower weights, reflecting their reduced reliability. Conversely, locations with lower error variance are deemed more reliable and thus are given greater weights in the grid level merging process.

Table 2 demonstrates the distribution of TC-based weights for CTP and HI across various global regions for the three datasets. The spatial maps in the supplementary material (Fig. S2) detail the weights for each reanalysis product for CTP and HI globally. These weights are directly proportional to the relative error variance; areas where MERRA2 and CFSR show larger variances (Fig. S3), especially in the northern hemisphere and South America, tend to favour ERA5 for the weighting of the merged CTP. Based on Table 2, ERA5 emerges as the leading dataset for CTP, being allocated the highest weight in most regions. However, in Europe and Africa, the weight distribution for CTP is almost similar among the three datasets, indicating a balanced reliance on each dataset within these continents.




Table 2: Average weight across North America (NAM), South America (SAM), Africa (AFR), Europe (EUR), Asia (ASA), and Australia (AUS). The color gradient is applied based on the minimum value (yellow color) to the maximum value (green color) for CTP and HI for AIRS overpass.

Weight distribution across the region for CTP (J/Kg)						
Region	NAM	SAM	AFR	EUR	ASA	AUS
MERRA2	0.27	0.27	0.32	0.35	0.28	0.30
CFSR	0.30	0.30	0.32	0.34	0.30	0.30
ERA5	0.43	0.43	0.35	0.31	0.42	0.40
Weight distribution across the region for HI (°C)						
Region	NAM	SAM	AFR	EUR	ASA	AUS
MERRA2	0.33	0.29	0.35	0.37	0.35	0.31
CFSR	0.31	0.31	0.33	0.34	0.30	0.34
ERA5	0.36	0.40	0.32	0.29	0.35	0.36

The weight allocation for the HI shows considerable regional variation. In South America, the MERRA2 dataset is assigned the lowest weight, implying that it contributes less to the combined HI product. Conversely, in Europe, it is the ERA5 dataset that receives the lowest weight, signifying its reduced contribution to the HI variable in this region. These regional differences in the weighting of datasets underscore the merging process, allowing for a location-specific approach to creating a merged product.

4.1.2 Performance of Merged CTP-HI

The data are merged following equations 10(a) and 10(b), and the resultant spatial distribution of average CTP and HI during the summer season (June, July, and August of JJA) for the year 2012 is portrayed in supplementary material (Fig S3 and S4). These figures compare the CTP and HI values derived from the MERRA2, CFSR, and ERA5 alongside the merged product.

The accuracy of the merged data, along the individual reanalysis datasets MERRA2, CFSR, and ERA5, is evaluated through a comparison with IGRA2 radiosonde and AIRSv7 satellite observations, as depicted in Fig. 4 and 5. The evaluation utilizes the Mean Absolute Error (MAE), bias, and Correlation Coefficient (CC) across major global regions. The results indicate that the merged dataset consistently achieves the lowest MAE for CTP and HI variables, outperforming the individual reanalysis datasets. This improvement in accuracy suggests that the merging process effectively consolidates the strengths of the individual datasets while mitigating their respective biases despite the correlated errors seen in Fig. 3.

The merged product compensates for these inaccuracies in regions where the MERRA2 dataset exhibits substantial discrepancies with observational data, demonstrating a practical approach that improves the overall metric. Furthermore, the merged dataset shows a considerable reduction in positive bias for CTP across South America, Europe, and Asia, as well as a decrease in negative bias for HI in most regions, except for Asia. These improvements highlight the efficacy of the merging process in yielding a more reliable dataset, which is especially beneficial for L-A coupling studies in regions challenged by less accurate reanalysis data.

Weight distribution across the region				
Region	NAM	SAM	AFR	EUR
MERRA2	0.2724	0.2745	0.3241	0.35
CFSR	0.2964	0.2992	0.3241	0.33
ERA5	0.4311	0.4262	0.3496	0.30
Weight distribution across the region				
Region	NAM	SAM	AFR	EUR
MERRA2	0.3289	0.2883	0.3471	0.36
CFSR	0.3125	0.3124	0.3318	0.33
ERA5	0.3584	0.3991	0.321	0.29

Deleted:

Formatted: Line spacing: 1.5 lines

Deleted: gets

Deleted: (

Deleted: is

Deleted: s

Deleted: datasets

Deleted: set

Deleted: alongside

Deleted: evaluation

Deleted: estimation

Deleted: performance metrics such as

Deleted: that perform best independently

Deleted: integration

Deleted: representation

Deleted: regions like





Figure 4: Bar plot of performance metric and intercomparison of merged data and reanalysis with radiosonde observation from IGRA2 in different region and globally for (a) CTP and (b) HI.



Deleted:

Deleted: each global region





Figure 5: Bar plot of performance metric and intercomparison of merged data and reanalysis with satellite remote sensing from AIRSv7 in different region and globally for (a) CTP and (b) HI.



Deleted:

Deleted: each global region for

Fig. 5 presents the same set of performance metrics as in Fig. 4 but compared against the AIRSv7 data. The result indicates a general trend of overestimation for CTP and HI by the reanalysis and merged datasets across all regions as compared to AIRSv7. It is essential to acknowledge the limitations in measurement capabilities and inherent biases of AIRSv7, particularly in terms of lower troposphere retrieval due to limited vertical resolution in AIRS, which is different from those in the reanalysis datasets. Specifically, the limited sensitivity and vertical resolution of AIRSv7 in the PBL might lead to positive biases, resulting in discrepancies with the reanalysis datasets vs. that observed with radiosonde data as the reference.

Regarding temporal correlation, the merged dataset and the ERA5 reanalysis score the highest values, reflecting a consistent signal with observations from IGRA2 and AIRSv7 (as shown in Fig. 4 and 5). When evaluated against both ground-based observations and satellite remote sensing, there is a noticeable improvement in the performance of the merged product compared to the individual reanalysis datasets. The merged product demonstrates a closer alignment with the ground truth, as represented by IGRA2, capturing the observed signal with greater accuracy. Overall, the outcome remains consistent across the globe when validating the merged product and individual reanalysis datasets with respect to IGRA2 and AIRSv7 for CTP and HI.

The previous evaluation of the CTP and HI products were done at the AIRS overpass time (~1:30AM local time), however, the theory of the CTP-HI framework for classifying coupling relies on early morning observations of the CTP and HI (Findell and Eltair, 2003) which also aligns with the SMAP overpass time (~6AM local time). Table S5 gives the weights for the merged product at sunrise. As compared to the AIRS overpass weights (Table 2), the sunrise weights show similar spatial patterns with marginal variations. This is consistent with previous research by Roundy and Santanello (2017) that demonstrated the application of CTP-HI framework with AIRS estimates and demonstrated similarity in the overarching pattern of L-A coupling classification. Thus, the remainder of the analysis will use the sunrise merged CTP and HI products to classify and analyse L-A coupling.

4.2 Coupling Strength in the Contiguous United States

The sunrise merged CTP and HI products and three different datasets of soil moisture (SMAPL3, SMAPL4 L3, and SMAPL4) are used to classify the CTP-HI space and create a timeseries of L-A coupling from 2003-2022. The timeseries is then used to quantify the L-A coupling strength based on the persistence probabilities from a 1st order three state Markov Chain. Fig. 6 examines the coupling strength over the contiguous United States using the same CTP-HI data but with different soil moisture datasets. Fig. 6(a) reveals a consistent pattern of a persistent dry regime in the inter-mountain west region and a persistent wet regime in the north-western and eastern parts of the country. A side-by-side evaluation shows that the SMAPL4 dataset displays a more persistent pattern under dry and wet regimes. This indicates a stronger representation of L-A coupling in SMAPL4 that may be due to the strong vertical coupling of soil moisture in the catchment assimilation processes. On the other hand, SMAPL4_L3 data shows less persistence than SMAPL4, suggesting that part of observed difference between SMAPL3 and SMAPL4 is due to sample size. Despite these differences, the overarching spatial pattern remains consistent across all datasets, except over the northern Great Lakes for SMAPL4 where it shows a persistent wet coupling regime.

Deleted: accuracy

Deleted: figure

Deleted: A

Deleted: ing

Deleted: may ...s different from those in the other ...analysis datasets, is essential... For example (... [56])

Deleted: The bar representing the global comparison reinforces the trend observed in individual regions, providing a comprehensive overview. This enhancement in accuracy emphasizes the merged product's value as a reliable and comprehensive dataset, which is especially beneficial for future research on L-A coupling.

Deleted: product of sunrise ...TP and HI products to classify and analyse (... [57])

Deleted: The previous section evaluated the merged CTP and HI products at the AIRS overpass time (~1:30AM local time), however, the theory of the CTP-HI framework for classifying coupling relies on early morning observations of the CTP and HI (Findell and Eltair, 2003) which also aligns with the SMAP overpass time (~6AM local time). Table S5 gives the weights for the merged product at sunrise. As compared to the AIRS overpass weights (Table 2 Fig. 4), the sunrise weights show similar spatial patterns with marginal variations. This is consistent with previous research by Roundy and Santanello (2017) that demonstrated some differences between the AIRS and sunrise overpasses estimates of CTP and HI, yet acknowledged the overarching similarity in their patterns. Thus, the remainder of the analysis will use a merged product of sunrise CTP and HI to classify the L-A coupling. (... [58])

Deleted: overpass ...erged CTP and, (... [58])

Deleted: a...nd (... [59])

Deleted:

Deleted: are combined with

Deleted: create a...lassify the ...TP-HI framework...pace and create a timeseries of . Using the sunrise overpass merged C (... [60])

Moved down [6]: Fig. 6 examines the L-A coupling strength of

Deleted: timeseries of ...-A coupling classification (... [61])

Deleted: timeseries classified ...rom 2003-2022. (... [62])

Deleted: t...e timeseries is then used to quantify the Fig. (... [63])

Deleted: of

Deleted: persistence ...three state Markov Chain. Fig. 6 ex (... [64])

Formatted: Superscript

Deleted: for the

Deleted: datasets

Deleted: simulation...atasets in CTP-HI framework... Fig (... [65])

Deleted: ent

Deleted: conditions

Deleted: n...stronger enhanced...representation of L-A co (... [66])

Deleted: for

To delve deeper into the noted differences in coupling strength, the lagged correlation between the three sets of soil moisture and the CTP and HI over 2015-2022 are analysed. Lag correlation is employed to identify the relationship between soil moisture and future CTP and HI and vice versa. While previous work has discussed the potential for soil moisture predictability out to 60 days (Dirmeyer et al., 2018), this analysis uses a 10-day lag to capture the role of soil moisture in predicting the atmospheric state (CTP and HI) and the atmospheric state in predicting soil moisture on time scales relative to typical weather predictability. Within this setup, the ability of soil moisture to predict future CTP and HI is given as a positive lag correlation and the ability of the CTP and HI to predict future soil moisture is given as a negative lag. For both the CTP and HI, the correlation with soil moisture is negative due to the relationship between SM-CTP and SM-HI. Wet soil typically results in surface cooling when solar radiation is limited, leading to a more stable temperature profile in the lower atmosphere. This stability restricts vertical movement and consequently leads to a lower CTP, thus creating a negative correlation. HI, on the other hand, measures atmospheric moisture content. Higher HI values signify drier air, while lower values indicate moisture-rich air closer to saturation. High soil moisture enhances evaporation, which adds water vapor to the atmosphere, reducing the gap between temperature and dew point and thus lowering the HI, resulting in a negative correlation.




Figure 6: Comparative analysis of L-A coupling strength using SMAPL3, SMAPL4 L3, and SMAPL4 data (a) coupling strength for the dry and wet coupling regime and (b and c) lag- correlation between soil moisture and CTP and HI averaged over the contiguous United States.

Fig. 6(b) and 6(c) show the average lag correlation out to 10 days over the contiguous US and indicate that soil moisture has a stronger predictive influence on CTP and HI as shown by the larger magnitudes of correlations over positive lags, which suggest that while CTP and HI are not as strong in predicting soil moisture (lower magnitude of correlations over negative lags), soil moisture is more effective in predicting changes in CTP and HI. This is particularly noticeable for shorter lags, suggesting a more immediate impact of soil moisture on atmospheric stability and humidity. Conversely, the decrease in correlation magnitude with longer lags highlights the diminishing influence of L-A interaction over time. For the different datasets, SMAPL4 consistently shows higher correlations at all lag intervals for both CTP and HI. However, the sample size

- Deleted: observed
- Deleted: and
- Deleted: with
- Deleted: d
- Deleted:
- Deleted: only extends to
- Deleted: to
- Deleted: to
- Moved down [7]: is shown in Fig. 6(b) and 6(c).
- Deleted: for the three data sets
- Deleted: The reason behind the lag-correlation study is to identify the
- Deleted: driving factors behind the observed differences in coupling strength. By analysing the lagged correlation between soil moisture and the CTP and HI metrics, we aim to understand the variable-dependent predictability of
- Deleted: and HI from soil moisture (
- Deleted:)
- Deleted: predictability
- Deleted: soil moisture informed by
- Deleted: (
- Deleted:)
- Deleted: (Dirmeyer et al., 2018) (Santanello et al., 2018)
- Formatted: Caption
- Formatted: Font color: Black
- Deleted: is shown in
- Deleted: .
- Deleted: ¶
- Reliable weather predictability is generally limited to 10 days Because reliable weather predictability is generally limited to 10 days (Krishnamurthy, 2019). Therefore,
- Deleted: 10-day
- Deleted: is analysed to identify variable-dependent predictability of CTP and HI from soil moisture (positive lag) and the (... [67])
- Deleted: for shorter lag periods,
- Deleted: e higher correlation
- Deleted: s
- Moved up [8]: and HI, the correlation with soil moisture is
- Moved (insertion) [8]
- Deleted: ¶
- Deleted: , the 10-day average lag correlation over the con (... [68])
- Deleted: For both the and HI, the correlation with soil mo (... [69])
- Deleted: soil moisture is a more reliable predictor of CTP (... [70])

725 does play a role in this assessment as noticed by a decrease in the magnitude of correlation for the SMAPL4 L3. Despite this, the SMAPL4 L3 dataset still shows a higher magnitude in lag correlation as compared SMAPL3, particularly for CTP. This suggests that the assimilation of SMAP observations into a model, as in SMAPL4, may yield a stronger relationship in the temporal dynamics between the land surface and the atmosphere. In contrast, the pattern of a stronger L-A connection for SMAPL4 is less evident for HI.

To further analyse the impact of soil moisture on coupling strength, Fig. 7 delves into a grid-scale analysis at a location in Nevada, USA (38.89, -115.59). Fig. 7(a) illustrates the classified CTP-HI space based on SMAPL3, SMAPL4 L3, and SMAPL4 datasets. The coupling regimes are clearly distinguished within the CTP-HI framework across the datasets,

730 highlighting the variations and interactions between soil moisture and atmospheric conditions. SMAPL4 indicates more bins assigned to wet and dry coupling regimes, thereby resulting in an increased coupling strength within the time series. In Fig. 7(b), the joint probability of CTP-HI-SM space illustrates the bin average SM within the CTP-HI space based on historical observations and helps to identify patterns in the CTP-HI-SM relationship. States of wet coupling are associated with higher soil moisture levels in the combined probability space of CTP-HI-SM, whereas dry states are linked to consistent lower

735 moisture levels within the CTP-HI space. While the overall disparities between SMAPL3 and SMAPL4 in the distribution of soil moisture are quite evident in Fig. 7(c), there is also very little difference in soil moisture distribution due to sample size (difference between SMAPL4 and SMAPL4 L3). Given that the classification algorithm accounts for climatological difference in the soil moisture datasets, the difference stems from the shape of the soil moisture distribution and its projection on to the CTP-HI space. The SMAPL3 dataset shows a tendency for observations to skew toward the lower end of the soil

740 moisture spectrum, while the SMAPL4 tends to exhibit a clustering of observations in the mid-range, between 0.4 to 0.8. These differences in both the soil moisture distribution and its projection in the CTP-HI space affect the classification of coupling regimes and therefore the coupling strength of the timeseries. Fig. 7(d) depicts the daily coupling classification for an arbitrary month (May 2010). The SMAPL4 dataset, with its higher number of wet regime classifications, demonstrates a greater likelihood of days being categorized as a wet regime. This is evidenced in the time series, where most days are classified under

745 wet conditions in SMAPL4, in contrast to the SMAPL3 dataset, which indicates more days in an atmospherically controlled regime. Sample size has a small impact on the classification, with only two days being different between SMAPL4 L3 and SMAPL4. This difference underscores the influence of soil moisture on the daily classification of L-A coupling within the CTP-HI framework, even though it is not directly used in creating daily timeseries.

- Deleted: , further
- Deleted: ing
- Deleted: assimilation process
- Deleted: s
- Deleted: is
- Deleted: with SMAPL3, especially for the
- Deleted: CTP
- Deleted: ¶
- Deleted: the specific coordinates (38.89, -115.59), located
- Deleted: , which shows notable differences in the dry and
- Deleted: shows
- Deleted: The outcome and clearly distinguishes t
- Deleted: in
- Deleted: space
- Deleted: .
- Deleted: Not surprisingly,
- Deleted: indicating
- Deleted: observed
- Deleted: the frequency of occurrences for various combinations of
- Deleted: ,
- Deleted: , and
- Deleted: . This
- Deleted: how
- Deleted: soil moisture influences the lower atmosphere
- Deleted: Fig. 7(b) illustrates the relationship between soil
- Deleted: The
- Deleted: .
- Deleted: . At the same time
- Deleted: ¶
- Deleted: There is also very little difference in soil moistu





Figure 7: Multifaceted assessment of coupling classification influenced by soil moisture across SMAPL3, SMAPL4_L3, and SMAPL4 data at a specific grid location (38.89, -115.59) situated in Nevada, USA (a) CTP-HI framework (b) A joint probability space of CTP-HI-SM (c) Saturate soil moisture distribution (d) Coupling classification time series for May 2010.

4.3 Global Coupling Strength

The previous section highlighted the connection between coupling strength and the lag-correlation between SM and the CTP and HI. To explore this further, Fig. 8 presents the average coupling strength (for both dry and wet regimes) on the x-axis and the average positive (soil moisture predicts CTP-HI) lag correlation out to 10 days on the y-axis for all land grid cells across the globe. Overall, all soil moisture datasets show a non-linear relationship between coupling strength and lag correlation, with little relationship between the variables for low coupling strength that transitions into a stronger relationship as coupling strength increases. To help quantify this, the data is fit to a quadratic model, and the R^2 for this relationship is shown. The R^2 value is calculated to measure how well the variance in the observed data can be explained by the quadratic model. As indicated in Fig. 8(a), all soil moisture datasets show a weaker relationship between average coupling strength and average lag correlation for CTP as indicated by a lower R^2 (explained variance by the regression line) and a shallower slope in the regression line. In contrast, Fig. 8(b) shows the SM-HI relationship is stronger, with higher R^2 values and more pronounced nonlinear relationship. This indicates that persistency as a measure of coupling strength is predominantly driven by the SM-HI relationship, suggesting a direct influence of soil moisture on lower-level humidity and its correlation over time. In contrast, the SM-CTP relations is more complex due to the indirect influence on atmospheric stability, which may be more influenced by larger-scale atmospheric conditions. Furthermore, SMAPL4 shows a stronger relationship as compared to SMAPL3, that is only slightly impacted by the difference in sample size between SMAPL3 and SMAPL4.



- Deleted: ¶
- Formatted: Line spacing: 1.5 lines
- Deleted: These differences in both the soil moisture distribution and its projection in the CTP-HI space affect the classification (... [75])
- Deleted: (persistence)
- Deleted: soil moisture
- Deleted: XX
- Deleted: over
- Deleted:
- Deleted: mentioned
- Deleted: For
- Deleted: , the soil moisture CTP relationship
- Deleted: is
- Deleted: as
- Deleted: the
- Deleted:
- Deleted: of
- Deleted: Fig. 8b highlights
- Deleted: soil moisture
- Deleted: soil moisture
- Deleted: , as opposed to the
- Deleted: s
- Deleted: stability
- Deleted: y (CTP)
- Deleted: ¶






Figure 8: Comparative analysis of average dry and wet coupling strength and 10-day lag correlation using SMAPL3, SMAPL4_L3, and SMAPL4 soil moisture datasets for (a) CTP and (b) HI.

The average coupling strength for dry and wet regimes across the different regions and globally using SMAPL3, SMAPL4_L3, and SMAPL4 is given in Fig. 9. All datasets are consistent in showing that Africa has the largest average coupling strength, while North America has the smallest average coupling strength. Yet, there are differences in the relative strength for other regions. Notably, the SMAPL4 dataset demonstrates a stronger coupling strength in both the dry and wet regimes, indicating a stronger temporal persistence in the coupling regime. The variation in coupling strength becomes noticeable when the sample size is considered. This is particularly true for Africa, where there is little difference between the average coupling strength between SMAPL3 and SMAPL4_L3. In contrast, North America shows the largest difference in coupling strength between SMAPL3 and SMAPL4_L3. As with other regions of the globe, there is little difference in the coupling strength for the dry regime, as differences are predominantly seen in the wet regime coupling strength. This outcome is consistent with Fig. 6, which shows the spatial variability of coupling strength over the contiguous United States.



Deleted:

Moved down [9]: Fig. 9 shows

Deleted: th

Deleted: global

Deleted: from

Moved (insertion) [9]

Deleted: shows

Deleted: taken into account

Deleted: However, when considering the impact of sample size, the difference in coupling strength is dimensioned.

Deleted: but is

Deleted: †

Formatted: Centered

Figure 9: Comparative analysis of Land-Atmosphere average coupling strength using different soil moisture from SMAPL3, SMAPL4_L3, and SMAPL4 data across various region and globally.

5. Discussion

An important aspect of this analysis is the development of the merged CTP and HI product. The merging process employs the triple collocation method based on the relative errors among the MERRA2, CFSR, and ERA5 datasets. The variations in the weight distributions for CTP and the HI reflect the inherent differences in the datasets. In the merged dataset, the ERA5 reanalysis had the highest weight which may be a result of improved representations of tropospheric temperature and humidity (Hersbach et al. 2020). Nonetheless, it is critical to understand that a higher weight for ERA5 does not mean that the merged product will closely resemble ERA5 in its characteristics. The merging process involves integrating information from multiple datasets, and the resulting merged product is a distinct and independent entity. Fig. 3 also reveals the presence of correlated errors when compared with radiosonde observations, suggesting the potential for a biased merged product. However, the merged product outperformed the individual reanalysis datasets when compared to radiosondes and satellite-based estimates of CTP and HI (Figs 4 and 5). This suggests that the merging process reduces the bias arising from the individual reanalysis products and provides a more accurate product. While this validation of the merged product is robust, it is not without its flaws. The differences in spatial and temporal resolutions between the merged product and the IGRA2 radiosonde and AIRSv7 observations is prone to uncertainty. Despite this, the merged dataset demonstrates a more accurate reflection of in-situ and satellite observations of CTP and HI compared to individual datasets, thus providing a temporal and spatially consistent dataset for analysing L-A coupling.

The merged CTP-HI product was used to investigate how soil moisture from SMAP contributes to quantifying L-A coupling strength globally. As depicted in Figs. 6 through 9, the findings underscore the pivotal role of soil moisture in the representation of L-A coupling strength within the CTP-HI framework developed by Roundy et al. (2013). The coupling strength is quantified as the persistence in the wet and dry coupling regimes defined using a three state Markov Chain model. Figs 6 and 8, demonstrate the ability of soil moisture to predict future CTP and HI through lag-correlation analysis and its direct relationship with coupling strength. The observed lag correlation is most pronounced for the SM-HI relation, indicating the direct influence of soil moisture on lower tropospheric humidity. In contrast, there is a weaker relationship between soil moisture and atmospheric stability as indicated by the SM-CTP lag-correlation. Reinforcing this point, Entekhabi et al. (1996) discussed the significant feedback mechanisms between soil moisture and atmospheric processes, highlighting how changes in soil moisture can directly impact the atmospheric environment. Wang et al. (2024) further substantiate this argument, which demonstrates the potential for soil moisture to predict the future coupling state and the utility of the persistence in the coupling regime as a measure of coupling strength.

The SMAPL4 dataset exhibits higher coupling strength in both the wet and dry regime across the globe (Fig. 9). This indicates that SMAPL4 would give stronger short-term predictions of future coupling states using the Coupling Stochastic Model (CSM) developed by Roundy and Wood, (2015), compared to the SMAPL3 product. However, it is unclear if the stronger coupling

- Deleted: [76]
- Deleted: over the various regions across the Globe.
- Deleted: r
- Deleted: linked to
- Deleted: '
- Deleted: behaves
- Deleted: as
- Deleted:
- Deleted: As such,
- Deleted: The goal of this analysis is to investigate
- Deleted: t
- Deleted: The goal of this analysis is to explore the role of (... [77])
- Deleted: our
- Deleted: using within
- Deleted: Roundy et al. (2013)
- Deleted: ,
- Deleted: process
- Deleted: as quantified in this study by the persistence in th (... [78])
- Deleted: shown
- Deleted: be related to
- Deleted: out to 10 days (Figs. 7 and 8)
- Deleted:
- Deleted: soil moisture
- Deleted: ship
- Deleted: and consistent
- Deleted: -level humidity in the lower atmosphere
- Deleted: across time
- Deleted: to
- Deleted: the more variable effects on influence
- Deleted: , this suggests that soil moisture has a more (... [79])
- Deleted: This relationship underscores the critical role of (... [80])
- Deleted: persiste
- Deleted: nce
- Deleted: The research conducted by Dong and Crow, (20 (... [81])
- Deleted: Furthermore, the
- Deleted: main
- Deleted: al regions
- Deleted:
- Deleted: the
- Deleted: has a greater potential to provide

predictions would lead to more skillful predictions and this topic is left to future research. One possible reason for the weaker coupling strength in SMAPL3 is that all available SMAPL3 data points were used to maintain a larger dataset without considering the quality flags. Dong and Crow (2019) highlights the importance of high signal-to-noise ratios in L-band SMAP soil moisture data for accurately quantifying L-A coupling strength. However, reducing the SMAPL3 sample size by filtering out data based on quality flags may introduce additional variability and noise, potentially impacting the coupling strength, especially in regions with variable data quality. This could potentially be teased out with a longer SMAP data record that would allow greater scrutiny of the retrievals without compromising the sample size.

The higher coupling strength observed in SMAPL4 is attributed to the reliance on a land surface model and assimilation process. The constraints of a deterministic model based on fixed equations, makes a land surface model less susceptible to variation and random noise and thus creates a higher autocorrelation compared to SMAPL3. While this is a unique characteristic of SMAPL4, it is important to remember that the strong coupling may not accurately mirror the complexities of real-world conditions (Van Vuuren et al., 2012). The accuracy of the stronger coupling in SMAPL4 is difficult to quantify due to the scarcity of in-situ observations across the globe where simultaneous atmospheric profile and soil moisture measurements can be obtained. The limitation of comprehensive ground-based observations poses a significant challenge in validating the true representation of coupling and understanding the intricate interplay between soil moisture and atmospheric conditions (Santanello et al. 2018; Beamesderfer et al. 2022). This conclusion also aligns with the insight provided by Findell et al. (2023), who emphasized the importance of high-frequency data for accurately assessing land-atmosphere coupling in climate models. Consequently, our ability to ascertain which dataset offers a more accurate representation of L-A coupling remains a subject of ongoing investigation.

Fig. 7c shows that differences between SMAPL3 and SMAPL4 coupling strength is mainly due to the shape of the distribution of soil moisture and its projection in the CTP-HI space. A minor difference in soil moisture distribution is observed in Fig. 7c when comparing SMAPL4 and SMAPL4 L3, though this difference is more notable in the joint probability space of CTP-HI-SM (Fig. 7a and b). Soil moisture estimates from SMAPL3 tend to skew towards drier values, likely due to the retrieval of the topsoil layer, which tends to dry quickly after rainfall events. This skew is influenced by fixed conditions in the retrieval process such as the prescribed freeze/thaw condition, as well as land surface characteristics like vegetation cover and soil properties. In contrast, SMAPL4, which uses observations from the same satellite, employs model-based soil hydraulic parameters, providing a greater range in the depiction of soil moisture variations across different landscapes. Studies by Reichle et al. (2017), Reichle et al. (2019), Reichle et al. (2021) have shown a reduced bias and expand dynamic range of surface soil moisture, in SMAPL4 as compared to in-situ and previous version of SMAPL4. These differences highlight the varying methodologies and characteristics of SMAPL3 and SMAPL4, resulting in distinct soil moisture estimates.

Recent research underscores the vital role of the SMAPL3 soil moisture product in agricultural applications, as demonstrated by Zhu et al. (2024). The accuracy of soil moisture measurement is crucial, and Tavakol et al. (2019) highlighted the SMAPL3 and SMAPL4 soil moisture products are at the forefront of soil moisture accuracy. For instance, Xu (2020) has concluded that SMAPL4 surface soil moisture product is more accurate, with lower errors (ubRMSE < 0.04 m³/m³), compared to the SMAPL3

- Deleted: is
- Deleted: Since the CSM model the probabilistic and stoch (... [82])
- Deleted: This indicates that the SMAPL4 has a stronger L (... [83])
- Deleted: However, the...ne possible reason for the weak (... [84])
- Deleted: The research conducted by ... ADDIN ZOTER (... [85])
- Deleted: this...educing the SMAPL3 sample size by filter (... [86])
- Deleted: during the dry coupling regimes ...n SMAPL4 is (... [87])
- Deleted: the increased temporal frequency used in the ana (... [88])
- Deleted: the
- Deleted: the
- Deleted: model itself and the associated
- Deleted: through the
- Deleted: SMAPL4 soil moisture data is partly derived from
- Deleted: Given the...he constraints of a deterministic mo (... [89])
- Deleted: ... land surface model, making it...less suscepti (... [90])
- Deleted: -
- Deleted: , which relies more directly on satellite observat (... [91])
- Deleted: s...of SMAPL4's assimilation process (... [92])
- Deleted: .
- Moved up [10]: SMAPL4 soil moisture data is partly derived
- Deleted: , which allows for a more detailed representation (... [93])
- Deleted: In contrast, the higher coupling strength observe (... [94])
- Deleted: However, it is important to consider that the (... [95])
- Moved (insertion) [11]
- Deleted: also ...bserved in Fig. 7c when comparing SMA (... [96])
- Deleted: In synthesizing the comparison between SMAPL (... [97])
- Deleted: s...to skew towards drier values, likely due to th (... [98])
- Deleted: and water phase in the retrieval process
- Deleted: used in the algorithm
- Deleted: also ...ses observations from the same satellite, (... [99])
- Deleted: dynamic approach by incorporating ...odel-bas (... [100])
- Deleted: Conversely, SMAPL4 offers a more detailed (... [101])
- Deleted: Also
- Deleted: Notably
- Deleted: , s
- Deleted: to
- Deleted:
- Deleted: effectively ...duced bias and expanded...the (... [102])
- Deleted: estimates
- Moved up [11]: A minor difference in soil moisture distribution
- Deleted: data assimilation techniques between
- Deleted: While acknowledging the presence of uncertain (... [103])

product ($\sim 0.06 \text{ m}^3/\text{m}^3$). Moreover, SMAP outperforms other soil moisture products, such as AMSR2 L3 and SMOS-IC, across most of the global land surface (Kim et al., 2021). This enhanced accuracy has been corroborated by Zhang et al. (2017), which showed that SMAPL4 captures spatial and temporal soil moisture variations more reliably as compared with Advanced Microwave Scanning Radiometer (AMSR2) across the United States. Reichle et al., (2017) have shown that version 4 SMAPL4's bias is significantly reduced as compared to version 3, exhibiting a 46% decrease in surface soil moisture uncertainty. Considering the above points, SMAP observation could improve drought monitoring and provide detailed insight of drought conditions, as demonstrated in studies by Mishra et al. (2017), Velpuri et al. (2016), Mladenova et al.(2020). This work further demonstrates the particular relevance of SMAP observations for studies of drought and L-A interaction, providing deeper insights into how land surface conditions influence atmospheric responses (Zhang et al., 2023b) and how it could potentially improve drought monitoring and prediction using statistical modelling (Roundy and Wood, 2015).

Conclusion

This work developed a merged reanalysis-based product for CTP and HI, which outperforms individual reanalysis products when validated against radiosonde and satellite observations. The merged CTP and HI product was used in combination with two different SMAP soil moisture products to analyse the coupling strength across the globe. Coupling strength is defined by the persistence probability for wet and dry coupling regimes as given by a first order three state Markov Chain model and is directly related to the ability of soil moisture to predict future atmospheric states out to 10 days. It is demonstrated that the measure of coupling strength is primarily driven by the SM-HI relationship, suggesting a direct influence of soil moisture on lower tropospheric humidity over time. In contrast, the SM-CTP relationship is more complex and likely influenced by larger-scale atmospheric conditions.

Another key conclusion is that SMAPL4 consistently presents a stronger representation of coupling strength compared to SMAPL3. While some of the difference in coupling strength can be attributed to the limited sample size of SMAPL3, analysing similar samples from both SMAP products still demonstrates stronger persistence in the wet coupling regimes in SMAPL4. The increased coupling strength in SMAPL4 may result from SMAPL4's reliance on a land surface model, which reduces susceptibility to random noise compared to SMAPL3. The difference in coupling strength in the two soil moisture datasets using the same CTP-HI underscores the importance of soil moisture data in estimating coupling strength using the CTP-HI framework. Nevertheless, it is not clear which of the two SMAP measures of coupling strength provide a better representation of the "real world coupling" as enhanced observation networks (i.e., co-located ground and atmospheric profile measurements) are needed to assess the accuracy of L-A interactions on a global scale.

Future work will investigate how the differences and similarities in the SMAP coupling strength may influence the ability to monitor and forecast the initiation, intensification, and abatement of drought conditions, using methods such as the L-A coupling based drought index (Roundy et al., 2014) and coupling stochastic model (Roundy and Wood, 2015). Such work may

Deleted: For instance, Xu (2020) has shown that SMAPL4's bias is significantly reduced, exhibiting a 46% decrease in surface soil moisture uncertainty. This enhanced accuracy has been corroborated by Zhang et al. (2017), which showed that SMAPL4 captu... [104]

Deleted:

Deleted: While,

Deleted: Importantly,

Deleted: e...SMAP observations for studies of drought a... [105]

Deleted: offering improved drought monitoring and wea... [106]

Deleted: However, it's crucial to recognize SMAPL4's p... [107]

Deleted: Another important aspect of this analysis is the... [108]

Deleted: ¶

Deleted: The development of...a merged reanalysis-base... [109]

Deleted: metric

Deleted: with

Deleted: datasets

Deleted: for ...o analysing [110]

Deleted: over

Deleted: the ...et and dry coupling regimes defined...s [111]

Deleted: his

Deleted: ¶ [112]

Deleted: -level

Deleted: This enhanced temporal consistency is evidenc... [113]

Deleted: the assimilation techniquesprocess and its

Deleted: ,

Deleted: The results indicate that SMAPL4 demonstrat... [114]

Deleted: se...insights ...n the two soil moisture datasets... [115]

Deleted: on

Deleted: the classification of L-A coupling and its subse... [116]

Deleted:

Deleted: ¶ [117]

Deleted: due...s enhanced observation networks (i.e., co... [118]

Deleted: Looking ahead, more research is needed into... [119]

Deleted: (Roundy et al., 2013)...Roundy et al., 2014) ... [120]

Deleted: to forecast short-term drought

Deleted: (Roundy et al., 2013).

Deleted: This drought index is anticipated to be instrum... [121]

Deleted: Moreover, a comparative analysis between of... [122]

Deleted: This research has the

Deleted: Thus

lead to potential improvements to drought monitoring and forecasting, strengthening the capacity for effective drought preparedness and response.

Data Availability

All data mentioned in the dataset section of this paper are publicly available on their respective websites. In addition, the merged dataset for CTP and HI can be accessed via the following: Makhasana, P., J. Roundy, J. A. Santanello, P. M. Lawston-Parker (2024). Triple Collocation based Merged Dataset for Convective Triggering Potential (CTP) and Humidity Index (HI), HydroShare, <https://doi.org/10.4211/hs.90bf9b575b684c849e617f620c2d63fb>

Author Contributions

Payal R. Makhasana led the development of the Triple Collocation based merging dataset, conducted the analysis on coupling strength, and primarily responsible for writing the original draft of the manuscript. Joseph A. Santanello and Patricia M. Lawston-Parker contributed to the refinement of the study design, provided critical reviews, and edits to the manuscript, enhancing its intellectual content. Joshua K. Roundy developed the code for revised CTP-HI framework for coupling classification, and participated in manuscript editing. Joseph A. Santanello, Patricia M. Lawston-Parker, and Joshua K. Roundy, played a significant role in shaping the research direction and methodology, ensuring the rigor and accuracy of the work presented.

Competing interests

The authors have no competing interests to declare.

Acknowledgment

We acknowledge the generous support and funding the NASA Soil Moisture Active-Passive Mission Science Team Program provided under grant NNH19ZDA001N-SMAP.

Reference

AIRS Project: Aqua/AIRS L3 Monthly Support Product (AIRS+AMSU) 1 degree x 1 degree V7.0, <https://doi.org/10.5067/GZT3YJIL1HUV>, 2020.

Alessi, M. J., Herrera, D. A., Evans, C. P., DeGaetano, A. T., and Ault, T. R.: Soil Moisture Conditions Determine Land-Atmosphere Coupling and Drought Risk in the Northeastern United States, JGR Atmospheres, 127, e2021JD034740, <https://doi.org/10.1029/2021JD034740>, 2022.

Deleted: ,

Deleted: ing

Deleted: to

Deleted: and thereby enhancing our understanding of drought dynamics using L-A interaction

Deleted: and

Deleted: was

Deleted: -based

Deleted: code

- Alexander, G. A., Holmes, H. A., Sun, X., Caputi, D., Faloona, I. C., and Oldroyd, H. J.: Simulating land-atmosphere coupling in the Central Valley, California: Investigating soil moisture impacts on boundary layer properties, *Agricultural and Forest Meteorology*, 317, 108898, <https://doi.org/10.1016/j.agrformet.2022.108898>, 2022.
- 2735 Anderson, W. B., Zaitchik, B. F., Hain, C. R., Anderson, M. C., Yilmaz, M. T., Mecikalski, J., and Schultz, L.: Towards an integrated soil moisture drought monitor for East Africa, *Hydrol. Earth Syst. Sci.*, 16, 2893–2913, <https://doi.org/10.5194/hess-16-2893-2012>, 2012.
- Aoying Zhou, Zhiyuan Cai, Li Wei, and Weining Qian: M-kernel merging: towards density estimation over data streams, in: Eighth International Conference on Database Systems for Advanced Applications, 2003. (DASFAA 2003). Proceedings., Proceedings Eighth International Conference on Database Systems for Advanced Applications (DASFAA 2003), Kyoto, Japan, 285–292, <https://doi.org/10.1109/DASFAA.2003.1192393>, 2003.
- 2740 Arshad, M., Ma, X., Yin, J., Ullah, W., Liu, M., and Ullah, I.: Performance evaluation of ERA-5, JRA-55, MERRA-2, and CFS-2 reanalysis datasets, over diverse climate regions of Pakistan, *Weather and Climate Extremes*, 33, 100373, <https://doi.org/10.1016/j.wace.2021.100373>, 2021.
- Beamesderfer, E. R., Buechner, C., Faiola, C., Helbig, M., Sanchez-Mejia, Z. M., Yáñez-Serrano, A. M., Zhang, Y., and Richardson, A. D.: Advancing Cross-Disciplinary Understanding of Land-Atmosphere Interactions, *JGR Biogeosciences*, 127, e2021JG006707, <https://doi.org/10.1029/2021JG006707>, 2022.
- Bell, B., Hersbach, H., Simmons, A., Berrisford, P., Dahlgren, P., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Radu, R., Schepers, D., Soci, C., Villaume, S., Bidlot, J., Haimberger, L., Woollen, J., Buontempo, C., and Thépaut, J.: The ERA5 global reanalysis: Preliminary extension to 1950, *Quart J Royal Meteor Soc*, 147, 4186–4227, <https://doi.org/10.1002/qj.4174>, 2021.
- 2750 Bennet, M. J., Kingston, D. G., and Cullen, N. J.: Extreme Compound and Seesaw Hydrometeorological Events in New Zealand: An Initial Assessment, *JGR Atmospheres*, 128, e2022JD038346, <https://doi.org/10.1029/2022JD038346>, 2023.
- Bocquet, M., Brajard, J., Carrassi, A., Bertino, L., ,CEREA, joint laboratory École des Ponts ParisTech and EDF R & D, Université Paris-Est, Champs-sur-Marne, France, ,Nansen Environmental and Remote Sensing Center, Bergen, Norway, and Sorbonne University, CNRS-IRD-MNHN, LOCEAN, Paris, France, ,Departement of Meteorology, University of Reading and NCEO, United-Kingdom, and Mathematical Institute, Utrecht University, The Netherlands, and ,Nansen Environmental and Remote Sensing Center, Bergen, Norway: Bayesian inference of chaotic dynamics by merging data assimilation, machine learning and expectation-maximization, *Foundations of Data Science*, 2, 55–80, <https://doi.org/10.3934/fods.2020004>, 2020.
- 2755 Chen, C., He, M., Chen, Q., Zhang, J., Li, Z., Wang, Z., and Duan, Z.: Triple collocation-based error estimation and data fusion of global gridded precipitation products over the Yangtze River basin, *Journal of Hydrology*, 605, 127307, <https://doi.org/10.1016/j.jhydrol.2021.127307>, 2022.
- 2760 Dirmeyer, P. A., Halder, S., and Bombardi, R.: On the Harvest of Predictability From Land States in a Global Forecast Model, *J. Geophys. Res. Atmos.*, 123, <https://doi.org/10.1029/2018JD029103>, 2018.
- Dong, J. and Crow, W. T.: L-band remote-sensing increases sampled levels of global soil moisture-air temperature coupling strength, *Remote Sensing of Environment*, 220, 51–58, <https://doi.org/10.1016/j.rse.2018.10.024>, 2019.
- 2765 Dong, X., Wang, Y., Hou, S., Ding, M., Yin, B., and Zhang, Y.: Robustness of the Recent Global Atmospheric Reanalyses for Antarctic Near-Surface Wind Speed Climatology, *Journal of Climate*, 33, 4027–4043, <https://doi.org/10.1175/JCLI-D-19-0648.1>, 2020.

- Durre, I. and Yin, X.: Enhanced Radiosonde Data For Studies of Vertical Structure, *Bull. Amer. Meteor. Soc.*, 89, 1257–1262, <https://doi.org/10.1175/2008BAMS2603.1>, 2008.
- 2770 Durre, I., Yin, X., Vose, R. S., Applequist, S., Arnfield, J., Korzeniewski, B., and Hundermark, B.: Integrated Global Radiosonde Archive (IGRA), Version 2, <https://doi.org/10.7289/V5X63K0Q>, 2016.
- Entekhabi, D., Rodriguez-Iturbe, I., and Castelli, F.: Mutual interaction of soil moisture state and atmospheric processes, *Journal of Hydrology*, 184, 3–17, [https://doi.org/10.1016/0022-1694\(95\)02965-6](https://doi.org/10.1016/0022-1694(95)02965-6), 1996.
- Entekhabi, D., Das, N., Njoku, E., Johnson, J., and Shi, J.: SMAP L3 Radar/Radiometer Global Daily 9 km EASE-Grid Soil Moisture, Version 3, <https://doi.org/10.5067/7KKNQ5UURM2W>, 2016.
- Feng, F. and Wang, K.: Merging Satellite Retrievals and Reanalyses to Produce Global Long-Term and Consistent Surface Incident Solar Radiation Datasets, *Remote Sensing*, 10, 115, <https://doi.org/10.3390/rs10010115>, 2018.
- Ferguson, C. R. and Wood, E. F.: Observed Land–Atmosphere Coupling from Satellite Remote Sensing and Reanalysis, *Journal of Hydrometeorology*, 12, 1221–1254, <https://doi.org/10.1175/2011JHM1380.1>, 2011.
- 2780 Findell, K. L. and Eltahir, E. A. B.: Atmospheric Controls on Soil Moisture–Boundary Layer Interactions. Part II: Feedbacks within the Continental United States, *J. Hydrometeorol.*, 4, 570–583, [https://doi.org/10.1175/1525-7541\(2003\)004<0570:ACOSML>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)004<0570:ACOSML>2.0.CO;2), 2003.
- Findell, K. L., Yin, Z., Seo, E., Dirmeyer, P. A., Arnold, N. P., Chaney, N., Fowler, M. D., Huang, M., Lawrence, D. M., Ma, P.-L., and Santanello Jr., J. A.: Accurate Assessment of Land–Atmosphere Coupling in Climate Models Requires High Frequency Data Output, *Climate and Earth system modeling*, <https://doi.org/10.5194/egusphere-2023-2048>, 2023.
- 2785 Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., Da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 2790 Gruber, A., Dorigo, W. A., Crow, W., and Wagner, W.: Triple Collocation-Based Merging of Satellite Soil Moisture Retrievals, *IEEE Trans. Geosci. Remote Sensing*, 55, 6780–6792, <https://doi.org/10.1109/TGRS.2017.2734070>, 2017.
- Gruber, A., De Lannoy, G., Albergel, C., Al-Yaari, A., Brocca, L., Calvet, J.-C., Colliander, A., Cosh, M., Crow, W., Dorigo, W., Draper, C., Hirschi, M., Kerr, Y., Konings, A., Lahoz, W., McColl, K., Montzka, C., Muñoz-Sabater, J., Peng, J., Reichle, R., Richaume, P., Rüdiger, C., Scanlon, T., Van Der Schalie, R., Wigneron, J.-P., and Wagner, W.: Validation practices for satellite soil moisture retrievals: What are (the) errors?, *Remote Sensing of Environment*, 244, 111806, <https://doi.org/10.1016/j.rse.2020.111806>, 2020.
- 2795 Hassler, B. and Lauer, A.: Comparison of Reanalysis and Observational Precipitation Datasets Including ERA5 and WFDE5, *Atmosphere*, 12, 1462, <https://doi.org/10.3390/atmos12111462>, 2021.
- 2800 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Q.J.R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 2805

- Hsu, H. and Dirmeyer, P. A.: Soil moisture–evaporation coupling shifts into new gears under increasing CO₂, *Nat Commun*, 14, 1162, <https://doi.org/10.1038/s41467-023-36794-5>, 2023.
- 2810 Huang, H., Liang, Z., Li, B., Wang, D., Hu, Y., and Li, Y.: Combination of Multiple Data-Driven Models for Long-Term Monthly Runoff Predictions Based on Bayesian Model Averaging, *Water Resour Manage*, 33, 3321–3338, <https://doi.org/10.1007/s11269-019-02305-9>, 2019.
- Jach, L., Schwitalla, T., Branch, O., Warrach-Sagi, K., and Wulfmeyer, V.: Sensitivity of land–atmosphere coupling strength to changing atmospheric temperature and moisture over Europe, *Earth Syst. Dynam.*, 13, 109–132, <https://doi.org/10.5194/esd-13-109-2022>, 2022.
- 2815 Kim, S., Dong, J., and Sharma, A.: A Triple Collocation-Based Comparison of Three L-Band Soil Moisture Datasets, SMAP, SMOS-IC, and SMOS, Over Varied Climates and Land Covers, *Front. Water*, 3, 693172, <https://doi.org/10.3389/frwa.2021.693172>, 2021.
- Kozubek, M., Krizan, P., and Lastovicka, J.: Homogeneity of the Temperature Data Series from ERA5 and MERRA2 and Temperature Trends, *Atmosphere*, 11, 235, <https://doi.org/10.3390/atmos11030235>, 2020.
- 2820 Liu, Y., Yao, L., Jing, W., Di, L., Yang, J., and Li, Y.: Comparison of two satellite-based soil moisture reconstruction algorithms: A case study in the state of Oklahoma, USA, *Journal of Hydrology*, 590, 125406, <https://doi.org/10.1016/j.jhydrol.2020.125406>, 2020.
- Lorenzo, A. T., Morzfeld, M., Holmgren, W. F., and Cronin, A. D.: Optimal interpolation of satellite and ground data for irradiance nowcasting at city scales, *Solar Energy*, 144, 466–474, <https://doi.org/10.1016/j.solener.2017.01.038>, 2017.
- 2825 Lu, J., Wang, G., Chen, T., Li, S., Hagan, D. F. T., Kattel, G., Peng, J., Jiang, T., and Su, B.: A Harmonized Global Land Evaporation Dataset from Reanalysis Products Covering 1980–2017, *Data, Algorithms, and Models*, <https://doi.org/10.5194/essd-2021-61>, 2021.
- Lyu, F., Tang, G., Behrangi, A., Wang, T., Tan, X., Ma, Z., and Xiong, W.: Precipitation Merging Based on the Triple Collocation Method Across Mainland China, *IEEE Trans. Geosci. Remote Sensing*, 59, 3161–3176, <https://doi.org/10.1109/TGRS.2020.3008033>, 2021.
- 2830 Massoud, E. C., Lee, H., Gibson, P. B., Loikith, P., and Waliser, D. E.: Bayesian Model Averaging of Climate Model Projections Constrained by Precipitation Observations over the Contiguous United States, *Journal of Hydrometeorology*, 21, 2401–2418, <https://doi.org/10.1175/JHM-D-19-0258.1>, 2020.
- 2835 Miranda Espinosa, M. T., Giuliani, G., and Ray, N.: Reviewing the discoverability and accessibility to data and information products linked to Essential Climate Variables, *International Journal of Digital Earth*, 13, 236–252, <https://doi.org/10.1080/17538947.2019.1620882>, 2020.
- Mishra, A., Vu, T., Veettil, A. V., and Entekhabi, D.: Drought monitoring with soil moisture active passive (SMAP) measurements, *Journal of Hydrology*, 552, 620–632, <https://doi.org/10.1016/j.jhydrol.2017.07.033>, 2017.
- 2840 Mladenova, I. E., Bolten, J. D., Crow, W., Sazib, N., and Reynolds, C.: Agricultural Drought Monitoring via the Assimilation of SMAP Soil Moisture Retrievals Into a Global Soil Water Balance Model, *Front. Big Data*, 3, 10, <https://doi.org/10.3389/fdata.2020.00010>, 2020.
- Mukherjee, S. and Mishra, A. K.: Global Flash Drought Analysis: Uncertainties From Indicators and Datasets, *Earth’s Future*, 10, <https://doi.org/10.1029/2022EF002660>, 2022.

- 2845 Nguyen, G. V., Le, X.-H., Van, L. N., Jung, S., Yeon, M., and Lee, G.: Application of Random Forest Algorithm for Merging Multiple Satellite Precipitation Products across South Korea, *Remote Sensing*, 13, 4033, <https://doi.org/10.3390/rs13204033>, 2021.
- ONEill, Peggy E., Chan, Steven, Njoku, Eni G, Jackson, Tom, Bindlish, Rajat, Chaubell, M. Julian, and Colliander, Andreas: SMAP Enhanced L3 Radiometer Global and Polar Grid Daily 9 km EASE-Grid Soil Moisture, Version 5, <https://doi.org/10.5067/4DQ54OUI9DL>, 2021.
- 2850 Park, S., Son, S.-W., Jung, M.-I., Park, J., and Park, S. S.: Evaluation of tropospheric ozone reanalyses with independent ozonesonde observations in East Asia, *Geosci. Lett.*, 7, 12, <https://doi.org/10.1186/s40562-020-00161-9>, 2020.
- Pratola, C., Barrett, B., Gruber, A., and Dwyer, E.: Quality Assessment of the CCI ECV Soil Moisture Product Using ENVISAT ASAR Wide Swath Data over Spain, Ireland and Finland, *Remote Sensing*, 7, 15388–15423, <https://doi.org/10.3390/rs71115388>, 2015.
- 2855 Qi, Y., Chen, H., and Zhu, S.: Influence of Land–Atmosphere Coupling on Low Temperature Extremes Over Southern Eurasia, *JGR Atmospheres*, 128, e2022JD037252, <https://doi.org/10.1029/2022JD037252>, 2023.
- Qiu, J., Dong, J., Crow, W. T., Zhang, X., Reichle, R. H., and De Lannoy, G. J. M.: The benefit of brightness temperature assimilation for the SMAP Level-4 surface and root-zone soil moisture analysis, *Hydrol. Earth Syst. Sci.*, 25, 1569–1586, <https://doi.org/10.5194/hess-25-1569-2021>, 2021.
- 2860 Reichle, Rolf, Lannoy, Gabrielle De, Koster, Randal, Crow, Wade, and Kimball, John: SMAP L4 Global 3-hourly 9 km Surface and Rootzone Soil Moisture Geophysical Data, Version 3, <https://doi.org/10.5067/B59DT1D5UMB4>, 2017.
- Reichle, R. H., De Lannoy, G. J. M., Liu, Q., Ardizzone, J. V., Colliander, A., Conaty, A., Crow, W., Jackson, T. J., Jones, L. A., Kimball, J. S., Koster, R. D., Mahanama, S. P., Smith, E. B., Berg, A., Bircher, S., Bosch, D., Caldwell, T. G., Cosh, M., González-Zamora, Á., Holifield Collins, C. D., Jensen, K. H., Livingston, S., Lopez-Baeza, E., Martínez-Fernández, J., McNairn, H., Moghaddam, M., Pacheco, A., Pellarin, T., Prueger, J., Rowlandson, T., Seyfried, M., Starks, P., Su, Z., Thibeault, M., Van Der Velde, R., Walker, J., Wu, X., and Zeng, Y.: Assessment of the SMAP Level-4 Surface and Root-Zone Soil Moisture Product Using In Situ Measurements, *Journal of Hydrometeorology*, 18, 2621–2645, <https://doi.org/10.1175/JHM-D-17-0063.1>, 2017.
- 2865 Reichle, R. H., Liu, Q., Koster, R. D., Crow, W. T., De Lannoy, G. J. M., Kimball, J. S., Ardizzone, J. V., Bosch, D., Colliander, A., Cosh, M., Kolassa, J., Mahanama, S. P., Prueger, J., Starks, P., and Walker, J. P.: Version 4 of the SMAP Level-4 Soil Moisture Algorithm and Data Product, *J Adv Model Earth Syst*, 11, 3106–3130, <https://doi.org/10.1029/2019MS001729>, 2019.
- Reichle, R. H., Liu, Q., Ardizzone, J. V., Crow, W. T., De Lannoy, G. J. M., Dong, J., Kimball, J. S., and Koster, R. D.: The Contributions of Gauge-Based Precipitation and SMAP Brightness Temperature Observations to the Skill of the SMAP Level-4 Soil Moisture Product, *Journal of Hydrometeorology*, 22, 405–424, <https://doi.org/10.1175/JHM-D-20-0217.1>, 2021.
- 2875 Roundy, J. K. and Santanello, J. A.: Utility of Satellite Remote Sensing for Land–Atmosphere Coupling and Drought Metrics, *Journal of Hydrometeorology*, 18, 863–877, <https://doi.org/10.1175/JHM-D-16-0171.1>, 2017.
- Roundy, J. K. and Wood, E. F.: The Attribution of Land–Atmosphere Interactions on the Seasonal Predictability of Drought, *Journal of Hydrometeorology*, 16, 793–810, <https://doi.org/10.1175/JHM-D-14-0121.1>, 2015.
- 2880 Roundy, J. K., Ferguson, C. R., and Wood, E. F.: Temporal Variability of Land–Atmosphere Coupling and Its Implications for Drought over the Southeast United States, *Journal of Hydrometeorology*, 14, 622–635, <https://doi.org/10.1175/JHM-D-12-090.1>, 2013.

- Roundy, J. K., Ferguson, C. R., and Wood, E. F.: Impact of land-atmospheric coupling in CFSv2 on drought prediction, *Clim Dyn*, 43, 421–434, <https://doi.org/10.1007/s00382-013-1982-7>, 2014.
- Saha, K., Dash, P., Zhao, X., and Zhang, H.: Error Estimation of Pathfinder Version 5.3 Level-3C SST Using Extended Triple Collocation Analysis, *Remote Sensing*, 12, 590, <https://doi.org/10.3390/rs12040590>, 2020.
- 2885 Saha, S., Moorthi, S., Pan, H.-L., Wu, X., Wang, J., Nadiga, S., Tripp, P., Kistler, R., Woollen, J., Behringer, D., Liu, H., Stokes, D., Grumbine, R., Gayno, G., Wang, J., Hou, Y.-T., Chuang, H., Juang, H.-M. H., Sela, J., Iredell, M., Treadon, R., Kleist, D., Van Delst, P., Keyser, D., Derber, J., Ek, M., Meng, J., Wei, H., Yang, R., Lord, S., Van Den Dool, H., Kumar, A., Wang, W., Long, C., Chelliah, M., Xue, Y., Huang, B., Schemm, J.-K., Ebisuzaki, W., Lin, R., Xie, P., Chen, M., Zhou, S., Higgins, W., Zou, C.-Z., Liu, Q., Chen, Y., Han, Y., Cucurull, L., Reynolds, R. W., Rutledge, G., and Goldberg, M.: The NCEP Climate Forecast System Reanalysis, *Bull. Amer. Meteor. Soc.*, 91, 1015–1058, <https://doi.org/10.1175/2010BAMS3001.1>, 2010.
- 2890 Saini, R., Wang, G., and Pal, J. S.: Role of Soil Moisture Feedback in the Development of Extreme Summer Drought and Flood in the United States, *Journal of Hydrometeorology*, 17, 2191–2207, <https://doi.org/10.1175/JHM-D-15-0168.1>, 2016.
- Santanello, J. A., Roundy, J., and Dirmeyer, P. A.: Quantifying the Land–Atmosphere Coupling Behavior in Modern Reanalysis Products over the U.S. Southern Great Plains, *Journal of Climate*, 28, 5813–5829, <https://doi.org/10.1175/JCLI-D-14-00680.1>, 2015.
- Santanello, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., Ek, M., Gentine, P., Guillod, B. P., Van Heerwaarden, C., Roundy, J., and Wulfmeyer, V.: Land–Atmosphere Interactions: The LoCo Perspective, *Bulletin of the American Meteorological Society*, 99, 1253–1272, <https://doi.org/10.1175/BAMS-D-17-0001.1>, 2018.
- 2900 Schwitalla, T., Jach, L., Wulfmeyer, V., and Warrach-Sagi, K.: Soil moisture-atmosphere coupling strength over Central Europe in the recent warming climate, *Atmospheric, Meteorological and Climatological Hazards*, <https://doi.org/10.5194/egusphere-2023-1725>, 2023.
- Seneviratne, S. I. and Stöckli, R.: The Role of Land-Atmosphere Interactions for Climate Variability in Europe, in: *Climate Variability and Extremes during the Past 100 Years*, edited by: Brönnimann, S., Luterbacher, J., Ewen, T., Diaz, H. F., Stolarski, R. S., and Neu, U., Springer Netherlands, Dordrecht, 179–193, https://doi.org/10.1007/978-1-4020-6766-2_12, 2008.
- 2905 Seo, E. and Dirmeyer, P. A.: Understanding the diurnal cycle of land–atmosphere interactions from flux site observations, *Hydrol. Earth Syst. Sci.*, 26, 5411–5429, <https://doi.org/10.5194/hess-26-5411-2022>, 2022.
- Seo, Y.-W. and Ha, K.-J.: Changes in land-atmosphere coupling increase compound drought and heatwaves over northern East Asia, *npj Clim Atmos Sci*, 5, 100, <https://doi.org/10.1038/s41612-022-00325-8>, 2022.
- 2910 Stoffelen, A.: Toward the true near-surface wind speed: Error modeling and calibration using triple collocation, *J. Geophys. Res.*, 103, 7755–7766, <https://doi.org/10.1029/97JC03180>, 1998.
- Sun, L. and Fu, Y.: A new merged dataset for analyzing clouds, precipitation and atmospheric parameters based on ERA5 reanalysis data and the measurements of the Tropical Rainfall Measuring Mission (TRMM) precipitation radar and visible and infrared scanner, *Earth Syst. Sci. Data*, 13, 2293–2306, <https://doi.org/10.5194/essd-13-2293-2021>, 2021.
- 2915 Tavakol, A., Rahmani, V., Quiring, S. M., and Kumar, S. V.: Evaluation analysis of NASA SMAP L3 and L4 and SPoRT-LIS soil moisture data in the United States, *Remote Sensing of Environment*, 229, 234–246, <https://doi.org/10.1016/j.rse.2019.05.006>, 2019.

- 2920 Teixeira, J., Chen, S., Clayton, C. A., Fridlind, A. M., Lebsock, M., McCarty, W., Salmun, H., Santanello, J. A., Turner, D. D., Wang, Z., and Zeng, X.: Toward a Global Planetary Boundary Layer Observing System: The NASA PBL Incubation Study Team Report, NASA PBL Incubation Study Team, 1–134, 2021.
- Velpuri, N. M., Senay, G. B., and Morissette, J. T.: Evaluating New SMAP Soil Moisture for Drought Monitoring in the Rangelands of the US High Plains, *Rangelands*, 38, 183–190, <https://doi.org/10.1016/j.rala.2016.06.002>, 2016.
- 2925 Wakefield, R. A., Basara, J. B., Furtado, J. C., Illston, B. G., Ferguson, Craig, R., and Klein, P. M.: A Modified Framework for Quantifying Land–Atmosphere Covariability during Hydrometeorological and Soil Wetness Extremes in Oklahoma, *Journal of Applied Meteorology and Climatology*, 58, 1465–1483, <https://doi.org/10.1175/JAMC-D-18-0230.1>, 2019.
- Wang, G., Fu, R., Zhuang, Y., Dirmeyer, P. A., Santanello, J. A., Wang, G., Yang, K., and McColl, K.: Influence of lower-tropospheric moisture on local soil moisture–precipitation feedback over the US Southern Great Plains, *Atmos. Chem. Phys.*, 24, 3857–3868, <https://doi.org/10.5194/acp-24-3857-2024>, 2024.
- 2930 Wilson, A. G. and Fronczyk, K. M.: Bayesian Reliability: Combining Information, *Quality Engineering*, 0–0, <https://doi.org/10.1080/08982112.2016.1211889>, 2016.
- Wu, X., Lu, G., Wu, Z., He, H., Scanlon, T., and Dorigo, W.: Triple Collocation-Based Assessment of Satellite Soil Moisture Products with In Situ Measurements in China: Understanding the Error Sources, *Remote Sensing*, 12, 2275, <https://doi.org/10.3390/rs12142275>, 2020.
- 2935 Xu, X.: Evaluation of SMAP Level 2, 3, and 4 Soil Moisture Datasets over the Great Lakes Region, *Remote Sensing*, 12, 3785, <https://doi.org/10.3390/rs12223785>, 2020.
- Yilmaz, M. T., Crow, W. T., Anderson, M. C., and Hain, C.: An objective methodology for merging satellite- and model-based soil moisture products: OBJECTIVELY MERGING SOIL MOISTURE PRODUCTS, *Water Resour. Res.*, 48, <https://doi.org/10.1029/2011WR011682>, 2012.
- 2940 Yingshan, W., Weijun, S., Lei, W., Yanzhao, L., Wentao, D., Jizu, C., and Xiang, Q.: How Do Different Reanalysis Radiation Datasets Perform in West Qilian Mountains?, *Front. Earth Sci.*, 10, 852054, <https://doi.org/10.3389/feart.2022.852054>, 2022.
- Yoon, D., Kang, T., Cha, D., Song, C., Lee, M., Min, K., Kim, J., Chun, J. A., and Seo, E.: Role of Land–Atmosphere Interaction in the 2016 Northeast Asia Heat Wave: Impact of Soil Moisture Initialization, *JGR Atmospheres*, 128, e2022JD037718, <https://doi.org/10.1029/2022JD037718>, 2023.
- 2945 Yuan, R., Li, F., and Ye, R.: Global diagnosis of land–atmosphere coupling based on water isotopes, *Sci Rep*, 13, 21319, <https://doi.org/10.1038/s41598-023-48694-1>, 2023.
- Zhang, L., Ding, M., Zheng, X., Chen, J., Guo, J., and Bian, L.: Assessment of AIRS Version 7 Temperature Profiles and Low-Level Inversions with GRUAN Radiosonde Observations in the Arctic, *Remote Sensing*, 15, 1270, <https://doi.org/10.3390/rs15051270>, 2023a.
- 2950 Zhang, L. N., Short Gianotti, D. J., and Entekhabi, D.: Land Surface Influence on Convective Available Potential Energy (CAPE) Change during Interstorms, *Journal of Hydrometeorology*, 24, 1365–1376, <https://doi.org/10.1175/JHM-D-22-0191.1>, 2023b.
- Zhang, S.-Q., Ren, G.-Y., Ren, Y.-Y., Zhang, Y.-X., and Xue, X.-Y.: Comprehensive evaluation of surface air temperature reanalysis over China against urbanization-bias-adjusted observations, *Advances in Climate Change Research*, 12, 783–794, <https://doi.org/10.1016/j.accre.2021.09.010>, 2021.

Zhang, X., Zhang, T., Zhou, P., Shao, Y., and Gao, S.: Validation Analysis of SMAP and AMSR2 Soil Moisture Products over the United States Using Ground-Based Measurements, *Remote Sensing*, 9, 104, <https://doi.org/10.3390/rs9020104>, 2017.

2960 Zhou, S., Williams, A. P., Berg, A. M., Cook, B. I., Zhang, Y., Hagemann, S., Lorenz, R., Seneviratne, S. I., and Gentile, P.: Land-atmosphere feedbacks exacerbate concurrent soil drought and atmospheric aridity, *Proc. Natl. Acad. Sci. U.S.A.*, 116, 18848–18853, <https://doi.org/10.1073/pnas.1904955116>, 2019.

Zhu, L., Tian, G., Wu, H., Ding, M., Zhu, A.-X., and Ma, T.: Regional Assessment of Soil Moisture Active Passive Enhanced L3 Soil Moisture Product and Its Application in Agriculture, *Remote Sensing*, 16, 1225, <https://doi.org/10.3390/rs16071225>, 2024.