

# Hydrological Controls on Temporal Contributions of Three Nested Forested Subcatchments to the Export of Dissolved Organic Carbon ~~DOC~~ Export

Katharina Blaurock<sup>1</sup>, Burkhard Beudert<sup>2</sup>, Luisa Hopp<sup>1</sup>

5 <sup>1</sup> Department of Hydrology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, 95447 Bayreuth, Germany

<sup>2</sup> Department of Nature Conservation and Research, Bavarian Forest National Park, 94481 Grafenau, Germany

*Correspondence to:* Luisa Hopp (luisa.hopp@uni-bayreuth.de)

**Abstract.** Assessing ~~DOC~~ the export of dissolved organic carbon (DOC) from terrestrial systems into inland waters reliably is of vital importance to understand the global carbon cycle in detail. In this context, hydrological connectivity within catchments is the prerequisite for transport processes from terrestrial systems into inland waters. Using high-frequency measurements of DOC concentrations via UV-Vis spectrometry, we quantified the DOC export at the outlets of three nested forested subcatchments within a 3.5 km<sup>2</sup> headwater catchment in the Bavarian Forest National Park, Germany, during a 12 month period. The subcatchments differ with respect to topography, elevation, vegetation and soil types. We observed a high flow-weighted DOC export from the entire headwater catchment during spring and autumn. In contrast, during snowmelt, summer and winter, DOC export was low due to either low runoff generated and/or relatively lower ~~the interplay between runoff generated and~~ DOC available for mobilization. Flow-weighted DOC export also varied between the three subcatchments. Flow-weighted DOC export was always higher in the lower, flat subcatchment than in the upper steep subcatchments, indicating a large DOC store that can be activated, whenever hydrological connectivity is established. This was particularly evident during autumn, when large precipitation events mobilized DOC which had accumulated during the dry summer period and was also delivered from fresh leaf litter of deciduous trees. Our data show the strong hydrological control on seasonal DOC export. However, the runoff-based contribution of subcatchments over time is modulated by the interplay of soils, vegetation, topography and microclimate, which can be seen as secondary controls. As hydrological connectivity varies with topography, the relative contribution of topographically different subcatchments varies seasonally. Since climate change is predicted to influence precipitation patterns, spatial and temporal DOC export patterns are likely to change depending on topography.

## 1 Introduction

Organic carbon in inland waters strongly influences the global carbon cycle due to the role it plays for carbon sequestration, transport and mineralization (Battin et al., 2009; Cole et al., 2007). A recent study estimated that the global carbon export from

30 terrestrial systems to inland waters amounts to 5.1 Pg C annually (Drake et al., 2018). Dissolved carbon represents a large part  
of the exported carbon and therefore influences the net ecosystem balance (Kindler et al., 2011). The majority of the exported  
dissolved carbon is inorganic (Chaplot and Mutema, 2021) and many uncertainties regarding the contribution of dissolved  
organic carbon (DOC) to carbon export from terrestrial systems remain. However, especially in catchments with wetter and  
35 cooler climates (due to latitude and/or elevation) where organic-rich soils have evolved, DOC is a major component of the  
exported carbon (Ciais et al., 2008). DOC plays an important role in the context of climate change. It can outgas to the  
atmosphere from water as the greenhouse gases CO<sub>2</sub> and CH<sub>4</sub> because of in-stream metabolic processes. The largest part of all  
organic and inorganic carbon compounds, which are transported from the terrestrial system to the aquatic system, is emitted  
to the atmosphere and not sequestered in the sediments of streams or oceans (Drake et al., 2018; Raymond et al., 2013). DOC  
is linked to nutrient cycling as it acts as electron donor in anaerobic respiration processes, e.g., denitrification (Lovley et al.,  
40 1999; Pang and Wang, 2021). In addition, DOC also affects drinking water quality (Sadiq and Rodriguez, 2004; Alarcon-  
Herrera et al., 1994; Ledesma et al., 2012) and the transport of pollutants (Doerr and Muennich, 1991; Hope et al., 1994;  
Ravichandran, 2004). Therefore, it is important to characterize the patterns and dynamics of DOC export as accurately as  
possible.

A meta-analysis of DOC export data found that DOC export from catchments worldwide ranged from 1.2 to over 50 000 kg C  
45 km<sup>-2</sup> yr<sup>-1</sup> and that it is negatively related to catchment area and positively to discharge (Alvarez-Cobelas et al., 2012). The main  
source of DOC in forested catchments is soil organic matter (Batjes, 2014; Borken et al., 2011; Kaiser and Kalbitz, 2012;  
Kalbitz et al., 2000), and it has been shown that a high proportion of wetlands in a catchment leads to a relatively higher DOC  
export because high groundwater levels support the accumulation of organic matter in soils due to limited mineralization under  
hypoxic conditions (Laudon et al., 2011; Zarnetske et al., 2018). Riparian zones are especially regarded to be of importance  
50 for DOC export due to the accumulation of soil organic matter and the often high groundwater levels that facilitate transport  
into the streams (Ledesma et al., 2018; Mei et al., 2014; Musolff et al., 2018; Ploum et al., 2020; Strohmeier et al., 2013). The  
largest amounts of DOC are commonly exported during precipitation events (Raymond et al., 2016; Raymond and Saiers,  
2010) and DOC concentrations generally increase with rising discharge (Hobbie and Likens, 1973; Meyer and Tate, 1983). In  
order to capture the DOC export dynamics precisely, it is necessary to measure DOC concentrations at a high temporal  
55 resolution, especially during events. Earlier DOC export estimates have mostly been based on weekly or bi-weekly grab  
samples leading to a large uncertainty in export numbers, especially in small catchments (Schleppi et al., 2006a; Schleppi et  
al., 2006b). By using novel techniques, high-frequency measurements over long periods lead to better estimates of DOC export  
(Ritson et al., 2022).

DOC export does not only vary with event characteristics but also during and between seasons as a result of varying  
60 temperature and wetness conditions (Blaurock et al., 2021; Wen et al., 2020; Werner et al., 2019). Although biogeochemical  
processes determine the production, quality, and availability of DOC within catchments, water flow processes at catchment  
scale eventually control the dynamics and patterns of mobilization and transport of DOC into streams and rivers. Here,  
hydrological connectivity, influenced by catchment wetness conditions (Detty and McGuire, 2010; McGuire and McDonnell,


2010; Penna et al., 2015), is of crucial importance for solute export (Covino, 2017; Kiewiet et al., 2020). However, patterns and dynamics of hydrological connectivity do not only depend on antecedent precipitation and event characteristics but are also controlled by catchment topography, e. g. shape and slope angle of hillslopes and riparian zones (Detty and McGuire, 2010; Tetzlaff et al., 2014), which, in turn, controls patterns of catchment wetness, i.e., where in a catchment moisture accumulates. Virtual experiments have suggested that the hillslope shape (straight vs. concave) influences the mobilization of nutrients (Weiler and McDonnell, 2006). Therefore, topography is a factor that can be expected to influence DOC mobilization from terrestrial to aquatic systems (Jankowski and Schindler, 2019).

In an earlier study [in a temperate forest headwater catchment](#), we showed that the contribution of topographically different subcatchments to event-based DOC export depended on the establishment of hydrological connectivity between the DOC source areas and the stream (Blaurock et al., 2021). The objective of this study was to improve our understanding of the temporal and spatial patterns of DOC export from a forested headwater catchment beyond the event timescale to assess the importance of seasons with differing hydrological conditions including low-flow periods for DOC export. We conducted high-frequency measurements of DOC concentrations over one year in the streams of three nested subcatchments to 1) quantify total DOC export from the entire study catchment in the course of 12 months and 2) explore temporal differences in DOC export among topographically distinct subcatchments.

## 2 Material and Methods

### 80 2.1 Study site

The study was conducted in the Hinterer Schachtenbach catchment ( $HS_{\text{tot}}$ , 3.5 km<sup>2</sup>), which is part of the Große Ohe catchment (19.2 km<sup>2</sup>) in the Bavarian Forest National Park (BFNP, 243 km<sup>2</sup>), Germany (Fig. 1).



85 **Figure 1.** The studied catchment Hinterer Schachtenbach (HS<sub>tot</sub>) is part of the Große Ohe catchment and located in southeast Germany (black dot). It includes the subcatchments Kaltenbrunner Seige (KS), Markungsraben (MG) and Hinterer Schachtenbach (HS<sub>sub</sub>). The high-frequency sampling sites were located at the outlets of the two subcatchments KS (yellow diamond) and MG (green diamond) and at the outlet of the entire catchment HS<sub>tot</sub> (black diamond). Precipitation was measured at the meteorological station RDH (red dot). Both the location of the stream network and the digital elevation model were provided by the Bavarian State Office for Environment. The map of Germany by Vemaps (2022).

90 Mean annual precipitation is 1323 mm and mean annual temperature is 6.5 °C (1990 – 2019). Measurements were conducted in three nested subcatchments of HS<sub>tot</sub> (Table 1): The streams of the upper subcatchments Markungsraben (MG, 1.1 km<sup>2</sup>) and Kaltenbrunner Seige (KS, 0.9 km<sup>2</sup>) join to form the stream of the lower subcatchment Hinterer Schachtenbach (HS<sub>sub</sub>, 1.5 km<sup>2</sup>). Elevation in the entire catchment ranges from 771 m to 1373 m. MG and KS represent the upper part of the catchment with incised streams, and HS<sub>sub</sub> represents the lower part of the catchment with a wide riparian zone. The geology of the

95 ~~Hinterer Schachtenbach~~ catchment (HS<sub>tot</sub>) is dominated by biotite granite and cordierite–sillimanite gneiss. KS and MG are dominated by Cambisols and Podzols with a larger proportion of Cambisols at KS. HS<sub>sub</sub> is dominated by Cambisols and hydromorphic soils. Almost 40 % of the area of MG is characterized by soils that are interspersed with rocks, mostly below the surface. Pleistocene solifluction processes created deeper soil layers in the lower catchment parts than in the upper catchment part. The entire catchment is almost entirely covered by forest. Dominant tree species are Norway spruce (*Picea abies* (L.) Karsten) and European beech (*Fagus sylvatica* L.). However, large parts of the catchment are in a stage of

100 rejuvenation of mostly Norway spruce due to bark beetle outbreaks in the mid-1990s and 2000s, especially in the subcatchment MG (Beudert et al., 2015). KS is characterized by a higher proportion of mature deciduous forest compared to MG.

**Table 1. Catchment characteristics of the entire catchment HS<sub>tot</sub> and the subcatchments KS, MG and HS<sub>sub</sub>.**

	HS <sub>tot</sub>	KS	MG	HS <sub>sub</sub>
Area (km <sup>2</sup> )	3.5	0.9	1.1	1.5
Area ratio of HS <sub>tot</sub> (%)	100	26	31	43
Elevation (m a.s.l.)	771 - 1373	877 - 1279	876 - 1373	771 - 1085
Mean slope (°)	12.0	14.5	15.8	7.4
Soils (%)	Cambisol	66	79	65
	Podzol	15	16	0
	Hydromorphic soil	18	5	35
	Other	1	0	0
Vegetation (%)	Rejuvenation	34	28	21
	Deciduous Forest	41	53	42
	Coniferous Forest	9	1	17
	Mixed Forest	15	17	19
	Other	1	0	3

105

## 2.2 Field measurements

### 2.2.1 Discharge measurements and meteorological data

During the study period from May 30<sup>th</sup> 2020 to May 29<sup>th</sup> 2021, the water level was measured at HS<sub>tot</sub> (Fig. 1) every 15 minutes using a pressure transducer (SEBA Hydrometrie GmbH & Co. KG, Kaufbeuren, Germany). Discharge was determined via tracer dilution (NaCl, TQ-S; Sommer Messtechnik, Koblach, Austria) at 36 occasions to establish a rating curve and generate 15-min discharge data. Due to a technical failure, no discharge data for HS<sub>tot</sub> was available from August 1st to September 3rd, 2020. This data gap was filled using the discharge data of a neighboring catchment within the Große Ohe Catchment of similar size (Vorderer Schachtenbach, 5.9 km<sup>2</sup>). Using the discharge values of one year prior to the data gap, a relationship between the discharge of the two catchments was established ( $R^2=0.94$ ) and calculated values were used to fill the data gap (Fig. S1).

For MG, the discharge data for the complete study period were taken from the database of the Bavarian State Office for Environment (2021) at a 15-min interval. The discharge at KS was measured via tracer dilution (TQ-S; Sommer Messtechnik, Koblach, Austria) at eight occasions between April and December 2021 and 15-min discharge values were derived using a relationship between the highly resolved discharge data of MG and the measured discharge at KS ( $R^2=0.95$ ; Fig. S2). Precipitation and temperature during the study period were measured close to the outlet of MG and KS (meteorological station RDH) and data were provided by the BFNP. Measuring resolution was one second and storage interval was 10 minutes. Precipitation was measured using a weighing precipitation gauge (Pluvio<sup>2</sup> S, OTT Hydromet, Kempten Germany). A ring heater keeps the orifice rim free from snow and ice. As no long-term temperature data was available for this station, temperature

115  
120

data from a meteorological station 3.8 km east of the catchment outlet (48°55.771' N, 13°27.890' E, 953 m a.s.l.; outside of the map section shown in Fig. 1) was used to calculate the mean annual temperature.

## 125 2.2.2 High-frequency measurements of DOC concentrations

DOC concentrations were measured in situ at the outlets of the three subcatchments during the study period using three UV-Vis (ultraviolet-visible) spectrophotometers (spectro::lyser, s::can GmbH, Vienna, Austria). Device 1 (D1) was installed at the outlet of HS<sub>tot</sub> over the whole study period, Device 2 (D2) started at the outlet of MG but had to be replaced by Device 3 (D3) at the beginning of July 2020 due to a technical failure (after lightning strike). D3 was initially installed at KS and was moved  
130 to MG following the technical failure of D2. After repair, D2 was installed at the outlet of KS in October 2020. This means that for MG, there is a gap in DOC concentration data for the period from June 23rd to July 8th, 2020, and for KS, there is a gap in DOC concentration data from July 9th to October 9th, 2020. The spectrometric devices recorded the absorption spectrum of stream water from 200 to 750 nm with a resolution of 2.5 nm every 15 min. DOC concentrations were quantified using the internal calibration based on the absorption values using the software ana::pro. In order to refine the internal calibration, the  
135 DOC concentrations measured by the three UV-Vis spectrophotometers were corrected using grab stream samples taken over the course of the study period at various discharge conditions ( $n_{D1} = 52$ ,  $n_{D2} = 22$ ,  $n_{D3} = 44$ ; Fig. S3-S5). Samples were filtered in the field using polyethersulfone membrane filters (0.45  $\mu\text{m}$ ) or polycarbonate track etched membrane filters (0.45  $\mu\text{m}$ ). All samples were stored until further analysis at 4°C. DOC concentrations of the grab samples were analyzed by thermo-catalytic oxidation (TOC-L analyzer; Shimadzu, Kyoto, Japan). For further analysis, the corrected values ( $R^2$  for D1 = 0.98,  $R^2$  for D2  
140 = 0.87 and  $R^2$  for D3 = 0.97) were used. As no drift of the DOC concentration could be identified in the measured signal, we decided against a correction for biofouling as done by Werner et al. (2019). However, the sensor optics were manually cleaned in the field every 2 weeks using cotton swabs.

## 2.2.2 Data analysis

For the evaluation of the temporal DOC dynamics during our study period, we defined “hydrological periods” based on changes  
145 in precipitation regime and discharge behavior. For this, mean daily precipitation for preceding 14 day intervals, frequency of events and changes in discharge (lag times and peak values), together with visual inspection of the time series of precipitation and discharge, were considered. Since these hydrological periods overlap with calendrical seasons to a large extent, we designate the hydrological periods by the commonly used names for calendrical seasons (Table 2). In addition, we also introduced the hydrological period “snowmelt” to account for the very different hydrological conditions with increased runoff  
150 generation. We prefer this definition of hydrological periods for evaluating DOC export over the usage of calendrical or meteorological seasons to avoid the arbitrary attribution of precipitation and runoff events. Spring started with the end of snowmelt and ended with the beginning of a dry period in July 2020, which was defined as summer. Summer ended with the end of a long dry period prior to the beginning of a rainy period, which was defined as autumn. Winter started after the last precipitation event that did not fall as snow and when the temperature fell below zero. Snowmelt was defined as the period

155 when temperatures rose above zero, a snow cover was present, and snowmelt was clearly visible in discharge data at HS<sub>tot</sub> as diurnal fluctuations. The snowmelt period ended when the catchments were completely free of snow at all elevations. The length of the hydrological periods is presented in days. However, these are rounded values as the start or end of a hydrological period did not automatically fall on the start or end of a day. For the calculation of mean daily DOC export, non-rounded values were used. All hydrological periods were defined using discharge data of HS<sub>tot</sub> and precipitation data from the meteorological station RDH. Runoff ratios were calculated as the ratio of cumulative area-normalized discharge to the amount of precipitation from RDH during each hydrological period. Data analysis was performed using MATLAB (The MathWorks, Inc.).

165 **Table 2. Hydrological periods and number of days with measured DOC data for HS<sub>tot</sub> and for all subcatchments. Due to the technical failure of device D2, there is no period with data for all three subcatchments available during summer.**

	Dates	Total number of days	Number of days with DOC data for HS <sub>tot</sub>	Number of days with DOC data for all subcatchments
Spring	30.05.2020 – 30.06.2020	60	60	52
	02.05.2021 – 29.05.2021			
Summer	01.07.2020 – 24.09.2020	87	83	0
Autumn	25.09.2020 – 21.11.2020	57	57	34
Winter	22.11.2020 – 21.12.2020	60	60	58
	30.12.2020 – 28.01.2021			
Snowmelt	22.12.2020 – 29.12.2020	101	91	87
	29.01.2021 – 01.05.2021			

DOC export from HS<sub>tot</sub> and subcatchments was calculated as follows for the entire study period and the hydrological periods:

$$DOC\ export = \sum_{i=1}^n c_i \times Q_i \times 15$$

170 where  $c_i$  is the DOC concentration [mg L<sup>-1</sup>] and  $Q_i$  is the discharge [L min<sup>-1</sup>] for  $n$  15-minute intervals. Small data gaps of several hours were interpolated linearly. For the year investigated, DOC data for HS<sub>tot</sub> was available for 351 days out of 365 days. Daily DOC export averages were calculated by dividing the DOC export measured during one hydrological period by the length of the respective hydrological period. Total DOC export from HS<sub>tot</sub> for each hydrological period was calculated as the sum of the measured DOC export and, for each day without data, the calculated daily average (for the corresponding hydrological period). Results from all subcatchments were available on 231 days during the study period. Flow-weighted DOC export for a period was calculated by dividing the measured and estimated DOC export during this period by the cumulated discharge generated in this period. When comparing the contribution of different subcatchments to total DOC

export, we used the ratio of the subcatchment area to the total catchment area as a benchmark ( $KS/HS_{tot} = 0.26$ ,  $MG/HS_{tot} = 0.31$ ,  $HS_{sub}/HS_{tot} = 0.43$ ). All results presented for  $HS_{sub}$  were derived from the difference of  $HS_{tot}$  and the upper subcatchments (MG and KS). For these calculations, we assumed that all DOC entering  $HS_{sub}$  was still present at the outlet of  $HS_{tot}$  and did not leave the stream by surface water-groundwater exchange or in-stream processes on its passage towards the catchment outlet. Studies have indicated that the relevance of in-stream metabolic processes influencing stream DOC concentrations is limited, particularly in headwater streams where residence times of stream water are expected to be short (e.g., Bernal et al. (2019); Singh et al. (2015)). Using DOM quality metrics, Da Silva et al. (2021) and Kothawala et al. (2015) found no indication for in-stream metabolic processing in [similar](#) forested headwater streams.

### 3 Results

#### 3.1 DOC concentrations, discharge and precipitation during the investigation period

With 1280 mm of precipitation at RDH, the investigated period was drier than the long-term annual average (1535 mm for RDH 1990 – 2019). The mean temperature during the study period was higher (6.8°C) than the long-term annual average (6.5°C, 1990 – 2019).

The spring period was characterized by frequent precipitation events. During the summer period, long dry periods were interrupted by a few precipitation events. In autumn, several large precipitation events occurred, and sharply [rising-increases](#) [in](#) DOC concentrations [of](#) up to 20 mg/L were observed (Fig. 2). During the winter period, precipitation fell as snow, which led to low discharge and low DOC concentrations. Winter was interrupted by one large snowmelt event induced by rising temperatures and rainfall towards the end of December. During snowmelt, diurnal fluctuations of discharge and DOC concentrations were observed following the amplitudes of air temperature. The highest discharge value measured during the study period reached  $0.75 \text{ m}^3 \text{ s}^{-1}$  at the onset of the snowmelt period. Additionally, some rain events enhanced snowmelt. DOC concentrations clearly increased with discharge during precipitation events (Fig. 2), observed during events in the summer and the autumn period and during the short snowmelt event in late December. During baseflow, DOC concentrations usually vary between 2 and 3  $\text{mg L}^{-1}$  at MG and  $HS_{tot}$  (Da Silva et al., 2021).




Figure 2: DOC concentrations at the outlets of the subcatchments KS (yellow), MG (green) and HS (black) and discharge (Q) at HS<sub>tot</sub> (blue). The shaded areas correspond to the delineated hydrological periods.


### 205 3.2 DOC export from the entire catchment HS<sub>tot</sub>

The annual DOC export of HS<sub>tot</sub> was 13760 kg or 39.31 kg ha<sup>-1</sup>, if including both the measured (13136 kg, 351 days) and the estimated (624 kg, 14 days) data (Table 2). The largest amounts of DOC were exported during snowmelt (40.5 %), spring (24.6 %) and autumn (20.3 %), whereas the export in summer (10.4 %) and winter (4.2 %) was markedly lower. In the following, all results refer only to periods with measured data. The highest mean daily DOC export was observed during the hydrological periods of spring and snowmelt, followed by autumn (>45 kg d<sup>-1</sup>), whereas winter and summer exhibited markedly lower mean daily DOC release (< 20 kg d<sup>-1</sup>) (Table 3).

Table 3. Total precipitation P<sub>tot</sub> [mm] at RDH, total DOC export [kg], number of days per hydrological period, and mean daily DOC export [kg d<sup>-1</sup>] at HS<sub>tot</sub> during the periods for which measured data were available (see Table 2).

	P <sub>tot</sub> [mm]	Total DOC export [kg]	Days [d]	Mean daily DOC export [kg d <sup>-1</sup> ]
Spring	327	3381	60	56.4
Summer	214	1378	83	16.7
Autumn	262	2794	57	48.8
Winter	129	576	60	9.5
Snowmelt	332	5007	91	55.1
All periods	1264	13136	351	37.4


As the hydrological periods had different lengths (Table 3) and DOC and discharge are positively related (Fig. 2), we also examined the flow-weighted DOC export normalized by the length of the hydrological periods to remove the influence of discharge (Fig. 3). Autumn proved to be the hydrological period with the highest daily flow-weighted DOC release, followed by the spring period. Although hydrologically being quite different, the summer, winter and snowmelt periods exhibited similar mean daily flow-weighted DOC export.



**Figure 3. Mean daily flow-weighted DOC export from the Hinterer Schachtenbach catchment ( $HS_{tot}$ ) during the different hydrological periods.**

225 Plotting discharge and DOC export time series as cumulative curves illustrated the DOC export behavior more clearly (Fig. 4). Cumulative discharge increased almost linearly during the study period with a break in slope with the onset of snowmelt at the end of January 2021, indicating an increased runoff generation during the snowmelt and spring period that lasted until the end of the study period.


In contrast, pronounced step-like increases were visible in the cumulative DOC export curve, particularly in autumn and during early snowmelt. The deviation between the two cumulative curves indicated that DOC export increased disproportionately relative to discharge during these periods. ~~In contrast, t~~During most of the winter and towards the end of the snowmelt period, however, DOC export decreased relative to discharge.



235 **Figure 4. Cumulative discharge (blue line) and cumulative DOC export (black dashed line) from  $HS_{tot}$  during the different hydrological periods (shaded background). Missing data (white background) are not accounted for.**

### 3.2 DOC export behavior of the subcatchments

The area-normalized cumulative export of the three subcatchments showed a similar pattern running parallel from late spring until the middle of the autumn period, when  $HS_{sub}$  began to export less DOC than MG and KS (Fig. 5). The cumulative export of  $HS_{sub}$  remained smaller than the export of MG and KS throughout the winter period. In winter, DOC export was generally  
 240 low. Export of area-normalized DOC from all three subcatchments during the autumn and the winter period occurred in a step-like pattern whereas with the onset of snowmelt, DOC export happened more continuously over time, with particularly strong increases in  $HS_{sub}$  and MG at the beginning of the snowmelt period. Strong increases of DOC export could also be seen in spring following precipitation events. At the end of the study period, MG showed the highest DOC export per unit area, whereas the area-normalized export of KS was lower by 32%.




245

**Figure 5. Cumulative DOC export per unit area from the subcatchments MG, KS and HS<sub>sub</sub> during the different hydrological periods (shaded background). Note: For periods when not all three sensors were working (white background), cumulative DOC export from the subcatchments was not calculated (dashed lines).**

Area-normalized daily DOC export was high during snowmelt, spring and autumn and low during winter (Fig. 6a). Flow-weighted DOC export was always highest in the HS<sub>sub</sub> catchment, particularly during autumn, underlining the relevance of HS<sub>sub</sub> for DOC export (Fig. 6b). However, the relative contribution of the subcatchments to total DOC export varied between the seasons (Fig. 6c).

250



255 **Figure 6. Contributions to DOC export and runoff of the subcatchments KS, MG and HS<sub>sub</sub> for spring, autumn, winter, and snowmelt: a) area-normalized daily mean DOC export in kg d<sup>-1</sup> km<sup>-2</sup>, b) DOC export weighted by discharge generated in each subcatchment in mg L<sup>-1</sup>, c) contribution to total DOC export in %, d) contribution to total runoff in %. For c) and d), the horizontal lines correspond to the value expected according to the area ratio (see Table 1).**

During snowmelt and spring, the order of contribution to total DOC export was as expected in terms of catchment area as HS<sub>sub</sub> contributed most, followed by MG and KS. In winter, all subcatchments contributed equally to total DOC export due to a larger contribution than expected of KS and a smaller contribution than expected of HS<sub>sub</sub>. In autumn, MG contributed most to total DOC export, whereas HS<sub>sub</sub> contributed less than during the other hydrological periods (Fig. 6c). The contribution of MG was larger than expected during the entire year and even exceeded the contribution of the larger subcatchment HS<sub>sub</sub> in autumn. MG also contributed more to runoff than expected during all hydrological periods, whereas HS<sub>sub</sub> contributed less than expected to runoff (Fig. 6d). KS contributed more to runoff than expected from proportional area in autumn and winter.

265 The runoff ratios were highest for MG during all hydrological periods except for winter and summer, when KS showed the highest runoff ratio (Table 4). The runoff ratios were lowest for HS<sub>sub</sub> during all hydrological periods. The highest runoff ratios could be found during snowmelt for all subcatchments. The lowest runoff ratio could be found in spring for KS, in summer for MG and HS<sub>tot</sub> and in autumn for HS<sub>sub</sub>. Overall, HS<sub>tot</sub> exhibited a mean runoff coefficient of 0.57 for the entire study period.

270

**Table 4. Runoff ratios for the entire catchment  $HS_{tot}$  and the subcatchments KS, MG and  $HS_{sub}$  during the periods for which measured discharge data was available for all subcatchments.**

	$HS_{tot}$	KS	MG	$HS_{sub}$
Spring	0.48	0.54	0.68	0.30
Summer	0.38	0.62	0.47	0.18
Autumn	0.40	0.56	0.63	0.13
Winter	0.53	0.82	0.64	0.29
Snowmelt	0.94	0.92	1.15	0.81
Entire study period	0.57	0.69	0.75	0.38

275 The contribution to DOC export and the contribution to runoff of the subcatchments differed (Fig. 6 and Table 5). The contribution to DOC export from KS and MG during the entire period was lower than their contribution to runoff. On the contrary, the contribution of  $HS_{sub}$  to DOC export was markedly larger than its contribution to runoff for all hydrological periods (Table 5).

280 **Table 5. Ratio between relative contribution to DOC and relative contribution to runoff generation for the three subcatchments during the hydrological periods (without summer)\*.**

	KS	MG	$HS_{sub}$
Spring	0.66	0.88	1.57
Autumn	0.79	0.74	2.77
Winter	0.85	0.86	1.51
Snowmelt	0.68	0.9	1.32

\*If the ratio is larger than 1, then the contribution to DOC release is relatively larger than the contribution to runoff generation.

## 4 Discussion

### 4.1 Variations of DOC export from the entire headwater catchment over time

285 The annual DOC export of the catchment of 39.31 kg ha<sup>-1</sup> is in the range of other forested catchments studied (15-99 kg ha<sup>-1</sup> in Ågren et al. (2007); 31 kg ha<sup>-1</sup> in Raymond and Saiers (2010)) with values, e.g., being higher than measured in a Mediterranean catchment with 1.8±1 kg C ha<sup>-1</sup> (Bernal and Sabater, 2012), but lower than in a wetland-dominated catchment (84 kg ha<sup>-1</sup> in Strohmeier et al. (2013)). DOC export of the entire Große Ohe catchment (which the Hinterer Schachtenbach catchment is a subcatchment of) was estimated to be very similar as well with 42 kg ha<sup>-1</sup> (Beudert et al., 2012). In general, it has been found that precipitation, resulting in runoff events, is a primary control for DOC export (Wei et al., 2021) and that  
290 wetter years result in a higher export of DOC than drier years (Jager et al., 2009).

The absolute export of DOC during the hydrological periods was highest during snowmelt and lowest during the winter period. Absolute DOC loads are the result of the interplay between runoff volumes and DOC concentration levels reached. However, these DOC amounts are affected by the lengths of the hydrological periods and by the runoff generated. Since discharge and DOC concentrations are positively correlated and discharge is the main driver for DOC export, the flow-weighted evaluation of DOC export is necessary to disentangle the relevance of factors other than discharge, such as topography, soils or vegetation, for the release of DOC into streams. The flow-weighted values can also be interpreted as the efficiency of a catchment to mobilize DOC per unit generated discharge.

When evaluating the flow-weighted export of DOC from the headwater catchment Hinterer Schachtenbach (Fig. 3), we observed that DOC export was low during the periods of summer, winter, and snowmelt, compared with the other two hydrological periods of spring and autumn. In summer, discharge was at baseflow with occasional short increases in discharge due to precipitation events. During these events, DOC concentrations rose up to 20 mg L<sup>-1</sup>, reaching the highest concentrations observed during the study period. This indicated that DOC was available and ready to be mobilized. However, since discharge was mostly low, the exported DOC load remained overall also low for the summer period. Less precipitation input and an increased water demand by vegetation led to decreasing catchment wetness and drier soils. We argue that hydrological connectivity between the DOC source areas and the stream was established less often and to a lesser spatial extent leading to the inhibition of DOC export as explained by Blaurock et al. (2021), derived from an analysis of four precipitation events. This analysis suggested that the lower flatter part of the headwater catchment with the wide riparian zone may be affected more often by missing hydrological connectivity. At the same time, previous work indicated that this catchment part can release substantial amounts of DOC that accumulates in microtopographic depressions into the stream once hydrological connectivity is established (Blaurock et al., 2022). Overall, this again points to DOC export being transport-limited in this catchment, as has been observed in many other catchments (Zarnetske et al., 2018). The relevance of this transport-limitation of DOC release into streams may increase in the future if summer droughts will occur more frequently with climate change.

During the winter period, with precipitation falling as snow and adding to the snowpack, discharge at the outlet of the Hinterer Schachtenbach was characterized by a constant low level, except for a brief snowmelt period following rising air temperatures and a rain event at the end of December 2020. Also, DOC remained at concentrations typical for baseflow conditions (2-3 mg L<sup>-1</sup>) in this catchment (Da Silva et al., 2021), resulting in a low DOC release from the catchment. DOC production is generally low during winter because of cold temperatures (Kalbitz et al., 2000) and snowfall. DOC produced during [summer and](#) autumn accumulates in the soil and can be readily mobilized, as evidenced by the strong increase in DOC concentrations during the short snowmelt event at the end of December 2020.

The snowmelt period led to higher discharge and an increase of DOC concentrations in streamflow. Several studies have shown that DOC export during snowmelt can be substantial (Jager et al., 2009; Seybold et al., 2019; Wilson et al., 2013). The diurnal fluctuations of discharge and DOC during snowmelt indicated that DOC export was strongly controlled by hydrological processes, which responded to the freeze-thaw cycle of the snowpack. The largest DOC export could be observed at the beginning of the snowmelt periods probably due to the flushing of topsoil DOC that had accumulated during winter (Ågren et

325 al., 2010; Boyer et al., 1997) as a result of the leaching of DOC from the autumnal litter input or DOC production in snow-  
covered soil (Brooks et al., 1999). However, neither discharge nor DOC concentrations showed any pronounced event  
responses, which again led to overall low flow-weighted DOC export (Fig. 3) and a more continuous release of DOC (Fig. 4).  
During the spring period at the beginning and at the end of our study period, the daily flow-weighted DOC almost doubled,  
330 compared to the summer, winter and snowmelt periods (Fig. 3). Soils were likely saturated after the snowmelt period and,  
therefore, hydrological connectivity between the DOC sources and the stream ~~existed~~facilitating the connection of distal  
DOC sources (Croghan et al., 2023). This connection contributed to ~~and~~ an increased DOC export (~~Fig. 4~~) during the spring  
events (Fig. 4). Furthermore, the increasing temperatures during spring had enhanced DOC production, thereby increasing the  
potential mobilizable DOC pool.

The autumn period showed the highest flow-weighted DOC release. The onset of larger precipitation events led to sharp  
335 increases in discharge and DOC concentrations between 15 and 20 mg L<sup>-1</sup>. In autumn, DOC export was clearly driven by the  
large precipitation events. During the hydrological period of autumn which accounted for 16% of the study period, 14% of the  
total discharge were generated but 21% of total DOC were mobilized. The ratio between percentages of DOC exported and of  
runoff generated was highest for autumn (Table S1), again underlining the pronounced DOC export in relation to discharge  
generation during this period. Moreover, it is possible that DOC had accumulated in the soils during summer due to 1) the  
340 warm temperatures, which enhance biological activity as well as evapotranspiration, and 2) the lack of precipitation events  
and a low hydrological connectivity, therefore limiting DOC flushing (Dawson et al., 2008; Kawasaki et al., 2005; Seybold et  
al., 2019; Wei et al., 2021). In autumn, leaf litter from deciduous trees may constitute an additional important DOC source  
potentially amplifying DOC export during this period (Hongve, 1999; Meyer et al., 1998). Long-term observations of  
phenology and litter fall in the Bavarian Forest National Park show that over 80 % of litter fall is likely to occur during the  
345 here defined autumn period at all elevations (unpublished data, Bavarian Forest National Park Administration, 2022). It has  
also been shown that leaf litter leads to an annual carbon input of 150,000 kg C/km<sup>2</sup> in a neighboring catchment (Forellenbach)  
and can therefore substantially contribute to DOC export (unpublished data, Bavarian Forest National Park Administration,  
2022).

To conclude, we saw pronounced temporal variations in DOC export from the Hinterer Schachtenbach catchment during our  
350 one-year study period. As has been observed elsewhere, DOC showed a chemodynamic behavior with a positive relation to  
discharge. Hence, precipitation regime and catchment wetness, both governing hydrological connectivity and runoff responses,  
were important factors for DOC export. Resulting runoff events contributed substantially to DOC release within short times.  
After normalizing DOC export with runoff, the hydrological period of autumn, after a dry summer period, showed the highest  
DOC release, caused by the often-missing hydrological connectivity in the preceding summer period and a subsequent  
355 accumulation of DOC in the soils of the catchment. The hydrological periods of summer, winter and snowmelt exhibited low  
flow-weighted DOC export, driven by the interplay between runoff generated and available DOC pool that can be mobilized.  
The DOC production in a catchment, in turn, is influenced by temperature and the organic matter from vegetation available  
for decomposition.

## 4.2 The relative importance of the subcatchments for runoff generation and DOC export

360 The one-year dataset of 15-min DOC concentrations and discharge values at three topographic positions within the Hintere Schachtenbach catchment illustrated the relative role of the subcatchments for runoff generation and the mobilization of DOC. The area-normalized DOC release of the subcatchments (Fig. 6a) changed over time and also in relation to each other. Snowfall and low temperatures led to a low DOC export in winter. The relatively high contribution of KS during autumn and winter could be due to the influence of litter fall on DOC export as KS is the subcatchment with the highest fraction of deciduous

365 trees (Table 1). The breakdown of this additional input of organic matter could lead to a constant DOC production even during the winter months (Hongve, 1999). Additionally, litter fall (a) induces a lower water loss via canopy interception in autumn and (b) leads to strongly reduced transpiration rates. Both factors would result in shifts in the water balance, leading to an increased availability of water in the soil profile and active flow pathways, as suggested by the high runoff ratio at KS during winter (Table 4). During the spring period, the relative importance of MG for DOC export increased. This could be a

370 consequence of a delayed snowmelt. The subcatchment MG extends up to 1373 m a.s.l., which is 100 m higher than the neighboring subcatchment KS, leading to colder temperatures and a longer lasting snowpack (unpublished data, Bavarian Forest National Park Administration, 2022). Therefore, a delayed snowmelt at MG could contribute to the higher DOC export in the spring period. Similar observations were made by Boyer et al. (1997) who found a delayed contribution of upslope catchment parts to the total DOC export due to an asynchronous melting of the snow in an alpine catchment.

375 Moreover, the contribution of MG to runoff from the entire Hintere Schachtenbach catchment was markedly higher than expected from the area proportion during all hydrological periods and except for the winter period always the highest among the three subcatchments (Fig. 6d). One reason for this could be the higher altitude of the subcatchment MG, resulting in a higher precipitation due to orographic effects. Observations in the Große Ohe catchment have shown that precipitation increases by around 100 mm per 100 m increase in altitude. MG is characterized by steep slopes with over 22 % of the area

380 having slopes with 21 – 25° (in comparison to 2 % in the neighboring subcatchment KS). MG possesses also a higher percentage of Podzol soils that usually have a higher fraction of sand and subsequently large pores, leading to a higher hydraulic conductivity (Scheffer-Schachtschabel, 2018). The permeable soils in MG are also interspersed with large rocks, facilitating the occurrence of preferential flow paths. These factors may result in a quick movement of infiltrated water towards the stream channel and a high runoff generation. In contrast, at KS, interspersed rocks are much less prominent leading to slower flow

385 processes. The subcatchment MG also has the highest area percentage in the state of forest rejuvenation. It is therefore likely that due to these differences in extent of mature trees, the evaporation component in the water balance of MG is markedly lower compared to KS, resulting in an increased availability of water for runoff generation. These differences in topography, soils as well as vegetation characteristics could explain the differences in runoff generation between the two upslope subcatchments KS and MG.

390 The lower subcatchment HS<sub>sub</sub> exhibited particularly low mobilized DOC loads per unit area in the autumn period (Fig. 6a). This period was also the one with the lowest runoff contribution of HS<sub>sub</sub> (Fig. 6d). In our previous event-based study, we had

identified the missing hydrologic connectivity in the subcatchment HS<sub>sub</sub> after summer drought as a dominant reason for inhibited DOC mobilization from this subcatchment towards the stream during autumn (Blaurock et al., 2021). The flat riparian zone of HS<sub>sub</sub> is dependent on a high level of catchment wetness to enable hydrological connectivity in the shallow subsurface, which is the prerequisite for DOC export, whereas the steeper upper subcatchments are able to maintain flow towards the stream even during dry conditions. The one-year dataset confirmed our event-based interpretation.

395 However, considering only the area-normalized export of DOC may skew our view of catchment functioning. Normalizing the DOC export values with the cumulative discharge that was generated in each subcatchment per hydrological period allows to consider mobilization factors other than discharge. HS<sub>sub</sub> always showed the highest flow-weighted DOC export (Fig. 6b), e.g., in HS<sub>sub</sub>, one unit discharge mobilized around four times as much DOC as in the other subcatchments during the autumn period. This suggests that the lower subcatchment possesses a large DOC store that can be activated whenever hydrological connectivity is established.

We propose that the high flow-weighted DOC export in the subcatchment HS<sub>sub</sub> is the result of the interplay between hydrological processes and DOC production and accumulation. In our previous work at the study site, we identified microtopographic depressions (“ponds”) in the riparian zone that fill regularly with water from the bottom during large events and periods of high catchment wetness as hot spots for DOC accumulation and DOC release towards the stream (Blaurock et al., 2022). After the warm and dry summer 2020 with low runoff generation in the Hinterer Schachtenbach catchment, the soils in the flat riparian zone around the stream in HS<sub>sub</sub> had dried out, as did the ponds in the riparian zone. The few events in August led to only small discharge increases but still high DOC concentrations, resulting in a low DOC load. The month of September was particularly dry. Shallow piezometers installed in the riparian zone of HS<sub>sub</sub> during the 2020 summer months (one of them being installed in a pond) showed that during July, August and September 2020, water levels in the soils did rise into the upper riparian soil layers following precipitation events but also quickly receded again (Fig. 2 in Blaurock et al. (2022)). This pointed to a quick connection of potentially DOC-rich layers and a subsequent drying-out of the soils in the riparian zone of HS<sub>sub</sub> and lost hydrologic connectivity. Particularly low water levels in the shallow groundwater were reached at the beginning of August and at the end of September. The first autumn events led to a more substantial wet-up of the soils and produced strong DOC concentration increases in the stream (see Fig. 2). However, discharge remained still low. The highest value of water level in the piezometer installed in a pond (P1 in Fig. 2 from Blaurock et al. (2022)) corresponds to spill-over of the pond; this is the occasion when the ponds are fully-completely filled, flow over and connect with each other and the stream via localized overland flow. This spill-over was reached only once during June to end of October 2020 during the end of September, and even though the pond remained filled during most of October (water level > 0 cm, Fig. 2 in Blaurock et al., 2022), it did not spill again. This indicated that wide-spread hydrological connectivity had not been reached yet, therefore runoff generation in HS<sub>sub</sub> remained still low. Towards the end of the autumn period, in November 2020, the baseflow level of the Hinterer Schachtenbach rose, suggesting that runoff generation had picked up.

In addition, the lower elevation of HS<sub>sub</sub> and resulting differences in mean air and soil temperatures may have contributed to an overall higher production and availability of DOC. Particularly in autumn, after the dry summer, DOC had accumulated

and was readily exported into the streams during the first large rainfall events, leading to the pronounced flow-weighted DOC export from  $HS_{sub}$ .

430 Comparing the daily flow-weighted DOC export per hydrological period from the entire catchment  $HS_{tot}$  (Fig. 3) and the subcatchments (Fig. S6) revealed that each subcatchment behaved as the entire catchment Hinterer Schachtenbach but that the differences between the hydrological periods varied for the three subcatchments (note: summer could not be evaluated). The difference between spring and autumn was particularly pronounced for  $HS_{sub}$  and only small for MG. As stated above, this could be explained with the missing hydrological connectivity between DOC source areas and the stream during the dry summer that led to an accumulation of DOC in the soils of  $HS_{sub}$ . In the transition from winter to snowmelt, all subcatchments seemed to contribute equally to the doubling of flow-weighted DOC export of  $HS_{tot}$ . As DOC export was limited during winter, 435 DOC which had accumulated in the topsoil during the winter months could now be flushed by meltwater and precipitation events (Ågren et al., 2010; Boyer et al., 1997). From snowmelt to spring, the strongest change was observed in  $HS_{sub}$ , suggesting that  $HS_{sub}$  with its large available DOC pool was most likely driving the increase in flow-weighted DOC export. The upper subcatchments mobilized substantial amounts of DOC because of their high runoff generation, as indicated by their runoff ratios that were always higher than for the lower subcatchment (Table 4). At the same time, their contribution to DOC 440 export was not as high as their contribution to runoff within the Hintere Schachtenbach catchment (Table 5). In contrast, despite the often low runoff ratio of the lower subcatchment  $HS_{sub}$  (except for the snowmelt period), it contributed disproportionately to DOC export in relation to runoff, illustrating the DOC source strength of this catchment part.

Analyzing area-normalized as well as flow-weighted DOC export, in combination with the relative contributions of the subcatchments to DOC export and runoff generation of the Hinterer Schachtenbach catchment, revealed different factors that 445 drive the release of DOC. The upper subcatchment MG contributed substantially to DOC export because of the pronounced runoff generation taking place in MG. Topography, soils and vegetation can explain the differences in runoff generation and DOC release between the two steep upslope subcatchments MG and KS. The lower subcatchment  $HS_{sub}$  was limited in its contribution to DOC export during times of low catchment wetness since it is highly dependent on the establishment of hydrological connectivity between the DOC source areas in the flat riparian zone and the stream. However, due to its deeper 450 soils and microtopographic features that act as DOC production and accumulation hot spots, high amounts of DOC are present that can be readily mobilized, whenever wetter hydrological conditions occur and hydrological connectivity is established.

The transport-limited nature of DOC mobilization has been shown before (Zarnetske et al., 2018). Our study demonstrates that hydrological processes strongly control the export of DOC but that the runoff-based contribution of subcatchments over time 455 is modulated by the interplay of soils, vegetation, topography and microclimate, which can be seen as secondary controls.

## 5 Conclusions and outlook

Our high-resolution data show that DOC export varies seasonally as a result of both varying DOC production and varying hydrological connectivity, which is influenced by topographical position. Topography, together with soil hydraulic properties and vegetation (differences in litter fall and dynamics of interception), governs the accumulation of water in the landscape, the formation of hydrological flow pathways and the establishment of hydrological connectivity towards the stream channel. We argue that these hydrological processes control the DOC export into streams and lakes. Although hydrological connectivity is a general state that needs to be present to enable the transport of solutes towards the stream, some catchments are particularly sensitive towards the establishment of hydrological connectivity, due to their combination of topographic, soil and vegetation characteristics. Different catchment parts can mobilize high DOC loads for different reasons, e.g., due to strong runoff generation and/or the presence of DOC production and accumulation hot spots that are activated upon occurrence of hydrological connectivity towards the stream. In next decades, temperate ecoregions are likely to be influenced strongly by climate change as a result of rising temperatures and changing precipitation patterns (Intergovernmental Panel on Climate Change, 2021), influencing terrestrial-aquatic linkages within the carbon cycle with potentially large consequences for DOC export (Wit et al., 2021). Rising temperatures could increase DOC production and, therefore, the potential DOC pool. Higher temperatures in winter would reduce the amount of precipitation that falls as snow. Consequently, more DOC could be exported during winter and the flushing effect during snowmelt would be reduced. DOC export could be distributed more evenly during the winter and spring months. However, more extreme precipitation events and prolonged drought periods could lead to a stronger contrast in DOC export patterns during summer and autumn. Drought periods would also limit DOC production and a low DOC export in summer and autumn could then lead to a higher DOC export during the following winter and snowmelt (Ågren et al., 2010). Droughts can lead to large-scale die-offs of trees, e.g., by facilitating bark beetle infestations and/or windthrow. This would affect the partitioning of water within the catchment water balance and provide a disproportionately high carbon pool for decomposition within a short period. Additionally, topography and vegetation seem to influence the contribution of the different catchment parts to DOC export throughout the year. Drought periods could limit DOC export from the lower catchment parts, which are dependent on hydrological connectivity, leading to an increasing importance of the upper catchment parts for DOC export. This effect could be enhanced by longer warm periods at higher altitudes.

Our study highlights that we need a thorough understanding of catchment hydrological processes to understand the patterns and dynamics of carbon export from terrestrial systems into inland waters (Catalán et al., 2016). High-resolution data is essential to capture the highly dynamic DOC export behavior and to include fast mobilization processes in DOC export calculations. Therefore, high-resolution data contributes substantially to a better understanding of the seasonally varying contribution of landscape parts to streamflow generation and associated DOC mobilization and export.

*Data availability.* The data set used in this study is available at Figshare via <https://doi.org/10.6084/m9.figshare.22770365>. During the peer-review process, we kindly ask the reviewers to use the following link to access the dataset:

https://figshare.com/s/1216d7fecbd3f0488d83. The official DOI will be activated upon publication. For data analysis,  
490 MATLAB (The MathWorks, Inc.) was used.

*Supplement.* The supplement related to this article is available online at:

*Author contributions.* Conceptualization by LH, KB and BB; investigation, methodology and formal analysis by KB; Writing  
495 – original draft preparation by KB; Writing – review & editing by LH and KB.

*Competing interests.* The contact author has declared that none of the authors has any competing interests.

*Acknowledgements.* This research was funded by the Rudolf and Helene Glaser Foundation and the Zempelin Foundation of  
500 the Stifterverband für die Deutsche Wissenschaft, as part of the project “Influence of natural factors on concentration, quality  
and impact of dissolved organic carbon in the Bavarian Forest National Park” (Project No. T0083\30771\2017\kg). The authors  
would like to thank the Bavarian Forest National Park (BFNP) administration for providing physiographic and meteorological  
data, and the BFNP staff for their helpful assistance with the installation and maintenance of the field equipment. Funded by  
the Open Access Publishing Fund of the University of Bayreuth.

505

## References

- 510 Ågren, A., Buffam, I., Jansson, M., and Laudon, H.: Importance of seasonality and small streams for the landscape  
regulation of dissolved organic carbon export, *J. Geophys. Res.*, 112, doi:10.1029/2006JG000381, 2007.
- Ågren, A., Haei, M., Köhler, S. J., Bishop, K., and Laudon, H.: Regulation of stream water dissolved organic carbon (DOC)  
concentrations during snowmelt; the role of discharge, winter climate and memory effects, *Biogeosciences*, 7, 2901–  
2913, doi:10.5194/bg-7-2901-2010, 2010.
- 515 Alarcon-Herrera, M. T., Bewtra, J. K., and Biswas, N.: Seasonal variations in humic substances and their reduction through  
water treatment processes, *Can. J. Civ. Eng.*, 21, 173–179, doi:10.1139/194-020, 1994.
- Alvarez-Cobelas, M., Angeler, D. G., Sánchez-Carrillo, S., and Almendros, G.: A worldwide view of organic carbon export  
from catchments, *Biogeochemistry*, 107, 275–293, doi:10.1007/s10533-010-9553-z, 2012.
- Batjes, N. H.: Total carbon and nitrogen in the soils of the world, *European J Soil Science*, 65, 10–21,  
520 doi:10.1111/ejss.12114\_2, 2014.

- Battin, T. J., Luysaert, S., Kaplan, L. A., Aufdenkampe, A. K., Richter, A., and Tranvik, L. J.: The boundless carbon cycle, *Nature Geosci*, 2, 598–600, doi:10.1038/ngeo618, 2009.
- Bavarian State Office for Environment: Current discharge values Rachel Diensthuetten / Markungsgraben, <https://www.gkd.bayern.de/de/fluesse/wasserstand/passau/rachel-diensthuetten-17418004>, 2021.
- 525 Bernal, S., Lupon, A., Wollheim, W. M., Sabater, F., Poblador, S., and Martí, E.: Supply, Demand, and In-Stream Retention of Dissolved Organic Carbon and Nitrate During Storms in Mediterranean Forested Headwater Streams, *Front. Environ. Sci.*, 7, doi:10.3389/fenvs.2019.00060, 2019.
- Bernal, S. and Sabater, F.: Changes in discharge and solute dynamics between hillslope and valley-bottom intermittent streams, *Hydrol. Earth Syst. Sci.*, 16, 1595–1605, doi:10.5194/hess-16-1595-2012, 2012.
- 530 Beudert, B., Bäessler, C., Thorn, S., Noss, R., Schröder, B., Dieffenbach-Fries, H., Foullois, N., and Müller, J.: Bark Beetles Increase Biodiversity While Maintaining Drinking Water Quality, *Conservation Letters*, 8, 272–281, doi:10.1111/conl.12153, 2015.
- Beudert, B., Spitzky, A., Kloeking, B., Zimmermann, L., Baessler, C., and Foullois, N.: DOC-Langzeitmonitoring im Einzugsgebiet der "Großen Ohe", Wasserhaushalt und Stoffbilanzen im naturnahen Einzugsgebiet Große Ohe, Bericht  
535 Nr. 9, Nationalparkverwaltung Bayerischer Wald, 78 pp., 2012.
- Blaurock, K., Beudert, B., Gilfedder, B. S., Fleckenstein, J. H., Peiffer, S., and Hopp, L.: Low hydrological connectivity after summer drought inhibits DOC export in a forested headwater catchment, *Hydrol. Earth Syst. Sci.*, 25, 5133–5151, doi:10.5194/hess-25-5133-2021, 2021.
- Blaurock, K., Garthen, P., Da Silva, M. P., Beudert, B., Gilfedder, B. S., Fleckenstein, J. H., Peiffer, S., Lechtenfeld, O. J.,  
540 and Hopp, L.: Riparian Microtopography Affects Event-Driven Stream DOC Concentrations and DOM Quality in a Forested Headwater Catchment, *J. Geophys. Res.*, 127, doi:10.1029/2022JG006831, 2022.
- Borken, W., Ahrens, B., Schulz, C., and Zimmermann, L.: Site-to-site variability and temporal trends of DOC concentrations and fluxes in temperate forest soils, *Global Change Biology*, 17, 2428–2443, doi:10.1111/j.1365-2486.2011.02390.x, 2011.
- 545 Boyer, E. W., Hornberger, G. M., Bencala, K. E., and McKnight, D. M.: Response characteristics of DOC flushing in an alpine catchment, *Hydrol. Process.*, 11, 1635–1647, doi:10.1002/(SICI)1099-1085(19971015)11:12<1635:AID-HYP494>3.0.CO;2-H, 1997.
- Brooks, P. D., McKnight, D. M., and Bencala, K. E.: The relationship between soil heterotrophic activity, soil dissolved organic carbon (DOC) leachate, and catchment-scale DOC export in headwater catchments, *Water Resources Research*,  
550 35, 1895–1902, doi:10.1029/1998WR900125, 1999.
- Catalán, N., Marcé, R., Kothawala, D. N., and Tranvik, L. J.: Organic carbon decomposition rates controlled by water retention time across inland waters, *Nature Geosci*, 9, 501–504, doi:10.1038/ngeo2720, 2016.
- Chaplot, V. and Mutema, M.: Sources and main controls of dissolved organic and inorganic carbon in river basins: A worldwide meta-analysis, *Journal of Hydrology*, 603, 126941, doi:10.1016/j.jhydrol.2021.126941, 2021.

- 555 Ciais, P., Borges, A. V., Abril, G., Meybeck, M., Folberth, G., Hauglustaine, D., and Janssens, I. A.: The impact of lateral carbon fluxes on the European carbon balance, *Biogeosciences*, 5, 1259–1271, doi:10.5194/bg-5-1259-2008, 2008.
- Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Downing, J. A., Middelburg, J. J., and Melack, J.: Plumbing the Global Carbon Cycle: Integrating Inland Waters into the Terrestrial Carbon Budget, *Ecosystems*, 10, 172–185, doi:10.1007/s10021-006-9013-8, 2007.
- 560 Covino, T.: Hydrologic connectivity as a framework for understanding biogeochemical flux through watersheds and along fluvial networks, *Geomorphology*, 277, 133–144, doi:10.1016/j.geomorph.2016.09.030, 2017.
- Croghan, D., Ala-Aho, P., Lohila, A., Welker, J., Vuorenmaa, J., Kløve, B., Mustonen, K.-R., Aurela, M., and Marttila, H.: Coupling of Water-Carbon Interactions During Snowmelt in an Arctic Finland Catchment, *Water Resources Research*, 59, doi:10.1029/2022WR032892, 2023.
- 565 Da Silva, M. P., Blaurock, K., Beudert, B., Fleckenstein, J. H., Hopp, L., Peiffer, S., Reemtsma, T., and Lechtenfeld, O. J.: Delineating Source Contributions to Stream Dissolved Organic Matter Composition Under Baseflow Conditions in Forested Headwater Catchments, *J. Geophys. Res.*, 126, doi:10.1029/2021JG006425, 2021.
- Dawson, J. J. C., Soulsby, C., Tetzlaff, D., Hrachowitz, M., Dunn, S. M., and Malcolm, I. A.: Influence of hydrology and seasonality on DOC exports from three contrasting upland catchments, *Biogeochemistry*, 90, 93–113, doi:10.1007/s10533-008-9234-3, 2008.
- 570 Detty, J. M. and McGuire, K. J.: Topographic controls on shallow groundwater dynamics: implications of hydrologic connectivity between hillslopes and riparian zones in a till mantled catchment, *Hydrol. Process.*, 24, 2222–2236, doi:10.1002/hyp.7656, 2010.
- Doerr, H. and Muennich, K. O.: Lead and cesium transport in european forest soils, *Water Air Soil Pollut*, 57-58, 809–818, doi:10.1007/BF00282944, 1991.
- 575 Drake, T. W., Raymond, P. A., and Spencer, R. G. M.: Terrestrial carbon inputs to inland waters: A current synthesis of estimates and uncertainty, *Limnol Oceanogr Letters*, 3, 132–142, doi:10.1002/lol2.10055, 2018.
- Hobbie, J. E. and Likens, G. E.: Output of Phosphorus, Dissolved Organic Carbon, and Fine Particulate Carbon from Hubbard Brook Watersheds, *Limnology & Oceanography*, 18, 734–742, doi:10.4319/lo.1973.18.5.0734, 1973.
- 580 Hongve, D.: Production of dissolved organic carbon in forested catchments, *Journal of Hydrology*, 224, 91–99, doi:10.1016/S0022-1694(99)00132-8, 1999.
- Hope, D., Billett, M. F., and Cresser, M. S.: A review of the export of carbon in river water: fluxes and processes, *Environmental pollution (Barking, Essex 1987)*, 84, 301–324, doi:10.1016/0269-7491(94)90142-2, 1994.
- Intergovernmental Panel on Climate Change (IPCC): Climate Change 2021: The Physical Science Basis: Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, U.K., 2391 pp., 2021.
- 585

- Jager, D. F., Wilmking, M., and Kukkonen, J. V. K.: The influence of summer seasonal extremes on dissolved organic carbon export from a boreal peatland catchment: evidence from one dry and one wet growing season, *The Science of the total environment*, 407, 1373–1382, doi:10.1016/j.scitotenv.2008.10.005, 2009.
- 590 Jankowski, K. J. and Schindler, D. E.: Watershed geomorphology modifies the sensitivity of aquatic ecosystem metabolism to temperature, *Scientific reports*, 9, 17619, doi:10.1038/s41598-019-53703-3, 2019.
- Kaiser, K. and Kalbitz, K.: Cycling downwards – dissolved organic matter in soils, *Soil Biology and Biochemistry*, 52, 29–32, doi:10.1016/j.soilbio.2012.04.002, 2012.
- Kalbitz, K., Solinger, S., Park, J.-H., Michalzik, B., and Matzner, E.: CONTROLS ON THE DYNAMICS OF DISSOLVED  
595 ORGANIC MATTER IN SOILS: A REVIEW, *Soil Science*, 165, 277–304, doi:10.1097/00010694-200004000-00001, 2000.
- Kawasaki, M., Ohte, N., and Katsuyama, M.: Biogeochemical and hydrological controls on carbon export from a forested catchment in central Japan, *Ecological Research*, 20, 347–358, doi:10.1007/s11284-005-0050-0, 2005.
- Kiewiet, L., van Meerveld, I., Stähli, M., and Seibert, J.: Do stream water solute concentrations reflect when connectivity  
600 occurs in a small, pre-Alpine headwater catchment?, *Hydrol. Earth Syst. Sci.*, 24, 3381–3398, doi:10.5194/hess-24-3381-2020, 2020.
- Kindler, R., Siemens, J. A., Kaiser, K., Walmsley, D. C., Bernhofer, C., Buchmann, N., Cellier, P., Eugster, W., Gleixner, G., Grünwald, T., Heim, A., Ibrom, A., Jones, S. K., Jones, M., Klumpp, K., Kutsch, W., Larsen, K. S., Lehuger, S., Loubet, B., McKenzie, R., Moors, E., Osborne, B., Pilegaard, K. I., Reibmann, C., saunders, M., Schmidt, M. W. I.,  
605 Schrumpf, M., Seyfferth, J., Skiba, U. T., Soussana, J.-F., Sutton, M. A., Tefs, C., Vowinckel, B., Zeeman, M. J., and Kaupenjohann, M.: Dissolved carbon leaching from soil is a crucial component of the net ecosystem carbon balance, *Global Change Biology*, 17, 1167–1185, doi:10.1111/j.1365-2486.2010.02282.x, 2011.
- Kothawala, D. N., Ji, X., Laudon, H., Ågren, A. M., Futter, M. N., Köhler, S. J., and Tranvik, L. J.: The relative influence of land cover, hydrology, and in-stream processing on the composition of dissolved organic matter in boreal streams, *J.  
610 Geophys. Res.*, 120, 1491–1505, doi:10.1002/2015JG002946, 2015.
- Laudon, H., Berggren, M., Ågren, A., Buffam, I., Bishop, K., Grabs, T., Jansson, M., and Köhler, S.: Patterns and Dynamics of Dissolved Organic Carbon (DOC) in Boreal Streams: The Role of Processes, Connectivity, and Scaling, *Ecosystems*, 14, 880–893, doi:10.1007/s10021-011-9452-8, 2011.
- Ledesma, J. L. J., Köhler, S. J., and Futter, M. N.: Long-term dynamics of dissolved organic carbon: implications for  
615 drinking water supply, *The Science of the total environment*, 432, 1–11, doi:10.1016/j.scitotenv.2012.05.071, 2012.
- Ledesma, J. L. J., Kothawala, D. N., Bastviken, P., Maehder, S., Grabs, T., and Futter, M. N.: Stream Dissolved Organic Matter Composition Reflects the Riparian Zone, Not Upslope Soils in Boreal Forest Headwaters, *Water Resources Research*, 54, 3896–3912, doi:10.1029/2017WR021793, 2018.
- Lovley, D. R., Fraga, J. L., Coates, J. D., and Blunt-Harris, E. L.: Humics as an electron donor for anaerobic respiration,  
620 *Environmental microbiology*, 1, 89–98, doi:10.1046/j.1462-2920.1999.00009.x, 1999.

- McGuire, K. J. and McDonnell, J. J.: Hydrological connectivity of hillslopes and streams: Characteristic time scales and nonlinearities, *Water Resources Research*, 46, doi:10.1029/2010WR009341, 2010.
- Mei, Y., Hornberger, G. M., Kaplan, L. A., Newbold, J. D., and Aufdenkampe, A. K.: The delivery of dissolved organic carbon from a forested hillslope to a headwater stream in southeastern Pennsylvania USA, *Water Resources Research*, 625 50, 5774–5796, doi:10.1002/2014WR015635, 2014.
- Meyer, J. L. and Tate, C. M.: The Effects of Watershed Disturbance on Dissolved Organic Carbon Dynamics of a Stream, *Ecology*, 64, 33–44, doi:10.2307/1937326, 1983.
- Meyer, J. L., Wallace, J. B., and Eggert, S. L.: Leaf Litter as a Source of Dissolved Organic Carbon in Streams, *Ecosystems*, 1, 240–249, doi:10.1007/s100219900019, 1998.
- 630 Musolff, A., Fleckenstein, J. H., Opitz, M., Büttner, O., Kumar, R., and Tittel, J.: Spatio-temporal controls of dissolved organic carbon stream water concentrations, *Journal of Hydrology*, 566, 205–215, doi:10.1016/j.jhydrol.2018.09.011, 2018.
- Pang, Y. and Wang, J.: Various electron donors for biological nitrate removal: A review, *The Science of the total environment*, 794, 148699, doi:10.1016/j.scitotenv.2021.148699, 2021.
- 635 Penna, D., van Meerveld, H. J., Oliviero, O., Zuecco, G., Assendelft, R. S., Dalla Fontana, G., and Borga, M.: Seasonal changes in runoff generation in a small forested mountain catchment, *Hydrol. Process.*, 29, 2027–2042, doi:10.1002/hyp.10347, 2015.
- Ploum, S. W., Laudon, H., Peralta-Tapia, A., and Kuglerová, L.: Are dissolved organic carbon concentrations in riparian groundwater linked to hydrological pathways in the boreal forest?, *Hydrol. Earth Syst. Sci.*, 24, 1709–1720, 640 doi:10.5194/hess-24-1709-2020, 2020.
- Ravichandran, M.: Interactions between mercury and dissolved organic matter--a review, *Chemosphere*, 55, 319–331, doi:10.1016/j.chemosphere.2003.11.011, 2004.
- Raymond, P. A., Hartmann, J., Lauerwald, R., Sobek, S., McDonald, C., Hoover, M., Butman, D., Striegl, R., Mayorga, E., Humborg, C., Kortelainen, P., Dürr, H., Meybeck, M., Ciais, P., and Guth, P.: Global carbon dioxide emissions from 645 inland waters, *Nature*, 503, 355–359, doi:10.1038/nature12760, 2013.
- Raymond, P. A. and Saiers, J. E.: Event controlled DOC export from forested watersheds, *Biogeochemistry*, 100, 197–209, doi:10.1007/s10533-010-9416-7, 2010.
- Raymond, P. A., Saiers, J. E., and Sobczak, W. V.: Hydrological and biogeochemical controls on watershed dissolved organic matter transport: pulse-shunt concept, *Ecology*, 97, 5–16, doi:10.1890/14-1684.1, 2016.
- 650 Ritson, J. P., Kennedy-Blundell, O., Croft, J., Templeton, M. R., Hawkins, C. E., Clark, J. M., Evans, M. G., Brazier, R. E., Smith, D., and Graham, N. J. D.: High frequency UV-Vis sensors estimate error in riverine dissolved organic carbon load estimates from grab sampling, *Environmental monitoring and assessment*, 194, 831, doi:10.1007/s10661-022-10515-9, 2022.

- Sadiq, R. and Rodriguez, M. J.: Disinfection by-products (DBPs) in drinking water and predictive models for their occurrence: a review, *The Science of the total environment*, 321, 21–46, doi:10.1016/j.scitotenv.2003.05.001, 2004.
- 655 Scheffer-Schachtschabel: *Lehrbuch der Bodenkunde*, 17.th ed., Springer Spektrum, 749 pp., 2018.
- Schleppi, P., Waldner, P. A., and Fritschi, B.: Accuracy and precision of different sampling strategies and flux integration methods for runoff water: comparisons based on measurements of the electrical conductivity, *Hydrol. Process.*, 20, 395–410, doi:10.1002/hyp.6057, 2006a.
- 660 Schleppi, P., Waldner, P. A., and Stähli, M.: Errors of flux integration methods for solutes in grab samples of runoff water, as compared to flow-proportional sampling, *Journal of Hydrology*, 319, 266–281, doi:10.1016/j.jhydrol.2005.06.034, 2006b.
- Seybold, E., Gold, A. J., Inamdar, S. P., Adair, C., Bowden, W. B., Vaughan, M. C. H., Pradhanang, S. M., Addy, K., Shanley, J. B., Vermilyea, A., Levia, D. F., Wemple, B. C., and Schroth, A. W.: Influence of land use and hydrologic variability on seasonal dissolved organic carbon and nitrate export: insights from a multi-year regional analysis for the northeastern USA, *Biogeochemistry*, 146, 31–49, doi:10.1007/s10533-019-00609-x, 2019.
- 665 Singh, S., Inamdar, S., and Mitchell, M.: Changes in dissolved organic matter (DOM) amount and composition along nested headwater stream locations during baseflow and stormflow, *Hydrol. Process.*, 29, 1505–1520, doi:10.1002/hyp.10286, 2015.
- 670 Strohmeier, S., Knorr, K.-H., Reichert, M., Frei, S., Fleckenstein, J. H., Peiffer, S., and Matzner, E.: Concentrations and fluxes of dissolved organic carbon in runoff from a forested catchment: insights from high frequency measurements, *Biogeosciences*, 10, 905–916, doi:10.5194/bg-10-905-2013, 2013.
- Tetzlaff, D., Birkel, C., Dick, J., Geris, J., and Soulsby, C.: Storage dynamics in hydrogeological units control hillslope connectivity, runoff generation, and the evolution of catchment transit time distributions, *Water Resources Research*, 50, 969–985, doi:10.1002/2013WR014147, 2014.
- 675 The MathWorks, Inc.: Versions 2019–2024, <https://www.mathworks.com>.
- Vemaps: Outline Map of Germany, <https://vemaps.com>, 2022.
- Wei, X., Hayes, D. J., Fernandez, I., Zhao, J., Fraver, S., Chan, C., and Diao, J.: Identifying Key Environmental Factors Explaining Temporal Patterns of DOC Export From Watersheds in the Conterminous United States, *J. Geophys. Res.*, 126, doi:10.1029/2020JG005813, 2021.
- 680 Weiler, M. and McDonnell, J. J.: Testing nutrient flushing hypotheses at the hillslope scale: A virtual experiment approach, *Journal of Hydrology*, 319, 339–356, doi:10.1016/j.jhydrol.2005.06.040, 2006.
- Wen, H., Perdrial, J., Abbott, B. W., Bernal, S., Dupas, R., Godsey, S. E., Harpold, A., Rizzo, D., Underwood, K., Adler, T., Sterle, G., and Li, L.: Temperature controls production but hydrology regulates export of dissolved organic carbon at the catchment scale, *Hydrol. Earth Syst. Sci.*, 24, 945–966, doi:10.5194/hess-24-945-2020, 2020.
- 685

- Werner, B. J., Musolff, A., Lechtenfeld, O. J., Rooij, G. H. de, Oosterwoud, M. R., and Fleckenstein, J. H.: High-frequency measurements explain quantity and quality of dissolved organic carbon mobilization in a headwater catchment, *Biogeosciences*, 16, 4497–4516, doi:10.5194/bg-16-4497-2019, 2019.
- 690 Wilson, H. F., Saiers, J. E., Raymond, P. A., and Sobczak, W. V.: Hydrologic Drivers and Seasonality of Dissolved Organic Carbon Concentration, Nitrogen Content, Bioavailability, and Export in a Forested New England Stream, *Ecosystems*, 16, 604–616, doi:10.1007/s10021-013-9635-6, 2013.
- Wit, H. A. de, Stoddard, J. L., Monteith, D. T., Sample, J. E., Austnes, K., Couture, S., Fölster, J., Higgins, S. N., Houle, D., Hruška, J., Krám, P., Kopacek, J., Paterson, A. M., Valinia, S., van Dam, H., Vuorenmaa, J., and Evans, C. D.: Cleaner air reveals growing influence of climate on dissolved organic carbon trends in northern headwaters, *Environmental research letters ERL [Web site]*, 16, 1–13, doi:10.1088/1748-9326/ac2526, 2021.
- 695 Zarnetske, J. P., Bouda, M., Abbott, B. W., Saiers, J., and Raymond, P. A.: Generality of Hydrologic Transport Limitation of Watershed Organic Carbon Flux Across Ecoregions of the United States, *Geophysical Research Letters*, 45, doi:10.1029/2018GL080005, 2018.