

Impacts of climate and land-surface change on catchment evapotranspiration and runoff from 1951-2020 in Saxony, Germany

Maik Renner^{1,2} and Corina Hauffe¹

¹Institute of Hydrology and Meteorology, Technische Universität Dresden, Dresden, 01069, Germany

²also at, Landesamt für Umwelt, Brandenburg

Correspondence: Maik Renner <maik.renner@gmail.com>, Corina Hauffe <corina.hauffe@tu-dresden.de>

Abstract. This paper addresses the question how catchment scale water and energy balance have responded to climatic and land-surface changes over the last 70 years in the federal state of Saxony, in eastern Germany. Therefore observational data of hydrological and meteorological monitoring sites from 1951-2020 across 71 catchments are examined in a relative water and energy partitioning framework to put the recent drought induced changes in a historical perspective. A comprehensive visualization method is used to analyze the observed time series. The study focuses on changes in decadal time scale and finds the largest decline in observed runoff in the last decade (2011-20). The observed decline can be explained by the significant increase in aridity, caused by the reduction of annual mean rainfall and an increase in potential evaporation at the same time. In a few mainly forested head water catchments the observed decline in runoff was even stronger than predicted by climate conditions alone. These catchments are still on the recovery from past widespread forest damages in the 1970-80s resulting in a continuous increase of actual evapotranspiration due to forest regrowth. In contrary runoff stayed almost constant in other catchments despite an increase in aridity.

These results highlight that water budgets in Saxony are in an unstable, non-stationary regime due to significant climatic changes and regional impacts of land-surface changes such as forest health. The recent decreases in the mean annual runoff are substantial and must be taken into account by the authorities for fresh water management.

1 Introduction

Annual mean runoff is an important variable in water management as it determines the available fresh water of a catchment. Runoff is governed by climatic conditions which describe the available water by precipitation and the atmospheric demand for water and by catchment properties such as topography, land-cover, soils and geology (Gentine et al., 2012).

Anthropogenic warming and increasing human pressure on ecosystem services is expected to impact water balance components (Gudmundsson et al., 2021). Recent hydroclimatic trend analysis across Europe shows continental scale diverging trends for annual mean precipitation with increases in northern Europe and declines in the Mediterranean region (Masseroni et al., 2021). The signal follows the trend pattern of the wet getting wetter and the dry getting drier (Held and Soden, 2006) which was early related to global warming. Although streamflow trends tend to follow the changes in precipitation, there are regions with diverse trends even opposing the meteorological pattern (Masseroni et al., 2021). Reasons for the diverging pattern of

25 streamflow trends could be manifold: different climatic sensitivities, changes in land-surface conditions (Teuling et al., 2019),
seasonal changes with earlier snow melt (Renner and Bernhofer, 2011; Berghuijs et al., 2014), compensating effects, decadal
variability influencing trend analysis (Hannaford et al., 2013) but also methodological reasons such as different time windows
of available observations.

The eastern part of Germany is such a region where trends in streamflow did not follow trends in climate and where previous
30 research for the federal state of Saxony found large decadal variations, including catchments where major land-cover changes
occurred that influenced catchment evapotranspiration and runoff (Renner et al., 2014).

Saxony has a long history of hydrological and meteorological observations which allows hydro-climatic analysis such as
the famous work by Schreiber (1904), who established the first type of a Budyko curve also with data from Saxony. Also
Saxony has a wide range of different topographic characteristics from the forested Ore mountains in the south to the flat and
35 arable regions in the northern part. The climate is shaped by the transition of the maritime climate in the west towards more
continental climate in the east. Over the last 70 years mean annual air temperature rose significantly by about 1.5 K (Franke
and Rühle, 2022) due to global warming by anthropogenic greenhouse gas emissions. While greenhouse gas emissions are still
increasing, the emissions of sulphur dioxide were successfully reduced since its peak in 1980s (Maas and Grennfelt, 2016).
Sulfur dioxide has strong environmental impacts. High atmospheric concentrations affected the transmission of visible solar
40 radiation leading to a reduction of surface solar radiation towards the 1990s and an trend reversal since then (Wild et al., 2005).
Air pollution was also the main cause of the tree dieback in the mountainous regions of the Ore and Izer mountains (Mazurski,
1986; Šrámek et al., 2008; Maas and Grennfelt, 2016). The forested higher regions were hit more severely since the needle
leaf trees comb out fog quite effectively. The fog, which was enriched by sulfuric acid, was intercepted by needle leaf trees
and then infiltrated into soils leading to soil acidification. Over the years the acidic conditions strongly reduced tree growth and
45 caused widespread tree damages (Pitelka and Raynal, 1989; Maas and Grennfelt, 2016). With the reduction of emissions in the
1990s and reforestation activities the forests started to regrow (SMUL, 2006; Šrámek et al., 2008). In recent years forests were
also hit by a series of large storms, the 2018-2020 droughts and bark beetle bloom. Here especially monoculture spruce tree
forest plantations were hit severely, leading to yet another forest disturbance (Otto et al., 2022).

Renner et al. (2014) analyzed decadal variations in water and energy balance components from 1950 to 2009 and separated
50 the climatic from the land-cover related impacts on evapotranspiration. They found significant impacts of forest disturbance
on catchment evapotranspiration and consistent but not significant impacts of climate changes. After 2009 climate change
continued and a record breaking drought in Europe occurred (Rakovec et al., 2022; Büntgen et al., 2021). It becomes natural
to ask how runoff responded to this drought and to put this change into relation to the earlier changes.

This study extends and updates the work of Renner et al. (2014). Therefore, the time series for 71 catchments in the federal
55 state of Saxony are updated and extended to cover the period from 1951 to 2020. The methods section introduces the coupling
of the water and energy balances and explains how this coupling can be used to separate climatic from land-cover impacts.
Then the dataset is introduced and in the results section the last 70 years are analyzed. The study focuses on the changes from
the 2001-10 into the 2011-20 decade which actually showed the largest climatic shift observed in this region. The paper closes

with a discussion of the limitations and the impacts of the drought as well as recent land-cover changes are discussed which
60 were partly induced by the drought itself.

2 Methods

2.1 Catchment water and energy balance

The catchment water balance describes how precipitation P received over the catchment area is partitioned into actual evapo-
transpiration E_T , runoff R at the catchment outlet and changes in water storage ΔS_w :

$$65 \quad P = E_T + R + \Delta S_w \quad (1)$$

Precipitation, evapotranspiration and water storage in soils and groundwater vary spatially and river discharge integrates runoff
processes over the catchment towards the outlet. With the simple water balance equation in (1), we assume that (i) precipitation
which is spatially averaged over the catchment area is the only water input into the catchment and that (ii) runoff observed at the
outlet comprises all outflow components. The latter two assumptions require that there are no water fluxes, e.g. by groundwater
70 or water management across the catchment area (Fan, 2019). When integrating over time the change in water storage in
catchments ΔS_w should converge to zero (Dyck and Peschke, 1995) and one can estimate catchment evapotranspiration by
 $E_T = P - R$. However, 5–10 years of data are recommended to reduce water storage effects (Zhang et al., 2001).

Evapotranspiration E_T is also part of the surface energy balance, here normalized by the latent heat of vaporisation $L =$
 $2.5 \text{ MJ kg}^{-1} \text{ K}^{-1}$:

$$75 \quad R_n/L = E_T + H/L + \Delta S_e, \quad (2)$$

with net radiation R_n , sensible heat H and an energy storage change term ΔS_e . Since available energy usually expressed as
 R_n/L is not well observed, we assume that it can be described by potential or reference evapotranspiration denoted as E_0
(Choudhury, 1999; Arora, 2002).

E_T couples the water and energy balances. As first consequence of the coupling we can state that E_T is limited by both
80 precipitation and available energy at the same time (Budyko, 1948). Thereby P and E_0 become the two main predictors of
the water and energy balance, which is well documented by the Budyko curve (Budyko, 1948). It describes how the relative
water balance, here E_T/P , can be well predicted by the aridity index E_0/P . The semi-empirical functions of Schreiber (1904);
Ol'Dekop (1911); Budyko (1948); Pike (1964) and others show good agreement with catchment water budget data from humid
to arid climates. The relative water balances of most catchments are close to the physical limits. However, variation in regional
85 catchment properties may affect the partitioning. For example steep orography or seasonality of precipitation may decrease E_T ,
while density and type of vegetation, plant accessible soil water tend to increase E_T under the same meteorological forcing.
These effects were summarized in a catchment parameter from which parametric Budyko curves were developed (Turc, 1961;
Mezentsev, 1955; Fu, 1981).

The well-known way to visualize the water and energy coupling is the Budyko plot. The aridity index E_0/P is plotted on the x-axis and the relative water partitioning E_T/P on the y-axis. The latter ratio should range between 0 and 1. Low values indicate that most of the rainfall is converted into runoff and the upper limit of 1 would indicate that all of the received precipitation evaporates.

2.2 Decomposition of climatic and land-surface impacts

The second consequence of the complementarity of water and energy balances is that changes in the aridity of a catchment will not only change the relative water balance $q = E_T/P$ but also the relative energy balance $f = E_T/E_0$, only in the opposite direction. This is illustrated in Fig.1 which plots the relative water balance on the x-axis and the relative energy partitioning on the y-axis. Any point in the range between 0 and 1 is a reasonable state of the relative water and energy partitioning. The diagonal represents the condition when the aridity index $E_0/P = 1$, with humid conditions above and arid conditions below the diagonal line.

The complementarity becomes apparent under a change in aridity. For example more precipitation will decrease E_T/P but increase E_T/E_0 , if the catchment follows a Budyko type of curve. In contrast, if there is solely a change in the catchment attributes without variation in the aridity index there will be a parallel shift to the line of constant aridity through the origin. An observed change in E_T (from point q_0, f_0 to point q_1, f_1) can be decomposed into a land surface change part (from point q_b, f_b to q_0, f_0 , red arrow in Figure 1) and a climate change part (from point q_1, f_1 to q_b, f_b , blue arrow). Renner et al. (2014) assumed that climatic changes lead to a shift in q and f perpendicular to the line of constant aridity. With this assumption the amount of evapotranspiration $E_{T,b}$ without any climatic changes can be computed from data before (index 0) and after the change (index 1):

$$E_{T,b} = P_0(f_0 f_1 q_0 + q_0^2 q_1) / (f_0^2 + q_0^2). \quad (3)$$

Then the climate-induced change in evapotranspiration can be calculated by $\Delta E_{T,C} = E_{T,1} - E_{T,b}$, as well as the remaining part denoted as the the land-surface induced change $\Delta E_{T,L} = E_{T,b} - E_{T,0}$, see Renner et al. (2014) and subsection B1 in the appendix for a detailed derivation.

By applying the water budget equation to the changes, i.e. $\Delta P = \Delta E_T + \Delta R$, we can also compute the amount of runoff change attributed to climatic changes:

$$\Delta R_C = R_1 - P_0 + E_{T,b}, \quad (4)$$

where, R_1 is the mean annual runoff in the second period and P_0 is the mean annual precipitation of the first period. The land-surface induced changes in runoff are derived by

$$\Delta R_L = P_0 - E_{T,b} - R_0. \quad (5)$$

where R_0 is the mean annual runoff of the first period. Note, that changes in runoff are mainly driven by changes in precipitation (Dooge, 1992) and secondly by changes in evaporative demand and land-surface conditions.

120 Other approaches to decompose climate and land-surface impacts on runoff were proposed by Wang and Hejazi (2011); Jaramillo et al. (2013); Jaramillo and Destouni (2015). These methods use Budyko functions to estimate the climatic change part. However, the uncertainties in quantifying catchment precipitation, runoff and energy demand are more relevant than the methodological differences.

2.3 Comprehensive visualization method

125 The decadal changes in catchment evapotranspiration are related to precipitation and potential evapotranspiration and plotted together in the joint water-energy balance diagram. This allows a visual interpretation of separate climatic and land-surface impacts on catchment evapotranspiration as illustrated in Fig. 1. Therefore decadal averages of the relative water (E_T/P) and energy (E_T/E_0) partitioning for each of the 71 catchments are calculated. Subsequently the decadal averages of both ratios for a given decade and each catchment are plotted in the joint water-energy balance diagram. The position of a catchment within
130 the plot is depicted as pie-chart. It holds information about the land-use types (see chapter 3.3) of each catchment.

The change of the relative water (E_T/P) and energy (E_T/E_0) partitioning towards the following decade is integrated as an arrow. The direction of the arrow reveals, if only climatic or land-surface changes occurred or if both types of changes were present. The arrow length refers to the magnitude of the change. In order to assess if a change is statistically significant a two sample Hotellings T2 test (Todorov and Filzmoser, 2009) is performed by sampling the annual data of each decade. Black
135 bold arrows highlight if the magnitude of a change in the decadal mean between two decades is larger than the year-to-year variability using a significance level of $\alpha = 0.1$. Gray thin arrows visualize the direction of an insignificant change.

Furthermore, a map is added with the outline of the analyzed Saxonian catchments to illustrate where the changes occurred. The arrows are also included in the map and start at the river gauge location. This is combined with spatial information of forest damage data (see chapter 3.3).

140 If several of these comprehensive visualization diagrams are plotted as sequence over a number of decades, it provides information about temporal as well as spatial changes in the catchment water balance. It allows the identification of general behaviour within a large catchment sample and may also reveal differences for certain catchment groups. At the same time the analysis of probable causes is possible.

3 Data

145 In this section details about the used datasets for the hydro-climatological analysis are provided. Note that this study updates the analysis presented by Renner et al. (2014) by adding data from 2010-2020. Thereby the same catchments and methods are used as in Renner et al. (2014). However, the hydrological and meteorological input data were updated containing data corrections and additional meteorological data due to digitization efforts.

3.1 Hydrological data

150 The study covers data of 71 river gauging stations with long term, high quality data records. The sites were chosen according to continuity, homogeneity, small direct impacts by water management due to dam operation or large mining activities such as in the Lausitz region in the East of Saxony. The sampling of the catchments thus slightly over-represents the higher regions of Saxony. It is also important to note that the sample includes nested catchments which share parts of the catchment area and are thus not independent. All catchment attribute data such as river names, catchment polygons etc were used from Renner
155 et al. (2014). The map in Fig. 2 provides an overview. All time series data were updated. Key data provider is the Saxon State Department for Environment, Agriculture and Geology (Landesamt für Umwelt, Landwirtschaft und Geologie, LfULG), which provided daily runoff time series data via the webportal <https://ida.sachsen.de> and by e-Mail. The river gauge Greiz/Weiße Elster is operated by the Thuringian State Department of Environment, Mining and Nature Conservation (TLUBN) and time-series data were obtained directly from the authority. All data were checked graphically for correctness. Data screening for
160 homogeneity and dominant influences like water management was performed in Renner et al. (2014) and such catchments were removed from the dataset. This was necessary because we employ the simple water balance equation 1 to estimate E_T .

At first daily discharge data were aggregated to monthly averages with at least 20 non-missing values per month. Then annual averages were computed using the hydrological year from November to October. Finally decadal averages and statistics were derived when at least 6 years of complete data per decade were available. Runoff yield ($mm\ yr^{-1}$) was calculated from
165 average discharge records ($m^3\ s^{-1}$) by multiplying with the number of seconds per year and dividing by the catchment area.

The study period covers 1951-2020. For computing the long-term average runoff all available annual data per site were used. Note that not all runoff time series cover the whole period. In a few catchments river gauge relocation occurred, leading to a change of the station-id. At the following sites the time series of the prior gauge and the current station were merged for this study: Buschmühle/Kirnitzsch, Piskowitz 2/Ketzerbach and Annaberg/Sehma. Table A1 in the appendix provides an overview
170 of the river gauges and the time series availability.

3.2 Meteorological data

Precipitation and potential evaporation were derived from meteorological station data, interpolated on a spatial grid, from which catchment averages were extracted. Precipitation is well observed in the study area with 845 sites used in the regionalisation and 366 sites with more than 40 years of data, shown as circles in Fig. 3a. The size of the circles reflects the available observation
175 years. Key operators are the German weather service, the Landestalsperrenverwaltung Sachsen (river dam authority of the Federal State of Saxony) as well as the Czech and Polish Hydrometeorological Institutes. All meteorological station data were acquired at daily resolution from the REKIS (rekis.org) data service, which is a platform for regional meteorological data service covering the Federal States of Saxony, Thuringia and Saxony-Anhalt.

The original daily precipitation data was added up to annual totals using the hydrological year from November to October.
180 The annual station data were then interpolated on a 1.5 km grid using an automatic universal Kriging procedure provided as

R package automap (Hiemstra et al., 2009). Thereby station altitude is used as the dependent variable. The procedure was validated by Renner and Bernhofer (2011) and Renner et al. (2014).

To model potential evaporation (E_0) we use the FAO-56 method for grass reference evapotranspiration (Allen et al., 1994). This is a simplification of the Penman-Monteith equation and Allen et al. (1994) provide methods to substitute net radiation data and ground heat flux which are not commonly available. Here, the daily station data was first aggregated to monthly averages to compute E_0 from temperature (daily mean, minimum, maximum), sun shine duration, relative humidity and wind speed data. The locations of the 115 climate stations (40 sites have more than 40 years of data) are shown as circles in Fig. 3b. Similar to precipitation spatial interpolation was performed on available annual (hydrological year) station data and interpolated onto a 1.5 km grid using universal Kriging with altitude as dependent variable.

190 3.3 Land-use and forest damage data

The remote-sensing based Corine Land Cover data set (© European Union, Copernicus Land Monitoring Service, European Environment Agency (EEA)) was used for land-use classification of the catchments and as a proxy for forest damage. Snapshots for different years are available.

From the Corine snapshot of the year 2000 we classified the land-use types of forested and near natural vegetation classes as “forest”, merged all agricultural and grass lands in the class “agricultural” and grouped the remaining as “others”, mainly built-up areas. The first Corine snapshot from the year 1990 was used to estimate the relative area of forest damage per catchment, following Renner et al. (2014). The Corine land cover class Transitional Scrub-Forest (324) includes areas of damaged forests (Bossard et al., 2000). This snapshot of 1990 captures the period of maximal forest damage extent in Central Europe. Based on this data Renner et al. (2014) calculated the relative affected area for each catchment and classified the catchments in groups with different magnitudes of forest damage. The class major forest damages with a threshold of $> 20\%$ of catchment area was selected following Bosch and Hewlett (1982) who reported significant streamflow increases in paired catchment studies after clear-cuts when more than 20% were affected. The threshold category of 2%-20% was applied to select catchments where a measurable area was affected. Renner et al. (2014) found that 38 % of the catchments have no forest damage ($< 0.01\%$), 21 % have minor affected areas (0.01–2 %), 31 % have considerable damaged areas (2–20 %), and 10 % of the catchments have damaged areas larger than 20 %. The respective catchments are shown in the map in Fig. 4.

In order to get an idea of the forest damage over time we also used the Corine snapshots of the year 2000 and 2012. For the pre-satellite era maps of canopy damage data available for the years 1960, 70, 80 and 1990 were used. Canopy damage was assessed by needle and leaf losses in the canopy (SMUL, 2006) and follows a measurement protocol defined in the former German Democratic Republic (Forstprojektierung, 1970).

4.1 Long-term hydro-climatology of Saxony

In this section the long-term average (1951-2020) hydro-climatology of Saxony is illustrated based on the established dataset, see also Table A1 in the appendix.

Annual average precipitation is 830 mm yr^{-1} across the 71 catchments and ranges between 607 and 1151 mm yr^{-1} . For the whole of Saxony an average of 712 mm yr^{-1} is estimated. The difference is due to the sampling of the catchments, which over-represent higher regions of Saxony. Precipitation shows a distinct north to south increasing gradient which is linked to altitude. Another, although weaker gradient is induced by the transition of maritime to continental climate which results in decreased precipitation from west to east (Fig. 3a and Fig. 5a).

The FAO grass reference potential evapotranspiration E_0 across the study catchments ranges between 572 and 738 mm yr^{-1} and is on average 669 mm yr^{-1} . E_0 is negatively correlated with elevation (see Fig. 3b).

The spatial pattern of runoff is dominated by topography and precipitation with most catchments receiving a large amount of its water from the headwater catchments in the Ore Mountains, the Lausitzer Bergland, and Izer mountains, see Fig. 3c. The lowest annual runoff values are found in the lowland catchments in the North (minimum 59 mm yr^{-1}) and highest in the headwater catchments in the South (maximum 799 mm yr^{-1}). On average runoff is 372 mm yr^{-1} across the 71 catchments.

The difference of $P - R$ is used to estimate actual catchment evapotranspiration E_T (Fig. 3d). E_T is on average of all catchments 461 mm yr^{-1} and 572 mm yr^{-1} on maximum. The lowest values of annual E_T are found in the high headwater basins (minimum 219 mm yr^{-1}). In general E_T has a smaller spatial variation than runoff. All long-term average data is summarized in detail in Table A1.

The average water and energy partitioning of the catchments in Saxony is illustrated in Fig. 5a) and Fig. 5b), which is the Budyko plot. Each catchment is depicted as pie chart, representing the relative portion of forested and agricultural area. Furthermore contour lines are included in Fig. 5a), derived from the mean catchment altitude.

The relative water partitioning E_T/P is placed on the x-axis in Fig. 5a) as well as on the y-axis in Fig. 5b). The long-term averages of E_T/P vary between 0.22 and 0.91 and are on average 0.57 .

On the y-axis of Fig. 5a) the relative energy partitioning E_T/E_0 is plotted, also ranging between 0 and 1 . This ratio varies between 0.37 and 0.83 , with most catchments clustering around the average value of $E_T/E_0 = 0.69$. Plotting both ratios together combines the two key meteorological forcing variables P and E_0 . Four general examples for the aridity index $\phi = E_0/P$ are included in Fig. 5a) as lines through the origin. When precipitation equals potential evaporation $P = E_0$ this would fall on the diagonal. More humid conditions would be above the diagonal of $\phi = 1$, more arid conditions fall below this diagonal. All together catchments at higher altitudes (see altitude contour lines in Fig. 5a) are more humid with $E_T/P < 1$ and have a larger portion of forested area. Catchments with $E_T/P > 1$ are generally found at lower altitudes and dominated by agriculture land use.

The aridity index ϕ is also visualized on the x-axis in the Budyko plot (see Fig. 5b) and ranges between 0.50 at the highest catchment and 1.21 at the lowest catchment. On average we find a value of $E_0/P = 0.82$. The Budyko function (Budyko,

1948) is plotted as reference allowing the prediction of catchment E_T from the aridity index. The distribution of the data generally agrees with the Budyko function. However, there is an overestimation of E_T/P at the more humid catchments and an underestimation for the drier catchments.

4.2 Annual and decadal dynamics in joint water and energy balance

4.2.1 Variability of meteorological forcing

The variability of the meteorological forcings P and E_0 will be analyzed using the annual mean values as well as the decadal means (see figure 6). Generally there is a larger variability of precipitation P both across catchments and from year to year than E_0 . The decadal mean precipitation over all basins was always higher than of E_0 . However, the last decade (2011–20) shows a remarkable increase in E_0 , being the highest decadal average so far (710 mm/yr) and getting closer to P (814 mm/yr). At annual time scale about 17% of the years had a negative climatic water balance, where catchment annual average E_0 was larger than P : 1951, 1953, 1959, 1963, 1964, 1976, 1982, 1990, 1991, 2003, 2015, and 2018. The occurrence of these negative values does not have a clear trend, but indicates that there have already been dry conditions in the 1950s and 1960s.

4.2.2 Variability of catchment evapotranspiration

The water budget residual $P - R$ is analyzed as proxy for basin scale E_T , which may include the unknown influence of past water storage changes ΔS_w . To reduce the effect of storage changes decadal averages of E_T are derived, presented in Fig. 7. In the figure catchments are grouped by the percentage of areal forest damage extent as shown in Fig. 4.

The boxplots in Fig. 7 depict variation of the decadal average of E_T of all catchments in each group. During the first decade 1951–60 there is no large difference between these four groups. Between 1961–80 the heavily damaged group experienced a large decline in E_T , while the two groups with low damage effects show little variation. The catchments with moderately damaged forest are in between and show a minor decrease in E_T during the decades from 1971–90. From 1991–2010 the effect is reversed for all catchments with moderately as well as heavily damaged forest and is reflected in a pronounced increase in E_T (see Fig. 7). And while the positive trend in E_T continues for the group with considerable forest damages, the low damaged catchments show a small but consistent decline during the last decade.

4.3 Decomposition of climate and land-surface effects

The changes in runoff are decomposed in a climate and a land-surface change component which together sum up to the observed change. The decomposition method by Renner et al. (2014) introduced in the methods section quantifies the contributions of climatic changes in P and E_0 as well as land-surface changes in E_T in retrospective. Figure 8 presents the results of the decomposition for catchment runoff. The gray box-plots depict the changes in the observed runoff from one decade to the next for all the catchments. There is not much variation of the median of the changes from 1951 to 2010 and no clear tendency for the direction of change. The observed changes in runoff varied between -50 and 50 mm/yr across the Saxonian catchments.

In the last decade observed runoff decreased the most across the majority of catchments with a median decline of -70 mm/yr, which is 8% of mean annual precipitation.

The decomposition quantifies the contributions of climatic changes in P and E_0 on runoff, visualized as blue box-plots in Fig. 8. Thereby the climatic driven runoff changes were relatively small in the first five transitions with median changes between -13 mm/yr towards the 80s and +23 mm/yr towards the 2000s. The climate driven changes became the dominant cause in the last transition with a median change of -65 mm/yr and thus explaining the observed decline in runoff in most catchments in Saxony. However note, that the transitions from the 80s to the 90s and from the 90s to the 2000s showed two consecutive increases in the climatic runoff change component due to increases in precipitation.

The red box-plots in Fig. 8 represent the land-surface change component. The outliers show that in a few catchments it dominated the change in runoff during the past decades. There have been two consecutive, but small increases of 14 mm/yr in runoff due to land-surface conditions towards the 70s and 80s. They were followed by a strong and more widespread decline in the 1990s with a median decrease of -38 mm/yr in the land-surface component.

4.4 Decadal dynamics of water-energy partitioning from 1951 to 2000

The decadal changes within the 70 year period are depicted by six panels in Fig. 9. In each panel the changes in the water and energy partitioning ratios from one decade to the next is visualized by the direction of the arrows as well as the length. The displayed pie-charts always refer to the same land-use types derived from the Corine dataset of the year 2000. The number of catchments taken into account varies due to the different availability of data.

Panel a) shows a set of 28 catchments in the decade 1951-60 with transition into the next decade. No significant changes in E_T/E_0 and E_T/P are apparent. In Fig. 9b) the transition from the 60s to the 70s is visualized. The number of available catchments increases to 45. The changes within the catchments are either pointing towards the origin of the diagram or in the opposite direction, but are not significant for most of the catchments. Only two neighboring head water catchments in the upper Ore mountains (Rothenthal/Natschung and Zöblitz/Schwarze Pockau) have significant decreases in both ratios indicating an impact of land-surface changes on E_T . Also other neighboring catchments show decreases which hints at a similar cause. These declines in E_T are mostly in the more humid head water catchments and are part of the group with dominant forest damage, as presented in Fig. 7.

The extent and severity of forest damages increased in the 1980s, see inlay maps of forest damage in Fig. 9c). The transition from the 1970s to the 80s was analyzed for 69 catchments and can be seen in Fig. 9c). The development of the two ratios appears to be dominated by this ongoing spread of forest damages, with many of the arrows showing directions of land-surface changes. For two catchments in the upper Ore mountains (Rehefeld/Wilde Weisseritz and Wolfsgrund/Chemnitzbach) the declines in E_T/E_0 and E_T/P are significant, while four catchments experience a significant increase in the two ratios.

In contrast to the prior transitions the changes are almost consistent for all of the catchments in the transition from the 80s to the 90s depicted in Fig. 9d). It reveals the most drastic and coherent shifts observed in the whole period, with dominant and significant increases in both ratios in 11 out of 69 catchments. For some of the catchments this even means a reversed direction of changes for E_T/E_0 and E_T/P compared to earlier periods. In 1990 forest damage reached its largest extent in Saxony.

In addition increases in precipitation and potential evaporation led to a further increase in catchment evapotranspiration, see Fig. 6. The following decadal change from 1991–2000 into 2001–2010 is visualized in Fig. 9e). Now there are two groups with a different development in the water and energy partitioning. While the mountainous catchments, which were highly effected by the vast forest damages, still have a continuing increase in E_T/E_0 and E_T/P , the lower-lying regions generally show less changes with no clear direction. Fig. 9e) also reveals five mountainous catchments with significant recovery of both ratios. In contrary the lowland catchment Seerhausen/Jahna has a significant decrease in the aridity index due to higher precipitation.

In general the land-surface attributed impacts were dominating the development of the two ratios E_T/E_0 and E_T/P for the time period between 1951 and 2010. This analysis has been part of the previous work of Renner et al. (2014). The picture changes, when it comes to the transition from 2001–10 to the last fully observed decade of 2011–20, which is presented in the next section.

4.5 Decadal dynamics of water-energy partitioning from the 2000s to the 2010s

The last decadal transition is dominated by the meteorological conditions of 2011–20, which have been much drier than the two decades before. This results in the highest number of significant changes (16 out of 64, 25%) in catchment water and energy partitioning, presented in Fig. 9f). The direction of change is now mainly orthogonal to the prior transitions, meaning that climate attributed impacts effecting the ratios of E_T/E_0 and E_T/P the most.

Two more figures are included to illustrate the noticeable changing conditions from 2001–2010 to 2011–20 in Saxony. Fig. 10 visualizes the decadal changes as bar plot for all analyzed water balance components and all 71 catchments. The reduction in precipitation (blue bars in Fig. 10) on the one hand and the uptake in the potential evaporation (red bars in Fig. 10) on the other hand are apparent. The widespread and large decline in annual precipitation resulted in a median decrease of -59 mm/yr. It was rather pronounced in the Ore mountains reaching up to -143 mm/yr and the Izer mountains, which feed the upper Lausitzer Neiße river, see Fig. 11a). The maps in Fig. 11a) combine precipitation changes (filled polygons of the catchment area) with changes in observed runoff (filled triangles at the river gauge locations). The same color scale is used to highlight, how much the regional precipitation decrease affected runoff. This combination shows that catchment runoff broadly followed precipitation. In 56 out of 64 catchments the observed runoff decreased with a median decline of -73 mm/yr.

Focusing at the significant changes three different types can be observed: (i) five catchments in the Ore mountains show continuing recovery effects with significant increases in both ratios, marked by letter 'r' in Fig. 10. These catchments have had large areal extents of forest damages in which the recovery led to catchment evapotranspiration rates now being even higher than during the 1951–60 decade, as can be seen in Fig. 7. Type (ii) are catchments, especially in the Mulde river basin, which experienced quite significant decreases in annual mean precipitation. Nine of the catchments show significant movements in the climatic change direction, marked by letter 'c' in Fig. 10. These catchments follow the Budyko curve and retain high evapotranspiration leading to increases in the relative water balance. Type (iii) are the lowland catchments in northern and eastern Saxony recognizable by the downward pointing arrows in Fig. 9f) and marked by letter 'd' in Fig. 10. They represent a decrease in the relative energy partitioning without a change in the relative water balance. Only one catchment (Zescha/Hoyerswerdaer Schwarzwasser) experienced a significant change. Here the increase in potential evaporation did not

lead to an increase catchment evapotranspiration. This highlights that both, climate changes (i.e. higher aridity) and land-surface changes, co-occurred in this transition period. A similar behavior was also found by Pluntke et al. (2023) for a small, forested research catchment in the south of Dresden, Saxony.

345 The map in Fig. 11a) also highlights notable exceptions: (i) there are a few catchments in the Upper Ore mountains with larger decreases in runoff than in precipitation, and (ii) four catchments located in the East of Saxony having notable increases in runoff despite increasing aridity.

In addition to less precipitation amounts there was a substantial increase in FAO reference potential evaporation across all regions with a median increase of 36 mm/yr, see red bars in Fig. 10. The map in Fig. 11b) presents a rather uniform spatial
350 distribution of the change in potential evapotranspiration with slightly larger increases in lower parts of Saxony. In this map the changes in actual catchment evapotranspiration E_T (derived by $P - R$) are added with the same color scale. The response of actual evapotranspiration is rather mixed. 27 out of 64 catchments depict a reduced E_T with -10 to -67 mm/yr, see also the orange bars in Fig. 10. These catchments are found in the lower regions of Saxony. Especially the smaller tributaries to the Elbe River and catchments in the Lausitz show these reductions (triangles in Fig. 11b). Furthermore, 16 catchments experienced
355 increases in E_T larger than 10 mm/yr. In five of these catchments in the upper mountain regions the increase even exceeded 50 mm/yr. The remaining 21 catchments have only smaller changes in E_T .

The already discussed decomposition method was applied to quantify the climatic and the land-surface impact on runoff changes. The climatic impact of the observed changes in precipitation and potential evaporation on runoff is visualized in the map in Fig. 11c). The catchments in the Mulde river basin show the largest decreases in the middle of Saxony and there are
360 also larger effects seen in the upper Lausitzer Neiße river basin. Note that these catchments have a more humid climate and higher runoff ratios, which generally leads to a higher sensitivity of absolute runoff amount to changes in precipitation (Dooge, 1992; Renner et al., 2012). In the decade of 2011-2020 the increasing dryness has had the strongest climate impact on runoff in Saxony in the whole period analyzed here (see Fig. 8).

Fig. 11d) shows the spatial distribution of the land-surface impacts on runoff. Many catchments reveal only minor effects
365 of land-surface changes. However, there are interesting exceptions. The group of catchments with the large increases in E_T and corresponding pronounced declines in runoff is the one with the most land-surface impacts. The catchments are all located in forested regions, where the recovery of past forest damages is still ongoing (Ore Mountains, the Lausitzer Bergland) as presented in Fig. 11d). There are also catchments with increasing runoff attributed to land-surface impacts as shown in Fig. 11d). These are predominantly lowland catchments with higher aridity indices and no decrease in runoff as expected from
370 the climatic conditions, resulting in downward arrows in Fig. 9f). The discussion of the potential causes is left for the next section.

5 Discussion

In this study we identified and estimated impacts of climate and land-surface conditions on catchment water balance in the Federal State of Saxony. First we discuss the attributed climatic and land-cover impacts on catchment water balance and close
375 with an explanation of the potential limitations of the study.

5.1 Climate change becomes a significant driver of water balance

The drought from 2018–20 in Europe was exceptional in terms of spatial extent, duration and intensity (Rakovec et al., 2022; Büntgen et al., 2021). Compared to previous drought events, it was exceptionally warm (Rakovec et al., 2022) and was probably caused by anthropogenic warming (Büntgen et al., 2021). The drought assessment of Rakovec et al. (2022) used a modeled
380 soil moisture index based on meteorological data reconstructions from 1766 to 2020 for Europe. According to Rakovec et al. (2022) northern Saxony and the Lausitz region exhibit a soil moisture drought of 24 months. Low flow statistics showed long duration of hydrological drought conditions which were also exceptionally low in Saxony. The Saxon State Department of Environment, Agriculture and Geology reported that during summertime of 2018–19 and 2020 40 % to 70 % of 146 runoff gauges in Saxony showed values below the mean annual low flow (Franke and Rühle, 2022). Due to the long duration also
385 mean annual flows of 2018–20 were lower than the long term mean (Franke and Rühle, 2022).

The widespread runoff decline in Saxony represents the largest and most coherent decadal shift in runoff and the decomposition methods highlight that this reduction is caused by climatic changes - that is the reduction of rainfall and the increase in atmospheric demand for water (see Fig. 6 to Fig. 8). For the majority of catchments the reduction of runoff can thus be predicted by knowing the changes in annual mean precipitation and potential evaporation. However, there were notable exceptions
390 in certain regions which are discussed next.

5.2 Land-surface impacts on water and energy partitioning

Any change in runoff which cannot be predicted with the climatic signal is attributed to land-surface changes. In the last decade with the shift towards more arid conditions we observed two different reactions in catchment runoff which were attributed to land-surface impacts. While some catchments showed stable or even increasing runoff, others experienced very strong declines
395 in runoff which could not be explained by the dryness, see section 4.5.

5.2.1 Hydrological response to forest recovery under dry conditions

Large scale forest dieback due to effects of air pollution occurred in Central Europe with the industrial use of lignite coal burning with significant regional impacts starting in the 1960s (Mazurski, 1986; Maas and Grennfelt, 2016). Renner et al. (2014) documented on the one hand that these forest damages led to strong reductions in catchment actual evapotranspiration
400 lasting over two to three decades and on the other hand that an increase in E_T correlates with recovery of forests. This link of forest damage and substantial declines in catchment evapotranspiration is updated in this study and clearly revealed by decadal averages of E_T (see Fig. 7). The effect becomes apparent when the catchments are grouped by the percentage of forest damage.

Since 1990 the recovery of these areas led to increases in catchment evapotranspiration and reduced the gap between disturbed and undisturbed catchments in Saxony. Nevertheless, the analysis of climatic and land-surface attributed changes on runoff, visualized in Fig. 9f) and Fig. 11d), point out that in a few mountainous catchments recovery of E_T is still ongoing. The largest increases were seen in the Eastern Ore Mountains at the neighboring catchments Baerenfels/Pöbelbach and Rehefeld/Wilde Weißeritz, followed by the Middle Ore Mountains (Rauschenbach/Rauschenfluß, Deutschgeorghthal/Rauschenbach) and a head water catchment in the Western Ore mountains (Sachsengrund/Große Pyra). At these catchments the strong increases in E_T correlate to exceptional decreases in annual mean runoff. Hereby, it is noticeable that the change in runoff is larger than the observed decrease in precipitation. Furthermore, the increase in E_T is higher than in E_0 (see Fig. 10). This strongly hints at the recovery of transpiration and evaporation from interception due to vegetation regrowth. The long-term time series of catchment evapotranspiration presented in Fig. 7 shows that the recovery in terms of E_T in these catchments is almost completed. On the one hand the magnitude of E_T is now at the same level or even higher than prior to the period of large scale forest damages. On the other hand E_T is now in a similar range compared to catchments without severe forest damages.

5.2.2 Potential causes of constant or increasing runoff

At first sight it seems contradictory that runoff remained stable or increased even though precipitation decreased and potential evaporation increased. A potential hypothesis explaining this behavior may be the reduced transpiration and evaporation from interception due to reduced growth or plant damage. This hypothesis is supported by forest inventory data from Saxony (Otto et al., 2022) showing that by far the largest amount of damaged wood due to bark beetle infestations occurred in 2018-20 within more than 70 years of observations. The combination of warm and dry conditions reduced the fitness of needle leaf trees, which then led to an outbreak of bark beetles in 2018 continuing to 2020 throughout Saxony. Thereby the forests in eastern Saxony were most affected, which is inline with regions where we observed stable or increasing runoff conditions and thus decreases in catchment actual evapotranspiration (see Fig. 11).

Apart from forests, there is evidence of reduced crop yields during the drought year 2018 (Statistisches Landesamt Freistaat Sachsen, 2022). This year showed the highest aridity index in the analyzed period. The reduction in yields corresponds to the depletion of soil moisture during the long and severe drought, especially in northern and eastern Saxony (Rakovec et al., 2022). Water limitation causes reduced plant growth as well as reduced transpiration via stomata closure (Teuling et al., 2010). However, more research is required to quantify the contribution of reduced transpiration of short vegetation such as grass and crops to reduced catchment evapotranspiration. In either case a future recovery of vegetation, especially of forests, will likely increase E_T and thus lead to less runoff from such catchments if precipitation will remain low.

An alternative hypothesis for the stable runoff under increasing aridity could be the contribution of groundwater release as baseflow. This would be a change in the water storage term in equation 1, which we tried to avoid by using the decadal averaging. This alternative hypothesis is supported by the fact that the catchments with nearly constant runoff are all located in regions with highly conductive aquifers, see the map in Figure A1 in the appendix. Studies of groundwater levels also found a record breaking groundwater drought in the last decade beginning in 2013 (Müller et al., 2023). Here we recommend

further research to analyze runoff components and the integration of groundwater levels into the analysis to actually test these hypothesis.

5.3 Limitations

440 5.3.1 Catchment selection and influences by water management

The selection of catchments aimed to capture a large part of Saxony, while excluding catchments with direct hydrological impacts such as reservoirs with storage volumes large enough to influence the annual water balance. Also catchments which are impacted by large water extractions and inputs have been excluded. These exclusions were done using meta data as well as homogeneity tests as explained in Renner et al. (2014). Nevertheless, there may still be artifacts in single catchments due to these issues. For example the catchments Rehefeld/Wilde Weisseritz, Baerenfels/Pöbelbach and Geising/Rotes Wasser are adjacent to the catchment of the drinking water reservoir Altenberg, which was established in the 1980s. Thus, the operation of trenches could directly influence the water budget of these catchments. Since the reservoir is not directly part of these three catchments we left them in the dataset, but excluded downstream river gauges whose catchments are dominated by reservoirs, such as the Rote Weisseritz river. For a full treatment of catchments, which are affected by water management there is the need to collect and share data on reservoir operation, water abstractions and transitions.

5.3.2 Catchment integration and groundwater transfers

Closing the water balance is a key assumption to estimate catchment evapotranspiration. Water fluxes across catchment boundaries, for example by groundwater flows contributing to downstream parts of the river basin (Fan, 2019), were not considered in this study. Both natural and anthropogenic cross boundary fluxes are often difficult to quantify and thus lead to biases in $E_T = P - R$ and especially absolute values must be taken with care. However, a reasonable assumption is that unmeasured water fluxes across catchment boundaries may not change over time. Thus the decadal change signals in runoff and catchment evapotranspiration are likely unaffected by this type of uncertainty. Nevertheless, we recommend to address this issue in further studies which explicitly include groundwater fluxes and water transfers across surface catchment boundaries.

5.3.3 Runoff estimation

460 Runoff was computed from daily discharge time series data obtained from the monitoring authorities. Discharge is usually estimated from continuous water level records and a water level – discharge relationship. It is established by point measurements usually done every one or two months. These relationships can change over time due to changes in the profile, such as weed growth and weed control but also by flood deposits or ice conditions. Hence, runoff estimates can be quite uncertain under low flow conditions as well as under flood conditions when the river leaves its profile (Coxon et al., 2015). By averaging the impact of these uncertainties they can be reduced but not completely removed. Again the large sample of catchments analyzed here shows that there are consistent changes across catchments. It highlights that this uncertainty is not a major issue in interpretation of the results.

5.3.4 Temporal resolution

The analysis is based on annual averages and thus neglects any subseasonal changes such as hydro-meteorological shifts in spring season (Bernhofer et al., 2008; Renner and Bernhofer, 2011; Berghuijs et al., 2014; Ionita et al., 2020). These seasonal shifts are out of scope of this study and should be investigated in further studies. Further decadal averages are used for the decomposition of climate and land-surface impacts to reduce the potential impact of interannual storage changes, which can be large in groundwater dominated catchments. Changes in storage, if present at the interdecadal time-scale, would certainly influence the climate and land-surface decomposition, since a storage change would be mistaken for a change in evapotranspiration (see Eq. 1). This potential misinterpretation must be kept in mind when applying the decomposition framework. However, the results of this study, providing spatially and temporally consistent trajectories in the relative water and energy balances, indicate that storage influences are a minor issue.

5.4 Comparison of decomposition methods

Last but not least the decomposition method itself is discussed. For validation of this method the results are compared with the algorithm presented by Wang and Hejazi (2011). They employ a parametric Budyko function, here the form of Mezentsev (1955), to estimate the climatic and the land-surface change part. First they estimate the catchment parameter n using the data of the first period. Then the parametric Budyko function is used to derive E_T based on precipitation and potential evaporation of the second period. The difference to the observed change in E_T is the land-surface induced change on E_T . The input data to both methods is the same. One key difference is that the method of this study uses a linear approximation whereas Wang and Hejazi (2011) employ a non-linear Budyko function, which addresses problems in this non-linear space for example when approaching the physical water and energy limits. A second, probably more relevant difference is the order of the decomposition parts. While Wang and Hejazi (2011) first computes the climatic part this study derives the land-surface part first as depicted in Fig. 1. The differences between the two methods are illustrated in Fig.B1 (see chapter B2 in the appendix). In most cases the differences are minor with a high correlation ($R^2 > 0.93$) and no obvious bias of the attributed changes in runoff and E_T . Hence, uncertainties in the input data are arguably more relevant than methodological issues.

6 Conclusions

Here we employed a coupled water and energy balance framework to analyze changes in the hydro-climatology of the federal state of Saxony in Germany. The large set of catchments with long term hydrological monitoring data in combination with a dense network of meteorological stations allowed a comprehensive study on how climate and land-surface changes impacted runoff. During the last decade from 2011-20 a long and extreme drought event led to significant changes in catchment runoff, which were quantified using a decomposition method. The decomposition highlighted the impacts of land-surface changes in a number of catchments revealing two main and spatially distinct patterns. First, recovery from past forest disturbance leads to a reduced runoff and intensifies the already existing negative trend due to a lack in precipitation. Second, impacts of the drought

on vegetation health are severe and affect the water balance decisively. This is especially relevant for forested regions which
500 have experienced huge amounts of damaged wood due to bark beetle infestations reducing catchment E_T .

We conclude that the hydro-climatology of Saxony has become rather non-stationary leading to methodological and practical
challenges for water resources management. Saxony is located in a region where climate models predict no clear trend. It is in
the transition of the "dry getting drier and the wet getting wetter" paradigm. The past years point out, that this does not mean
there are no changes. Instead we may prepare for more variation with extended phases of dry and hot conditions and increasing
505 number of intense rainfall events, which will affect water redistribution and thereby the average fresh water availability.

Code and data availability. Aggregated data and code to perform analysis and diagrams can be acquired from the first author.

Author contributions. MR conceived the study, acquired data and performed analysis. MR and CH discussed the results and drafted the
outline of the manuscript. CH provided information on anthropogenic influences on catchment water budgets and hydrogeology. MR wrote
the first draft which was reviewed and edited by CH.

510 *Competing interests.* The authors have no competing interests.

Acknowledgements. We thank all data providers for sharing their data. In particular the REKIS.org initiative which collects meteorological
data from different services for Saxony and bordering countries. MR is grateful to Rico Kronenberg (TU Dresden) for providing the daily
data and Philipp Körner (IAMK GmbH, Dresden) for discussion on reference evapotranspiration. For the hydrological data we thank the
Landesamt für Umwelt, Landwirtschaft und Geologie (LfULG) for providing most of their discharge data on the website as well as Heike
515 Mitzschke (Landeshochwasserzentrum, LfULG) for sending additional data and explanations on river gauge location data. Further we thank
Ralf Haupt (Hochwassernachrichtenzentrale, Thüringer Landesamt für Umwelt, Bergbau und Naturschutz) for providing the data of gauge
Greiz/Weiße Elster. Finally MR thanks Christian Bernhofer and Thomas Plunkte (TU Dresden) for discussions on ongoing changes related
to the Wernersbach research catchment, which raised the motivation to extend the previous study from 2014.

References

- 520 Allen, R., Smith, M., Pereira, L., and Perrier, A.: An update for the calculation of reference evapotranspiration, *ICID bulletin*, 43, 35–92, 1994.
- Arora, V.: The use of the aridity index to assess climate change effect on annual runoff, *Journal of Hydrology*, 265, 164–177, 2002.
- Berghuijs, W. R., Woods, R. A., and Hrachowitz, M.: A precipitation shift from snow towards rain leads to a decrease in streamflow, *Nature Climate Change*, 4, 583–586, <https://doi.org/10.1038/nclimate2246>, 2014.
- 525 Bernhofer, C., Goldberg, V., Franke, J., Häntzschel, J., Harmansa, S., Pluntke, T., Geidel, K., Surke, M., Prasse, H., Freydank, E., Hänsel, S., Mellentin, U., and Küchler, W.: Klimamonographie für Sachsen (KLIMOSA) – Untersuchung und Visualisierung der Raum- und Zeitstruktur diagnostischer Zeitreihen der Klimatelemente unter besonderer Berücksichtigung der Witterungsextreme und der Wetterlagen., Sächsisches Staats-Ministerium für Umwelt und Landwirtschaft (Hrsg.), Sachsen im Klimawandel, Eine Analyse, 211, 2008.
- Bosch, J. M. and Hewlett, J. D.: A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration, *Journal of Hydrology*, 55, 3–23, [https://doi.org/10.1016/0022-1694\(82\)90117-2](https://doi.org/10.1016/0022-1694(82)90117-2), 1982.
- 530 Bossard, M., Feranec, J., and Otahel, J.: CORINE land cover technical guide: Addendum 2000, European Environment Agency Copenhagen, 2000.
- Budyko, M.: Evaporation under natural conditions, *Gidrometeorizdat*, Leningrad, English translation by IPST, Jerusalem, 1948.
- Büntgen, U., Urban, O., Krusic, P. J., Rybníček, M., Kolář, T., Kyncl, T., Ač, A., Koňasová, E., Čáslavský, J., Esper, J., Wagner, S., Saurer, M., Tegel, W., Dobrovolný, P., Cherubini, P., Reinig, F., and Trnka, M.: Recent European drought extremes beyond Common Era background variability, *Nature Geoscience*, 14, 190–196, <https://doi.org/10.1038/s41561-021-00698-0>, number: 4 Publisher: Nature Publishing Group, 2021.
- 535 Choudhury, B.: Evaluation of an empirical equation for annual evaporation using field observations and results from a biophysical model, *Journal of Hydrology*, 216, 99–110, 1999.
- 540 Coxon, G., Freer, J., Westerberg, I. K., Wagener, T., Woods, R., and Smith, P. J.: A novel framework for discharge uncertainty quantification applied to 500 UK gauging stations, *Water Resources Research*, 51, 5531–5546, <https://doi.org/10.1002/2014WR016532>, 2015.
- Dooge, J.: Sensitivity of runoff to climate change: A Hortonian approach, *Bulletin of the American Meteorological Society*;(United States), 73, 2013–2024, 1992.
- Dyck, S. and Peschke, G.: *Grundlagen der Hydrologie*, 1995.
- 545 Fan, Y.: Are catchments leaky?, *WIREs Water*, 6, e1386, <https://doi.org/10.1002/wat2.1386>, 2019.
- Forstprojektierung: Aufnahme von Schadstufen bei Rauchschäden Betriebsregelungsanweisung - BRA IV / 320, VEB Forstprojektierung, 1970.
- Franke, J. and Rühle, K.: *Wetter trifft auf Klima, Jahresrückblick 2021*, Tech. rep., Landesamt für Umwelt, Landwirtschaft und Geologie, Freistaat Sachsen, 2022.
- 550 Fu, B.: On the calculation of the evaporation from land surface, *Scientia Atmospherica Sinica*, 5, 23–31, 1981.
- Gentine, P., D’Odorico, P., Lintner, B. R., Sivandran, G., and Salvucci, G.: Interdependence of climate, soil, and vegetation as constrained by the Budyko curve, *Geophysical Research Letters*, 39, L19 404, <https://doi.org/10.1029/2012GL053492>, 2012.
- Gudmundsson, L., Boulange, J., Do, H. X., Gosling, S. N., Grillakis, M. G., Koutroulis, A. G., Leonard, M., Liu, J., Müller Schmied, H., Papadimitriou, L., Pokhrel, Y., Seneviratne, S. I., Satoh, Y., Thiery, W., Westra, S., Zhang, X., and Zhao, F.: Globally observed trends in

- 555 mean and extreme river flow attributed to climate change, *Science*, 371, 1159–1162, <https://doi.org/10.1126/science.aba3996>, publisher: American Association for the Advancement of Science, 2021.
- Hannaford, J., Buys, G., Stahl, K., and Tallaksen, L. M.: The influence of decadal-scale variability on trends in long European streamflow records, *Hydrology and Earth System Sciences*, 17, 2717–2733, <https://doi.org/10.5194/hess-17-2717-2013>, publisher: Copernicus GmbH, 2013.
- 560 Held, I. M. and Soden, B. J.: Robust responses of the hydrological cycle to global warming, *Journal of Climate*, 19, 5686–5699, <http://journals.ametsoc.org/doi/abs/10.1175/JCLI3990.1>, 2006.
- Hiemstra, P., Pebesma, E., Twenhöfel, C., and Heuvelink, G.: Real-time automatic interpolation of ambient gamma dose rates from the dutch radioactivity monitoring network, *Computers & Geosciences*, 35, 1711–1721, 2009.
- Ionita, M., Nagavciuc, V., Kumar, R., and Rakovec, O.: On the curious case of the recent decade, mid-spring precipitation deficit in central Europe, *npj Climate and Atmospheric Science*, 3, 1–10, <https://doi.org/10.1038/s41612-020-00153-8>, number: 1 Publisher: Nature Publishing Group, 2020.
- 565 Jaramillo, F. and Destouni, G.: Local flow regulation and irrigation raise global human water consumption and footprint, *Science*, 350, 1248–1251, <https://doi.org/10.1126/science.aad1010>, 2015.
- Jaramillo, F., Prieto, C., Lyon, S. W., and Destouni, G.: Multimethod assessment of evapotranspiration shifts due to non-irrigated agricultural development in Sweden, *Journal of Hydrology*, 484, 55–62, <https://doi.org/10.1016/j.jhydrol.2013.01.010>, 2013.
- 570 Maas, R. and Grennfelt, P.: Towards cleaner air, Arctic Monitoring and Assessment Programme, 2016.
- Masseroni, D., Camici, S., Cislighi, A., Vacchiano, G., Massari, C., and Brocca, L.: The 63-year changes in annual streamflow volumes across Europe with a focus on the Mediterranean basin, *Hydrology and Earth System Sciences*, 25, 5589–5601, <https://doi.org/10.5194/hess-25-5589-2021>, publisher: Copernicus GmbH, 2021.
- 575 Mazurski, K. R.: The destruction of forests in the polish Sudetes Mountains by industrial emissions, *Forest Ecology and Management*, 17, 303–315, [https://doi.org/10.1016/0378-1127\(86\)90158-1](https://doi.org/10.1016/0378-1127(86)90158-1), 1986.
- Mezentsev, V.: More on the calculation of average total evaporation, *Meteorol. Gidrol*, 5, 24–26, budyko02, 1955.
- Müller, U., Börke, P., and Mellentin, U.: Wie entwickeln sich die Wasserdargebote in Sachsen?, *Wasserbau und Wasserwirtschaft im 'Stresstest'*, pp. 10–18, 2023.
- 580 Ol'Dekop, E.: On evaporation from the surface of river basins, *Transactions on meteorological observations University of Tartu*, 4, 200, 1911.
- Otto, L.->, Matschulla, F., and Sonnemann, S.: Waldschutzsituation in Sachsen 2021, *AFZ/Der Wald*, 2022.
- Pike, J.: The estimation of annual run-off from meteorological data in a tropical climate, *Journal of Hydrology*, 2, 116–123, 1964.
- Pitelka, L. F. and Raynal, D. J.: Forest Decline and Acidic Deposition, *Ecology*, 70, 2–10, <https://doi.org/10.2307/1938405>, articleType: research-article / Full publication date: Feb., 1989 / Copyright © 1989 Ecological Society of America, 1989.
- 585 Pluntke, T., Bernhofer, C., Grünwald, T., Renner, M., and Prasse, H.: Long-term climatological and ecohydrological analysis of a paired catchment – flux tower observatory near Dresden (Germany). Is there evidence of climate change in local evapotranspiration?, *Journal of Hydrology*, 617, 128 873, <https://doi.org/10.1016/j.jhydrol.2022.128873>, 2023.
- Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., Hanel, M., and Kumar, R.: The 2018–2020 Multi-Year Drought Sets a New Benchmark in Europe, *Earth's Future*, 10, e2021EF002 394, <https://doi.org/10.1029/2021EF002394>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021EF002394>, 2022.
- 590 Renner, M. and Bernhofer, C.: Long term variability of the annual hydrological regime and sensitivity to temperature phase shifts in Saxony/Germany, *Hydrology and Earth System Sciences Discussions*, 8, 811–853, <https://doi.org/doi:10.5194/hessd-8-811-2011>, 2011.

- Renner, M., Seppelt, R., and Bernhofer, C.: Evaluation of water-energy balance frameworks to predict the sensitivity of streamflow to climate change, *Hydrology and Earth System Sciences*, 16, 1419–1433, <https://doi.org/10.5194/hess-16-1419-2012>, 2012.
- 595 Renner, M., Brust, K., Schwärzel, K., Volk, M., and Bernhofer, C.: Separating the effects of changes in land cover and climate: a hydro-meteorological analysis of the past 60 yr in Saxony, Germany, *Hydrology and Earth System Sciences*, 18, 389–405, <https://doi.org/10.5194/hess-18-389-2014>, 2014.
- Schreiber, P.: Über die Beziehungen zwischen dem Niederschlag und der Wasserführung der Flüsse in Mitteleuropa, *Zeitschrift für Meteorologie*, 21, 441–452, 1904.
- 600 SMUL: Waldzustandsbericht 2006 Freistaat Sachsen, in: Waldzustandsbericht, Sachsenforst, Pirna, <https://publikationen.sachsen.de/bdb/>, 2006.
- Statistisches Landesamt Freistaat Sachsen: Bodennutzung und Ernte im Freistaat Sachsen, Statistischer Bericht C II 2 - j/21, https://www.statistik.sachsen.de/download/statistische-berichte/statistik_sachsen_c-II2_bodennutzung-ernte.xlsx, 2022.
- Teuling, A. J., Seneviratne, S. I., Stöckli, R., Reichstein, M., Moors, E., Ciais, P., Luysaert, S., van den Hurk, B., Ammann, C., Bern-
605 hofer, C., Dellwik, E., Gianelle, D., Gielen, B., Grünwald, T., Klumpp, K., Montagnani, L., Moureaux, C., Sottocornola, M., and Wohlfahrt, G.: Contrasting response of European forest and grassland energy exchange to heatwaves, *Nature Geoscience*, 3, 722–727, <https://doi.org/10.1038/ngeo950>, 2010.
- Teuling, A. J., de Badts, E. A. G., Jansen, F. A., Fuchs, R., Buitink, J., Hoek van Dijke, A. J., and Sterling, S. M.: Climate change, reforestation/afforestation, and urbanization impacts on evapotranspiration and streamflow in Europe, *Hydrology and Earth System Sciences*, 23,
610 3631–3652, <https://doi.org/10.5194/hess-23-3631-2019>, publisher: Copernicus GmbH, 2019.
- Todorov, V. and Filzmoser, P.: An Object-Oriented Framework for Robust Multivariate Analysis, *Journal of Statistical Software*, 32, 1–47, <https://doi.org/10.18637/jss.v032.i03>, 2009.
- Turc, L.: Evaluation des besoins en eau d’irrigation, évapotranspiration potentielle, *Ann. agron.*, 12, 13–49, 1961.
- Wang, D. and Hejazi, M.: Quantifying the relative contribution of the climate and direct human impacts on mean annual streamflow in the
615 contiguous United States, *Water Resources Research*, 47, W00J12, <https://doi.org/10.1029/2010WR010283>, 2011.
- Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis, A., Russak, V., and Tsvetkov, A.: From Dimming to Brightening: Decadal Changes in Solar Radiation at Earth’s Surface, *Science*, 308, 847–850, <https://doi.org/10.1126/science.1103215>, 2005.
- Zhang, L., Dawes, W., and Walker, G.: Response of mean annual evapotranspiration to vegetation changes at catchment scale, *Water Resources Research*, 37, 701–708, 2001.
- 620 Šrámek, V., Slodičák, M., Lomský, B., Balcar, V., Kulhavý, J., Hadaš, P., Pulkráb, K., Šišák, L., Pěnička, L., and Sloup, M.: The Ore Mountains: Will successive recovery of forests from lethal disease be successful, *Mountain Research and Development*, 28, 216–221, 2008.

Figures




Figure 1. Illustration of the separation of climate and land surface changes using a relative energy to relative water balance diagram. The example shows two hydroclimatic states before (q_0, f_0) and after transition (q_1, f_1) . The position of point (q_b, f_b) is determined by using the described geometric approach (3). The bold arrows depict the climatic and the land surface components of this transition. For illustration we used case conditions as a reference: $P_0 = 1000$ mm/yr, $E_{0,0} = 800$ mm/yr, $E_{T,0} = 500$ mm/yr and a state after hypothetical climatic and land-surface change with $P_1 = 1400$ mm/yr, $E_{0,1} = 800$ mm/yr, $E_{T,1} = 350$ mm/yr. Thereby E_T decreased by 30%. The figure is taken from Renner et al. (2014).




Figure 2. Map of the federal State of Saxony, its topography, catchment boundaries and river discharge measurement stations analyzed in this study. Land-use fraction depicted as pie chart of each catchment is derived from CORINE Land-cover data from the year 2000. Panel b) shows the location of Saxony with Germany and Europe for orientation using polygon data from NUTS 2021 (ec.europa.eu). The figure is adapted from Renner et al. (2014).




Figure 3. Maps of long term (1951–2020) annual mean (a) precipitation, (b) annual potential evapotranspiration (FAO-56 grass-reference evapotranspiration (Allen et al., 1994)), (c) annual runoff and (d) the residual water budgets ($E_T = P - R$). Observation stations which have been used to derive spatial fields of P and E_0 are depicted as circles with diameter corresponding to the number of years of available data. The maps of mean annual precipitation P and the FAO reference potential evapotranspiration E_0 have been derived by averaging the individual annual raster maps used to calculate the basin averages. Note, that panels (a) and (c) as well as (b) and (d) use the same color scale respectively.




Figure 4. Map of forest damage using Corine 1990 class 324 (transitional scrub forest). Catchments are grouped by the relative area of forest damage.




Figure 5. Long term catchment water and energy partitioning of the catchments. Panel a) plots the relative energy vs. water balance. The average basin elevation is used to predict the contour lines using local polynomial (LOESS) regression. This demonstrates the general height dependency of the catchments climate and land-use (Corine dataset of 2000) as well as the transition from wet basins with high runoff ratio to lower values. The inset in panel a) shows a map of the catchments, its gauge id number and altitude as background raster. Panel b) visualizes the same data but in the classic Budyko diagram with relative water balance vs. aridity index. The catchments are depicted as pie-charts which show the areal percentage of land use of each catchment.




Figure 6. Time series of annual P and E_0 for all basins (thin dashed lines), cross basin annual average time series (full lines) and the all basin decadal averages (dot-dashed lines).




Figure 7. Boxplots of decadal averages of E_T for four groups of catchments with different ranges of damaged forest area per catchment area using the Corine 1990 class 324 (transitional scrub forest).




Figure 8. Decadal changes in annual mean catchment runoff across all catchments. Box-whisker plots are shown for observed (gray), climate (blue) and land-surface attributed runoff changes using the decomposition method introduced in the methods section. The boxes show the inter-quartile range (IQR) with the bold horizontal line being the median of the distribution. The whiskers show the outer ranges, with points marking outliers outside a 1.5 IQR range.




Figure 9. Annual averages per decade of the water and energy partitioning ratios for each basin analyzed. The arrows denote the change from one decade into the following decade. There is one panel for each decadal change, starting in 1950s in panel (a) to 2000s to 2010s in panel (f). Significant changes are marked with bold arrows and a red border of the respective catchment in the map inset. The inlay maps show the location of the river gauges, the spatial extent of forest damages and since 1990 blue pixels show the Corine transitional forest class.




Figure 10. Change in decadal water balance components from the decade 2001-10 to 2011-20 in mm/yr for each catchment.




Figure 11. Maps of decadal changes (2001-10 to 2011-20) in (a) precipitation and runoff, (b) potential evaporation (FAO-56 grass-reference evapotranspiration, (Allen et al., 1994)) and catchment evapotranspiration ($E_T = P - R$), (c) climate attributed changes in runoff, and (d) land-surface attributed changes in runoff. While panels (a) and (b) visualize the changes on the catchment level and at the gauging stations (triangles), panels (c) and (d) use only the catchment area to present changes.

625 Appendix A: Catchment data

	station / river	major basin	gaugeID	elev	area	forest	damage	P	E_0	R	E_T	miss
2	Buschmuehle/Kirnitzsch	Upper Elbe	550090	397	98	78	1	854	683	296	565	20
3	Kirnitzschtal/Kirnitzsch	Upper Elbe	550110	379	154	77	0	848	688	289	559	0
4	Porschdorf/Lachsbad	Upper Elbe	550190	378	267	39	0	850	691	362	487	0
5	Sebnitz/Sebnitz	Upper Elbe	550302	422	102	47	1	877	681	438	450	19
6	Neustadt/Polenz	Upper Elbe	550390	394	40	32	0	859	687	417	451	21
7	Markersbach/Bahra	Upper Elbe	550710	545	49	45	10	805	657	364	439	20
8	Bischofswerda/Wesenitz	Upper Elbe	550801	363	69	30	0	823	692	354	470	19
9	Elbersdorf/Wesenitz	Upper Elbe	550810	315	227	17	0	793	703	297	496	0
10	Dohna/Müglitz	Upper Elbe	550940	557	198	35	6	840	652	403	437	1
11	Geising/Rotes Wasser	Upper Elbe	550961	780	26	53	35	957	602	537	417	16
12	Kreischa/Lockwitzbach	Upper Elbe	551000	380	44	23	0	766	692	252	517	14
13	Klotzsche/Prießnitz	Upper Elbe	551100	261	40	60	0	732	720	267	472	20
14	Rehefeld/Wilde Weißeritz	Upper Elbe	551302	808	15	91	61	974	597	754	218	12
15	Ammelsdorf/Wilde Weißeritz	Upper Elbe	551310	734	49	64	20	966	614	599	374	2
16	Beerwalde/Wilde Weißeritz	Upper Elbe	551320	663	81	53	12	940	630	519	422	6
17	Baerenfels/Pöbelbach	Upper Elbe	551510	742	6	75	12	983	611	746	261	20
18	Freital/Poisenbach	Upper Elbe	551561	295	12	42	0	701	715	216	490	26
19	Piskowitz/Ketzerbach	Upper Elbe	552011	214	157	0	0	660	727	122	542	25
20	Seerhausen/Jahna	Upper Elbe	552110	178	153	0	0	627	734	58	571	16
21	Merzdorf/Döllnitz	Upper Elbe	552210	167	211	7	0	611	738	135	475	0
22	Zescha/Hoyersw. Schwarzwasser	S. Elster	554220	226	180	13	0	704	721	181	527	15
23	Pietzschwitz/Langes Wasser	S. Elster	554260	255	42	12	0	726	716	232	498	12
24	Koenigsbrueck/Pulsnitz	S. Elster	554420	278	92	30	0	755	713	272	484	10
25	Grossdittmannsdorf/Große Röder	S. Elster	554520	247	300	31	0	739	721	238	500	0
26	Kleinraschuetz/Große Röder	S. Elster	554550	198	679	25	0	683	731	187	494	11
27	Golzern/Vereinigte Mulde	Mulde	560021	481	5442	31	4	840	671	354	486	0
28	Bad Dübren/Vereinigte Mulde	Mulde	560051	439	6171	30	3	811	679	319	490	10
29	Zwickau/Zwickauer Mulde	Mulde	562070	633	1030	56	2	953	637	434	519	0
30	Wechselburg/Zwickauer Mulde	Mulde	562115	491	2107	35	1	851	669	392	459	0
31	Muldenberg/Rote Mulde	Mulde	563000	769	5	99	0	986	609	628	337	26
32	Sachsengrund/Große Pyra	Mulde	563290	927	7	100	12	1151	571	798	340	22
33	Aue/Schwarzwasser	Mulde	563790	744	362	68	6	1014	615	548	466	0
34	Niedermuelzen/Mülsenbach	Mulde	564201	364	50	20	0	753	693	276	482	22
35	Niederlungwitz/Lungwitzbach	Mulde	564300	353	138	11	0	744	700	310	445	18
36	Harthau/Würschnitz	Mulde	564620	435	136	17	0	819	688	336	491	19
37	Berthelsdorf/Freiberger Mulde	Mulde	566010	597	243	27	1	928	646	444	484	0
38	Nossen/Freiberger Mulde	Mulde	566040	486	585	20	0	858	670	368	489	0
39	Erlin/Freiberger Mulde	Mulde	566100	504	2983	30	6	851	666	376	482	19
40	Wolfsgrund/Chemnitzbach	Mulde	567000	627	37	13	0	946	639	556	389	0
41	Tannenberg/Zschopau	Mulde	567400	663	91	38	0	960	632	509	448	9
42	Hopfgarten/Zschopau	Mulde	567420	703	529	46	11	933	624	468	465	0
43	Lichtenwalde/Zschopau	Mulde	567451	619	1575	44	11	907	642	431	476	0
44	Annaberg/Sehma	Mulde	567590	753	49	47	2	959	615	499	458	17
45	Pockau/Flöha	Mulde	568140	689	385	63	27	920	625	469	454	8
46	Borstendorf/Flöha	Mulde	568160	662	644	57	19	918	631	438	480	0
47	Deutschgeorghenthal/Rauschenbach	Mulde	568200	743	10	84	14	983	616	549	443	16
48	Neuwernsdorf/Wernsbach	Mulde	568250	757	7	94	68	983	610	589	401	17
49	Rauschenbach/Rauschenfluß	Mulde	568300	741	7	96	48	958	622	642	328	15

50	Roenthal/Natzschung	Mulde	568350	768	75	81	47	911	610	566	344	0
51	Zoeblitz/Schwarze Pockau	Mulde	568400	706	129	58	15	924	623	531	390	1
1	Klingenthal/Zwota (Svatava)	W. Elster	530020	721	59	78	2	990	616	605	386	11
52	Adorf/Weiße Elster	W. Elster	576400	599	171	57	0	793	643	303	485	20
53	Oelsnitz/Weiße Elster	W. Elster	576410	574	328	50	0	792	649	297	496	10
54	Magwitz/Weiße Elster	W. Elster	576420	558	376	44	0	780	652	281	499	0
55	Strassberg/Weiße Elster	W. Elster	576421	532	611	37	0	742	657	258	491	15
56	Greiz/Weiße Elster	W. Elster	576470	474	1255	31	0	732	670	260	471	0
57	Hasenmühle/Trieb	W. Elster	577100	532	100	39	0	816	659	297	525	17
58	Mylau/Göltzsch	W. Elster	577220	515	155	34	0	839	661	374	465	0
59	Streitwald/Wyhra	W. Elster	577901	242	178	8	0	662	717	153	510	4
60	Leipzig/Parthe	W. Elster	578110	144	315	14	0	606	735	93	512	0
61	Schirgiswalde/Spree	Spree	582010	382	179	34	2	806	688	350	460	12
62	Wuischke/Wuischker Wasser	Spree	583140	403	3	91	0	759	689	305	447	28
63	Kotitz/Kotitzer Wasser	Spree	583170	252	29	4	0	686	714	191	490	22
64	Oehlich/Schwarzer Schöps	Spree	583230	275	35	10	0	681	714	186	495	27
65	Holtendorf/Weißer Schöps	Spree	583280	272	54	12	0	685	715	189	498	5
66	Saerichen/Weißer Schöps	Spree	583290	243	135	9	0	678	722	187	494	12
67	Hartau/Lausitzer Neiße	Neiße	660010	483	376	45	4	896	663	488	419	7
68	Zittau/Lausitzer Neiße	Neiße	660100	443	686	36	3	844	671	406	445	5
69	Seifhennersdorf/Mandau	Neiße	662001	419	75	22	4	815	676	378	445	17
70	Rennersdorf/Pließnitz	Neiße	663000	331	79	25	0	709	698	211	494	20
71	Rennersdorf1/Petersbach	Neiße	663100	328	64	25	0	707	698	219	483	20

Table A1: River stations analyzed over the period 1951 - 2020. The column *elev* denotes the mean catchment elevation in meters above sea level, *area* denotes catchment area in km^2 , *forest* gives the relative coverage of forest land use (%) based on Corine 1990 data. Forest damage in percent is given in the “damage” column, using the Corine class 324. The columns P , E_0 , R and E_T denote average annual water balance components for the basins in mm/yr. *miss* gives the number of missing years.

A1 Groundwater aquifers in Saxony




Figure A1. Map of the Saxon hydrogeology categorized into highly conductive aquifers as strong indicator for large groundwater influences, low conductive aquifers for minor groundwater influence, and low conductive bedrocks for regions without groundwater influences. Data source is the LfULG portal (<https://ida.sachsen.de>). On top the river gauge locations and the catchment borders are plotted.

Appendix B: Methods

B1 Derivation of the decomposition for runoff

To get to equation 4 we start with the climate-induced change in evapotranspiration:

$$630 \quad \Delta E_{T,C} = E_{T,1} - E_{T,b} \quad (B1)$$

Next we apply the water budget equation $\Delta P = \Delta E_T + \Delta R$, which yields for the climate part:

$$\Delta R_C = \Delta P - \Delta E_{T,C} \quad (B2)$$

Then we can substitute $\Delta E_{T,C}$ to yield:

$$\Delta R_C = \Delta P - E_{T,1} + E_{T,b} \quad (B3)$$

635 You can rewrite $E_{T,1} = P_1 - R_1$ and $\Delta P = P_1 - P_0$ and insert both:

$$\Delta R_C = P_1 - P_0 - (P_1 - R_1) + E_{T,b} \quad (B4)$$

to yield equation 4:

$$\Delta R_C = R_1 - P_0 + E_{T,b} \quad (B5)$$

To get to equation 5 we start with the land cover induced change in evapotranspiration as noted in chapter 2.2:

$$640 \quad \Delta E_{T,L} = E_{T,b} - E_{T,0} \quad (B6)$$

and also apply the water balance equation to get to the runoff changes.

$$\Delta R_L = \Delta P - \Delta E_{T,L} \quad (B7)$$

Then we can substitute $\Delta E_{T,L}$ to yield:

$$\Delta R_L = \Delta P - E_{T,b} + E_{T,0} \quad (B8)$$

645 By replacing $E_{T,0}$ and ΔP we get to the following equation:

$$\Delta R_L = P_1 - P_0 - E_{T,b} + P_0 - R_0 \quad (B9)$$

and simplify to receive equation 5:

$$\Delta R_L = P_1 - E_{T,b} - R_0 \quad (B10)$$

B2 Comparison of decomposition methods




Figure B1. Comparison of the decomposition method of this study with the method of Wang and Hejazi (2011) using the Mezentsev (1955) parametric Budyko function to separate climate and land-surface induced changes on E_T (panel a) and runoff (b). Here all decadal transitions of all catchments with sufficient data are used for the comparison.